

A SET OF POSTULATES FOR GENERAL PROJECTIVE GEOMETRY*

BY

MEYER G. GABA

Since Klein promulgated his famous *Erlangen Programme*† it has been known that the various types of geometry are such that each is characterized by a group of transformations. In view of the importance of the concept of transformation in nearly all mathematics and perhaps especially in geometry, geometers may properly seek to develop the various types of geometry in terms of point and transformation. For euclidean geometry this has been done by Pieri.‡

This paper is devoted to a similar treatment of general projective geometry.§ One would naturally lay such postulates on the system of transformations so as to make the system form the group associated with the geometry. This was the scheme that Pieri used. His postulates make his transformations form the group of motions. In general projective geometry, however, this method is not necessary. If we are given the group of all projective transformations we can deduce the geometry from it but it will be shown in the sequel that we can also do that from a properly chosen semi-group belonging to that group.

Our basis, to repeat, is a class of undefined elements called points and a class of undefined functions on point to point|| or transformations called collineations. For notation we will use small *Roman* letters to designate

* Read before the American Mathematical Society, April 26, 1913.

† F. Klein, *Vergleichende Betrachtungen über neuere geometrische Forschungen*, Erlangen, 1872. English translation by M. W. Haskell, *Bulletin of the American Mathematical Society*, vol. 2 (1893).

‡ M. Pieri, *Della geometria elementare come sistema ipotetico-deduttivo; monografia del punto e del moto*, Memorie delle Scienze di Torino (1899).

§ General projective geometry is defined by Veblen and Young as a geometry associated (analytically) with a general number field; that is, its theorems are valid, not alone in the ordinary real and the ordinary complex projective spaces but also in the ordinary rational spaces and in the finite spaces. This paper connects closely with the postulates for general projective geometry given by O. Veblen and J. W. Young in the *American Journal of Mathematics*, vol. 30 (1908), and in their *Projective Geometry*, Ginn and Company (1910).

|| By point to point is meant that to every point p there corresponds a single point p' and no point p' is the correspondent of two distinct points p_1 and p_2 .

points and small Greek letters for collineations. Thus

$$\tau(p_1, p_2, p_3) = p'_1, p'_2, p'_3$$

means that the collineation τ transforms the points p_1, p_2, p_3 into p'_1, p'_2, p'_3 respectively. Line will be defined in terms of points and collineations. If we should interpret our undefined collineations as the group of all projective collineations, our defined line will satisfy the Veblen-Young assumptions for their undefined line and the postulates I to VIII that we will soon give are theorems in general projective geometry. This proves the consistency of our postulates. On the other hand, leaving collineation as undefined and using postulates I to VI* we are able to prove as theorems the Veblen-Young assumptions $A_1, A_2, A_3, E_0, E_1, E_2, \dots, E_n, E_n'$, and P .† This shows that our six postulates are sufficient to establish the general projective geometry of n -dimensions. The undefined collineation will be proven to be a projective collineation which justifies the notation. If we desire that our class of collineations should be the group of all projective collineations we add postulate VII to the preceding six. To the Veblen-Young postulate H_0 corresponds our postulate VIII.

The independence of our assumptions is proven by the set of independence examples given at the end of this paper.

POSTULATE I. *There are at least $n + 2$ distinct points.*

POSTULATE II. *If τ_1 is a collineation and τ_2 is a collineation then the resultant of operating first with τ_1 and then with τ_2 (in notation $\tau_2 \tau_1$) is also a collineation.*

DEFINITION. A linear set is a class of points such that:

(a) every collineation that leaves two distinct points of the class invariant leaves the class invariant,

(b) every collineation that leaves three distinct points of the class invariant leaves every point of the class invariant.

DEFINITION. Points belonging to the same linear set are called collinear.

DEFINITION. A linear set that contains at least three distinct points is called a line.‡

POSTULATE III. *If p_1, p_2, p_3 are three distinct collinear points and p'_1, p'_2, p'_3 are three distinct collinear points then a collineation exists that transforms p_1, p_2, p_3 into p'_1, p'_2, p'_3 respectively.*

POSTULATE IV. *If p_1, p_2, p_3, p_4 are four distinct points such that no three*

* Postulates I to VI explicitly require that our set of collineations form a semi-group but all sets of transformations that we found satisfying I-VI were groups. The question whether they necessarily form a group in the general case has not as yet been proven.

† The precise statement of these postulates is given later in this paper.

‡ It will be proved that every pair of distinct points is contained in one and only one line.

are such that each is collinear with the same two distinct points then a collineation exists leaving p_1 and p_2 invariant and interchanging p_3 and p_4 .*

DEFINITION. If p_1, p_2 are two distinct points, a line containing p_1 and p_2 or, in case no line exists containing p_1 and p_2 , then the pair of points p_1, p_2 is called a one-space (P_1) containing p_1, p_2 .

DEFINITION. If P_{k-1} is a $(k-1)$ -space and p_0 is a point not contained in P_{k-1} , the class $P_k \equiv [P_{k-1}, p_0]$ of all points p collinear with the point p_0 and some point of P_{k-1} is called the k -space determined by P_{k-1} and p_0 .

DEFINITION. k points are called independent of each other if there exists no $(k-2)$ -space that contains them all.

POSTULATE V. If p_1, p_2, \dots, p_{n+1} are $n+1$ distinct points of the same k -space, $k < n$, then a collineation distinct from the identity exists leaving p_1, p_2, \dots, p_{n+1} invariant.†

POSTULATE VI. If p_1, p_2, \dots, p_{n+2} are $n+2$ distinct points then there exists a k -space, $k \leq n$, that contains them all.

POSTULATE VII. If p_1, p_2, \dots, p_{n+2} are $n+2$ points of the same n -space such that every $n+1$ are independent, and $p'_1, p'_2, \dots, p'_{n+2}$ are $n+2$ points of the same n -space such that every $n+1$ are independent then a collineation exists that transforms p_1, p_2, \dots, p_{n+2} into $p'_1, p'_2, \dots, p'_{n+2}$ respectively.

DEFINITION. A complete quadrangle is a figure consisting of four distinct coplanar points such that no three are collinear, called its vertices and six distinct lines containing the vertices in pairs called its sides. Two sides having no vertex in common are called opposite and points common to two opposite sides are called diagonal points.

POSTULATE VIII. The diagonal points of a complete quadrangle are non-collinear.

THEOREM 1. If p_1 and p_2 are distinct points, there is not more than one line containing both p_1 and p_2 .

Let us assume that two lines P and P' exist such that each contains p_1 and p_2 . If the two lines are distinct then at least one of the lines must contain a point not in the other. Let us assume that p'_3 is a point of P' and not of P . Since P is a line it contains in addition to p_1 and p_2 a third point p_3 . From Postulate III we know that a collineation τ exists such that $\tau(p_1, p_2, p_3) = p_1, p_2, p'_3$. But τ leaves two points of P invariant, therefore it leaves P invariant and p_3 cannot be transformed into p'_3 which is a point not of P . Since we are led to a contradiction our assumption that p_1 and p_2 are contained in two distinct lines must be false.

THEOREM 2. Two distinct lines cannot have more than one common point.

* Compare Postulate IV with Theorem 4. Postulate IV is weaker than Theorem 4 as independence example — IV will show.

† Postulate V with Theorem 14 shows that all points lie in no $P_k, k < n$.

THEOREM 3. *If three points are such that each is collinear with the same two points, they are collinear and conversely.*

THEOREM 4. *If p_1, p_2, p_3, p_4 are four distinct points such that no three are collinear then a collineation τ exists such that $\tau(p_1, p_2, p_3, p_4) = p_1, p_2, p_4, p_3$. The theorem follows from Postulate IV and Theorem 3.*

THEOREM 5. *If p_1, p_2 are two distinct points and q_1, q_2 are two distinct points then a collineation τ exists such that $\tau(p_1, p_2) = q_1, q_2$.*

If p_1, p_2 are on a line and q_1, q_2 are on a line, the theorem follows from Postulate III. If no three of the points p_1, p_2, q_1, q_2 are collinear and they are all distinct then by Theorem 4 we know that a collineation τ_1 exists such that $\tau_1(p_1, q_1, p_2, q_2) = p_1, q_1, q_2, p_2$ and a collineation τ_2 exists such that $\tau_2(p_1, q_1, q_2, p_2) = q_1, p_1, q_2, p_2$. Therefore by Postulate II a collineation $\tau_2 \tau_1$ exists such that $\tau_2 \tau_1(p_1, p_2) = q_1, q_2$.

There are, notation apart, two possible cases remaining which are: (1) q_2 collinear with p_1 and p_2 but q_1, q_2 on no line; (2) p_2 collinear with q_1 and q_2 , but p_1, p_2 on no line. Let us first suppose that a point r exists such that r is non-collinear with every two of the points p_1, p_2, q_1, q_2 . In the first case, collineations τ_3 and τ_4 exist such that $\tau_3(p_1, p_2, r, q_1) = q_1, p_2, r, p_1$ and $\tau_4(q_1, p_2, r, q_2) = q_1, q_2, r, p_2$, and therefore $\tau_4 \tau_3$ is the required collineation. In the second case, collineations τ_5 and τ_6 exist such that $\tau_5(p_1, p_2, r, q_2) = p_1, q_2, r, p_2$ and $\tau_6(p_1, q_2, r, q_1) = q_1, q_2, r, p_1$ and in this case $\tau_6 \tau_5$ is a collineation that transforms p_1, p_2 into q_1, q_2 respectively.

If no such point r exists then every point is contained in some one of the one-spaces determined by two of the four points p_1, p_2, q_1, q_2 . Let us consider the case where p_1 and p_2 are on no line and where p_1 is collinear with q_1, q_2 and let us further suppose that p_1 is distinct from q_1 and from q_2 . All points are in the two-space or plane determined by the line q_1, q_2 and the point p_2 is not in the line q_1, q_2 . If $n = 2$ then by Postulate V a collineation not the identity exists having p_1, q_1, q_2 as invariant points and there must therefore be at least one additional point r_1 on the one-space q_1, p_2 or one additional point s_1 on the one-space q_2, p_2 . If r_1 exists then collineations τ_7 and τ_8 exist such that $\tau_7(p_1, p_2, r_1, q_2) = p_1, q_2, r_1, p_2$ and $\tau_8(q_1, q_2, p_1) = p_1, q_2, q_1$. If s_1 exists we have collineations τ_9 and τ_{10} such that $\tau_9(p_1, p_2, s_1, q_1) = p_1, q_1, s_1, p_2$ and

$$\tau_{10}(q_1, q_2, p_1) = q_2, p_1, q_1.$$

Hence $\tau_8 \tau_7$ or $\tau_{10} \tau_9$ will be the required collineation according as r_1 or s_1 exists. If $n > 2$ there are by Postulate I at least $n + 2 > 4$ distinct points; hence the additional point r_1 or s_1 exists as before, and the argument is completed as in the case $n = 2$.

Let us now suppose that p_1 coincides with q_1 . Since $n \geq 2$, at least four

distinct points exist. Let us assume q_3 , distinct from q_1 and q_2 , exists on the one-space $q_1 q_2$. A collineation distinct from the identity exists leaving invariant q_1, q_2, q_3 if $n = 2$ and, if $n > 2$, q_1, q_2, q_3 , and $n - 2$ other points; therefore a point r_1 distinct from q_2, p_2 must lie on the one-space $q_2 p_2$. If the existence of r_1 had been assumed, then we could have proven in a similar manner that q_3 existed and since one or the other must exist, both exist. Since the four points $p_1 = q_1, p_2, r_1, q_3$ are such that no three are collinear a collineation τ_{11} exists such that $\tau_{11}(p_1, p_2, r_1, q_3) = p_1, q_3, r_1, p_2$ and a collineation τ_{12} exists such that $\tau_{12}(p_1, q_2, q_3) = q_1, q_3, q_2$. The collineation $\tau_{12} \tau_{11}$ is the collineation that transforms $p_1 p_2$ into $q_1 q_2$. For all the other possible cases the proofs are very similar to the preceding and therefore need not be repeated.

THEOREM 6. *A line exists.*

If all linear sets contained but two points there would be but $k + 1$ points in a k -space. This would make Postulates I and VI contradictory and therefore at least one linear set contains more than two points and hence a line exists.

THEOREM 7. *Every collineation transforms lines into lines.*

Let τ_1 be any collineation that transforms the line $P = [p]$ into a set of points $Q = [q]$. We are to prove that the set of points Q constitute a linear set. Let τ_2 be any collineation that leaves two of the q 's, say q_1 and q_2 invariant. The points q_1 and q_2 are the transforms under τ_1 of two points of P which we will call p_1 and p_2 . By Theorem 5 there is a collineation τ_3 that transforms q_1, q_2 into p_1, p_2 . Then $\tau_3 \tau_2 \tau_1(P) = P$ since the points p_1 and p_2 of P are left invariant. For the same reason $\tau_3 \tau_1(P) = P$. Therefore $\tau_3(Q) = P$, that is to say every p is the transform under τ_3 of some q and that for every q $\tau_3(q)$ is a p . But $\tau_3 \tau_2(Q) = P$ therefore $\tau_2(Q) = Q$. Hence any collineation that leaves two points of Q fixed leaves Q invariant.

Let τ_4 be any collineation that leaves three of the q 's invariant, say q_1, q_2, q_3 , where q_1, q_2, q_3 are the transforms under τ_1 of p_1, p_2, p_3 respectively. $\tau_3 \tau_4 \tau_1[P] = P$ since p_1 and p_2 are left invariant. $\tau_3 \tau_4 \tau_1(p_1, p_2, p_3) = p_1, p_2, p_3'$. Since p_1, p_2, p_3 and p_1, p_2, p_3' are sets of collinear points a collineation τ_5 exists such that $\tau_5(p_1, p_2, p_3) = p_1, p_2, p_3'$. Then

$$\tau_5 \tau_3 \tau_4 \tau_1(p_1, p_2, p_3) = p_1, p_2, p_3$$

and

$$\tau_5 \tau_3 \tau_1(p_1, p_2, p_3) = p_1, p_2, p_3.$$

Therefore the collineations $\tau_5 \tau_3 \tau_1$ and $\tau_5 \tau_3 \tau_4 \tau_1$ leave every point of P invariant. If τ_1 transforms p_i into q_i , $\tau_5 \tau_3$ must transform q_i into p_i and hence τ_4 must leave every point of Q invariant. We have shown that properties (a) and (b) of a linear set hold for Q and since P was a line (containing at least three points) Q is a line.

THEOREM 8. *If p_1 and p_2 are distinct points, there is at least one line containing both p_1 and p_2 .*

We know that at least one line exists from Theorem 6. That line has two points q_1 and q_2 . By Theorem 5 a collineation τ exists transforming q_1, q_2 into p_1, p_2 respectively. The line containing q_1, q_2 is transformed by τ into a line which contains p_1, p_2 .

THEOREM 9. *Every collineation transforms a k -space into a k -space.*

THEOREM 10. *All points are not on the same line.*

THEOREM 11. *If p_1, p_2, p_3, p_4, p_5 are five distinct points such that p_1, p_2, p_3 are non-collinear, p_1, p_2, p_4 are collinear and p_1, p_3, p_5 are collinear, then there exists a point p_6 such that p_2, p_3, p_6 are collinear and p_4, p_5, p_6 are collinear.**

No three of the points p_2, p_3, p_4, p_5 are collinear for if they were p_1, p_2, p_3 would be collinear. By Theorem 4 a collineation τ exists such that

$$\tau(p_2, p_3, p_4, p_5) = p_2, p_4, p_3, p_5.$$

The collineation τ transforms the lines $p_3 p_5$ and $p_2 p_4$ into the lines $p_4 p_5$ and $p_2 p_3$ respectively. The point p_1 common to the lines $p_3 p_5$ and $p_2 p_4$ will therefore be transformed into a point p_6 common to the lines $p_4 p_5$ and $p_2 p_3$.

We have already proven as theorems the Veblen-Young postulates A_1, A_2, A_3, E_0, E_1 , and E_2 . The postulate A_1 is our Theorem 8; A_2 is our Theorem 1; A_3 (if p_1, p_2, p_3 are points not all on the same line and p_4 and p_5 ($p_4 \neq p_5$) are points such that p_1, p_2, p_4 are on a line and p_1, p_3, p_5 are on a line, there is a point p_6 such that p_2, p_3, p_6 are on a line and p_4, p_5, p_6 are on a line) is in content equivalent to Theorem 11; E_0 (there are at least three points on every line) is true from definition of line; E_1 (there exists at least one line) is Theorem 6; and E_2 is Theorem 9. We therefore know that our line satisfies the six preceding postulates that Veblen and Young lay down for their undefined line, hence all theorems that they derive from the six assumptions listed will hold in our geometry. One such theorem is:

THEOREM 12. *Let the k -space P_k be defined by the point p_0 and the $(k-1)$ -space P_{k-1} , then*

(a) *There is a k -space on any $k+1$ independent points.*

(b) *Every line on two points of P_k has one point in common with P_{k-1} and is in P_k .*

(c) *Every P_g ($g < k$) on $g+1$ independent points of P_k is in P_k .*

* Theorem 11 is essentially equivalent to the Veblen-Young postulate A_3 : If p_1, p_2, p_3 are points not all on the same line and p_4 and p_5 ($p_4 \neq p_5$) are points such that p_1, p_2, p_4 are on a line and p_1, p_3, p_5 are on a line, there is a point p_6 such that p_2, p_3, p_6 are on a line and p_4, p_5, p_6 are on a line. The form of statement for Theorem 11 was suggested by Professor E. H. Moore as a substitute for A_3 since the latter is redundant in that it includes the obvious cases where p_4 is coincident with p_1 or p_2 or where p_5 is coincident with p_1 or p_3 .

(d) Every P_g ($g < k$) on $g + 1$ independent points of P_k has a P_{g-1} in common with P_{k-1} provided all $g + 1$ points are not in P_{k-1} .

(e) Every line P_1 on two points of P_k has one point in common with every P_{k-1} in P_k .

(f) If q_0 and Q_{k-1} (q_0 not in Q_{k-1}) are any point and any $(k - 1)$ -space respectively of the k -space determined by p_0 and P_{k-1} , the latter space is the same as that determined by q_0 and Q_{k-1} .

Another important theorem that Veblen and Young prove is:

THEOREM 13. On $k + 1$ independent points there is one and but one k -space.

THEOREM 14. If $k + 1$ points of a P_{k-1} , such that every k are independent, are left invariant by a collineation τ then τ leaves every point of P_{k-1} invariant.

This theorem clearly is true for a line or P_1 . Let us assume that the theorem is true for a P_{g-1} . In a P_g if every $g + 1$ of $g + 2$ points are to be independent, then if p_1, \dots, p_g determine a P_{g-1} , p_{g+1} and p_{g+2} cannot lie in P_{g-1} , nor can the line $p_{g+1} p_{g+2}$ contain any of the points p_1, \dots, p_g . The line $p_{g+1} p_{g+2}$ has a single point p_0 in common with P_{g-1} by Theorem 12 (e). When the $g + 2$ points are left invariant p_0 , being the intersection of P_{g-1} and $p_{g+1} p_{g+2}$, is left invariant. The line P_{g-1} and the line $p_{g+1} p_{g+2}$ are each therefore left identically invariant. Let p be any point in P_g not in P_{g-1} nor on the line $p_{g+1} p_{g+2}$. The lines pp_{g+1} and pp_{g+2} each meet the P_{g-1} by Theorem 12 (e). When the given $g + 2$ points are left invariant, these lines and consequently their intersection is left invariant. The theorem being true for a g -space if true for a $(g - 1)$ -space and holding for a one-space is therefore true for a k -space.

THEOREM 15. All points are not on the same k -space if $k < n$.

It can easily be shown that every k -space has $k + 2$ points such that every $k + 1$ are independent. If these $k + 2$ points and $n - k - 1$ other points, which exist by Postulate I, are not in the k -space the theorem is true. If these $n + 1$ points are in the k -space then by Postulate V a collineation distinct from the identity exists leaving these $n + 1$ points invariant and therefore by Theorem 14 that collineation leaves the k -space identically invariant. Hence not all points can be in the k -space.

THEOREM 16. There exist $n + 2$ points such that every $n + 1$ are independent.

The definitions of perspectivity, projectivity, etc., can now be given exactly as Veblen and Young give them. We will now proceed to identify what we call a collineation with what they call a projective collineation. To do this we will first prove:

THEOREM 17. Every central perspective correspondence between points of two lines can be secured by a collineation.

Let the perspectivity be defined by p_1, p_2 having as their correspondents q_1, q_2 . We have by Theorem 4, since no three of p_1, p_2, q_1, q_2 are collinear,

that the two collineations τ_1 and τ_2 exist such that

$$\tau_1(p_1, p_2, q_1, q_2) = p_1, q_2, q_1, p_2$$

and

$$\tau_2(p_1, q_2, q_1, p_2) = q_1, q_2, p_1, p_2,$$

therefore $\tau = \tau_2 \tau_1$ exists by Postulate II and is such that $\tau(p_1, p_2, q_1, q_2) = q_1, q_2, p_1, p_2$. The lines $p_1 q_1$ and $p_2 q_2$ will be left invariant by τ and so will their point of intersection o . The lines $p_1 p_2$ and $q_1 q_2$ are interchanged by τ as well as the lines $p_1 q_2$ and $p_2 q_1$. Let the intersection of $p_1 p_2$ and $q_1 q_2$ be r_3 and of $p_1 q_2$ and $p_2 q_1$ be r_4 . The collineation τ must leave r_3 and r_4 invariant. There are two cases possible,* the first where o , r_3 , and r_4 are non-collinear and the second where these three points are collinear. Let us first consider case 1. The line $r_3 r_4$ meets the lines $p_1 q_1$ and $p_2 q_2$ in points r_1 and r_2 . The collineation τ leaves r_1 , r_2 , r_3 , and r_4 invariant; hence every point of the line r_3, r_4 is left invariant by τ . Let p be any point on the line $p_1 p_2$ and call r the point of intersection of the line op with the line $r_1 r_2$. The line op is left invariant by τ since that collineation leaves two of its points, o and r , invariant. Since the line $p_1 p_2$ is transformed into the line $q_1 q_2$ by τ , p is transformed by τ into the intersection of the lines op and $q_1 q_2$ which we will call q .

For the second case, where o , r_3 , and r_4 are collinear, the points o and r are coincident. The collineation $\tau^2 = \tau\tau$ leaves p_1, p_2 , and r_3 invariant and therefore leaves every point of $p_1 p_2$ invariant. Hence if q denotes the point of $q_1 q_2$ into which τ transforms a point p of $p_1 p_2$ then τ must transform q into p . Hence the line pq is left invariant by τ . Since o , r_3 , and r_4 are collinear the lines $p_1 q_1$, $p_2 q_2$, and $r_3 r_4$ are concurrent at o . If the line pq did not pass through o it would intersect the three lines $p_1 q_1$, $p_2 q_2$, and $r_3 r_4$ at three distinct points. But this would make the line pq identically invariant under the collineation τ and hence p would be invariant and coincide with q , but this is possible only if p is r_3 and we assumed that p was any point on the line $p_1 p_2$ therefore pq passes through o . Therefore the collineation τ makes correspond to the points of the line (p) the points of line (q) perspective to (p) with center of perspectivity o .

THEOREM 18. *If a projective correspondence exists between the points of two lines, then a collineation exists that transforms the points of the first line into the projectively corresponding points of the second line.*

A projective correspondence between the points of two lines is the resultant of a sequence of central perspectivities. By Theorem 16, each central perspective correspondence has associated with it a collineation. The resultant of the sequence of collineations corresponding to the sequence of perspectivities that define the projectivity is the required collineation.

* If Postulate VIII were assumed, the first case only would arise.

THEOREM 19. *If a projectivity leaves each of three distinct points of a line invariant it leaves every point of the line invariant.*

THEOREM 20. *If P is a P_n all points are in P .*

This theorem follows readily from Postulate VI and Theorem 12.

In addition to the Veblen-Young postulates A_1, A_2, A_3, E_0 , and E_1 we have proven E_n , which is Theorem 15, E'_n , which is Theorem 19, and P , which is Theorem 18. We have therefore proven all of the postulates that Veblen and Young assume for general projective geometry of n -space and consequently all the theorems that can be derived from their postulates hold in our geometry.

We can now prove

THEOREM 21. *Every collineation is a projective collineation.*

Let τ be any collineation. From Theorem 20 we know that there exist $n + 2$ points such that every $n + 1$ are independent. The collineation τ will transform these $n + 2$ points which we will call p_1, p_2, \dots, p_{n+2} into $p'_1, p'_2, \dots, p'_{n+2}$ respectively. There exists a projective collineation π (from the Veblen-Young geometry which is at our disposal) such that $\pi(p_1, p_2, \dots, p_{n+2}) = p'_1, p'_2, \dots, p'_{n+2}$. The points p_1, p_2, \dots, p_n determine an $(n - 1)$ -space which will meet the line $p_{n+1} p_{n+2}$ in a point p_0 . Let p'_0 denote the intersection of the $(n - 1)$ -space determined by p'_1, p'_2, \dots, p'_n with the line $p'_{n+1} p'_{n+2}$. This point, p'_0 , will correspond to p_0 both by transformation τ and transformation π . The line $p_{n+1} p_{n+2}$ having three of its points p_{n+1}, p_{n+2} , and p_0 transformed into p'_{n+1}, p'_{n+2} , and p'_0 both by τ and π , will have every one of its points transformed into the same corresponding point of $p'_{n+1} p'_{n+2}$ both by τ and π . This is true for every line $p_i p_j$ ($i, j = 1, 2, 3, \dots, n + 2; i \neq j$). Every plane $p_i p_j p_k$ will go into the plane $p'_i p'_j p'_k$ and every point in the first plane will have the same correspondent in the second plane both by τ and π . Let p_{ijk} be any point in the plane $p_i p_j p_k$. Draw the lines $p_i p_{ijk}$ and $p_j p_{ijk}$. These lines will intersect the lines $p_j p_k$ and $p_i p_k$ in points that we can call p_{jk} and p_{ik} . Transformations τ and π will transform p_i, p_j, p_{jk} , and p_{ik} into the same points, that is into p'_i, p'_j, p'_{jk} , and p'_{ik} . To p_{ijk} will correspond the intersection of $p'_i p'_{jk}$ with $p'_j p'_{ik}$ by either τ or π . By continuing this process we can prove that to any point p there corresponds by either τ or π the same point p' . Hence τ and π are identical.

If we desire the set of collineations from which we start to be the group of all projective collineations we add Postulate VII to the postulates we have used and we have the theorem.

THEOREM 22. *Every projective collineation is a collineation.*

INDEPENDENCE EXAMPLES, $n \geq 3$

In the following a Roman numeral preceded by a minus sign denotes an example of a system in which the postulate denoted by that numeral is false but all the other postulates of the set I–VIII are true.*

– I. Let the class of points consist of a single element and the class of collineations of the identity transformation.

– II. Case 1, n odd. Let the class of points consist of $2(n+1)$ elements in $(n+1)/2$ sets of four. Let the collineations be all the transformations that permute the points of each set amongst themselves, the identity transformation excepted, together with all the transformations that permute the points of each of all but two sets amongst themselves but leaves one point of each of the two remaining sets invariant and transforms the other three points of each of these sets into the remaining three points of the other set. Each set is a line and there are no other lines.

Case 2, n even. Let the class of points consist of $2n+1$ points in $n/2$ sets of four and one single point. Let the collineations be transformations on the $n/2$ sets like those of Case 1 leaving the extra point invariant together with the transformations that permute the points of each of all but one of the sets amongst themselves, interchange the single point with one of the points of the remaining set and leaves none of the three other points of that set invariant. Each set is again a line and there are no other lines.

– III. Let the class of points be $2(n+1)$ elements in $n+1$ sets of two. The collineations are the 2^{n+1} transformations on the points leaving each pair invariant. The lines are the $n(n+1)/2$ tetrads of points each consisting of two pairs.

– IV. Case 1, n odd. Let the class of points consist of $2(n+1)$ elements in $(n+1)/2$ sets of four. Let the class of collineations be all the transformations that permute the points of each set or interchange the sets. Each set is a line.

Case 2, n even. Let the class of points consist of $2n+1$ elements in $n/2$ sets of four and one single point. Let the collineations be like those of case 1 on the sets and leave the extra point invariant. Each set is a line.

– V. Ordinary projective geometry of $(n-1)$ -space.

– VI. Ordinary projective geometry of $(n+1)$ -space.

– VII. Let the class of points be the class of all sets of $n+1$ rational numbers $(X_1, X_2, \dots, X_{n+1})$, the set $(0, 0, \dots, 0)$ excepted. The sets $(X_1, X_2, \dots, X_{n+1})$ and $(kX_1, kX_2, \dots, kX_{n+1})$ are understood to be equivalent for all rational values of k distinct from zero. Let the collineations be the class of all linear homogeneous transformations on $n+1$ variables,

*In – I, Postulates III–VIII are satisfied *vacuously*. The same is true of Postulates VII and VIII in – I and – III and of Postulate VII in – IV.

having rational coefficients and whose determinants of transformation are $(n + 1)$ th powers of rational numbers not zero.

— VIII. The finite projective geometry of n -space having three points on a line.*

* For finite projective geometries see O. Veblen and W. H. Bussey, *Finite projective geometries*, these Transactions, vol. 7 (1906), pp 241–259.