

IDEALS IN BIRKHOFF LATTICES

BY

R. P. DILWORTH⁽¹⁾

Introduction. In previous papers by the author (Dilworth [1, 2])⁽²⁾ methods were developed for studying the arithmetical properties of Birkhoff lattices, that is, the properties of irreducibles and decompositions into irreducibles. These methods, however, required the assumption of both the ascending and descending chain conditions. In this paper we give a new technique which is applicable in general and which under the assumption of merely the ascending chain condition gives results quite as good as those of the previous work. Now the descending chain condition is equivalent to the requirement that every ideal⁽³⁾ be principal. Hence if the descending chain condition does not hold we find it convenient to relate the arithmetical properties of the lattice to the structure of its lattice of ideals. Furthermore since the Birkhoff condition itself may lose much of its force if the descending chain condition does not hold, a lattice is defined to be a Birkhoff lattice if every element satisfies the Birkhoff condition⁽⁴⁾ in the lattice of ideals. Hence if the descending chain condition holds, this definition reduces to that used in the previous papers. In the lattice of ideals, the existence of sufficient covering ideals to make the Birkhoff conditions effective can be proved.

In D1 and D2 it was shown that the arithmetical behavior of an element a was closely related to the structure of the quotient lattice \mathfrak{S}_a generated by the elements covering a . Here we make a similar correlation with the structure of the quotient lattice of ideals \mathfrak{I}_a generated by the ideals covering a . The important properties of \mathfrak{S}_a follow from its finite dimensionality. \mathfrak{I}_a on the other hand is in general *not* finite dimensional and thus one of the essential problems of the present treatment is the proof of the archimedean character of \mathfrak{I}_a in the cases of interest.

If the descending chain condition holds, the Birkhoff condition is equivalent to Mac Lane's point-free exchange axiom E_6 (Mac Lane [1]). Now E_6 is independent of covering conditions, which suggests that it should be closely related to the Birkhoff condition in the lattice of ideals. We show that the Birkhoff condition in the lattice of ideals always implies E_6 and, if each principal ideal is covered by only a finite number of ideals, the two conditions are equivalent.

Presented to the Society, December 27, 1939; received by the editors May 11, 1940.

⁽¹⁾ Sterling Research Fellow, Yale University.

⁽²⁾ These papers will be referred to as D1 and D2.

⁽³⁾ An ideal is a sublattice which contains with each element all of its divisors. G. Birkhoff (Birkhoff [1]) uses the term *dual ideal* for such a sublattice.

⁽⁴⁾ See §1, Conditions B1 and B1'.

In D1 it was shown that a lattice of finite dimensions has unique irreducible decompositions if and only if it is a Birkhoff lattice in which every modular sublattice is distributive. This result no longer holds if we drop the descending chain condition as we show by an example. However, by strengthening slightly the condition that every modular sublattice be distributive, we have the following theorem:

THEOREM 6.6. *Let \mathfrak{S} satisfy the ascending chain condition. Then every element of \mathfrak{S} is uniquely expressible as a reduced crosscut of irreducibles if and only if the following conditions hold.*

E_5 . (Mac Lane's point-free exchange axiom.) $a \supset b \supset a \cap c$, $c \neq a \cap c$ implies that $c_1 \neq a \cap c$ exists such that $c \supset c_1 \supset a \cap c$ and $b = a \cap (b \cup c_1)$.

A. $a \cup b \supset x \supset a \cap b$, $a \cap x = b \cap x = a \cap b$ implies $x = a \cap b$.

If we go over to the lattice of ideals, E_5 may be replaced by the condition that \mathfrak{S} be a Birkhoff lattice, and A, by the requirement that the ideals covering a principal ideal generate a Boolean algebra.

In D2, Birkhoff lattices in which the number of components in the irreducible decompositions of each element is unique were characterized in terms of the structure of the quotient lattices \mathfrak{S}_a . We prove here:

THEOREM 5.1. *Let \mathfrak{S} be a Birkhoff lattice satisfying the ascending chain condition and let \mathfrak{I} denote its lattice of ideals. Then the number of components in the irreducible decompositions of each element of the lattice \mathfrak{S} is unique if and only if the ideals covering any principal ideal of the lattice \mathfrak{I} generate a dense, modular sublattice of \mathfrak{I} .*

By means of ideal methods we give a new proof of the Kurosch-Ore decomposition theorem for modular lattices in its most general form. The proof rests on the fact that if an element of a modular lattice has a decomposition into irreducibles then the sublattice generated by the ideals covering the element is of finite dimensions.

Finally §§7 and 8 contain examples which show the complications which may arise when the descending chain condition does not hold.

1. Notation and definitions. The fixed lattice of elements a, b, c, \dots will be denoted by \mathfrak{S} . \cup and \cap will denote union and cross-cut in place of the symbols $(,)$ and $[,]$ used in D1 and D2. \supset denotes lattice division. $a = b$ is defined by the two formulas $a \supset b$, $b \supset a$. If $a \supset b$, $a \neq b$ and $a \supset x \supset b$ implies $a = x$ or $x = b$, we say that a covers b and write $a > b$. Elements which cover the null element z of a lattice are called *points* and elements covered by the unit element u are said to be *simple*.

A lattice \mathfrak{S} satisfies the ascending (descending) chain condition if every chain $a_1 \subset a_2 \subset a_3 \subset \dots$ ($a_1 \supset a_2 \supset a_3 \supset \dots$) has only a finite number of distinct elements. If both the ascending and descending chain conditions hold, \mathfrak{S} is said to be *archimedean* or of *finite dimensions*.

Throughout the paper we shall be particularly interested in lattices which satisfy the following weak form of the modular axiom.

B1. $a > a \cap b \rightarrow a \cup b > b^{(5)}$.

Another form of B1 is the following:

B1'. $b > a, c \supset a, c \not\supset b \rightarrow b \cup c > c$.

If B1' is satisfied for a given a and any b and c we say that a satisfies the *Birkhoff condition* in \mathfrak{S} . Hence B1 holds in \mathfrak{S} if and only if each element of \mathfrak{S} satisfies the Birkhoff condition.

We state now some lemmas on elements satisfying the Birkhoff condition which are refinements of Lemmas 3.1–3.3 of D2.

LEMMA 1.1. *Let a satisfy the Birkhoff condition in \mathfrak{S} and let $a_1, \dots, a_k > a$. Then each union independent⁽⁶⁾ set of the a_i is contained in a maximal independent set.*

The usual proof is valid under the weaker hypotheses of the lemma.

LEMMA 1.2. *Let a satisfy the Birkhoff condition and let $a_1, \dots, a_k > a$. Then each union independent set of the a_i generates a Boolean algebra.*

We note that the usual proof (for example Theorem 2.3 of D1) is not valid in this case since it depends upon the existence of a rank function. Under the hypotheses of the lemma, complete chains need not have the same length and hence a rank function will in general not exist.

Now let A and B be two arbitrary subsets of the set $\{a_1, \dots, a_k\}$. Let $\Sigma(A)$ denote the union of the elements of A and denote the set-theoretic union and cross-cut of A and B by $A \cup B$ and $A \cap B$ respectively. We shall show that

$$(1) \quad \Sigma(A) \cap \Sigma(B) = \Sigma(A \cap B).$$

Let $\mu(A)$ denote the number of elements in A and set $\nu(A) = k - \mu(A)$. If $\nu(A \cap B) = 0$, then $\mu(A \cap B) = k$ and $A = B$. Hence (1) holds. If $\nu(A \cap B) = 1$, then either $A \supset B$ or $B \supset A$ and again (1) holds. Now let (1) hold for all A and B such that $\nu(A \cap B) < l$. Let $\nu(A \cap B) = l$ for some A and B . Then $\mu(A \cap B) = k - l = r$. Hence $A = \{a_1, \dots, a_r, a_{r+1}, \dots, a_s\}$ and $B = \{a_1, \dots, a_r, a'_{r+1}, \dots, a'_t\}$. Since (1) is trivial if $B \supset A$, we may assume that $s > r$. Let $B' = \{a_1, \dots, a_r, a_{r+1}, a'_{r+1}, \dots, a'_t\}$. Now $\mu(A \cap B') = r + 1$ and hence $\nu(A \cap B') = k - (r + 1) = l - 1 < l$. By the induction assumption $\Sigma(A \cap B') = \Sigma(A) \cap \Sigma(B')$. Thus $\Sigma(A \cap B') = \Sigma(A) \cap \Sigma(B') \supset \Sigma(A) \cap \Sigma(B) \supset \Sigma(A \cap B)$.

⁽⁵⁾ \rightarrow denotes formal implication.

⁽⁶⁾ A set of elements x_1, \dots, x_n is said to be *union independent* or simply *independent* if $x_1 \cup \dots \cup x_{i-1} \cup x_{i+1} \cup \dots \cup x_n \not\supset x_i$, $i = 1, \dots, n$. Similarly the set is said to be *cross-cut independent* if $x_i \not\supset x_1 \cap \dots \cap x_{i-1} \cap x_{i+1} \cap \dots \cap x_n$, $i = 1, \dots, n$.

Since a_1, \dots, a_k are independent we have $\Sigma(A \cap B) \not\supset a_{r+1}$ and hence $\Sigma(A \cap B') = a_{r+1} \cup \Sigma(A \cap B) > \Sigma(A \cap B)$. If $\Sigma(A \cap B') = \Sigma(A) \cap \Sigma(B)$, then $\Sigma(B) \supset a_{r+1}$ contrary to the independence of a_1, \dots, a_k . Hence $\Sigma(A) \cap \Sigma(B) = \Sigma(A \cap B)$. Thus (1) holds for $\nu(A \cap B) = l$ and by induction (1) holds for all A and B . Clearly $\Sigma(A) \cup \Sigma(B) = \Sigma(A \cup B)$. If $\Sigma(A) = \Sigma(B)$, then $A = B$ by the independence of a_1, \dots, a_k . Hence the elements which can be expressed as a union of the a_i are isomorphic to the subsets of a_1, \dots, a_k under union and cross-cut and thus a_1, \dots, a_k generate a Boolean algebra. This completes the proof of the lemma.

LEMMA 1.3. *Let a satisfy the Birkhoff condition and let $a_1, \dots, a_k > a$. Then any two maximal union independent sets of the a_i have the same number of elements and any element of one set may be replaced by a suitably chosen element of the other without altering the maximal property.*

The usual proof is valid in this case.

LEMMA 1.4. *Let a satisfy the Birkhoff condition and let $a_1, \dots, a_k > a$. Then any chain joining $a_1 \cup \dots \cup a_k$ to a has not more than $k+1$ distinct members.*

We may clearly suppose that a_1, \dots, a_k are independent. Let $a = b_0 \subset b_1 \subset b_2 \subset \dots \subset b_{l-1} \subset b_l = a_1 \cup \dots \cup a_k$ be a chain joining $a_1 \cup \dots \cup a_k$ to a having $l+1$ distinct members and let us assume that $l > k$. Clearly $b_0 < b_0 \cup a_1 < \dots < b_0 \cup a_1 \cup \dots \cup a_{k-1} < a_1 \cup \dots \cup a_k$ by the Birkhoff condition. Now suppose that it has been shown that $a \subset b_1 \subset \dots \subset b_i \subset b_i \cup a_1 < \dots < b_i \cup a_1 \cup \dots \cup a_{k_i-1} < a_1 \cup \dots \cup a_k$ where $k_i \leq k-i$ and $i < k$. Consider the chain $a \subset b_1 \subset \dots \subset b_i \subset b_{i+1} \subset b_{i+1} \cup a_1 \subset \dots \subset b_{i+1} \cup a_1 \cup \dots \cup a_{k_i-1} \subset a_1 \cup \dots \cup a_k$. Let us assume that all of the members of this chain are distinct. If $b_i \cup a_1 \cup \dots \cup a_{k_i-1} \not\supset b_{i+1}$, then $a_1 \cup \dots \cup a_k \supset b_{i+1} \cup a_1 \cup \dots \cup a_{k_i-1} \supset b_i \cup a_1 \cup \dots \cup a_{k_i-1}$ and $b_{i+1} \cup a_1 \cup \dots \cup a_{k_i-1} \neq b_i \cup a_1 \cup \dots \cup a_{k_i-1}$. But $a_1 \cup \dots \cup a_k > b_i \cup a_1 \cup \dots \cup a_{k_i-1}$ and hence $a_1 \cup \dots \cup a_k = b_{i+1} \cup a_1 \cup \dots \cup a_{k_i-1}$ contrary to our assumption. Thus $b_i \cup a_1 \cup \dots \cup a_{k_i-1} \supset b_{i+1}$. If $b_i \cup a_1 \cup \dots \cup a_{k_i-2} \not\supset b_{i+1}$, we have $b_i \cup a_1 \cup \dots \cup a_{k_i-1} \supset b_{i+1} \cup a_1 \cup \dots \cup a_{k_i-2} \supset b_i \cup a_1 \cup \dots \cup a_{k_i-2}$. But $b_i \cup a_1 \cup \dots \cup a_{k_i-1} > b_i \cup a_1 \cup \dots \cup a_{k_i-2}$ and hence $b_i \cup a_1 \cup \dots \cup a_{k_i-1} = b_{i+1} \cup a_1 \cup \dots \cup a_{k_i-2}$ contrary to our assumption. Thus $b_i \cup a_1 \cup \dots \cup a_{k_i-2} \supset b_{i+1}$. Continuing in this manner we eventually have $b_i \cup a_1 \supset b_{i+1}$. But then $b_i \cup a_1 \supset b_{i+1} \supset b_i$ and $b_{i+1} \neq b_i$. Hence $b_{i+1} = b_i \cup a_1 = b_{i+1} \cup a_1$ which contradicts our assumption. We conclude, then, that at least two members of the above chain are equal. Thus (renumbering the a 's if necessary) using the Birkhoff condition we have $a \subset b_1 \subset \dots \subset b_i \subset b_{i+1} < b_{i+1} \cup a_1 < \dots < b_{i+1} \cup a_1 \cup \dots \cup a_{k_{i+1}-1} < a_1 \cup \dots \cup a_k$ where $k_{i+1} \leq k_i - 1 \leq k - (i+1)$. By induction, we get $a = b_0 \subset b_1 \subset \dots \subset b_{r-1} < a_1 \cup \dots \cup a_k$ where $r \leq k$. But then $b_r = a_1 \cup \dots \cup a_k$ and hence $r = l$ which contradicts $l > k$. Thus $l \leq k$ and the lemma follows.

The dual of condition B1 is the condition

$$\text{B2. } a \cup b > b \rightarrow a > a \cap b.$$

G. Birkhoff (Birkhoff [2]) has proved the following lemma which relates B1 and B2 to modularity.

LEMMA 1.5. *An archimedean lattice \mathfrak{S} is modular if and only if B1 and B2 are satisfied.*

2. **Lattice ideals.** A sublattice \mathfrak{a} of \mathfrak{S} is said to be an *ideal* if $x \supset a$, $a \in \mathfrak{a}$ implies $x \in \mathfrak{a}$. If \mathfrak{a} consists of all elements x such that $x \supset a$ for a fixed a , then \mathfrak{a} is said to be a *principal* ideal and we write $\mathfrak{a} = (a)$. Now suppose that \mathfrak{S} satisfies the descending chain condition. Then the set of elements in \mathfrak{a} has a cross-cut which can be expressed as a cross-cut of a finite number of them and hence belongs to \mathfrak{a} . Thus \mathfrak{a} consists of all divisors of a fixed element of \mathfrak{S} and hence is principal. Conversely, if every ideal of \mathfrak{S} is principal, then a descending chain $a_1 \supset a_2 \supset \cdots$ generates an ideal \mathfrak{a} which consists of all x such that $x \supset a_k$ for some k . But then $\mathfrak{a} = (a)$ and $a \supset a_k$ for some k . Hence $a = a_k = a_{k+1} = \cdots$ and every descending chain has only a finite number of distinct elements. We thus have

LEMMA 2.1. *\mathfrak{S} satisfies the descending chain condition if and only if every ideal is principal.*

The set of ideals of \mathfrak{S} will be denoted by \mathfrak{I} .

DEFINITION 2.1. *The union $\mathfrak{a} \cup \mathfrak{b}$ of two ideals \mathfrak{a} and \mathfrak{b} is the set of all elements x such that $x \supset a \cup b$ for some $a \in \mathfrak{a}$ and $b \in \mathfrak{b}$. Similarly the cross-cut $\mathfrak{a} \cap \mathfrak{b}$ is the set of all elements y such that $y \supset a \cap b$ for some $a \in \mathfrak{a}$ and $b \in \mathfrak{b}$.*

It is readily verified that the union and cross-cut so defined are ideals and that \mathfrak{I} is a lattice under these operations. The union $\mathfrak{a} \cup \mathfrak{b}$ is simply the set-theoretic cross-cut of \mathfrak{a} and \mathfrak{b} .

The definition of cross-cut may be readily extended to any subset S of \mathfrak{I} . $\Pi(S)$ consists of all elements of \mathfrak{S} which belong to the cross-cut of a finite number of ideals of S . If \mathfrak{S} has a unit element u , the union $\Sigma(S)$ is also defined and is simply the set-theoretic cross-cut of the ideals of S .

If \mathfrak{a} and \mathfrak{b} are principal ideals $\mathfrak{a} = (a)$ and $\mathfrak{b} = (b)$, then by Definition 2.1 $\mathfrak{a} \cup \mathfrak{b} = (a \cup b)$ and $\mathfrak{a} \cap \mathfrak{b} = (a \cap b)$. Hence the set of principal ideals forms a sublattice of \mathfrak{I} which is isomorphic to \mathfrak{S} and we may thus consider \mathfrak{S} as a sublattice of \mathfrak{I} .

LEMMA 2.2. *\mathfrak{I} is a modular (distributive) if and only if \mathfrak{S} is modular (distributive).*

Since \mathfrak{S} is a sublattice of \mathfrak{I} , the modularity (distributivity) of \mathfrak{I} implies the modularity (distributivity) of \mathfrak{S} .

Now let \mathfrak{S} be distributive and let $x \in a \cup (b \cap c)$. Then $x \supset a \cup (b \cap c)$ where $a \in \mathfrak{a}$, $b \in \mathfrak{b}$ and $c \in \mathfrak{c}$ by Definition 2.1. But then $x \supset a \cup (b \cap c) \supset (a \cup b) \cap (a \cup c)$ since \mathfrak{S} is distributive and hence $x \in (a \cup b) \cap (a \cup c)$. Thus $a \cup (b \cap c) \supset (a \cup b) \cap (a \cup c)$. But $(a \cup b) \cap (a \cup c) \supset a \cup (b \cap c)$ trivially. Hence \mathfrak{L} is distributive. Now let \mathfrak{S} be modular. Suppose $a \supset b$ and $x \in b \cup (a \cap c)$. Then $x \supset b \cup (a \cap c)$ where $a \in \mathfrak{a}$, $b \in \mathfrak{b}$ and $c \in \mathfrak{c}$ by Definition 2.1. Now since $a \supset b$ we have $a \supset b_1$ where $b_1 \in \mathfrak{b}$. But then $x \supset (b \cap b_1) \cup (a \cap c)$ and $a \supset b \cap b_1$ where $b_1 \cap b \in \mathfrak{b}$. Hence $x \supset a \cap ((b \cap b_1) \cup c)$ since \mathfrak{S} is modular and $x \in a \cap (b \cup c)$. Thus $b \cup (a \cap c) \supset a \cap (b \cup c)$ and since $a \cap (b \cup c) \supset b \cup (a \cap c)$ trivially, \mathfrak{L} is modular. This completes the proof.

LEMMA 2.3. *Let $a \supset b \supset \cdots \supset u \supset \cdots$ be a chain of ideals such that $u \supset (a)$ and $u \neq (a)$ for all ideals of the chain. Then if \mathfrak{p} is the cross-cut of the ideals of the chain, $\mathfrak{p} \supset (a)$ and $\mathfrak{p} \neq (a)$.*

We note that \mathfrak{p} is the set-theoretic union of the elements of the ideals a, b, \cdots, u, \cdots . For if $x \in \mathfrak{p}$, then x divides a finite cross-cut of the ideals of the chain and hence divides some ideal of the chain. Now suppose $\mathfrak{p} = a$. Then $a \in \mathfrak{p}$ and $a \in u$ for some u . But then $u = (a)$ contrary to assumption. Hence $\mathfrak{p} \neq a$.

The results so far have been independent of the well ordering hypothesis. However, to prove the fundamental property of the ideals we must assume that the elements of \mathfrak{S} can be well ordered. This will be assumed through the remainder of the paper.

THEOREM 2.1. *Let $b \supset (a)$ and $b \neq (a)$. Then there exists an ideal \mathfrak{p} such that $b \supset \mathfrak{p} > (a)$.*

Proof. Let U be the set of all elements x such that $x \supset a$. Let U be well ordered, $U = \{x_\nu\}$, $\nu < \sigma$. Define $a_0 = b$. Now suppose that a_μ has been defined for all $\mu < \nu$ in such a way that $a_\mu \neq (a)$, $a_\mu \supset a_{\mu'}$ if $\mu \leq \mu'$, and $a_\mu \cap x_\mu = a_\mu$ or $a_\mu \cap x_\mu = a$. Let c_ν be the cross-cut of all a_μ with $\mu < \nu$. Then $c_\nu \neq (a)$ by Lemma 2.3. If $c_\nu \cap x_\nu \neq (a)$, let $a_\nu = c_\nu \cap x_\nu$; otherwise let $a_\nu = c_\nu$. Then $a_\nu \neq (a)$ and $a_\mu \supset a_\nu$, all $\mu < \nu$. Clearly $a_\nu \cap x_\nu = a$ or a_ν . Now let $\mathfrak{p} = \Pi_{\nu < \sigma} a_\nu$. Then $\mathfrak{p} \neq a$ by Lemma 2.3 and $b \supset \mathfrak{p}$. If $\mathfrak{p} \supset a \supset (a)$ and $\mathfrak{p} \neq a$, there exists an element $x \in \mathfrak{p}$ such that $x \notin \mathfrak{p}$. Since $x \supset a$ we have $x = x_\nu$ for some ν . But then $a_\nu \cap x = a$ since otherwise $x_\nu \supset a_\nu \supset \mathfrak{p}$ which contradicts $x \notin \mathfrak{p}$. Thus $a = a_\nu \cap x \supset \mathfrak{p} \cap a = a \supset a$ and $a = (a)$. Hence $\mathfrak{p} > a$.

In the special instances of Boolean algebras and distributive lattices, Theorem 2.1 gives respectively the existence of the prime ideals of Stone (Stone [1]) and the maximal collections of Wallman (Wallman [1]).

We next prove a theorem which enables us to pass from ideal relations to the corresponding element relations. The following lemma is required.

LEMMA 2.4. *Let $\mathfrak{a} = \mathfrak{a}(a_1, \cdots, a_n)$ be an ideal obtained from the ideals*

a_1, \dots, a_n by forming a finite number of unions and cross-cuts. Then if $x \in a$, there exist elements a_1, \dots, a_n , $a_i \in a_i$, such that $x \supset a(a_1, \dots, a_n)$.

For let $n(a)$ denote the number of union and cross-cut symbols in the expression $a(a_1, \dots, a_n)$. Suppose that the lemma is true for all expressions a for which $n(a) < k$. Let $n(a) = k$. Then $a = a_1 \circ a_2$ where \circ is either \cap or \cup and $n(a_1) < k$, $n(a_2) < k$. Now if $x \in a$ we have $x \supset x_1 \circ x_2$ where $x_1 \in a_1$ and $x_2 \in a_2$ by the definition of union and cross-cut. But then by the induction assumption elements a'_1, \dots, a'_n and a''_1, \dots, a''_n exist such that $x_1 \supset a_1(a'_1, \dots, a'_n)$, $x_2 \supset a_2(a'_1, \dots, a'_n)$. Let $a_i = a'_i \cap a''_i$. Then $x \supset x_1 \circ x_2 \supset a_1(a'_1, \dots, a'_n) \circ a_2(a'_1, \dots, a'_n) \supset a_1(a_1, \dots, a_n) \circ a_2(a_1, \dots, a_n) = a(a_1, \dots, a_n)$ and a_i is clearly in a_i . Since the lemma is trivially true when $n(a) = 1$ by Definition 2.1, the proof is complete.

THEOREM 2.2. Let $(a) = a(a_1, \dots, a_n)$ where a is obtained from a_1, \dots, a_n by forming a finite number of union and cross-cuts. Then $(a) = a(a_1, \dots, a_n)$ where $a_i \in a_i$.

Proof. By Lemma 2.4 $a \supset a(a_1, \dots, a_n)$ where $a_i \in a_i$. But then $a(a_1, \dots, a_n) \supset a(a_1, \dots, a_n) = (a)$. Hence $(a) = a(a_1, \dots, a_n)$.

As an example, if $a = a_1 \cap \dots \cap a_n$ then elements $a_i \in a_i$ exist such that $a = a_1 \cap \dots \cap a_n$.

We conclude this section with two useful lemmas on irreducibles⁽⁷⁾.

LEMMA 2.5. If q is irreducible in \mathfrak{S} , then q is irreducible in \mathfrak{L} .

For if q is reducible in \mathfrak{L} , then $q = a \cap b$, $a, b \neq q$. But then $q = a \cap b$, $a \notin a$, $b \in b$ by Theorem 2.2. Clearly $a \neq q$ and $b \neq q$. Hence q is reducible in \mathfrak{S} . Inverting the logic gives the lemma.

LEMMA 2.6. Let every element of \mathfrak{S} be expressible as a cross-cut of irreducibles. Then if $a \supset b$, $a \neq b$, there exists an irreducible q of \mathfrak{S} such that $q \supset b$, $q \not\supset a$.

For since $a \neq b$, b exists such that $b \in b$, $b \notin a$. Let $b = q_1 \cap \dots \cap q_k$. If $q_i \in a$ for every i then $b \in a$ contrary to assumption. Hence $q_i \notin a$ for some i . But then $q_i \supset b \supset b$.

3. Birkhoff lattices. In D1 and D2 a lattice satisfying B1 was defined to be a Birkhoff lattice. Since both the ascending and descending chain conditions were assumed to hold, B1 was never satisfied trivially. Now in a sufficiently general lattice no covering relations may exist and B1 will hold vacuously. Hence we formulate a more general definition which reduces to that used in D1 and D2 if the descending chain condition holds.

DEFINITION 3.1. A lattice \mathfrak{S} is said to be a Birkhoff lattice if each element of \mathfrak{S} satisfies the Birkhoff condition in the lattice of ideals.

(7) An element q is said to be *cross-cut irreducible* or simply *irreducible* if $q = a \cap b \rightarrow q = a$ or $q = b$. q is said to be *union irreducible* if $q = a \cup b \rightarrow q = a$ or $q = b$.

A lattice \mathfrak{S} is never vacuously a Birkhoff lattice since by Theorem 2.1 covering ideals always exist. Furthermore if the descending chain condition holds, then every ideal is principal and \mathfrak{S} is a Birkhoff lattice if and only if B1 holds in \mathfrak{S} .

Now if \mathfrak{S} has a unit element u and a is any element of \mathfrak{S} , then the union of the ideals covering a exists and will be denoted by u_a . Let \mathfrak{L}_a denote the quotient lattice of all ideals of \mathfrak{L} which are divisible by u_a and which divide a . Then \mathfrak{L}_a is a dense sublattice of \mathfrak{L} and every proper divisor of a in \mathfrak{L} divides some point ideal of \mathfrak{L}_a by Theorem 2.1. Clearly \mathfrak{L}_a reduces to the sublattice \mathfrak{S}_a of the previous papers if the descending chain condition holds. The essential properties of \mathfrak{S}_a followed from its finite dimensionality. But \mathfrak{L}_a is in general *not* finite dimensional. However we now prove a theorem which insures the archimedean character of \mathfrak{L}_a in most cases of arithmetical interest. We need the following lemma:

LEMMA 3.1. *Let \mathfrak{S} be a Birkhoff lattice. Then if p_1, \dots, p_k is a maximal independent set of point ideals of \mathfrak{L}_a , the length of any chain of \mathfrak{L}_a is not greater than k .*

Since the length of any chain is one less than the number of distinct members of the chain, the lemma follows immediately from Lemma 1.4 and Definition 3.1.

According to Lemma 3.1, \mathfrak{L}_a is archimedean if and only if u_a can be expressed as a union of a finite number of point ideals of \mathfrak{L}_a .

THEOREM 3.1. *Let \mathfrak{S} be a Birkhoff lattice in which every element may be represented as a cross-cut of irreducibles. Then \mathfrak{L}_a is archimedean if and only if the number of components in the irreducible decompositions of a is bounded.*

Proof. Let the number of components in the irreducible decompositions of a be bounded, say less than n . Then if \mathfrak{L}_a is not archimedean, by Lemmas 1.2 and 3.1 there are n union independent point ideals p_1, \dots, p_n of \mathfrak{L}_a which generate a Boolean algebra. Let $a_i = p_1 \cup \dots \cup p_{i-1} \cup p_{i+1} \cup \dots \cup p_n$. Then $a = a_1 \cap \dots \cap a_n$. Hence by Theorem 2.2 $a = a_1 \cap \dots \cap a_n$ where $a_i \in a_i$. Now let $a_i = q_{i1} \cap \dots \cap q_{ik_i}$ where q_{i1}, \dots, q_{ik_i} are irreducibles of \mathfrak{S} . Then $a = q_{11} \cap q_{12} \cap \dots \cap q_{nk_n}$ and this representation may be reduced⁽⁸⁾ by dropping our superfluous irreducibles. However not all of the irreducibles belonging to any one a_i may be dropped out since otherwise $a = q_{11} \cap \dots \cap q_{nk_n} \supset a_1 \cap \dots \cap a_{i-1} \cap a_{i+1} \cap \dots \cap a_n \supset a_1 \cap \dots \cap a_{i-1} \cap a_{i+1} \cap \dots \cap a_n \supset p_i$ contrary to $p_i > a$. Hence a has a decomposition having at least n components. But this contradicts our assumption that the number of components is less than n . Hence \mathfrak{L}_a is archimedean and of length less than n .

On the other hand let the number of components be unbounded. Then for

⁽⁸⁾ A representation $a = a_1 \cap a_2 \cap \dots \cap a_n$ is said to be reduced if a_1, \dots, a_n are cross-cut independent.

every k there is an irreducible decomposition $a = q_1 \cap \cdots \cap q_n$ with $n \geq k$. Let $q'_i = q_1 \cap \cdots \cap q_{i-1} \cap q_{i+1} \cap \cdots \cap q_n$. Then $q'_i \supset a$ and $q'_i \neq a$ since the representation is reduced. Hence $q'_i \supset p_i > a$ by Theorem 2.1. Suppose $p_1 \cup \cdots \cup p_{i-1} \cup p_{i+1} \cup \cdots \cup p_n \supset p_i$. Then $q_i \supset q'_1 \cup \cdots \cup q'_{i-1} \cup q'_{i+1} \cup \cdots \cup q'_n \supset p_1 \cup \cdots \cup p_{i-1} \cup p_{i+1} \cup \cdots \cup p_n \supset p_i$ and $a = q_i \cap q'_i \supset p_i$ which contradicts $p_i > a$. Thus p_1, \cdots, p_n are union independent. Hence for every k there are more than k union independent point ideals of \mathfrak{L}_a and \mathfrak{L}_a is *not* archimedean.

If \mathfrak{L}_a is archimedean it has some simple structure properties which follow from the Birkhoff condition.

THEOREM 3.2. *Let \mathfrak{S} be a Birkhoff lattice. Then if \mathfrak{L}_a is archimedean, it is complemented and every ideal can be expressed as a cross-cut of simple ideals.*

Proof. Let $a \in \mathfrak{L}_a$ and let p_1, \cdots, p_k be a maximal independent set of point ideals of \mathfrak{L}_a divisible by a . Imbed p_1, \cdots, p_k in a maximal independent set p_1, \cdots, p_n . Let $a' = p_{k+1} \cup \cdots \cup p_n$. Then $a \cup a' \supset p_1 \cup \cdots \cup p_n \supset u_a$. Hence $a \cup a' = u_a$. Now suppose that $a \cap a' \neq a$. Then $a \cap a' \supset p > a$ by Theorem 2.1. Since $a \supset p$ we have $p_1 \cup \cdots \cup p_k \supset p$ by the maximal property of p_1, \cdots, p_k and $a = (p_1 \cup \cdots \cup p_k) \cap a' \supset p$, which contradicts $p > a$. Hence $a \cap a' = a$ and \mathfrak{L}_a is complemented.

Now let q be irreducible in \mathfrak{L}_a . Let p_1, \cdots, p_k be a maximal independent set of point ideals of \mathfrak{L}_a divisible by q and let this set be imbedded in a maximal independent set $p_1, \cdots, p_k, \cdots, p_n$. Then $q \not\supset p_{k+1}, \cdots, p_n$ and hence $q \cup p_i > q$, $i = k+1, \cdots, n$, by B1'. But since q is irreducible in \mathfrak{L}_a we have $q \cup p_{k+1} = \cdots = q \cup p_n$. Hence $u_a = q \cup u_a = q \cup p_{k+1} \cup \cdots \cup q \cup p_n = q \cup p_{k+1} > q$. Thus each ideal which is irreducible in \mathfrak{L}_a is a simple ideal of \mathfrak{L}_a and since \mathfrak{L}_a is archimedean each ideal of \mathfrak{L}_a can be represented as a cross-cut of simple ideals.

If \mathfrak{L}_a is not archimedean it will in general neither be complemented nor will every ideal be expressible as a cross-cut of simple ideals⁽⁹⁾. In the archimedean case an arbitrary complement of a in \mathfrak{L}_a will be denoted by a' .

DEFINITION 3.2. *An ideal $c \neq u_a$ of \mathfrak{L}_a is said to be characteristic if there exists an irreducible q of \mathfrak{S} which divides exactly the same point ideals of \mathfrak{L}_a as c .*

THEOREM 3.3. *An element $a \in \mathfrak{S}$ has a reduced representation $a = q_1 \cap \cdots \cap q_n$ where q_1, \cdots, q_n are irreducibles if and only if a has a reduced representation $a = c_1 \cap \cdots \cap c_n$ where c_1, \cdots, c_n are characteristic ideals of \mathfrak{L}_a such that $q_i \supset c_i$.*

Proof. Let $a = q_1 \cap \cdots \cap q_n$ be a reduced representation of a as a cross-cut of irreducibles. If $q_i \supset u_a$ for some i , then $q_1 \cap \cdots \cap q_{i-1} \cap q_{i+1} \cap \cdots \cap q_n \supset p_i > a$ and hence $a = q_1 \cap \cdots \cap q_n \supset p_i > a$, which is impossible. Thus $q_i \not\supset u_a$. Let c_i be a characteristic ideal associated with q_i . There is always at least one

⁽⁹⁾ See §7 for an example.

such ideal, namely, the union of the point ideals of \mathfrak{L}_a divisible by q_i . Now $a = q_1 \cap \cdots \cap q_n \supset c_1 \cap \cdots \cap c_n \supset a$ implies $a = c_1 \cap \cdots \cap c_n$. Suppose $c_i \supset c_1 \cap \cdots \cap c_{i-1} \cap c_{i+1} \cap \cdots \cap c_n$. Then $q_i \cap \cdots \cap q_{i-1} \cap q_{i+1} \cap \cdots \cap q_n \supset p_i \supset a$ implies $c_1 \cap \cdots \cap c_{i-1} \cap c_{i+1} \cap \cdots \cap c_n \supset p_i$. But then $a = c_i \cap c_1 \cap \cdots \cap c_{i-1} \cap c_{i+1} \cap \cdots \cap c_n \supset p_i$ which is impossible. Hence the representation $a = c_1 \cap \cdots \cap c_n$ is reduced.

Now let $a = c_1 \cap \cdots \cap c_n$ where c_1, \dots, c_n are characteristic ideals and the representation is reduced. Let q_1, \dots, q_n be associated irreducibles. Suppose $q_1 \cap \cdots \cap q_n \supset p \supset a$. Then $a = c_1 \cap \cdots \cap c_n \supset p \supset a$ which is impossible. Hence $a = q_1 \cap \cdots \cap q_n$. It follows easily that this representation is reduced.

The characteristic ideals of \mathfrak{L}_a can be characterized in terms of the structure of \mathfrak{L} as follows:

THEOREM 3.4. *Let \mathfrak{S} be a Birkhoff lattice in which each element can be expressed as a cross-cut of irreducibles. Then if \mathfrak{L}_a is archimedean, c is characteristic if and only if there exists an ideal $\mathfrak{r} \in \mathfrak{L}$ such that $\mathfrak{r} \supset c$, $c' \cup \mathfrak{r} > \mathfrak{r}$ and $c' \cap \mathfrak{r} = a$ for every c' .*

Proof. Let us first assume that such an ideal \mathfrak{r} exists. Then $u_a \cup \mathfrak{r} = c \cup c' \cup \mathfrak{r} = c' \cup \mathfrak{r}$. Let q be an irreducible such that $q \supset \mathfrak{r}$, $q \nmid u_a \cup \mathfrak{r}$ (Lemma 2.6). Since $q \supset \mathfrak{r} \supset c$, q divides every point ideal of \mathfrak{L}_a which c divides. Now let $q \supset p$. Then if $\mathfrak{r} \nmid p$ we have $c' \cup \mathfrak{r} = u_a \cup \mathfrak{r} \supset p \cup \mathfrak{r} \supset \mathfrak{r}$ and $p \cup \mathfrak{r} \neq \mathfrak{r}$. Hence $c' \cup \mathfrak{r} = p \cup \mathfrak{r}$ and $q \supset p \cup \mathfrak{r} \supset c' \cup \mathfrak{r}$ which contradicts the definition of q . Hence $\mathfrak{r} \supset p$. Now if $c \nmid p$, then $c' \supset p$ for some c' . But then $a = c' \cap \mathfrak{r} \supset p$ which is impossible. Hence $q \supset p$ implies $c \supset p$ and c is thus characteristic.

On the other hand let c be characteristic and let q be an irreducible associated with c . Then $q \cup c' > q$ for every c' . For there is a point ideal p such that $c' \supset p$, $c \nmid p$ since otherwise we would have $c' = a$ and $c = u_a$ contrary to the definition of a characteristic ideal. Now $q \cup p = q \cup u_a > q$ since q is irreducible in \mathfrak{L} by Lemma 2.5. Hence $q \cup u_a = q \cup c' = q \cup p > q$. Now if $c' \cap q \neq a$, then $c' \cap q \supset p > a$ and hence $c' \supset p$, $q \supset p$ by Theorem 2.2. But then $c \supset p$ and hence $a = c \cap c' \supset p$ which is impossible. Thus $c' \cap q = a$ for every c' .

COROLLARY 3.1. *Each simple ideal of \mathfrak{L}_a is characteristic.*

We may take \mathfrak{r} to be the simple ideal itself.

THEOREM 3.5. *Let \mathfrak{S} be a Birkhoff lattice in which each element can be expressed as a cross-cut of irreducibles. Then if \mathfrak{L}_a is archimedean, each characteristic ideal c of \mathfrak{L}_a occurs in a reduced representation $a = c \cap c_1 \cap \cdots \cap c_k$ where k is the number of maximal independent point ideals divisible by c and c_1, \dots, c_k are characteristic ideals of \mathfrak{L}_a .*

Proof. Let p_1, \dots, p_k be a maximal independent set of point ideals of \mathfrak{L}_a divisible by c . Imbed p_1, \dots, p_k in a maximal independent set $p_1, \dots, p_k, \dots, p_n$. Let $c_i = p_1 \cup \cdots \cup p_{i-1} \cup p_{i+1} \cup \cdots \cup p_k \cup \cdots \cup p_n$, $i = 1, \dots, k$. If $c \cap c_1$

$\cap \cdots \cap c_k \neq a$ we have $c \cap c_1 \cap \cdots \cap c_k \supset p > a$ and $c \supset p$ implies $p_1 \cup \cdots \cup p_k \supset p$. But then $a = (p_1 \cup \cdots \cup p_k) \cap c_1 \cap \cdots \cap c_k \supset p$ which is impossible. Hence $a = c \cap c_1 \cap \cdots \cap c_k$. Also since $c \cap c_1 \cap \cdots \cap c_{i-1} \cap c_{i+1} \cap \cdots \cap c_k \supset p_i$ the representation is reduced. Since c_1, \cdots, c_k are simple ideals of \mathfrak{L}_a , they are characteristic by Corollary 3.1.

COROLLARY 3.2. *Let \mathfrak{S} be a Birkhoff lattice in which every element can be expressed as a cross-cut of irreducibles. Then if \mathfrak{L}_a is archimedean of length k , a has a reduced decomposition into irreducibles with k components.*

For by Lemma 1.2 and Theorem 3.4, a has a reduced representation as a cross-cut of k characteristic ideals of \mathfrak{L}_a .

LEMMA 3.2. *Let \mathfrak{S} be a Birkhoff lattice and let \mathfrak{L}_a be archimedean for some a . Then \mathfrak{L}_a is modular if and only if it satisfies B2.*

For let \mathfrak{L}_a satisfy B2 and let q be a union irreducible ideal of \mathfrak{L}_a . If $\mathfrak{s} \nmid q$ and \mathfrak{s} is a simple ideal of \mathfrak{L}_a we have $q > q \cap \mathfrak{s}$ by B2. Hence since q is union irreducible we have $q \cap \mathfrak{s} = q \cap \mathfrak{s}'$ for any two simple ideals \mathfrak{s} and \mathfrak{s}' which do not divide q . Let $a = \mathfrak{s}_1 \cap \cdots \cap \mathfrak{s}_n$ where $\mathfrak{s}_1, \cdots, \mathfrak{s}_i \supset q; \mathfrak{s}_{i+1}, \cdots, \mathfrak{s}_n \nmid q$. Then $a = q \cap a = q \cap \mathfrak{s}_1 \cap \cdots \cap \mathfrak{s}_n = (q \cap \mathfrak{s}_{i+1}) \cap \cdots \cap (q \cap \mathfrak{s}_n) = q \cap \mathfrak{s}_{i+1} < q$. Hence q is a point of \mathfrak{L}_a and every ideal of \mathfrak{L}_a is a union of point ideals. Now let $a > a \cap b$ in \mathfrak{L}_a . Then since every ideal is a union of point ideals, there exists a point ideal p such that $a \supset p, a \cap b \nmid p$. But then $a = (a \cap b) \cup p$. Hence $a \cup b = (a \cap b) \cup p \cup b = p \cup b > b$ since \mathfrak{S} is a Birkhoff lattice. Thus B1 and B2 hold in \mathfrak{L}_a and \mathfrak{L}_a is modular by Lemma 1.5. Conversely, if \mathfrak{L}_a is modular, then B2 is satisfied by Lemma 1.5. This completes the proof.

According to Theorem 3.1, if every element of a lattice \mathfrak{S} has a decomposition into irreducibles and the number of components in the decompositions of a is bounded, then \mathfrak{L}_a is archimedean. This result can be sharpened considerably if \mathfrak{S} is modular.

LEMMA 3.3. *Let \mathfrak{S} be a modular lattice. Then if an element a has a decomposition into irreducibles, \mathfrak{L}_a is archimedean.*

For let $a = q_1 \cap \cdots \cap q_k$ where q_1, \cdots, q_k are irreducible. Since \mathfrak{S} is modular, \mathfrak{L} is modular by Lemma 2.2. Now if $q_i \nmid p$ where $p > a$, we have $q_i \cup p > q_i$ and hence $q_i \cup u_a > q_i$ since q_i is irreducible. But then $u_a > u_a \cap q_i$ since \mathfrak{L} is modular. Thus each irreducible q_i divides a simple characteristic ideal $c_i = q_i \cap u_a$. Since \mathfrak{L} is modular, we have $u_a > c_1 > c_1 \cap c_2 > \cdots > c_1 \cap \cdots \cap c_k = a$. Hence \mathfrak{L}_a is archimedean and the lemma is proved.

If \mathfrak{S} is modular and a has two reduced decompositions into irreducibles, then by Lemma 3.3, \mathfrak{L}_a is archimedean and a has two reduced representations as a cross-cut of simple ideals. Now by Lemma 3.2, B2 holds in \mathfrak{L}_a and hence by the dual of Lemma 1.3 any two reduced representations of a as a cross-cut of simple ideals have the same number of components and any simple ideal

of one decomposition may be replaced by a suitably chosen simple ideal of the other. Thus by Theorem 3.3 and Corollary 3.1 we have the

KUROSCH-ORE DECOMPOSITION THEOREM. *Let an element of a modular lattice have two reduced decompositions into irreducibles. Then the number of components in the two decompositions is the same and any component in one decomposition may be replaced by a suitably chosen component of the other.*

4. Lattices with unique decompositions. This section will be devoted to the proof of the following theorem:

THEOREM 4.1. *Let \mathfrak{S} satisfy the ascending chain condition. Then each element of \mathfrak{S} has a unique representation as a reduced cross-cut of irreducibles if and only if \mathfrak{S} is a Birkhoff lattice and \mathfrak{L}_a is a Boolean algebra for each a .*

We begin with a series of lemmas, the first of which proves the necessity of the conditions of the theorem.

LEMMA 4.1. *Let \mathfrak{S} satisfy the ascending chain condition and let each element have a unique representation as a reduced cross-cut of irreducibles. Then \mathfrak{S} is a Birkhoff lattice and \mathfrak{L}_a is a Boolean algebra for each a .*

For let $b > a$, $c \supset a$ and $c \not\supset b$. If $b \cup c \not\supset c$ we have $b \cup c \supset b \supset c$ where $b \cup c \neq b \neq c$. Since $b \not\supset b$, there exists a $d \in b$ such that $d \not\supset b$. Since $b \neq c$, there exists a c such that $c \in c$, $d \supset c$, and $c \not\supset b$. Furthermore since $c \not\supset b$ there exists an irreducible q_c such that $q_c \supset c$, $q_c \not\supset b$ (Lemma 2.6). But then $b \supset b \cap q_c \supset a$ and if $b = b \cap q_c$ we have $q_c \supset b \cup c \supset b$ which contradicts $q_c \not\supset b$. Hence $a = b \cap q_c$. Similarly there exists an irreducible q_d such that $q_d \supset d$ and $a = b \cap q_d$. By Theorem 2.2 we have $a = b_c \cap q_c$ and $a = b_d \cap q_d$ where $b_c, b_d \in b$. Let $b = b_c \cap b_d$. Then $b \in b$ and $a = b \cap q_c = b \cap q_d$. Let $b = q_{i_1} \cap \cdots \cap q_{i_r}$. Then a has two reduced representations $a = q_{i_1} \cap \cdots \cap q_{i_r} \cap q_c = q_{i_1} \cap \cdots \cap q_{i_m} \cap q_d$. Now $q_c \neq q_d$ since otherwise $q_c \supset b$ and $q_c \neq q_{i_r}$ since otherwise $q_c \supset b \cup c \supset b$ contrary to $q_c \not\supset b$. Hence a has two distinct reduced representations as a cross-cut of irreducibles which contradicts our hypothesis. Thus $b \cup c > c$ and hence each element of \mathfrak{S} satisfies the Birkhoff condition in the lattice of ideals.

Now since each element has a unique decomposition into irreducibles, the number of components is obviously bounded and hence \mathfrak{L}_a is archimedean by Theorem 3.1. Let p_1, \dots, p_k be a maximal independent set of point ideals of \mathfrak{L}_a . Then p_1, \dots, p_k generate a Boolean algebra with simple ideals $\mathfrak{s}_1, \dots, \mathfrak{s}_k$. $\mathfrak{s}_1, \dots, \mathfrak{s}_k$ are clearly simple ideals of \mathfrak{L}_a and hence are characteristic ideals by Corollary 3.2. Thus a has a decomposition $a = q_1 \cap \cdots \cap q_k$ where $q_i \supset \mathfrak{s}_i$ (Theorem 3.3). Now suppose there is a simple ideal \mathfrak{s} distinct from $\mathfrak{s}_1, \dots, \mathfrak{s}_k$. Let $q \supset \mathfrak{s}$, $q \not\supset u_a$. Then q is a component of a by Theorem 3.5 and hence $q = q_i$ for some i since a has but one reduced decomposition into irreducibles. But then $q \supset \mathfrak{s} \cup \mathfrak{s}_i = u_a$ which is impossible. Hence $\mathfrak{s}_1, \dots, \mathfrak{s}_k$ are all of the simple

ideals of \mathfrak{L}_a and since each ideal of \mathfrak{L}_a can be expressed as a union cross-cut of simple ideals, \mathfrak{L}_a is simply the Boolean algebra generated by p_1, \dots, p_k .

LEMMA 4.2. *If \mathfrak{L}_a is a Boolean algebra, then it is archimedean.*

For if \mathfrak{L}_a has an infinite number of point ideals, let p_1, p_2, p_3, \dots be a denumerable sequence of point ideals. Let $p'_i = p_1 \cup p_2 \cup \dots \cup p_{i-1} \cup p_{i+1} \cup \dots$. Then since \mathfrak{L}_a is a Boolean algebra we have $a = p'_1 \cap p'_2 \cap \dots$. But since the cross-cut of an infinite number of ideals consists of all elements contained in finite cross-cuts $a = p'_1 \cap p'_2 \cap \dots \cap p'_k$ for some k . Then $a \supset p_{k+1}$ which contradicts $p_{k+1} > a$. Hence \mathfrak{L}_a has only a finite number of point ideals and thus is archimedean.

LEMMA 4.3. *Let \mathfrak{S} be a Birkhoff lattice in which each \mathfrak{L}_a is archimedean. Then if every three ideals covering a principal ideal generate a Boolean algebra of order eight, \mathfrak{L}_a is a Boolean algebra for each a .*

For let the hypotheses of the lemma be satisfied and let every three ideals covering a principal ideal generate a Boolean algebra. We show first that the ideals of any finite set of ideals covering a principal ideal are independent. Suppose that for any a every $k-1$ ideals covering a are independent. Let p_1, \dots, p_k be k distinct ideals covering a . If p_1, \dots, p_k are not independent let $p_1 \cup p_2 \cup \dots \cup p_{k-1} \supset p_k$ say. Now $p_1 \cup p_i \not\supset p_2, \dots, p_{i-1}, p_{i+1}, \dots, p_k$ ($i=2, \dots, k$) since every three ideals covering a generate a Boolean algebra. Hence elements $x_{ij} \in p_i$ exist such that $x_{ij} \cup p_i \not\supset p_j$ ($j=2, \dots, i-1, i+1, \dots, k$; $i=2, \dots, k$). Let $x = x_{23} \cap x_{24} \cap \dots \cap x_{k, k-1}$. Then $x \in p_1$ and $x \cup p_i \not\supset p_2, \dots, p_{i-1}, p_{i+1}, \dots, p_k$ ($i=2, \dots, k$). Clearly $x \not\supset p_2, \dots, p_k$. Hence $p'_2 = x \cup p_2 > x, \dots, p'_k = x \cup p_k > x$ and p'_2, \dots, p'_k are distinct. Thus by the induction assumption p'_2, \dots, p'_k are independent. But $p'_2 \cup \dots \cup p'_{k-1} \supset x \cup p_2 \cup \dots \cup p_{k-1} \supset x \cup p_1 \cup \dots \cup p_{k-1} \supset x \cup p_k = p'_k$ which is contrary to the independence. Hence the independence of any finite set of covering elements follows by induction.

Now let $a \in \mathfrak{L}_a$ and let p_1, \dots, p_k be a maximal independent set of point ideals of \mathfrak{L}_a divisible by a . Imbed p_1, \dots, p_k in a maximal independent set $p_1, \dots, p_k, \dots, p_n$. Set $b = p_1 \cup \dots \cup p_k$. Then $a \supset b$. If $b \not\supset a$, there exists an element $b_1 \in b$ such that $b_1 \not\supset a$. Now $b \cup p_{k+i} \not\supset p_{k+j}$ ($j=k+1, \dots, k+i-1, k+i+1, \dots, n$; $i=1, \dots, n-k$). Hence as above there exists an element $b_2 \in b$ such that $b_2 \cup p_{k+i} \not\supset p_{k+j}$, $i \neq j$. Also $a = a \cup b \not\supset p_{k+1}, \dots, p_n$. Hence an element $b_3 \in b$ exists such that $a \cup b_3 \not\supset p_{k+1}, \dots, p_n$. Set $b = b_1 \cap b_2 \cap b_3$. Then $b \in b, b \not\supset a, b \cup p_{k+i} \not\supset p_{k+j}$, $i \neq j$, and $a \cup b \not\supset p_{k+1}, \dots, p_n$. Clearly $b \not\supset p_{k+1}, \dots, p_n$. Hence $p'_{k+1} = b \cup p_{k+1} > b, \dots, p'_n = b \cup p_n > b$ and p'_{k+1}, \dots, p'_n are distinct. Let $b \cup a \supset p > b$. Then p is distinct from p'_{k+1}, \dots, p'_n . For if $p = p'_{k+i}$, then $b \cup a \supset p_{k+i}$ contrary to the definition of b . Thus by the result of the above paragraph p, p'_{k+1}, \dots, p'_n are independent. But $p'_{k+1} \cup \dots \cup p'_n = b \cup p_{k+1} \cup \dots \cup p_n = b \cup p_1 \cup \dots \cup p_n \supset b \cup a \supset p$ which is impossible. Hence $a = b$ and

\mathfrak{L}_a is a point lattice. But then the point ideals of \mathfrak{L}_a are independent and generate \mathfrak{L}_a . Thus \mathfrak{L}_a is a Boolean algebra by Lemma 1.2.

LEMMA 4.4. *Let \mathfrak{S} be a Birkhoff lattice satisfying the ascending chain condition in which every three ideals covering a principal ideal generate a Boolean algebra. Let q be an irreducible of \mathfrak{S} such that $q \supset a$; $b, c > a$ and $b \neq c$. Then either $q \supset b$ or $q \supset c$.*

Let us suppose that for some a we have $q \supset a$; $b, c > a$, $b \neq c$, $q \not\supset b$ and $q \not\supset c$. We shall show that a proper divisor a' of a exists with the same properties and hence the lemma follows from the ascending chain condition. Now $q \neq a$ since otherwise $q = b \cap c$ contrary to the irreducibility of q . Hence $q \supset p > a$ by Theorem 2.1. Clearly $p \neq b, c$ since otherwise $q \supset b$ or $q \supset c$. Hence p, b and c generate a Boolean algebra. Since $p \cup b \not\supset c$ there exists an element $p \in p$ such that $p \cup b \not\supset c$. Since $q \supset p$, there exists an element $p' \in p$ such that $q \supset p'$. Let $a' = p \cap p'$. Then $a' \in p$ and hence $a' \neq a$. Clearly $q \supset a'$. Let $b' = a' \cup b$, $c' = a' \cup c$. Then $b' > a'$ and $c' > a'$ by the Birkhoff condition. If $b' = c'$, then $p \cup b \supset a' \cup b \supset c$, which contradicts $p \cup b \not\supset c$. Hence $b' \neq c'$. Since $q \not\supset b$, $q \not\supset c$ we have $q \not\supset b'$, $q \not\supset c'$. Thus a' is a proper divisor of a with the desired properties.

LEMMA 4.5. *Let \mathfrak{S} be a Birkhoff lattice satisfying the ascending chain condition in which every three ideals covering a principal ideal generate a Boolean algebra. Then if a has a reduced representation $a = q_1 \cap \cdots \cap q_k$, \mathfrak{L}_a is archimedean of length k and each q_i divides a simple ideal of \mathfrak{L}_a .*

For let α_i be the union of the point ideals of \mathfrak{L}_a which are divisible by q_i . Then $\alpha_i \neq u_a$ since α_i is a characteristic ideal of \mathfrak{L}_a . Now let p, p' be any two point ideals of \mathfrak{L}_a which are not divisible by α_i . Then $\alpha_i \cup p > \alpha_i$ and $\alpha_i \cup p' > \alpha_i$ by the Birkhoff condition. Now suppose that $\alpha_i \cup p \not\supset p'$. Then there exists an element $a_1 \in \alpha_i$ such that $a_1 \cup p \not\supset p'$. Since $q_i \supset \alpha_i$, there exists an element $a_2 \in \alpha_i$ such that $q_i \supset a_2$. Let $a_i = a_1 \cap a_2$. Then $q_i \supset a_i$ and $\alpha_i \cup p \not\supset p'$. Clearly $\alpha_i \not\supset p'$. If $\alpha_i \supset p$, then $q_i \supset p$ and $\alpha_i \supset p$ contrary to assumption. Hence $\alpha_i \cup p > \alpha_i$, $\alpha_i \cup p' > \alpha_i$ and $\alpha_i \cup p \neq \alpha_i \cup p'$. Since $q_i \supset \alpha_i$ by Lemma 4.4 we have either $q_i \supset \alpha_i \cup p$ or $q_i \supset \alpha_i \cup p'$. Hence $q_i \supset p$ or $q_i \supset p'$. But then $\alpha_i \supset p$ or $\alpha_i \supset p'$ contrary to assumption. Thus $\alpha_i \cup p \supset p'$ and $\alpha_i \cup p = \alpha_i \cup p'$ for every pair of point ideals of \mathfrak{L}_a not divisible by α_i . But then $\alpha_i \cup p = \alpha_i \cup u_a = u_a$ and $u_a > \alpha_i$. Hence α_i is simple and each q_i divides a simple ideal of \mathfrak{L}_a .

Now let $b_0 = u_a$ and let b_i denote the union of the point ideals of \mathfrak{L}_a which are divisible by q_1, \dots, q_i . Then $b_1 = \alpha_1$ and $b_0 > \alpha_1$ by the result we have just obtained. Clearly $b_{l-1} \supset b_l$. If $b_{l-1} = b_l$, let $q_1 \cap \cdots \cap q_{l-1} \cap q_{l+1} \cap \cdots \cap q_k \supset p_l > a$. p_l exists since the representation is reduced. Now $q_1 \cap \cdots \cap q_{l-1} \supset p_l$ and hence $b_{l-1} \supset p_l$. But then $b_l \supset p_l$ and hence $q_l \supset p_l$. Thus $a = q_1 \cap \cdots \cap q_k \supset p_l$ which is impossible. Hence $b_{l-1} \neq b_l$. Now let p and p' be two point ideals divisible by b_{l-1} but not by b_l . If $b_l \cup p \neq b_l \cup p'$ there exists an element $b_l \in b_l$ such that $q_l \supset b_l$, $b_l \cup p > b_l$, $b_l \cup p' > b_l$ and $b_l \cup p \neq b_l \cup p'$. But then $q_l \supset b_l \cup p$

or $q_l \supset b_l \cup p'$ by Lemma 4.4. Hence either $b_l \supset p$ or $b_l \supset p'$ which is contrary to assumption. Hence $b_l \cup p = b_l \cup p'$ for every two point ideals of b_{l-1} which are not divisible by b_l . Thus $b_l \cup p = b_l \cup b_{l-1} = b_{l-1}$ and $b_{l-1} > b_l$ by the Birkhoff condition. Hence we have the chain $u_a > b_1 > b_2 > \cdots > b_k$. But $b_k = a$ and the lemma follows from Lemma 3.1.

LEMMA 4.6. *Let \mathcal{S} be a Birkhoff lattice satisfying the ascending chain condition in which every three ideals covering a principal ideal generate a Boolean algebra. Then each $a \in \mathcal{S}$ has a unique reduced representation $a = q_1 \cap \cdots \cap q_k$ where q_1, \cdots, q_k are irreducibles. \mathcal{L}_a is a Boolean algebra of order 2^k and each q_i divides a simple ideal of \mathcal{L}_a .*

It follows from Lemmas 4.3 and 4.5 that \mathcal{L}_a is a Boolean algebra of order 2^k . q_i divides a simple ideal \mathfrak{s}_i of \mathcal{L}_a by Lemma 4.5. Now let $a = q'_1 \cap \cdots \cap q'_l$ be a reduced decomposition of a . By Lemma 4.5, $l = k$ and q'_i divides a simple ideal \mathfrak{s}_j . Let $b = q'_i \cap q_j$. Then $b \supset \mathfrak{s}_j$, and $b \not\supset u_a$. Let $\mathfrak{s}_j \not\supset p > a$. Then $u_a = \mathfrak{s}_j \cup p$ and $b \cup p = b \cup \mathfrak{s}_j \cup p = b \cup u_a > b$ by the Birkhoff condition. If $q_j \neq b$, we have $q_j \supset p_j > b$ and $p_j \neq p \cup b$ since otherwise $q_j \supset p \cup \mathfrak{s}_j = u_a$. Hence by Lemma 4.4, either $q'_i \supset p_j$ or $q'_i \supset p$. But if $q'_i \supset p_j$, then $b = q'_i \cap q_j \supset p_j > b$ which is impossible. Hence $q'_i \supset p$ and $q'_i \supset u_a$ which is impossible. Thus $q_j = b$ and similarly $q'_i = b$. Hence q'_i is equal to q_j and the two representations are identical. This completes the proof of the lemma.

Lemma 4.1 and Lemma 4.6 together give Theorem 4.1.

In view of Lemma 4.6, lattices with unique irreducible decompositions may be characterized in terms of the local properties of the lattice of ideals as follows:

THEOREM 4.2. *Let \mathcal{S} satisfy the ascending chain condition. Then each element of \mathcal{S} has a unique reduced decomposition into irreducibles if and only if \mathcal{S} is a Birkhoff lattice in which every three ideals covering an element of \mathcal{S} are independent.*

As a corollary to Lemma 4.6 we have

COROLLARY 4.1. *Let \mathcal{S} satisfy the ascending chain condition and let every element of \mathcal{S} have a unique reduced decomposition into irreducibles. Then the number of irreducible components of a is equal to the number of ideals covering a .*

COROLLARY 4.2. *Let \mathcal{S} be a Birkhoff lattice satisfying the ascending chain condition. Then if \mathcal{S} contains a modular, non-distributive sublattice, the lattice of ideals of \mathcal{S} contains a complete⁽¹⁰⁾ modular, non-distributive sublattice of order five.*

For if \mathcal{S} contains a modular, non-distributive sublattice of order five, at

⁽¹⁰⁾ A sublattice \mathcal{L}' of \mathcal{L} is said to be *complete* if $a > b$ in \mathcal{L}' implies $a > b$ in \mathcal{L} .

least one element of \mathfrak{S} does *not* have a unique decomposition into irreducibles. But then there are three ideals covering a principal ideal which are dependent. These three ideals generate a complete, modular, non-distributive sublattice of \mathfrak{L} of order five.

5. Unicity of the number of components. In the previous section lattices with unique irreducible decompositions were completely characterized as Birkhoff lattices with certain special properties. Simple examples show that a similar characterization of lattices in which the *number* of components is unique will require lattices that are considerably more general than Birkhoff lattices. Hence we shall restrict ourselves to the characterization of Birkhoff lattices having the number of components unique. We prove the following theorem:

THEOREM 5.1. *Let \mathfrak{S} be a Birkhoff lattice satisfying the ascending chain condition. Then the number of components in the reduced decompositions of each element into irreducibles is unique if and only if \mathfrak{L}_a is modular for each a .*

As in §4, the proof rests on a series of lemmas.

LEMMA 5.1. *Let \mathfrak{S} be a Birkhoff lattice satisfying the ascending condition. Then the number of components in the irreducible decompositions of a is unique if and only if \mathfrak{L}_a is archimedean, modular, and every characteristic ideal of \mathfrak{L}_a is simple.*

Since the ascending chain condition holds each element of \mathfrak{S} has a decomposition into irreducibles. Now if the number of components in the irreducible decompositions of a is unique it is certainly bounded and hence \mathfrak{L}_a is archimedean by Theorem 3.1. Now let c be a characteristic ideal of \mathfrak{L}_a and let p_1, \dots, p_k be a maximal independent set of point ideals divisible by c . Imbed p_1, \dots, p_k in a maximal independent set $p_1, \dots, p_k, \dots, p_n$. By Theorem 3.5 and Theorem 3.3, a has an irreducible decomposition having $k+1$ components. But by Corollary 3.2 a has a decomposition having n components. Hence if the number of components is unique we have $n=k+1$. But then $p_1 \cup \dots \cup p_k$ is a simple ideal of \mathfrak{L}_a and $u_a \supset c \supset p_1 \cup \dots \cup p_k$, $u_a \neq c$. Hence $c = p_1 \cup \dots \cup p_k$ and c is a simple ideal of \mathfrak{L}_a .

Now let \mathfrak{s} be an arbitrary simple ideal of \mathfrak{L}_a and let a be any ideal of \mathfrak{L}_a such that $\mathfrak{s} \nsubseteq a$. By Theorem 3.2, a has a reduced representation $a = \mathfrak{s}_1 \cap \dots \cap \mathfrak{s}_l$ where $\mathfrak{s}_1, \dots, \mathfrak{s}_l$ are simple ideals of \mathfrak{L}_a . If $a \cap \mathfrak{s} \neq a$, by Theorem 3.2 there exists a simple ideal \mathfrak{s}_{l+2} such that $\mathfrak{s}_{l+2} \nsubseteq a \cap \mathfrak{s}$. Similarly if $a \cap \mathfrak{s} \cap \mathfrak{s}_{l+2} \neq a$, there exists a simple ideal \mathfrak{s}_{l+3} such that $\mathfrak{s}_{l+3} \nsubseteq a \cap \mathfrak{s} \cap \mathfrak{s}_{l+2}$. Thus we eventually have $a \cap \mathfrak{s} \cap \mathfrak{s}_{l+2} \cap \dots \cap \mathfrak{s}_m = a$. Then $a = \mathfrak{s}_1 \cap \dots \cap \mathfrak{s}_l \cap \mathfrak{s} \cap \mathfrak{s}_{l+2} \cap \dots \cap \mathfrak{s}_m$ and since each simple ideal is characteristic this decomposition gives a decomposition into irreducibles with the same number of terms. Hence if the number of components in the irreducible decompositions of a is unique we have $m \geq n$ where n is the length of \mathfrak{L}_a . But $u_a \supset \mathfrak{s}_1 \supset \mathfrak{s}_1 \cap \mathfrak{s}_2 \supset \dots \supset \mathfrak{s}_1 \cap \dots \cap \mathfrak{s}_l \supset \mathfrak{s}_1$

$\cap \cdots \cap \mathfrak{s}_l \cap \mathfrak{s} \supset \mathfrak{s}_1 \cap \cdots \cap \mathfrak{s}_l \cap \mathfrak{s} \cap \mathfrak{s}_{l+2} \supset \cdots \supset \mathfrak{s}_1 \cap \cdots \cap \mathfrak{s}_m = a$ and the ideals of this chain are distinct. Hence $m \leq n$ by Lemma 3.1. Thus $m = n$ and each ideal of the chain covers the ideal which immediately follows. Hence $a > a \cap \mathfrak{s}$. Now let a and b be any two ideals of \mathfrak{L}_a such that $a \cup b > b$. By Theorem 3.2 and ideal \mathfrak{s} exists such that $\mathfrak{s} \supset b$, $\mathfrak{s} \not\supset a \cup b$. But then $b = (a \cup b) \cap \mathfrak{s}$. Hence $a > a \cap \mathfrak{s} = a \cap (a \cup b) \cap \mathfrak{s} = a \cap b$. Thus $a \cup b > b$ implies $a > a \cap b$ and B2 holds in \mathfrak{L}_a . But then \mathfrak{L}_a is modular by Lemma 3.2.

On the other hand let \mathfrak{L}_a be archimedean, modular, and every characteristic ideal be simple. Let $a = q_1 \cap \cdots \cap q_k$ be a reduced decomposition into irreducibles. By Theorem 3.3, a has a reduced representation $a = c_1 \cap \cdots \cap c_k$ where c_i is a characteristic ideal of \mathfrak{L}_a . But then c_i is a simple ideal of \mathfrak{L}_a by assumption. Thus $u_a > c_1 > c_1 \cap c_2 > \cdots > c_1 \cap \cdots \cap c_k = a$ since B2 holds in \mathfrak{L}_a by Lemma 1.5. Hence k is simply the length of \mathfrak{L}_a and every reduced decomposition of a into irreducibles has the same number of components. This completes the proof of the lemma.

LEMMA 5.2. *Let \mathfrak{S} be a Birkhoff lattice satisfying the ascending chain condition. Then if \mathfrak{L}_a is modular for each a , every characteristic ideal of \mathfrak{L}_a is simple.*

Let every characteristic ideal of \mathfrak{L}_b be simple for every proper divisor b of a . We shall show that every characteristic ideal of \mathfrak{L}_a is simple and the lemma follows by the ascending chain condition.

If c is a characteristic ideal of \mathfrak{L}_a which is not simple, let q be an associated irreducible. If x is any element of \mathfrak{S} divisible by q , let q_x denote the union of the point ideals of \mathfrak{L}_x divisible by q . Then since c is a characteristic ideal associated with q we have $c \supset q_a$ and hence q_a is not a simple ideal of \mathfrak{L}_a . Now suppose that for every two point ideals p and p' such that $q_a \not\supset p$, p' we have $q_a \cup p = q_a \cup p'$. Then $u_a = q_a \cup u_a = q_a \cup p > q_a$ and q_a is simple contrary to assumption. Hence there are two point ideals p and p' such that $q_a \not\supset p$, $q_a \not\supset p'$, and $q_a \cup p \neq q_a \cup p'$. Now $q \cap (q_a \cup p \cup p') = (q \cap u_a) \cap (q_a \cup p \cup p') = q_a \cup (q \cap u_a \cap (p \cup p')) = q_a \cup (q \cap (p \cup p'))$ since \mathfrak{L}_a is modular. If $q \cap (p \cup p') \neq a$, we have $q \cap (p \cup p') \supset p_1 > a$. If $p' = p_1$, we have $q \supset p'$ and hence $q_a \supset p'$ contrary to hypothesis. Thus $p_1 \neq p$ and $p_1 \neq p'$. Now $p \cup p' \supset p \cup p_1 \supset p$ and $p \cup p_1 \neq p$. Hence $p \cup p' = p \cup p_1$ by the Birkhoff condition. Since $q \supset p_1$ we have $q_a \supset p_1$ and hence $q_a \cup p \supset p_1 \cup p \supset p'$. But then $q_a \cup p = q_a \cup p'$ which contradicts the definition of p and p' . Thus $q \cap (p \cup p') = a$ and $q \cap (q_a \cup p \cup p') = q_a$.

Now suppose that q_a is not principal. Let X be the set of all elements x such that $q \supset x \supset q_a$, $q \neq x$. If $x \in X$, let $p_x = q \cap (x \cup p \cup p')$. Clearly X generates q_a . We shall show

- (1) There exists an $x_0 \in X$ such that $x \cup p \cup p' > p_x > x$ for all $x \in X$, $x_0 \supset x$.
- (2) The set of ideals p_x , $x \in X$, $x_0 \supset x$, generates q_a .

(1) Since $q_a \cup p \not\supset p'$ and X generates q_a , there exists an element $x_0 \in X$ such that $x_0 \cup p \not\supset p'$. Let $x_0 \supset x$, $x \in X$ and suppose that $x = p_x$. Since x is a proper divisor of a we have $u_x > q_x$. By the Birkhoff condition $x \cup p > x$,

$x \cup p' > x$ and hence $x \cup p, x \cup p'$ belong to \mathfrak{L}_x . Now $q_x \nsubseteq x \cup p \cup p'$ since otherwise $q \supset p$. Hence by the modularity of \mathfrak{L}_x we have $x \cup p \cup p' > q_x \cap (x \cup p \cup p')$. Then $x \cup p \cup p' \supset q \cap (x \cup p \cup p') \supset q_x \cap (x \cup p \cup p')$ and $q \cap (x \cup p \cup p') \neq x \cup p \cup p'$. Thus $x = p_x = q \cap (x \cup p \cup p') = q_x \cap (x \cup p \cup p')$ and hence $x \cup p \cup p' > x$. But then $x \cup p \cup p' \supset x \cup p \supset x$ and if $x = x \cup p$ we have $q \supset p$ which is impossible. Thus $x \cup p = x \cup p \cup p' \supset p'$ and $x_0 \cup p \supset x \cup p \supset p'$ contrary to the definition of x_0 . Hence $x \neq p_x$. Let $p_x \supset p'_x > x$. Clearly $x \cup p \cup p' \supset x \cup p \cup p'_x \supset x \cup p$ and $x \cup p \cup p' > x \cup p$ by the Birkhoff condition. If $x \cup p \cup p'_x = x \cup p$ we have $x \cup p \supset p'_x$ and $x = q \cap (x \cup p) \supset p'_x$ which contradicts $p'_x > x$. Hence $x \cup p \cup p' = x \cup p \cup p'_x$. But then $x \cup p \cup p'_x \supset p_x \supset p'_x$ and $x \cup p \cup p'_x > p'_x$. If $x \cup p \cup p'_x = p_x$, then $q \supset x \cup p \cup p'$ which is impossible. Hence $p_x = p'_x$ and $x \cup p \cup p' > p_x > x$.

(2) Clearly $p_x \supset q_a$ for every x since $p_x \supset x \supset q_a$. Now let $a_1 \in q_a$. Then $a_1 \supset q_a = q \cap (q_a \cup p \cup p')$ and hence $a_1 \supset q \cap (a_2 \cup p \cup p')$ where $a_2 \in q_a$ by Theorem 2.2. Let $x = x_0 \cap a_2$. Then $x \in X$ and $a_1 \supset q \cap (x \cup p \cup p') = p_x$. Hence each element of q_a divides some p_x and thus the ideals p_x generate q_a .

Now let y be an arbitrary element of $q_a \cup p$. Then $y \supset q_a$ and hence $y \supset p_x$ where $x_0 \supset x$ by (2). But then by (1) $x \cup p \cup p' > p_x > x$ and $p_x \nsubseteq p$ since otherwise $q \supset p$. Now $x \cup p \cup p' \supset p_x \cup p \supset p_x$ and $p_x \cup p \neq p_x$. Hence $x \cup p \cup p' = p_x \cup p$ which gives $p_x \cup p \supset p'$. Thus $y \cup p \supset p'$ for every y and hence $q_a \cup p \supset p'$. The assumption that q_a is *not* principal has thus led to a contradiction and we conclude that q_a is principal, say $q_a = (a_1)$. Since a_1 is a proper divisor of a , by hypothesis we have $u_{a_1} > q_{a_1}$. Hence $q_{a_1} \cup (a_1 \cup p \cup p') > q_{a_1}$. But $q_{a_1} \cap (a_1 \cup p \cup p') = q \cap (a_1 \cup p \cup p') = a_1$ and $a_1 \cup p \cup p' \nsubseteq a_1$. Hence \mathfrak{L}_{a_1} is non-modular contrary to assumption. Thus q_a is simple and hence c is a simple ideal of \mathfrak{L}_a .

LEMMA 5.3. *Let \mathfrak{S} be a Birkhoff lattice satisfying the ascending chain condition. Then if \mathfrak{L}_a is modular for every a , \mathfrak{L}_a is archimedean.*

For let $a = q_1 \cap \dots \cap q_n$ be a reduced decomposition of a into irreducibles. Then a has the reduced representation $a = c_1 \cap \dots \cap c_k$ where c_i is a characteristic ideal associated with q_i . By Lemma 5.2, c_i is a simple ideal of \mathfrak{L}_a . Hence since \mathfrak{L}_a is modular we have $u_a > c_1 > c_1 \cap c_2 > \dots > c_1 \cap \dots \cap c_k = a$. Thus \mathfrak{L}_a is archimedean of length k .

Lemmas 5.1–5.3 together give Theorem 5.1.

COROLLARY 5.1. *Let \mathfrak{S} be a Birkhoff lattice satisfying the ascending chain condition. Let the number of components in the reduced decompositions of an element a be unique. Then in any two reduced decompositions of a , each component of one decomposition may be replaced by a suitably chosen component of the other.*

For by Lemma 5.1, the two decompositions give two reduced representations of a as a cross-cut of simple ideals of \mathfrak{L}_a . However, since \mathfrak{L}_a is modular, B2 is satisfied and the replacement property follows from the dual of Lemma 1.3.

COROLLARY 5.2. *Let \mathfrak{S} be a Birkhoff lattice satisfying the ascending chain condition. Then if the number of components in the decompositions of an element a is unique, that number is simply the length of \mathfrak{L}_a .*

COROLLARY 5.3. *Let \mathfrak{S} be a complemented Birkhoff lattice in which every element can be expressed as a cross-cut of a finite number of irreducibles. Then the number of components in the reduced decompositions of the null element z is unique if and only if \mathfrak{S} is a complemented modular lattice of finite dimensions.*

For since \mathfrak{S} is complemented, \mathfrak{L}_z is simply \mathfrak{L} , the lattice of ideals.

6. The Mac Lane exchange axiom. In order to free condition B1 of the covering properties, Mac Lane (Mac Lane [1]) formulated the following axiom.

E₅. If $a \supset b \supset a \cap c$ and $c \neq a \cap c$, then there exists an element $c_1 \neq a \cap c$ such that $c \supset c_1 \supset a \cap c$ and $b = a \cap (b \cup c_1)$.

Mac Lane showed that E_5 is equivalent to a transposition property of chains and in case covering elements exist, that is, if $b \supset a$, $b \neq a$, implies b' exists such that $b \supset b' \supset a$, it reduces to B1. Thus both E_5 and the requirement that each element satisfy the Birkhoff condition in the lattice of ideals are generalizations of B1. We shall be particularly interested in the conditions under which they are equivalent.

THEOREM 6.1. *Every Birkhoff lattice satisfies E_5 .*

Proof. Let $a \supset b \supset a \cap c$ and $c \neq a \cap c$. Then by Theorem 2.1 and ideal \mathfrak{p} exists such that $c \supset \mathfrak{p} \supset a \cap b$. Now $b \not\supset \mathfrak{p}$ since otherwise $a \cap c \supset \mathfrak{p}$ which is impossible. Hence $b \cup \mathfrak{p} \supset b$ by the Birkhoff condition. But then $b \cup \mathfrak{p} \supset a \cap (b \cup \mathfrak{p}) \supset b$ and if $b \cup \mathfrak{p} = a \cap (b \cup \mathfrak{p})$ we have $a \supset \mathfrak{p}$ which is impossible. Hence $b = a \cap (b \cup \mathfrak{p})$. Thus by Theorem 2.2 an element $p \in \mathfrak{p}$ exists such that $b = a \cap (b \cup p)$. Let $c_1 = c \cap p$. Then $c \supset c_1 \supset \mathfrak{p} \supset a \cap c$ and hence $c \supset c_1 \supset a \cap c$, $c_1 \neq a \cap c$. Also $b = a \cap (b \cup p) \supset a \cap (b \cup c_1) \supset b$. Thus $b = a \cap (b \cup c_1)$ and c_1 satisfies the requirements of E_5 .

THEOREM 6.2. *Let \mathfrak{S} satisfy E_5 and have the property that each element is covered by only a finite number of covering ideals. Then \mathfrak{S} is a Birkhoff lattice.*

Proof. Let $a \supset a$, $\mathfrak{p} \supset a$ and $a \not\supset \mathfrak{p}$. Let $\mathfrak{p}, \mathfrak{p}_1, \dots, \mathfrak{p}_n$ be the finite number of ideals covering a . Now if $\mathfrak{p} \cup a \not\supset a$, we have $\mathfrak{p} \cup a \supset c \supset a$, $\mathfrak{p} \cup a \neq c \neq a$. Since $\mathfrak{p} \supset \mathfrak{p} \cap c \supset a$ and $c \not\supset \mathfrak{p}$ we have $\mathfrak{p} \cap c = a$ and hence by Theorem 2.2 elements $p' \in \mathfrak{p}$ and $c \in c$ exist such that $p' \cap c = a$. Since $c \neq a$, there exists an element $b' \in a$ such that $b' \not\supset c$. Let $b = c \cap b'$. Then $b \not\supset c$ and $c \supset b \supset c \cap p'$. Now since $\mathfrak{p} \not\supset \mathfrak{p}_i$ ($i=1, \dots, n$) elements p'_i exist such that $p'_i \in \mathfrak{p}$ and $p'_i \not\supset \mathfrak{p}_i$ ($i=1, \dots, n$). Set $p = p' \cap p'_1 \cap \dots \cap p'_n$. Then $p \in \mathfrak{p}$, $p' \supset p$ and $p \not\supset \mathfrak{p}_i$ ($i=1, \dots, n$). Clearly $a = c \cap p' \supset c \cap p \supset a$ implies $c \cap p = a$. Hence $c \supset b \supset c \cap p$ and $p \neq c \cap p$. Thus by E_5 an element p_1 exists such that $p_1 \neq a$, $p \supset p_1 \supset a$ and

$b = c \cap (b \cup p_1)$. Now $p_1 \supset p' > a$ and $p' \neq p_i$ ($i = 1, \dots, n$) since otherwise $p \supset p_1 \supset p_i$ contrary to the definition of p . Hence $p' = p$. But then $b = c \cap (b \cup p_1) \supset c \cap (a \cup p) \supset c$ which contradicts $b \not\supset c$. Hence $p \cup a > a$ and \mathcal{S} is a Birkhoff lattice.

Now by Lemma 4.6, if \mathcal{S} is a Birkhoff lattice in which the ascending chain condition holds and every three ideals covering a principal ideal generate a Boolean algebra, then each a is covered by only a finite number of ideals. However, Theorem 6.2 does not enable us to replace the Birkhoff condition in the lemma by E_s since the proof of the finiteness required the Birkhoff condition. To carry out this replacement we first replace the condition that every three ideals covering a principal ideal generate a Boolean algebra by an equivalent condition.

THEOREM 6.3. *Let \mathcal{S} be a Birkhoff lattice satisfying the ascending chain condition. Then every three ideals covering a principal ideal generate a Boolean algebra if and only if $a \cup b \supset q > a \cap b$ implies $a \supset q$ or $b \supset q$.*

Proof. Let every three ideals covering a principal ideal generate a Boolean algebra and suppose that $a \cup b \supset q > a \cap b$ but $a \not\supset q$, $b \not\supset q$. Then $a, b \neq a \cap b$. For if $a = a \cap b$, then $b = a \cup b \supset q$ contrary to assumption. Now with b fixed let a be maximal such that $a \cup b \supset q > a \cap b$ and $a \not\supset q$, $b \not\supset q$ for some q . Let $b \supset p > a \cap b$. Suppose $a = p \cup a \not\supset q$. Then $a \cup b = a \cup p \cup b = a \cup b \supset q$ and $a \cap b \supset a \cap b$. Now $a \neq a$ since otherwise $a \supset p$ and $a \cap b \supset p > a \cap b$ which is impossible. Let a_1' be an element of a such that $a_1' \not\supset q$. Now $a \cup b \supset b \supset p$ and hence $a \cup b \supset p \cup a = a$. Thus $a_1 = (a \cup b) \cap a_1' \supset a$ and $a \cup b \supset a_1$. But $a \cup b \supset a_1 \cup b \supset a \cup b$. Hence $a_1 \cup b = a \cup b \supset q$. Also $a_1 \not\supset q$ since otherwise $a_1' \supset q$. Now $q \supset (a_1 \cap b) \cap q \supset a \cap b$ and $q \neq (a_1 \cap b) \cap q$ since otherwise $b \supset q$. Hence $q > a \cap b = (a_1 \cap b) \cap q$. By the Birkhoff condition we have $q_1 = q \cup (a_1 \cap b) > a_1 \cap b$. Since $a_1 \cup b \supset q$ we have $a_1 \cup b \supset q_1 > a_1 \cap b$. Clearly $a_1 \not\supset q_1$, $b \not\supset q_1$, and $a_1 \neq a$. This contradicts the maximal property of a . Hence we have $p \cup a \supset q$.

Now let $a \supset p_1 > a \cap b$. $p_1 \neq q$ since otherwise $a \supset q$ and $p_1 \neq p$ since otherwise $a \cap b \supset p > a \cap b$. Hence $p_1 \cup p \not\supset q$ since every three ideals covering $a \cap b$ generate a Boolean algebra. Thus $p_1 \cup p \not\supset q$ for some $p_1 \in p_1$. Set $x = p_1 \cap a$. Then $a \supset x \supset a \cap b$ and $x \neq a \cap b$, $x \cup p \not\supset q$. Let a_2 be a maximal such x . Then $a_2 \neq a$ since $a \cup p \supset q$. Hence $a \supset p_2 > a_2$ for some ideal p_2 . Then if $p_2 \cup p \not\supset q$ we have $p_2 \cup p \not\supset q$ for some $p_2 \in p_2$. Let $a_2' = a \cap p_2$. Then $a \supset a_2' \supset a \cap b$, $a_2' \neq a \cap b$ and $a_2' \cup p \not\supset q$ contrary to the maximal property of a_2 . Hence $p_2 \cup p \supset q$. Let $q_2 = a_2 \cup q$ and $p_3 = a_2 \cup p$. We have $a_2 \not\supset p$ since $a \not\supset p$ and hence $p_2, q_2, p_3 > a_2$ by the Birkhoff condition. Clearly $p_2 \cup p_3 = p_2 \cup a_2 \cup p \supset a_2 \cup q = q_2$. Now $p_2 \neq q_2$ since otherwise $a \supset q$. Also $q_2 \neq p_3$ since otherwise $a_2 \cup p \supset q$ contrary to the definition of a_2 . Finally $p_2 \neq p_3$ since $a \not\supset p$. Thus p_2, q_2, p_3 do not generate a Boolean algebra, which contradicts our hypothesis. Hence either $a \supset q$ or $b \supset q$.

On the other hand if three ideals a, b, c covering d do not generate a Boolean algebra, then $a \cup b \supset c$ say. Since $d = a \cap b$, elements $a \in a$ and $b \in b$

exist such that $d = a \wedge b$. Hence $a \vee b \supset a \vee b \supset c > a \wedge b = d$. If $a \supset c$, then $a \supset a \vee c \supset b$ and $d = a \wedge b \supset b$ which is impossible. Hence $a \not\supset c$ and $b \not\supset c$. Thus $a \vee b \supset c > a \wedge b$ but $a \not\supset c$ and $b \not\supset c$. This completes the proof of the theorem.

We show now that if the ascending chain condition holds and the condition of Theorem 6.3 is satisfied, then E_5 is equivalent to the Birkhoff condition. A preliminary lemma is required.

LEMMA 6.1. \mathfrak{S} is a Birkhoff lattice if and only if $b \supset a$, $b \not\supset p > a$ implies $p \vee b > b$.

The necessity of the condition is obvious. To prove the sufficiency let $b > a$, $c \supset a$ and $c \not\supset b$. If $b \vee c \not\supset c$ we have $b \vee c \supset d \supset c$ where $b \vee c \neq d \neq c$. Since $b \not\supset b$ an element $d \in b$ exists such that $d \not\supset b$. Since $d \supset c$ and $c \not\supset b$ there exists an element $c \in c$ such that $d \supset c$ and $c \not\supset b$. Now $c \supset a$, $c \not\supset b > a$. Hence $c \vee b > c$. But $c \vee b \supset d \wedge (c \vee b) \supset c$ and $c \vee b \neq d \wedge (c \vee b)$ since $d \not\supset b$. Thus $c = d \wedge (c \vee b) \supset b \wedge (c \vee b) = b$ which contradicts $c \not\supset b$. Hence $b \vee c > c$ and \mathfrak{S} is a Birkhoff lattice.

THEOREM 6.4. Let \mathfrak{S} satisfy E_5 , the ascending chain condition, and let $a \vee b \supset q > a \wedge b$ imply $a \supset q$ or $b \supset q$. Then \mathfrak{S} is a Birkhoff lattice.

Proof. Let X be the set of all elements x such that $y \supset x$, $p > x$, $y \not\supset p$, and $p \vee y \not\supset y$ for some y and p . If \mathfrak{S} is not a Birkhoff lattice, then X is non-empty by Lemma 6.1. Let a be a maximal element of X . Then b and p exist such that $b \supset a$, $b \not\supset p > a$ and $b \vee p \supset c > b$, $b \vee p \neq c$. Now $p \supset p \wedge c \supset a$ and $p \neq p \wedge c$ since otherwise $b \vee p = c$. Hence $p \wedge c = a$. Let $p \in p$, $c \in c$ such that $p \wedge c = a$. Then $p \wedge c = p \wedge b = a$ and $c \neq b$. Hence by E_5 an element p_1 exists such that $p \supset p_1 \supset a$, $p_1 \neq a$ and $b = c \wedge (b \vee p_1)$. Now $b \vee p_1 \not\supset c$ since otherwise $b = c \wedge (b \vee p_1) \supset c$ which contradicts $c > b$. Now suppose that we have found p_1, \dots, p_k such that $b \vee p_1 \vee p_2 \vee \dots \vee p_k \not\supset c$ and $(b \vee p_1 \vee \dots \vee p_i) \wedge p_{i+1} = a$ ($i = 1, \dots, k-1$). Since $b \vee p_1 \vee p_2 \vee \dots \vee p_k \not\supset c$ we have $b \vee p_1 \vee \dots \vee p_k \not\supset p$ and hence $(b \vee p_1 \vee \dots \vee p_k) \wedge p = a$. Thus $p'_{k+1} \in p$ exists such that $(b \vee p_1 \vee \dots \vee p_k) \wedge p'_{k+1} = a$. Let $p'_{k+1} = p'_{k+1} \wedge p$. Then $(b \vee p_1 \vee \dots \vee p_k) \wedge p'_{k+1} = a$ and $c \wedge p_{k+1} = b \wedge p'_{k+1} = a$, $p'_{k+1} \neq a$. Hence by E_5 an element p_{k+1} exists such that $p'_{k+1} \supset p_{k+1} \supset a$, $p_{k+1} \neq a$ and $b = c \wedge (b \vee p_{k+1})$. Then $b \vee p_{k+1} \not\supset c$ since otherwise $b = c \wedge (b \vee p_{k+1}) \supset c$. Now $c \supset c \wedge ((b \vee p_1 \vee \dots \vee p_k) \wedge (b \vee p_{k+1})) \supset b$ and $c \neq c \wedge ((b \vee p_1 \vee \dots \vee p_k) \wedge (b \vee p_{k+1}))$ since otherwise $b \vee p_{k+1} \supset (b \vee p_1 \vee \dots \vee p_k) \wedge (b \vee p_{k+1}) \supset c$. Hence since $c > b$ we have $c > c \wedge ((b \vee p_1 \vee \dots \vee p_k) \wedge (b \vee p_{k+1})) = b$. But now since b is a proper divisor of a , by the maximal property of a we must have $c \vee ((b \vee p_1 \vee \dots \vee p_k) \wedge (b \vee p_{k+1})) > (b \vee p_1 \vee \dots \vee p_k) \wedge (b \vee p_{k+1})$. Now suppose that $b \vee p_1 \vee \dots \vee p_{k+1} \supset c$. Then $(b \vee p_1 \vee \dots \vee p_k) \vee (b \vee p_{k+1}) \supset c \vee ((b \vee p_1 \vee \dots \vee p_k) \wedge (b \vee p_{k+1})) > (b \vee p_1 \vee \dots \vee p_k) \wedge (b \vee p_{k+1})$. Hence by hypothesis either $b \vee p_1 \vee \dots \vee p_k \supset c$ or $b \vee p_{k+1} \supset c$ both of which are impossible. Thus $b \vee p_1 \vee \dots \vee p_{k+1} \not\supset c$. By induction we get an infinite chain $b \subset b \vee p_1 \subset b \vee p_1 \vee p_2 \subset \dots \subset b \vee p_1 \vee \dots$

$\cup p_i \subset \dots$ and $b \cup p_1 \cup \dots \cup p_i \neq b \cup p_1 \cup \dots \cup p_{i+1}$ since $(b \cup p_1 \cup \dots \cup p_i) \cap p_{i+1} = a$. This chain contradicts the ascending chain condition and hence \mathfrak{S} is a Birkhoff lattice.

The condition $a \cup b \supset q > a \cap b$ implies $a \supset q$ or $b \supset q$, may be given a purely combinatorial statement as follows:

THEOREM 6.5. *$a \cup b \supset q > a \cap b$ implies $a \supset q$ or $b \supset q$ if and only if*
 (A) *$a \cup b \supset x \supset a \cap b$, $a \cap x = b \cap x = a \cap b$ implies $x = a \cap b$.*

Proof. Let $a \cup b \supset x \supset a \cap b$, $a \cap x = b \cap x = a \cap b$. If $x \neq a \cap b$, let $x \supset q > a \cap b$. Then $a \cup b \supset q > a \cap b$ and hence $a \supset q$ say. But then $a \cap b = a \cap x \supset q > a \cap b$ which is impossible. Hence $x = a \cap b$.

On the other hand, let $a \cup b \supset q > a \cap b$. If $a \not\supset q$ and $b \not\supset q$ we have $a \cap q = b \cap q = a \cap b$. Hence for some $x \in q$ we have $a \cap x = b \cap x = a \cap b$ and $a \cup b \supset x$ by Theorem 2.2. But then $a \cup b \supset x \supset a \cap b$, $a \cap x = b \cap x = a \cap b$ and $x \neq a \cap b$ which contradicts condition A.

Lemma 4.6 with Theorems 6.1–6.5 give

THEOREM 6.6. *Let \mathfrak{S} satisfy the ascending chain condition. Then every element of \mathfrak{S} is uniquely expressible as a reduced cross-cut of irreducibles if and only if conditions E_5 and A are satisfied.*

Theorem 6.6 has the following interesting corollary:

COROLLARY 6.1. *Let \mathfrak{S} satisfy the ascending chain condition and let each element of \mathfrak{S} have a unique reduced decomposition into irreducibles. Then a sublattice \mathfrak{S}' of \mathfrak{S} has unique irreducible decompositions if and only if E_5 holds in \mathfrak{S}' .*

Axiom A is clearly a slightly stronger form of the requirement that every modular sublattice be distributive. In D1 it was shown that under the assumption of both the ascending and descending chain conditions, this weaker condition and B1 were necessary and sufficient for unique decomposition into irreducibles. But A *cannot* be replaced by the requirement that every modular sublattice be distributive in Theorem 6.6 as the example of Figure 1 shows.

The non-principal ideals of \mathfrak{S} are the ideals \mathfrak{a} , generated by a_1, a_2, a_3, \dots ; \mathfrak{b} , generated by b_1, b_2, b_3, \dots ; and \mathfrak{c} , generated by c_1, c_2, c_3, \dots . Clearly $\mathfrak{a} > \mathfrak{z}$, $\mathfrak{b} > \mathfrak{b}$, and $\mathfrak{c} > \mathfrak{z}$. Now $b \cup a_{2n} = d_n > a_{2n}$ and $b \cup a_{2n+1} = b_{2n+1} > a_{2n+1}$. Hence $b \cup a_i > a_i$. Similarly $b \cup c_i > c_i$. Now let x be any element of \mathfrak{S} not equal to b or \mathfrak{z} . Then b_1/x is an archimedean lattice and B1 is readily verified in b_1/x since each element has at most two covering elements. Thus we have only to verify the Birkhoff condition for non-principal ideals. Clearly $\mathfrak{b} > \mathfrak{a}$, $\mathfrak{b} > \mathfrak{c}$. Hence $\mathfrak{a} \cup \mathfrak{b} > \mathfrak{b}$, $\mathfrak{a} \cup \mathfrak{c} > \mathfrak{a}$, $\mathfrak{c} \cup \mathfrak{b} > \mathfrak{b}$, $\mathfrak{c} \cup \mathfrak{a} > \mathfrak{a}$. $\mathfrak{a} \cup c_{2n} = e_n > c_{2n}$ and $\mathfrak{a} \cup c_{2n+1} = b_{2n+2} > c_{2n+1}$. Hence $\mathfrak{a} \cup c_i > c_i$. Similarly $\mathfrak{c} \cup a_i > a_i$. Thus every element of \mathfrak{S} satisfies the Birkhoff condition in the lattice of ideals and hence \mathfrak{S} is a Birkhoff lattice. By Theorem 6.1, E_5 holds in \mathfrak{S} . Now if \mathfrak{S} contains a modular, non-distribu-

tive sublattice it also contains one of the form $\{u, v, w, x, y\}$ where $v \cup w = w \cup x = v \cup x = u$ and $v \cap w = w \cap x = v \cap x = y$. Since every element not equal to b or z is covered by at most two elements we must have $y = z$. But then $v = a_i$, $w = b$, $x = c_j$ and $v \cup w = a_i \cup b \neq b \cup c_j = w \cup x$ which contradicts $v \cup w$




FIG. 1

$= x \cup w$. Hence every modular sublattice of \mathfrak{S} is distributive. However \mathfrak{S} does not have unique irreducible decompositions since $z = a_1 \cap b = b \cap c_1 = a_1 \cap c_1$ and a_1, b, c are irreducibles of \mathfrak{S} . Axiom A does not hold since $a_i \cup c_j \supset b \supset a_i \cap c_j$ and $a_i \cap b = c_j \cap b = a_i \cap c_j$ but $b \neq a_i \cap c_j = z$.

Since a, b, c generate a modular lattice, \mathfrak{L}_x is modular for every x . Hence by Theorem 5.1, the number of components in the reduced decompositions of each element must be unique. This can be readily verified.

According to Theorem 6.2, if every element of a lattice \mathfrak{S} is covered by only a finite number of ideals, then E_5 implies that \mathfrak{S} is a Birkhoff lattice. We prove now an even stronger theorem, namely, under this restriction E_5 implies that B1 holds in the lattice of ideals. We begin with necessary lemmas.

LEMMA 6.2. *B1 holds in the lattice of ideals of \mathfrak{S} if and only if $a > x \cap a$ implies $x \cup a > x$ for every $a \in \mathfrak{I}$ and $x \in \mathfrak{S}$.*

For if B1 holds in \mathfrak{I} , then clearly $a > x \cap a$ implies $x \cup a > x$. Now let $a > x \cap a$ imply $a \cup x > x$ for each a and x . Suppose that B1 does not hold in \mathfrak{I} . Then ideals a and b exist such that $a > a \cap b$ but $a \cup b \not\supset c \supset b$, $a \cup b \neq c \neq b$. Let $x_1 \supset b$, $x_1 \not\supset a$. Such an x_1 always exists since $b \not\supset a$. Also since $b \not\supset c$, an element x_2 exists such that $x_2 \supset b$, $x_2 \not\supset c$. Finally since $b \cup c \not\supset a$ there is an element x_3 such that $x_3 \supset b$, $x_3 \cup c \not\supset a$. Let $x = x_1 \cap x_2 \cap x_3$. Then $x \supset b$, $x \not\supset a$, and $x \cup c \not\supset a$. Now $a \supset a \cap x \supset a \cap b$ and $a \neq a \cap x$. Hence $a \cap x = a \cap b$ and thus $a > a \cap x$. By

hypothesis then $x \cup a > x$. Now $x \cup a \supset c \cup x \supset x$ and $c \cup x \neq x$. Hence $x \cup a = c \cup x$ which implies $x \cup c \supset a$ contrary to the definition of x . Hence B1 holds in \mathfrak{L} .

LEMMA 6.3. *Let \mathfrak{S} be a Birkhoff lattice. Then if $a > x \cap a$ and $x \cup a \not> x$, each $x \cap a$, $a \in \mathfrak{a}$ is covered by an infinite number of ideals.*

For let $a > x \cap a$ and $x \cup a \not> x$. Then clearly $a \cap x \neq a$ for every $a \in \mathfrak{a}$ since otherwise $x \supset a \supset a$ and $a \not> x \cap a$. Hence $a \supset p_a > a \cap x$ for some ideal p_a by Theorem 2.1. Let S_a denote the set of all ideals p_a . Now $x \not\supset p_a$, since otherwise $a \cap x \supset p_a > a \cap x$ which is impossible. Thus $p_a \supset x \cap p_a \supset a \cap x$ and $p_a \neq x \cap p_a$. Hence $x \cap p_a = a \cap x$ and $p_a > x \cap p_a$ where $x \cap p_a$ is a principal ideal of \mathfrak{L} . Since \mathfrak{S} is a Birkhoff lattice we have $x \cup p_a > x$ for every p_a .

Now in S_a we set $p_a \sim p'_a$ if and only if $x \cup p_a = x \cup p'_a$. Then \sim is an equivalence relation which separates S_a into mutually exclusive sets of ideals. Let B_a denote an arbitrary equivalence class and let $b_a = \Sigma(B_a)$. If $p_a \in B_a$, then $x \cup p_a = x \cup p'_a$ for every other ideal p'_a of B_a and hence $x \cup p_a = x \cup b_a$. Thus $x \cup b_a > x$. Let T_a denote the set of ideals b_a . Now if $a \supset a_1 \supset a$ and $b_{a_1} \in T_{a_1}$, let $b_{a_1} \supset p_{a_1} > x \cap a_1$. Then $p_{a_1} \supset (x \cap a) \cap p_{a_1} \supset x \cap a_1$ and $p_{a_1} \neq (x \cap a) \cap p_{a_1}$ since otherwise $x \supset x \cap a \supset p_{a_1}$ which contradicts $x \not\supset p_{a_1}$. Hence $p_{a_1} > (x \cap a) \cap p_{a_1} = x \cap a_1$ and $(x \cap a) \cup p_{a_1} > x \cap a$ by the Birkhoff condition. Also $a \supset a \cup a_1 \supset (x \cap a) \cup p_{a_1} > x \cap a$ and hence $p_a = (x \cap a) \cup p_{a_1}$ belongs to S_a . Now $x \cup p_a = x \cup (x \cap a) \cup p_{a_1} = x \cup p_{a_1}$. Let $b_{a_1} \supset p'_{a_1} > x \cap a_1$. Then $x \cup p'_a = x \cup p'_{a_1}$ where $p'_a = (x \cap a) \cup p'_{a_1}$. Thus $x \cup p_a = x \cup p_{a_1} = x \cup p'_{a_1} = x \cup p'_a$ and $p_a \sim p'_a$ in S_a . Hence $b_{a_1} \subset b_a$ where b_a is an ideal of T_a . Now suppose that T_{a_1} contains a second ideal b'_{a_1} , which is divisible by b_a . Let $b'_{a_1} \supset p''_{a_1} > x \cap a_1$. Then $x \cup p_{a_1} = x \cup p_a = x \cup b_a \supset x \cup b'_{a_1} \supset x \cup p''_{a_1} \supset x$. Since $x \cup p''_{a_1} \neq x$, we have $x \cup p_{a_1} = x \cup p''_{a_1}$ and $p_{a_1} \sim p''_{a_1}$ in S_{a_1} contrary to assumption. Hence b_{a_1} is the only ideal of T_{a_1} divisible by b_a . Next suppose that T_a contains another ideal b'_a such that $b'_a \supset b_{a_1}$. Then $x \cup b'_a \supset x \cup b_{a_1} = x \cup b_a$ and hence $b_a = b'_a$. We thus conclude that *each ideal b_{a_1} of T_{a_1} is divisible by exactly one ideal b_a of T_a and b_{a_1} is the only ideal of T_{a_1} which is divisible by b_a .*

Let n_a denote the cardinal number of the set T_a . If $x \cap a$ is covered by only a finite number of ideals for some a , then n_a is finite for some a and hence has a minimal value for some a_0 . If $a_0 \supset a \supset a$, then $n_a \leq n_{a_0}$ since distinct ideals of T_a are divisible by distinct ideals of T_{a_0} . But since n_{a_0} is minimal we have $n_a = n_{a_0}$. Hence each ideal of T_{a_0} divides exactly one ideal of T_a . Let b_0 be an ideal of T_{a_0} and let b_a denote the ideal of T_a divisible by b_0 . In general, for any $a \in \mathfrak{a}$, let b_a be the ideal of T_a divisible by $x \cup b_0$. Such an ideal b_a always exists since $b_0 \supset b'_a$ where $b'_a \in T_{a \cap a_0}$ and hence $b_a \in T_a$ exists such that $b_a \supset b'_a$. Clearly $x \cup b_0 = x \cup b'_a = x \cup b_a \supset b_a$ and b_a is unique as shown above. If $a \supset a'$, we have $b_a \supset b_{a'}$ since $x \cup b_a = x \cup b_{a'}$. Hence $b_{a_1} \cap b_{a_2} \cap \dots \cap b_{a_n} \supset b_{a_1 \cap a_2 \cap \dots \cap a_n}$.

Now let $a_1 = \prod_{a \in \mathfrak{a}} (b_a)$. Since $a \supset b_a \supset a_1$, we have $a \supset a_1 \supset x \cap a$. If $a_1 = x \cap a_1$ then $x \cap a$ divides the cross-cut of a finite number of the ideals b_a and hence

$x \cap a \supset b_{a'}$ for some a' . But then $(x \cap a) \cap a' \supset b_{a'} \cap b_{a'} \supset b_{a'} \supset b_{a \cap a'} \supset x \cap (a \cap a')$ and $b_{a \cap a'} = x \cap (a \cap a')$ contrary to the definition of $b_{a \cap a'}$. Thus $a_1 \neq x \cap a$ and hence $a = a_1$ since $a > x \cap a$. But then $x \cup b_0 \supset x \cup a_1 \supset x \cup a \supset x$. Since $x \cup b_0 > x$ and $x \not\supset a$ we have $x \cup a = x \cup b_0 > x$ which contradicts $x \cup a \not\supset x$. Hence each




FIG. 2

$x \cap a$ is covered by an infinite number of ideals. The proof is thus complete.

Lemmas 6.2 and 6.3 and Theorem 6.2 give immediately

THEOREM 6.7. *Let each element of \mathfrak{S} be covered by at most a finite number of ideals. Then the following conditions are equivalent:*

- (1) E_8 holds in \mathfrak{S} .
- (2) \mathfrak{S} is a Birkhoff lattice.
- (3) B1 holds in the lattice of ideals.

If \mathfrak{S} is a Birkhoff lattice in which each element is *not* covered by at most a finite number of ideals, then even though the ascending chain condition holds in \mathfrak{S} B1 need not be satisfied in the lattice of ideals. For example, consider the lattice diagrammed in Figure 2.

All of the elements distinct from z form an ideal a which is generated by $a_1, a_2, a_3, \dots, b_1, b_2, b_3, \dots$ clearly form an ideal b which divides a . Now let $b \supset c \supset a$, $b \neq c$. Let $y \in c$, $y \notin b$. Then there exists a b_i such that $b_i > x \supset y$, $x \notin b$. But by the method of construction there exists an integer j such that

$x \cap b_k = a_k$ all $k \geq j$. Hence $a \supset x \cap b \supset y \cap b \supset c \cap b = c$. Thus $c = a$ and $b > a$. Clearly $b \cap a_1 = a$ and $b \cup a_1 = b_1$. But then $b > a_1 \cap b$ and $a_1 \cup b \not> a_1$. Hence B1 does *not* hold in \mathfrak{L} . On the other hand it is readily verified that \mathfrak{S} is a Birkhoff lattice since a is the only non-principal ideal which covers a principal ideal and every element distinct from z divides a .

The number of ideals covering an element of a lattice is closely related to the number of decompositions of the element into irreducibles. We prove

THEOREM 6.8. *Let \mathfrak{S} be a Birkhoff lattice in which each element can be represented as a cross-cut of irreducibles. Then if an element a has a finite number of decompositions into irreducibles, \mathfrak{L}_a is finite.*

Proof. Since a has only a finite number of decompositions into irreducibles, the number of components in the irreducible decompositions of a is bounded. Hence \mathfrak{L}_a is archimedean by Theorem 3.1. Let p_1, \dots, p_k be a maximal independent set of point ideals of \mathfrak{L}_a . Let $\mathfrak{s}_i = p_1 \cup \dots \cup p_{i-1} \cup p_{i+1} \cup \dots \cup p_k$ and suppose that \mathfrak{L}_a has an infinite sequence $\mathfrak{s}_1, \dots, \mathfrak{s}_k, \mathfrak{s}_{k+1}, \dots$ of simple ideals. Now if for each i there are only a finite number of simple ideals of the sequence which do not divide p_i , we have $\mathfrak{s}_n \supset p_i$ for all $n \geq l_i$ for some l_i . Let $n \geq \max(l_1, \dots, l_k)$. Then $\mathfrak{s}_n \supset p_i, i = 1, \dots, k$, and $\mathfrak{s}_n \supset u_a$, which is impossible. Hence for some p_i , say p_k , there are an infinity of ideals in the sequence $\mathfrak{s}_1, \mathfrak{s}_2, \dots$ which do not divide p_k . We may assume that $\mathfrak{s}_k, \mathfrak{s}_{k+1}, \dots$ do not divide p_k . Let $q_i \supset \mathfrak{s}_i, q_i \not\supset u_a$. Then $q_i \neq q_j, i \neq j$, since otherwise $q_i = q_j \cup q_j \supset \mathfrak{s}_i \cup \mathfrak{s}_j \supset u_a$. Now $a = p_k \cap \mathfrak{s}_{k+l} = \mathfrak{s}_1 \cap \dots \cap \mathfrak{s}_{k-1} \cap \mathfrak{s}_{k+l}$ ($l = 0, 1, 2, \dots$) implies $a = q_1 \cap q_2 \cap \dots \cap q_{k-1} \cap q_{k+l}$ ($l = 0, 1, 2, \dots$) by Theorem 3.3. If this representation is reduced for each l, q_{k+l} always remains since otherwise $a = q_1 \cap \dots \cap q_{k-1} \supset p_k$. Hence a has an infinite number of irreducible decompositions, which contradicts our hypothesis. Thus \mathfrak{L}_a has only a finite number of simple ideals. But by Theorem 3.2 every ideal of \mathfrak{L}_a can be expressed as a cross-cut of simple ideals. Hence \mathfrak{L}_a is finite.

Theorems 6.7 and 6.8 give

THEOREM 6.9. *Let \mathfrak{S} be a lattice in which every element has at least one and at most a finite number of decompositions into irreducibles. Then \mathfrak{S} is a Birkhoff lattice if and only if B1 is satisfied in the lattice of ideals⁽¹¹⁾.*

7. Example of a Birkhoff lattice. In §3 we have shown that the existence of a decomposition into irreducibles for an element a of a modular lattice implies that \mathfrak{L}_a is archimedean. Hence if the ascending chain condition holds,

⁽¹¹⁾ Various considerations suggest that the finiteness of the number of irreducible decompositions of an element always implies the finiteness of the number of ideals covering the element, in which case E_s and the finiteness of the number of decompositions would imply that \mathfrak{S} is a Birkhoff lattice. However, I have been unable to prove this. I have also been unable to prove that E_s is equivalent to the Birkhoff condition under the assumption of the ascending chain condition although this seems quite likely.

\mathfrak{L}_a is archimedean for each a . Also if the ascending chain condition holds in a Birkhoff lattice and the number of components is bounded for an element a , then \mathfrak{L}_a is archimedean. We shall construct in this section a Birkhoff lattice satisfying the ascending chain condition but containing an element a such that \mathfrak{L}_a is not archimedean. By the above remark the number of components in the irreducible decompositions of a must be unbounded.

Latin capitals A, B, C, \dots will denote finite subsets of the set of positive integers $1, 2, 3, \dots$. If A is such a set, let $n(A)$ denote the number of elements in A . Small Latin letters a, b, c, \dots will denote positive integers. $A \cup B$ and $A \cap B$ will denote set-theoretic union and cross-cut respectively and $a \cup b$, $a \cap b$ are respectively the maximum and minimum of a and b . Let \mathfrak{S} be the set of all ordered couples $\alpha = \{A, a\}$ where $n(A) < a$ together with the elements u and z . In \mathfrak{S} we define

$$\begin{aligned}\alpha \cup \beta &= \{A \cup B, a \cup b\} \text{ if } n(A \cup B) < a \cup b, \\ &= u \text{ if } n(A \cup B) \geq a \cup b, \\ \alpha \cap \beta &= \{A \cap B, a \cap b\} \text{ if } A \cap B \text{ is not null,} \\ &= z \text{ if } A \cap B \text{ is null,} \\ \alpha \cup z &= \alpha \cap u = \alpha, \quad \alpha \cup u = u, \quad \alpha \cap z = z.\end{aligned}$$

If $A \cap B$ exists, then $n(A \cap B) \leq n(A) < a \leq a \cap b$. Hence $\alpha \cap \beta$ is in \mathfrak{S} if α and β are in \mathfrak{S} . Now it can be readily verified that the union and cross-cut so defined in \mathfrak{S} are idempotent, commutative and associative. Consider $\alpha \cap (\alpha \cup \beta)$. If $\alpha \cup \beta = u$, then $\alpha \cap (\alpha \cup \beta) = \alpha$. If $\alpha \cup \beta \neq u$, then $\alpha \cap (\alpha \cup \beta) = \{A \cap (A \cup B), a \cap (a \cup b)\} = \{A, a\} = \alpha$. Hence $\alpha \cap (\alpha \cup \beta) = \alpha$ in all cases. Similarly $\alpha \cup (\alpha \cap \beta) = \alpha$. Hence \mathfrak{S} is a lattice under the union and cross-cut operations defined above. Clearly $\alpha \supset \beta$ if and only if $A \supset B$ and $a \leq b$.

\mathfrak{S} satisfies the ascending chain condition. For let $\alpha_1 \subset \alpha_2 \subset \alpha_3 \subset \dots$ be an infinite ascending chain. We may assume that $\alpha_1 \neq z$ so that $\alpha_1 = \{A_1, a_1\}$. But then $A_1 \subset A_2 \subset A_3 \subset \dots$ and $a_1 \geq a_2 \geq a_3 \geq \dots$. Now $n(A_i) < a_i \leq a_1$. Hence the chain $A_1 \subset A_2 \subset \dots$ has only a finite number of distinct sets. Clearly the chain $a_1 \geq a_2 \geq \dots$ has only a finite number of distinct members. Thus the chain $\alpha_1 \subset \alpha_2 \subset \dots$ has only a finite number of distinct members.

Now let \mathfrak{a} be an ideal of \mathfrak{S} with elements $\beta, \gamma, \delta, \dots$. Then the set of integers is either bounded or unbounded. If bounded, let a be the largest of them. Now suppose that $B \cap C \cap D \cap \dots$ is null. Then there exist a finite number of them B, C, \dots, L whose cross-cut is null. But then \mathfrak{a} contains z since \mathfrak{a} is closed with respect to finite cross-cut. Hence $\mathfrak{a} = \mathfrak{S}$ in this case. If $B \cap C \cap D \cap \dots$ is not null, let $A = B \cap C \cap D \cap \dots$. Then $A = B \cap C \cap \dots \cap L$ for a finite number of the sets and hence $\alpha' = \{A, a'\}$ and $\alpha'' = \{A', a\}$ are in \mathfrak{a} . But then \mathfrak{a} contains $\alpha = \alpha' \cap \alpha'' = \{A, a\}$ and $\beta \supset \alpha$ for every $\beta \in \mathfrak{a}$. Hence if the integers of the ideal are bounded, \mathfrak{a} is a principal ideal. If the integers of the ideal are unbounded, then as before either $\mathfrak{a} = \mathfrak{S}$

or there exists a set A such that $\{A, a\}$ is in \mathfrak{a} for each positive integer a and $\beta \supset \{A, a\}$ for some a if $\beta \in \mathfrak{a}$. Hence the ideals of \mathfrak{S} have the form $\mathfrak{a} = \{A, \infty\}$ if \mathfrak{a} is not principal. ∞ denotes the ideal of all positive integers. Clearly if a and b are positive integers or ∞

$$\begin{aligned} \mathfrak{a} \cap \mathfrak{b} &= \{A \cap B, a \cap b\} && \text{if } A \cap B \text{ is not null,} \\ &= z && \text{if } A \cap B \text{ is null,} \\ \mathfrak{a} \cup \mathfrak{b} &= \{A \cup B, a \cup b\} && \text{if } n(A \cup B) < a \cup b, \\ &= u && \text{if } n(A \cup B) \geq a \cup b. \end{aligned}$$

THEOREM 7.1. \mathfrak{S} is a Birkhoff lattice.

Proof. We shall show that B1 holds in the lattice of ideals. Let $\mathfrak{a} > \mathfrak{a} \cap \mathfrak{b}$. If $\mathfrak{a} \cap \mathfrak{b} = z$, then $A \cap B$ is null. Since $\mathfrak{a} > z$ we have $\mathfrak{a} = \{(a), \infty\}$. But then $\mathfrak{a} \cup \mathfrak{b} = \{(a) \cup B, \infty \cup b\} = \{(a) \cup B, b\}$ and $(a) \cup B > B$. Thus $\mathfrak{a} \cup \mathfrak{b} > \mathfrak{b}$. If $\mathfrak{a} \cap \mathfrak{b} \neq z$, then $\{A, a\} > \{A \cap B, a \cap b\}$ and hence either $A > A \cap B$, $a = a \cap b$ or $A = A \cap B$, $a = (a \cap b) - 1$. In the first case $A > A \cap B \rightarrow A \cup B > B$ and $a = a \cap b$ implies $b = a \cup b$. Hence $\mathfrak{a} \cup \mathfrak{b} = \{A \cup B, a \cup b\} = \{A \cup B, b\} > \{B, b\} = \mathfrak{b}$. If $n(A \cup B) \geq b$, then $n(B) = b - 1$ and hence $u > B$. In the second case $A = A \cap B \rightarrow A \cup B = B$ and $a > a \cap b \rightarrow a = b - 1$. Hence $\mathfrak{a} \cup \mathfrak{b} = \{B, a\} > \mathfrak{b}$. Thus $\mathfrak{a} > \mathfrak{a} \cap \mathfrak{b}$ implies $\mathfrak{a} \cup \mathfrak{b} > \mathfrak{b}$ and B1 holds in the lattice of ideals of \mathfrak{S} .

The point ideals of \mathfrak{L}_z are clearly the ideals $\mathfrak{p}_i = \{(i), \infty\}$. Now $\mathfrak{p}_1 \cup \mathfrak{p}_2 \cup \dots = u$ and hence $\mathfrak{L}_z = \mathfrak{L}$. The ascending chain $\mathfrak{p}_1 \subset \mathfrak{p}_1 \cup \mathfrak{p}_2 \subset \mathfrak{p}_1 \cup \mathfrak{p}_2 \cup \mathfrak{p}_3 \subset \dots$ has distinct members and \mathfrak{L}_z is thus *not* archimedean. The point ideals $\mathfrak{p}_1, \mathfrak{p}_2, \dots$ are *not* independent since the union of any infinite set is u . However, every finite set of the \mathfrak{p}_i is independent and thus generates a Boolean algebra. \mathfrak{L}_z is *not* complemented. For if $\mathfrak{p}_i \cup \mathfrak{a} = u$, then $\mathfrak{a} = u$ and $\mathfrak{a} \supset \mathfrak{p}_i$. The simple ideals of \mathfrak{L}_z are the elements of the form $\{A, a\}$ where $n(A) = a - 1$. Clearly \mathfrak{p}_i cannot be represented as a cross-cut of simple ideals. Hence it is *not* true that every ideal of \mathfrak{L}_z may be represented as a cross-cut of simple ideals. The irreducibles of \mathfrak{S} are the simple elements of \mathfrak{L}_z , namely, those elements $\{A, a\}$ with $n(A) = a - 1$. Now let A_i be the set $\{1, 2, \dots, i-1, i+1, \dots, k\}$ ($i = 1, \dots, k$). Then $\alpha_i = \{A_i, k\}$ is simple for each i and $z = \alpha_1 \cap \alpha_2 \cap \dots \cap \alpha_k$ since $A_1 \cap A_2 \cap \dots \cap A_k$ is null. This representation of z is clearly reduced. Hence for any positive integer $k > 1$, z has a reduced decomposition with k components.

This example clearly indicates the complications that may arise if \mathfrak{L}_a is *not* archimedean even though the ascending chain condition holds in \mathfrak{S} .

8. Example of a lattice satisfying E_5 which is not a Birkhoff lattice. Let S be the set of elements p_1, p_2, p_3, \dots . From the set of all subsets of S omit those infinite sets which contain either p_1 or p_2 but not both. Denote this set of subsets by \mathfrak{S} . \mathfrak{S} is clearly closed under infinite cross-cut. Since \mathfrak{S} contains a unit element, the union of any set of sets of \mathfrak{S} may be defined in terms

of the cross-cut operation. \mathfrak{S} is thus a continuous lattice in which every element is a union of points. Now consider $A \cup p$ where $A \in \mathfrak{S}$ and p is any element of S . If A is finite, then clearly $A \cup p = A + p$ where $+$ indicates set-theoretic union. Also if A is infinite and contains both p_1 and p_2 , $A \cup p = A + p$. Now if A is infinite and does not contain p_1 or p_2 , then $A \cup p = A + p$ if $p \neq p_1, p_2$ and $A \cup p = A + p_1 + p_2$ if $p = p_1$ or $p = p_2$. Now let $A + p_1 + p_2 \supset B \supset A$. If $B \neq A$, then B contains p_1 or p_2 and hence contains both p_1 and p_2 by the definition of \mathfrak{S} . Thus $B = A + p_1 + p_2$. Hence in every case $A \cup p > A$ if $p \notin A$.

Now let $A \supset B \supset A \cap C$ and $C \neq A \cap C$ where A, B, C are in \mathfrak{S} . Since every element of \mathfrak{S} is a union of points and $C \neq A \cap C$, there exists a point p such that $C \supset p$, $A \cap C \not\supset p$. Set $C_1 = (A \cap C) \cup p$. Then $C \supset C_1 \supset A \cap C$ and $C_1 \neq A \cap C$. Now $B \cup p \supset A \cap (B \cup C_1) \supset B$ and $B \cup p \neq A \cap (B \cup C_1)$ since otherwise $A \supset B \cup p \supset p$ and $A \cap C \supset p$ which contradicts $A \cap C \not\supset p$. Since $B \cup p > B$ we thus have $B = A \cap (B \cup C_1)$ and hence E_5 holds in \mathfrak{S} .

In \mathfrak{S} let \mathfrak{a} be the ideal generated by the sets $A_k = \{p_k, p_{k+1}, \dots\}$ ($k=3, 4, 5, \dots$). Then by Theorem 2.1, there exists a point ideal \mathfrak{p} such that $\mathfrak{a} \supset \mathfrak{p} > z$. Every set of \mathfrak{S} occurring in \mathfrak{p} contains an infinite number of elements. For suppose that $Q \in \mathfrak{p}$ and Q contains only a finite number of elements. Let k be the largest subscript occurring among the elements of Q . Then $z = Q \cap A_{k+1} \supset \mathfrak{p} \cap \mathfrak{a} \supset \mathfrak{p}$ which contradicts $\mathfrak{p} > z$. If $Q \in \mathfrak{p}$, then $Q \cup p_1 \supset p_2$ by the definition of \mathfrak{S} . Hence $\mathfrak{p} \cup p_1 \supset p_1 \cup p_2 \supset p_1$ where $\mathfrak{p} \cup p_1 \neq p_1 \cup p_2$ and $p_1 \cup p_2 \neq p_1$. Thus $\mathfrak{p} \cup p_1 \not\supset p_1$ and hence \mathfrak{S} is not a Birkhoff lattice. If $A \supset B$ and $A \neq B$, let $A \supset p$, $B \not\supset p$. Then $A \supset B_1 > B$ where $B_1 = B \cup p$. Thus \mathfrak{S} is an example of a continuous point lattice in which covering elements exist and E_5 holds, but which is not a Birkhoff lattice. Whether or not an exchange lattice, i.e., a continuous point lattice satisfying E_5 and a finite dependence axiom (Mac Lane [1]) is a Birkhoff lattice is an open question.

REFERENCES

G. BIRKHOFF

1. *Rings of sets*, Duke Mathematical Journal, vol. 3 (1937), pp. 443-454.
2. *On combination of subalgebras*, Proceedings of the Cambridge Philosophical Society, vol. 29 (1933), pp. 441-464.

R. P. DILWORTH

1. *Lattices with unique irreducible decompositions*, Annals of Mathematics.
2. *The arithmetical theory of Birkhoff lattices*, submitted to Duke Mathematical Journal.

S. MAC LANE

1. *A lattice formulation for transcendence degrees and p -bases*, Duke Mathematical Journal, vol. 24 (1938), pp. 455-468.

M. H. STONE

1. *The theory of representations of Boolean algebra*, these Transactions, vol. 40 (1936), pp. 37-111.

H. WALLMAN

1. *Lattices and topological spaces*, Annals of Mathematics, (2), vol. 39 (1938), pp. 112-126.

YALE UNIVERSITY,
NEW HAVEN, CONN.