

THE ϕ RECTIFIABLE SUBSETS OF THE PLANE

BY

A. P. MORSE AND JOHN F. RANDOLPH

CONTENTS

| | |
|--|-----|
| 1. Introduction..... | 236 |
| 2. Definitions..... | 238 |
| 3. Measure..... | 242 |
| 4. Connected sets..... | 246 |
| 5. Density theory..... | 250 |
| 6. Rectifiability and directionality..... | 262 |
| 7. Restrictedness and rectifiability..... | 271 |
| 8. Density ratios..... | 287 |
| 9. Density ratios and rectifiability..... | 293 |
| 10. Rectifiability and density ratios..... | 299 |
| 11. Interrelations..... | 302 |

1. Introduction. Throughout the paper, as will be reaffirmed in §2, we denote Euclidean space of two dimensions by E_2 and the family of all its subsets by \mathcal{E}_2 .

Among the functions on \mathcal{E}_2 to the extended non-negative real number system there are several that for some intuitive reason or another are designated as linear measure functions (see supplementary bibliography). Of these linear measure functions, that of Carathéodory, designated by L (see definitions 2.6 and 2.7), is without doubt the best known.

Besicovitch and others have made extensive studies of relations between rectifiable arcs and Carathéodory linearly measurable sets. These discussions have linear density properties as their central theme. It is natural to ask to what extent such results hold for the other linear measure functions. We answer such questions in this paper; but, rather than investigate each known linear measure function separately, we consider instead the measures of the class \mathfrak{U} (see 2.3) which satisfy the first four Carathéodory axioms, and introduce various notions in terms of density relations that suggest linearity.

For x a point of E_2 , A a set in E_2 , and ϕ a measure function, the ϕ densities

$$\mathfrak{D}_{\phi}^{\nabla}(A, x), \mathfrak{D}_{\phi}^{\Delta}(A, x), \mathfrak{D}_{\phi}^{\blacktriangle}(A, x)$$

of A at x are defined in 2.11. Our notions of directionality, ϕ directionality, restrictedness and ϕ restrictedness of A at x are also given among the definitions of §2. Also given are the notions of rectifiability and ϕ rectifiability of plane sets.

Presented to the Society, October 30, 1943; received by the editors March 2, 1943, and, in revised form, September 4, 1943.

If a set A lies on a rectifiable arc and ϕ is a measure which in any sense is a generalization of arc length, then it is rather to be expected that

$$0 < \mathfrak{D}_{\phi}^{\nabla}(A, x) = \mathfrak{D}_{\phi}^{\Delta}(A, x) = \mathfrak{D}_{\phi}^{\blacktriangle}(A, x)$$

for ϕ almost all x of A . On the other hand if E_2 contains A , $\phi \in \mathfrak{U}$, $\phi(A) < \infty$ and $\mathfrak{D}_{\phi}^{\nabla}(A, x) < \infty$ for ϕ almost all x in A , then (section 11.1)

$$\mathfrak{D}_{\phi}^{\Delta}(A, x) = \mathfrak{D}_{\phi}^{\nabla}(A, x)$$

for ϕ almost all x in A implies that A is essentially on a rectifiable arc; in fact if for ϕ almost all x in A either

$$\mathfrak{D}_{\phi}^{\Delta}(A, x) < 1.01 \mathfrak{D}_{\phi}^{\nabla}(A, x)$$

or

$$3\mathfrak{D}_{\phi}^{\blacktriangle}(A, x) < 4\mathfrak{D}_{\phi}^{\nabla}(A, x)$$

then there is a rectifiable arc which contains all of A except a subset of arbitrarily small ϕ measure.

A further conception of the nature of some of our results and their setting may be gained by reading 11.1, 11.2, 11.4 and 11.6.

If it is possible to inclose a non-measurable set A in a measurable set B with the same measure, then it is customary to elucidate the geometric properties of A by a consideration of the geometric properties of B . However, the generality of the measures which we consider precludes this method of attack.

In order to discuss the subsets of E_2 which have finite measure, but which are otherwise arbitrary, we make considerable use of the known techniques mentioned in 3.1 and 3.3. Because of these techniques it is in general relatively unimportant to us whether or not the sets we consider are measurable. In fact we are rather inclined to more than half believe the following overstatement: Measurability information is usually a luxury, rarely a convenience, never a necessity.

We are well aware that some of our results are old; in particular, our theorems which involve descriptive restrictedness and directionality are found in a different notation on pp. 262–269 of [7]⁽¹⁾ or pp. 28–30 of [6]. However, these known results are included in the paper partly for completeness and partly to make clearer the analogy and lack of analogy between the linear descriptive and the linear metric properties of plane sets. In this connection we feel we should point out that our notation $\text{dir}(A, x)$ has the same meaning which some other authors attach to the symbol $\text{contg}_A x$.

Now it is known and we also prove in 6.12 that

$$\mathfrak{D}_L^{\nabla}(A, x) = \mathfrak{D}_L^{\Delta}(A, x) = 1$$

(¹) Numbers in brackets refer to the bibliography at the end of the paper.

for L almost all x in a strictly rectifiable set A . From this, 5.15, and the known Theorem 3.4 we infer the known fact

$$\mathfrak{D}_L^\nabla(A, x) = \mathfrak{D}_L^\Delta(A, x) = 1$$

for L almost all x in an L rectifiable set A .

Supposing $L(A) < \infty$, we infer from 5.16 that

$$\mathfrak{D}_L^\Delta(A, x) \leq 1$$

for L almost all x ; we therefore conclude from 11.1(3) and 11.1(1) that the set

$$AE_x [3/4 < \mathfrak{D}_L^\nabla(A, x) < 1]$$

has L measure zero.

Accordingly we have the

THEOREM. *If $L(A) < \infty$, then the set*

$$AE_x [3/4 < \mathfrak{D}_L^\nabla(A, x) < 1]$$

has L measure zero.

This rather striking result of Besicovitch may be found on p. 324 of [1].

We also see at once, with the aid of 11.1(1) and 11.1(4), that the set

$$AE_x [\mathfrak{D}_L^\nabla(A, x) = \mathfrak{D}_L^\Delta(A, x) < 1]$$

has L measure zero whenever $L(A) < \infty$. This accords with chap. 11, pp. 324–327 of Besicovitch's [1].

Thus, if in 11.1 we replace " ϕ " by " L ," Theorem 11.1 becomes, in essence, a compendium of the results of Besicovitch in [1].

With three exceptions, namely, 7.5, 7.9 and 7.10, the fixed notations and definitions used in the paper are assembled in §2. In the beginning of §2 we list certain notations and definitions most of which are either suggestive or are in common use. Of the definitions enumerated in the remainder of §2 the reader will require for §3 only 2.3 through 2.8. The paper has been written with the idea in mind that the reader will make no more than a cursory examination of 2.1 through 2.27 before beginning his perusal of §3.

2. Definitions. Throughout the paper we use E_1 and E_2 for Euclidean space of one and two dimensions respectively, and \mathcal{E}_2 for the family of all subsets of E_2 . For x in E_2 we define the real numbers x_1, x_2 and $|x|$ so that $x = (x_1, x_2)$ and $|x| = (x_1^2 + x_2^2)^{1/2}$. We shall regard E_2 and the finite complex plane as the same with $i = (0, 1)$ and $i^2 = (-1, 0)$.

We regard the empty set and the number 0 as the same.

The notational phrase $x \in A$ is read x is a member of A ; $x \notin A$ is read x

is not a member of A ; and $A \subset B$ is read A is a subset of B . For $A \subset E_2$ the set $E_2 - A$ is denoted by \bar{A} .

The set $E_1 \cdot E_1[a \leq t \leq b]$ is denoted by $[a, b]$. We say J is a closed interval if and only if J is of the form $[a, b]$ where $-\infty < a < b < \infty$. We say J is an open interval if and only if J is of the form $E_1[a < t < b]$ where $-\infty < a < b < \infty$.

We use $\{x\}$ as an abbreviation for $E_y[y=x]$ and use $\{ \}$ only in this sense. Note that $\{x\}$ is the set whose sole element is x .

Whenever A is a nonvacuous subset of E_2 we define $\text{diam}(A)$ as the supremum of numbers of the form $|x-y|$ where $x \in A$ and $y \in A$; if $A = 0$ we define $\text{diam}(A) = 0$. Also we use $\text{clsr}(A)$ for the closure of A and $\text{bndr}(A)$ for the boundary of A . Thus $\text{bndr}(A) = (\text{clsr } A) \setminus (\text{clsr } \bar{A})$.

For $A \subset E_2$ and $B \subset E_2$ we define $\text{dist}(A, B)$ as the infimum of numbers of the form $|x-y|$ where $x \in A$ and $y \in B$.

We say A is disconnected if and only if $A \subset E_2$ and there exist non-empty subsets C_1 and C_2 of A such that

$$A = C_1 + C_2 \quad \text{and} \quad C_1(\text{clsr } C_2) + C_2(\text{clsr } C_1) = 0.$$

A set A is said to be connected if and only if $A \subset E_2$ and A is not disconnected. A closed connected set is said to be a continuum.

If a_x is a set for x in a set B , then

$$\sum_{x \in B} a_x \quad \text{and} \quad \prod_{x \in B} a_x$$

are so defined that

$$\begin{aligned} \sum_{x \in B} a_x &= E_y [y \in a_x \text{ for some } x \in B], \\ \prod_{x \in B} a_x &= E_y [y \in a_x \text{ for each } x \in B]. \end{aligned}$$

In connection with set theoretic sums we point out that

$$\mathfrak{F} = \sum_{s \in \mathfrak{F}} \{s\}$$

whereas

$$E_x [x \in s \text{ for some } s \in \mathfrak{F}] = \sum_{s \in \mathfrak{F}} s.$$

In fact we here and now define the more convenient notation $\sigma(\mathfrak{F})$ by

$$\sigma(\mathfrak{F}) = \sum_{s \in \mathfrak{F}} s.$$

It should be noted that $\sigma(\{y\}) = y$. Likewise

$$\sigma\left(\sum_{x \in B} a_x\right) = \sum_{x \in B} \sigma(a_x).$$

If $a_x \geq 0$ for x in a countable set B then

$$\sum_{x \in B} a_x$$

will also be used to denote the appropriate numerical sum. Whether numerical summation or point set summation is intended will be obvious from the context.

A family \mathfrak{F} is said to cover A if and only if $A \subset \sigma(\mathfrak{F})$.

In this paper we call a set A a Borel set if and only if it is a member of every subfamily \mathfrak{F} of \mathcal{E}_2 for which:

- (i) $B \in \mathfrak{F}$ whenever B is a closed subset of E_2 ;
- (ii) $\sigma(\mathfrak{G}) \in \mathfrak{F}$ whenever \mathfrak{G} is a countable subfamily of \mathfrak{F} ;
- (iii) $\tilde{B} \in \mathfrak{F}$ whenever $B \in \mathfrak{F}$.

We now enumerate some definitions and remarks specifically referred to later on.

2.1. DEFINITION. We say A and B are *disjoint* if and only if $AB = 0$. We say a family \mathfrak{F} of sets is *disjointed* if and only if $AB = 0$ whenever A and B are different members of \mathfrak{F} .

2.2. DEFINITION. We say A and B are *semi-disjoint* if and only if neither is a subset of the other. We say a family \mathfrak{F} is *semi-disjointed* if and only if no member of \mathfrak{F} is a subset of a different member.

2.3. DEFINITION. The family \mathfrak{U} is defined by $\phi \in \mathfrak{U}$ if and only if ϕ is a function on \mathcal{E}_2 to $E_t [0 \leq t \leq \infty]$ such that:

- I. $\phi(0) = 0$;
- II. $\phi(B) \leq \phi(A)$ whenever $B \subset A \subset E_2$;
- III. If \mathfrak{F} is a countable subfamily of \mathcal{E}_2 , then

$$\phi[\sigma(\mathfrak{F})] \leq \sum_{s \in \mathfrak{F}} \phi(s);$$

- IV. $\phi(A+B) = \phi(A) + \phi(B)$ whenever $\text{dist}(A, B) > 0$.

2.4. DEFINITION. The family \mathfrak{M} is defined by $\phi \in \mathfrak{M}$ if and only if $\phi \in \mathfrak{U}$ and $\phi(E_2) < \infty$.

Due to Carathéodory is the following

2.5. DEFINITION. We say A is ϕ measurable if and only if $A \subset E_2$, $\phi \in \mathfrak{U}$,

$$\phi(W) = \phi(WA) + \phi(W\tilde{A})$$

for each set $W \subset E_2$.

2.6. DEFINITION. For $0 < p$ we define L_p as the function on \mathcal{E}_2 such that $A \subset E_2$ implies $L_p(A)$ is the infimum of numbers of the form

$$\sum_{s \in \mathfrak{G}} (\text{diam } s)$$

where \mathfrak{G} is such a countable family of nonvacuous open subsets of E_2 that

$$\sigma(\mathfrak{G}) \supset A$$

and

$$\text{diam } s < \rho \text{ whenever } s \in \mathfrak{G}.$$

2.7. DEFINITION. We define L as the function on \mathcal{E}_2 such that $A \subset E_2$ implies

$$L(A) = \lim_{\rho \rightarrow 0+} L_\rho(A).$$

Definitions 2.6 and 2.7 are due to Carathéodory [2]. The number $L(A)$ is commonly known as the Carathéodory linear measure of A .

2.8. DEFINITION. By *counting measure* we mean the function ϕ on \mathcal{E}_2 such that $\phi(s)$ is the number of points in s whenever $s \subset E_2$.

2.9. ABBREVIATION. We use the symbol \mathfrak{R}_x^r as an abbreviation for

$$E_2 \cap \{ |y - x| < r \}$$

and the symbol \mathfrak{C}_x^r as an abbreviation for

$$E_2 \cap \{ |y - x| \leq r \}.$$

2.10. DEFINITION. A set Γ will be called a *circle* if and only if there is an $x \in E_2$ and a number r with $0 < r < \infty$ such that $\Gamma = \mathfrak{R}_x^r$.

2.11. DEFINITION. For $\phi \in \mathfrak{U}$, $A \subset E_2$ and $x \in E_2$ we define

$$\mathfrak{D}_\phi^\nabla(A, x) = \lim_{r \rightarrow 0+} \frac{\phi(A \cap \mathfrak{R}_x^r)}{2r},$$

$$\mathfrak{D}_\phi^\Delta(A, x) = \overline{\lim}_{r \rightarrow 0+} \frac{\phi(A \cap \mathfrak{R}_x^r)}{2r},$$

$$\mathfrak{D}_\phi^\Delta(A, x) = \overline{\lim}_{(\text{diam } \Gamma) \rightarrow 0, \Gamma \in \mathfrak{C}_x} \frac{\phi(A \cap \Gamma)}{(\text{diam } \Gamma)},$$

where $\mathfrak{C}_x = \{ \Gamma \mid \Gamma \text{ is a circle and } x \in \Gamma \}$ and call $\mathfrak{D}_\phi^\nabla(A, x)$ and $\mathfrak{D}_\phi^\Delta(A, x)$ respectively the lower and upper ϕ density of A at x .

2.12. DEFINITION. For $\phi \in \mathfrak{U}$ and $x \in E_2$ we define

$$\mathfrak{D}_\phi^\nabla(x) = \mathfrak{D}_\phi^\nabla(E_2, x), \quad \mathfrak{D}_\phi^\Delta(x) = \mathfrak{D}_\phi^\Delta(E_2, x) \quad \text{and} \quad \mathfrak{D}_\phi^\Delta(x) = \mathfrak{D}_\phi^\Delta(E_2, x).$$

2.13. DEFINITIONS. A set A is said to be *strictly rectifiable* if and only if $A \subset E_2$ and there is: a number $M > 0$; a closed interval J ; a function f , with domain J and range A , such that f satisfies the Lipschitz condition

$$|f(t'') - f(t')| \leq M |t'' - t'| \quad \text{for } t' \text{ and } t'' \text{ in } J.$$

2.14. DEFINITION. A set A is said to be *rectifiable* if and only if A is a subset of a strictly rectifiable set.

We shall use the obvious and elementary properties of rectifiable sets listed in

2.15. REMARK. (i) If A is rectifiable then $L(A) < \infty$.

(ii) Any subset of a rectifiable set is rectifiable.

(iii) If \mathfrak{F} is a finite family of rectifiable sets then $\sigma(\mathfrak{F})$ is rectifiable.

(iv) If A and A' are congruent and A' is rectifiable then A is rectifiable.

(v) If A is rectifiable then $\text{clsr } A$ is rectifiable.

2.16. DEFINITION. A set A is said to be ϕ rectifiable if and only if $A \subset E_2$, $\phi \in \mathbb{U}$ and corresponding to each $\epsilon > 0$ there is a strictly rectifiable set B with $\phi(A \setminus B) < \epsilon$.

2.17. DEFINITION. For $\phi \in \mathbb{U}$, $A \subset E_2$ and $x \in E_2$ we define:

(i) $\text{Sgn}(A, x) = \text{clsr} [\sum_{y \in B} \{(y-x)/|y-x|\}]$ where $B = A - \{x\}$;

(ii) $\text{dir}(A, x) = \prod_{\beta \in \mathfrak{F}} \text{Sgn}(A\beta, x)$ where $\mathfrak{F} = E_\beta[\beta \mathfrak{R}_x^r = 0 \text{ for some } r > 0]$;

(iii) $\text{dir}_\phi(A, x) = \prod_{\beta \in \mathfrak{G}} \text{Sgn}(A\beta, x)$ where $\mathfrak{G} = E_\beta[\mathfrak{D}_\phi^\Delta(\beta, x) = 0]$.

2.18. DEFINITION. For $\phi \in \mathbb{U}$ and $x \in E_2$ we define

$$\text{dir}_\phi(x) = \text{dir}_\phi(E_2, x)$$

2.19. DEFINITION. A set is said to be *diametral* if and only if it is of the form $\{-z\} + \{z\}$ where $z \in E_2$ and $|z| = 1$.

A set is said to be *nondiametral* if and only if it is a subset of E_2 and is not diametral.

2.20. DEFINITION. We say A is *directional* at x if and only if $\text{dir}(A, x)$ is diametral.

2.21. DEFINITION. We say A is ϕ *directional* at x if and only if $\text{dir}_\phi(A, x)$ is diametral.

2.22. DEFINITION. We say A is *restricted* at x if and only if there is a point z in E_2 with $|z| = 1$ for which $\{z\} \text{ dir}(A, x) = 0$.

2.23. REMARK. If $A \subset B$ and B is restricted at x then so is A .

2.24. DEFINITION. We say A is ϕ *restricted* at x if and only if

$$\mathfrak{D}_\phi^\Delta(A, x) > 0$$

and there is a diametral set Z for which

$$Z \text{ dir}_\phi(A, x) = 0.$$

2.25. ABBREVIATION. We employ \hat{A} as an abbreviation for

$$E_2 E_x [|x - y| \leq 2 \text{ diam } A \text{ for some } y \text{ in } A].$$

2.26. REMARK. We note that $\hat{\Gamma} = \mathfrak{R}_x^{5r}$ whenever $\Gamma = \mathfrak{R}_x^r$.

2.27. ABBREVIATION. We use $p * q$ as an abbreviation for

$$\mathfrak{R}_p^{|p-q|} \mathfrak{R}_q^{|p-q|}.$$

3. Measure. Consulting 2.3, 2.4, 2.5 for nomenclature we recall certain

general properties of measure.

If A is ϕ measurable and W and B are arbitrary subsets of E_2 , then

$$\phi(WA + WB) + \phi(WAB) = \phi(WA) + \phi(WB).$$

If A is ϕ measurable and $\phi(B) < \infty$ then

$$\phi(B - A) = \phi(B) - \phi(BA).$$

If \mathfrak{F} is a countable family of ϕ measurable sets then $\sigma(\mathfrak{F})$ is ϕ measurable.

If A is ϕ measurable then \bar{A} is ϕ measurable.

If $W \subset E_2$ and \mathfrak{F} is a countable disjointed family ϕ of measurable sets then

$$\phi\left(\sum_{s \in \mathfrak{F}} WS\right) = \sum_{s \in \mathfrak{F}} \phi(WS).$$

If B is a Borel set and $\phi \in \mathfrak{U}$ then B is ϕ measurable.

Several comments, which may incidentally help explain some peculiarities of certain subsequent proofs, are incorporated in

3.1. REMARK. Suppose $A_1 \subset A_2 \subset A_3 \subset \dots$ with $A = \sum_{n=1}^{\infty} A_n \subset E_2$.

If $\phi \in \mathfrak{U}$, W is an arbitrary subset of E_2 , and A_n is ϕ measurable for each positive integer n , then as is well known

$$(i) \quad \phi(WA) = \lim_{n \rightarrow \infty} \phi(WA_n).$$

Furthermore if $\phi \in \mathfrak{U}$ and for each positive integer n , A_n is contained in a ϕ measurable set with the same ϕ measure as A_n , then

$$\phi(A) = \lim_{n \rightarrow \infty} \phi(A_n).$$

However, even with $\phi \in \mathfrak{M}$, it may happen that

$$\lim_{n \rightarrow \infty} \phi(A_n) < \phi(A),$$

as may be shown by an example. Accordingly certain techniques, available to us in handling such a measure (see 2.6 and 2.7) as L , are unemployable in dealing with a general measure ϕ in \mathfrak{M} or \mathfrak{U} . Nevertheless if $\phi \in \mathfrak{U}$ and $\phi(A) > 0$ it is obvious from 2.3 III that

$$\lim_{n \rightarrow \infty} \phi(A_n) > 0.$$

We state without proof

3.2. THEOREM. If $\phi \in \mathfrak{U}$, $A \subset E_2$, and Φ is the function on E_2 such that $\Phi(s) = \phi(As)$ for $s \subset E_2$, then $\Phi \in \mathfrak{U}$ and s is Φ measurable whenever s is ϕ measurable; furthermore if in addition $\phi(A) < \infty$, then $\Phi \in \mathfrak{M}$.

3.3. CLICHÉ. If $\phi \in \mathfrak{U}$ and $\phi(A) < \infty$ and some statement involving ϕ and

A is readily proved by the introduction of the Φ of 3.2, we indicate this state of affairs by the use of some phrase containing the words "intersecting measure."

Our first use of intersecting measure occurs in 3.7.

After referring to 2.6 and 2.7 it is easy to check the following well known

3.4. THEOREM. $L \in \mathfrak{U}$ and corresponding to each set $A \subset E_2$ there is a G_δ set B for which

$$A \subset B \quad \text{and} \quad L(A) = L(B).$$

In this section and throughout the paper the measure functions in which we are primarily interested are rather special in that they are on \mathcal{E}_2 and belong either to \mathfrak{U} or its subclass \mathfrak{M} . Nevertheless the results of this section and their proofs are valid when we regard E_2 as an abstract metric space. In this sense 3.7 and 3.13 are apparently [7, Lemma 15.1, p. 152] heretofore unknown, the latter being an extension of a result, due to Hahn [3, Theorem IV, p. 447], which lessens the importance in measure theory of such special Borel sets as G_δ 's.

The following lemma is easy to prove without transfinite induction (see [4, p. 159]) and is rather well known.

3.5. LEMMA. A set $A \subset E_2$ is a Borel set if and only if it is a member of every family \mathfrak{F} such that:

- (i) $B \in \mathfrak{F}$ implies $B \subset E_2$;
- (ii) $B \in \mathfrak{F}$ whenever B is a closed subset of E_2 ;
- (iii) \mathfrak{G} is a countable nonvacuous subfamily of \mathfrak{F} implies

$$\prod_{s \in \mathfrak{G}} s \in \mathfrak{F} \quad \text{and} \quad \sum_{s \in \mathfrak{G}} s \in \mathfrak{F}.$$

We are indebted to Alfred Horn for pointing out that the use of the above lemma materially simplifies the proof of the following

3.6. THEOREM. If $\phi \in \mathfrak{M}$, A is a Borel set, and $\epsilon > 0$, then there is a closed subset C of A with $\phi(A \setminus C) < \epsilon$.

Proof. Let \mathfrak{F} be the family defined by the conditions: $B \in \mathfrak{F}$ if and only if $B \subset E_2$ and corresponding to each number $\epsilon > 0$ there is a closed set C such that $\phi(B \setminus C) < \epsilon$. The members of \mathfrak{F} are therefore subsets of E_2 and clearly $B \in \mathfrak{F}$ whenever B is a closed subset of E_2 . Also $B \in \mathfrak{F}$ implies B is ϕ measurable.

Let A_1, A_2, A_3, \dots be sets in \mathfrak{F} . Let $\eta > 0$. For $n = 1, 2, 3, \dots$ choose a closed set C_n such that

$$C_n \subset A_n, \quad \phi(A_n - C_n) < \eta 2^{-n}.$$

Thus

$$\prod_{j=1}^{\infty} A_j - \prod_{j=1}^{\infty} C_j \subset \sum_{j=1}^{\infty} (A_j - C_j)$$

and

$$(1) \quad \phi\left(\prod_{j=1}^{\infty} A_j - \prod_{j=1}^{\infty} C_j\right) \leq \phi\left[\sum_{j=1}^{\infty} (A_j - C_j)\right] \leq \sum_{j=1}^{\infty} \phi(A_j - C_j) < \eta$$

where $\prod_{j=1}^{\infty} C_j$ is clearly a closed subset of $\prod_{j=1}^{\infty} A_j$. On the other hand

$$\begin{aligned} \lim_{n \rightarrow \infty} \phi\left(\sum_{j=1}^{\infty} A_j - \sum_{j=1}^n C_j\right) &= \phi\left(\sum_{j=1}^{\infty} A_j - \sum_{j=1}^{\infty} C_j\right) \leq \phi\left[\sum_{j=1}^{\infty} (A_j - C_j)\right] \\ &\leq \sum_{j=1}^{\infty} \phi(A_j - C_j) < \eta, \end{aligned}$$

and n may be chosen so large that

$$(2) \quad \phi\left(\sum_{j=1}^{\infty} A_j - \sum_{j=1}^n C_j\right) < \eta$$

where $\sum_{j=1}^n C_j$ is of course a closed subset of $\sum_{j=1}^{\infty} A_j$. Accordingly the definition of \mathfrak{F} , the arbitrariness of η , (1), and (2), combine to tell us that

$$\sum_{j=1}^{\infty} A_j \in \mathfrak{F}, \quad \prod_{j=1}^{\infty} A_j \in \mathfrak{F}.$$

Thus using 3.5 we see that each Borel set is a member of \mathfrak{F} .

The theorem now follows from the definition of \mathfrak{F} .

3.7. THEOREM. *If $\phi \in \mathfrak{U}$, A is a Borel set with $\phi(A) < \infty$, and $\epsilon > 0$, then there is a closed subset C of A with $\phi(A \setminus C) < \epsilon$.*

Proof. In the spirit of 3.3 we introduce intersecting measure; that is we let Φ be the function on \mathcal{E}_2 such that

$$\Phi(s) = \phi(As) \quad \text{for } s \subset E_2,$$

invoke 3.2, note that $\Phi \in \mathfrak{M}$ and $\Phi(A\tilde{C}) = \phi(A\tilde{C})$ for $C \subset E_2$ and use 3.6.

3.8. COROLLARY. *If $\phi \in \mathfrak{U}$ and A is a Borel set with $\phi(A) < \infty$, then A is an \mathfrak{F}_σ plus a set of ϕ measure zero.*

3.9. COROLLARY. *If $\phi \in \mathfrak{U}$, $\psi \in \mathfrak{U}$, B is a Borel set, $\phi(B) + \psi(B) < \infty$, and if $\phi(C) = \psi(C)$ for each closed set $C \subset B$, then $\phi(B) = \psi(B)$.*

3.10. THEOREM. *If $\phi \in \mathfrak{M}$, $\psi \in \mathfrak{M}$ and $\phi(C) = \psi(C)$ for each closed set C , then $\phi(B) = \psi(B)$ for each Borel set B .*

Proof. This theorem is an immediate consequence of Corollary 3.9.

3.11. COROLLARY. If $\phi \in \mathfrak{M}$, $\psi \in \mathfrak{M}$, and $\phi(G) = \psi(G)$ for each open set G , then $\phi(B) = \psi(B)$ for each Borel set B .

3.12. THEOREM. If $\phi \in \mathfrak{U}$ and ψ is the function on E_2 such that $s \subset E_2$ implies $\psi(s)$ is the infimum of numbers of the form $\phi(S)$ where S is an open set containing s , then $\psi \in \mathfrak{U}$; furthermore if in addition $\phi \in \mathfrak{M}$, then $\psi \in \mathfrak{M}$ and $\phi(B) = \psi(B)$ for each Borel set B .

Proof. Clearly $\psi \in \mathfrak{U}$ whenever $\phi \in \mathfrak{U}$. We let $\phi \in \mathfrak{M}$, make a simple check to show that $\psi \in \mathfrak{M}$, note that $\phi(G) = \psi(G)$ for each open set G and apply Corollary 3.11.

3.13. THEOREM. If $\phi \in \mathfrak{U}$, $\epsilon > 0$, $\phi(\alpha) < \infty$, α is ϕ measurable, and if each subset A of E_2 is contained in a Borel set with the same ϕ measure, then there is a closed set β such that $\beta \subset \alpha$ and $\phi(\alpha \setminus \beta) < \epsilon$.

Proof. Let β_1 be a Borel set containing α with the same ϕ measure as α . Since $\phi(\beta_1 \setminus \alpha) = 0$ we may inclose $\beta_1 \setminus \alpha$ in a Borel set of ϕ measure zero and subtract this set from β_1 , in order to obtain a Borel set β_2 with $\beta_2 \subset \alpha$ and $\phi(\alpha) = \phi(\beta_1) = \phi(\beta_2)$. Application of Corollary 3.7 to the Borel set β_2 completes the proof.

A glance at 2.8 makes easy the proof of

3.14. THEOREM. If ϕ is counting measure, then $\phi \in \mathfrak{U}$ and every subset of E_2 is ϕ measurable.

4. **Connected sets.** Recalling the Definition 2.2 of a semi-disjointed family we state and prove

4.1. THEOREM. If \mathfrak{F} is any family of sets and

$$\mathfrak{F} \not\subseteq \left[s \subset S \right]$$

is a finite family whenever $s \in \mathfrak{F}$, then there is a semi-disjointed subfamily \mathfrak{G} of \mathfrak{F} such that

$$\sigma(\mathfrak{G}) = \sigma(\mathfrak{F}).$$

Proof. Let

$$\mathfrak{G} = \mathfrak{F} \setminus \left[s \subset S \in \mathfrak{F} \text{ implies } s = S \right].$$

It is obvious that \mathfrak{G} is semi-disjointed and from the hypothesis it is readily checked that every member of \mathfrak{F} is a subset of some set in \mathfrak{G} . Consequently

$$\sigma(\mathfrak{F}) \subset \sigma(\mathfrak{G}) \subset \sigma(\mathfrak{F}).$$

4.2. THEOREM. If A is a connected set, $x \in A$, and $0 \leq r \leq 2^{-1} (\text{diam } A)$, then $r \leq L(A \cap \mathbb{R}_2^r)$.

Proof. Since A is connected and $0 \leq r \leq 2^{-1} (\text{diam } A)$ it follows that $0 \leq r' < r$ implies there is a $y \in A$ for which $|y - x| = r'$. Let \mathfrak{F} be any countable family of nonvacuous open sets such that $\sigma(\mathfrak{F}) \supset A \setminus \mathfrak{R}_x$. For $s \in \mathfrak{F}$ let $M(s)$ and $m(s)$ be the supremum and infimum respectively of $\sum_{y \in s} \{|y - x|\}$ and note that $(\text{diam } s) \geq M(s) - m(s)$. Clearly $0 \leq r' < r$ implies there is an $s \in \mathfrak{F}$ for which $m(s) \leq r' \leq M(s)$, and therefore

$$\sum_{s \in \mathfrak{F}} (\text{diam } s) \geq \sum_{s \in \mathfrak{F}} [M(s) - m(s)] \geq r.$$

Consequently, the arbitrariness of \mathfrak{F} guarantees $L(A \setminus \mathfrak{R}_x) \geq r$.

For use in the next theorem and its proof the number $T_\alpha^\beta f$ is defined, whenever $-\infty < \alpha < \beta < \infty$ and f is a function whose domain includes $[\alpha, \beta]$ and whose range is contained in E_1 , as the supremum of numbers of the form

$$\sum_{i=1}^N |f(t_i) - f(t_{i-1})|$$

where N is a positive integer and $\alpha = t_0 < t_1 < t_2 < \cdots < t_N = \beta$.

We read $T_\alpha^\beta f$ as the total variation from α to β of f and recall, incidentally, that $L(S) = T_\alpha^\beta f$ whenever f is a continuous univalent function whose domain is the closed interval $[\alpha, \beta]$ and whose range is $SC E_2$.

4.3. THEOREM. *If \mathfrak{F} is a nonvacuous finite semi-disjointed family of n circles with $\sigma(\mathfrak{F})$ connected, then there is a number $a > 0$ and a function f such that:*

- (i) *the domain of f is $[0, a]$;*
- (ii) *the range of f is $\sum_{s \in \mathfrak{F}} (\text{bndr } s)$;*
- (iii) *f satisfies a Lipschitz condition on $[0, a]$;*
- (iv) $T_0^a f = \pi \sum_{s \in \mathfrak{F}} (\text{diam } s)$.

Proof. Assume the theorem is not true and let k be the smallest value of n for which the theorem is false. Since the theorem is obvious if $n = 1$ we see that $k \geq 2$.

Let \mathfrak{G} be a semi-disjointed family of k circles, with $\sigma(\mathfrak{G})$ connected, corresponding to which there is no function of the required type.

Since $\sigma(\mathfrak{G})$ is open and connected, it cannot be split into two non-empty open sets without points in common. Accordingly we select a circle of \mathfrak{G} and next select a different circle of \mathfrak{G} which intersects the first circle and proceeding step by step arrive at a family \mathfrak{H} of $k - 1$ circles of \mathfrak{G} such that $\sigma(\mathfrak{H})$ is connected. Letting s_0 be the sole member of $\mathfrak{G} - \mathfrak{H}$, we use the definition of k to ascertain a positive number a_0 and a function f_0 such that:

- (1) the domain of f_0 is $[0, a_0]$;
- (2) the range of f_0 is $\sum_{s \in \mathfrak{H}} (\text{bndr } s)$;
- (3) f_0 satisfies a Lipschitz condition;
- (4) $T_0^{a_0} f_0 = \pi \sum_{s \in \mathfrak{H}} (\text{diam } s)$.

Since $\sigma(\mathfrak{F})$ is connected, the set ss_0 is nonvacuous for some $s \in \mathfrak{H}$; and,

since s_0 and *this* s are semi-disjoint, the intersection of their boundaries is also nonvacuous. In view of this we select a point z_0 such that

$$z_0 \in (\text{bndr } s_0) \sum_{s \in \mathfrak{F}} (\text{bndr } s)$$

and choose t_0 so that $0 \leq t_0 \leq a_0$ and $f_0(t_0) = z_0$. Let x_0 and r_0 be such that $s_0 = \mathfrak{R}_{x_0}^{r_0}$, take α in E_1 so that $z_0 - x_0 = r_0 e^{i\alpha}$, and define the function f on $[0, a_0 + 2\pi]$ so that

$$f(t) = \begin{cases} f_0(t) & \text{for } 0 \leq t \leq t_0, \\ x_0 + r_0 e^{i(\alpha + t - t_0)} & \text{for } t_0 < t < t_0 + 2\pi, \\ f_0(t - 2\pi) & \text{for } t_0 + 2\pi \leq t \leq a_0 + 2\pi. \end{cases}$$

A check shows this function f corresponds to \mathfrak{G} as prescribed by our theorem, in contradiction to our previous statement that there was no such function.

By using the device known as parameterizing according to arc length we first obtain the

4.4. COROLLARY. *If \mathfrak{F} is a nonvacuous finite semi-disjointed family of open circles with $\sigma(\mathfrak{F})$ connected and if $l = \pi \sum_{s \in \mathfrak{F}} (\text{diam } s)$, then there is a function f on $[0, l]$ with range $\sum_{s \in \mathfrak{F}} (\text{bndr } s)$ such that*

$$|f(t'') - f(t')| \leq (t'' - t') \quad \text{for } 0 \leq t' \leq t'' \leq l.$$

Next by a transformation we obtain the

4.5. COROLLARY. *If \mathfrak{F} is a nonvacuous finite semi-disjointed family of open circles with $\sigma(\mathfrak{F})$ connected and if $l = \pi \sum_{s \in \mathfrak{F}} (\text{diam } s)$, then there is a function f on $[0, 1]$ with range $\sum_{s \in \mathfrak{F}} (\text{bndr } s)$ such that*

$$|f(t'') - f(t')| \leq l(t'' - t') \quad \text{for } 0 \leq t' \leq t'' \leq 1.$$

4.6. THEOREM. *If $A \subset E_2$ is a nonvacuous continuum with $L(A) < \infty$ then A is strictly rectifiable.*

Proof. We infer from 4.2 that A is bounded, and, relying on the fact that a bounded open set of diameter d may be included in a circle of diameter $2d$, we use Definition 2.7 of L and the Heine-Borel theorem to secure, for each positive integer n , a finite family \mathfrak{F}_n of circles such that

$$A \subset \sigma(\mathfrak{F}_n), \quad \sum_{s \in \mathfrak{F}_n} (\text{diam } s) < 2L(A) + 1$$

and such that, in addition,

$$\text{diam } s < 1/n, \quad sA \neq \emptyset \quad \text{for } s \in \mathfrak{F}_n.$$

Now for each positive integer n we select from \mathfrak{F}_n , according to Theorem 4.1, a semi-disjointed subfamily \mathfrak{G}_n covering A ; note that $s \in \mathfrak{G}_n$ implies

$$\sum_{S \in \mathfrak{G}_n} (\text{diam } S) < 2L(A) + 1, \quad \text{diam } s < 1/n, \quad sA \neq 0;$$

and, employing the connectedness of A , infer that $\sigma(\mathfrak{G}_n)$ is likewise connected. Upon setting

$$\lambda = \pi(2L(A) + 1)$$

we are now justified by 4.5 in ascertaining a sequence f such that: corresponding to each positive integer n , f_n is a function on $[0, 1]$; its range is $\sum_{S \in \mathfrak{G}_n} (\text{bndr } S)$; and it satisfies the Lipschitz condition

$$|f_n(t'') - f_n(t')| \leq \lambda(t'' - t') \quad \text{for } 0 \leq t' \leq t'' \leq 1.$$

Thus using the equi-continuity principle found, for example, in Theorem V on page 304 of [3], we extract a uniformly convergent subsequence g of f , and denote the limit function by \bar{g} . Clearly

$$|\bar{g}(t'') - \bar{g}(t')| \leq \lambda(t'' - t') \quad \text{whenever } 0 \leq t' \leq t'' \leq 1,$$

and a check reveals

$$A = \sum_{t \in [0,1]} \{\bar{g}(t)\}.$$

Consequently A is strictly rectifiable according to 2.13.

4.7. LEMMA. *If C is a continuum and \mathfrak{F} is a semi-disjointed family of circles such that*

(i) $\sigma(\mathfrak{F}) \subset C$,

(ii) $\epsilon > 0$ implies $\mathfrak{F}E_S[\text{diam } S > \epsilon]$ is finite,

then

$$D = [C - \sigma(\mathfrak{F})] + \sum_{S \in \mathfrak{F}} (\text{bndr } S)$$

is a continuum.

Proof. We first prove D is closed. Let p_0 be a cluster point of

$$\sum_{S \in \mathfrak{F}} (\text{bndr } S)$$

which is not a point of the closed set $C - \sigma(\mathfrak{F})$. Since C is closed, p_0 is a point of the open set $\sigma(\mathfrak{F})$ and we let $S_0 \in \mathfrak{F}$ be such that $p_0 \in S_0$ and note that

$$\delta = \text{dist}(\{p_0\}, \bar{S}_0)$$

is positive. Letting

$$\mathfrak{F} = \mathfrak{F} \setminus \mathfrak{F}_\delta [(\text{bndr } S) \mathfrak{R}_{p_0}^{3/2} \neq 0]$$

we see from the semi-disjointedness of \mathfrak{F} that $S \mathfrak{R}_{p_0}^{3/2} \neq 0$ and $S \bar{S}_0 \neq 0$ whenever

$S \in \mathfrak{S}$. Accordingly each member of \mathfrak{S} has diameter greater than $2^{-1}\delta$ and thus from (ii) it is clear that \mathfrak{S} is finite. Therefore p_0 is a cluster point of the closed set $\sum_{S \in \mathfrak{S}} (\text{bndr } S)$ with

$$p_0 \in \sum_{S \in \mathfrak{S}} (\text{bndr } S) \subset \sum_{S \in \mathfrak{F}} (\text{bndr } S) \subset D.$$

Hence D is closed.

Next the set D is connected as we prove by contradiction. Suppose $D = D_1 + D_2$ where D_1 and D_2 are non-empty closed sets and $D_1 D_2 = 0$.

We define

$$\mathfrak{F}_1 = \mathfrak{F} \overset{E}{\underset{\cdot}{\cap}} [(\text{bndr } s) \subset D_1], \quad \mathfrak{F}_2 = \mathfrak{F} \overset{E}{\underset{\cdot}{\cap}} [(\text{bndr } S) \subset D_2].$$

Since the boundary of a circle is connected we have

$$(1) \quad \mathfrak{F} = \mathfrak{F}_1 + \mathfrak{F}_2.$$

Moreover

$$(2) \quad \sigma(\mathfrak{F}_1)\sigma(\mathfrak{F}_2) = \sum_{s \in \mathfrak{F}_1} \sum_{S \in \mathfrak{F}_2} sS = 0$$

because the boundaries of two semi-disjoint intersecting circles intersect. Using the definition of D we conclude

$$\begin{aligned} \sigma(\mathfrak{F}_2)D_1 &= \sigma(\mathfrak{F}_2)D_1 \sum_{s \in \mathfrak{F}} (\text{bndr } s) = \sigma(\mathfrak{F}_2) \sum_{s \in \mathfrak{F}} [D_1(\text{bndr } s)] \\ &= \sigma(\mathfrak{F}_2) \sum_{s \in \mathfrak{F}_1} (\text{bndr } s) = \sum_{S \in \mathfrak{F}_2} \sum_{s \in \mathfrak{F}_1} S(\text{bndr } s) = 0; \end{aligned}$$

the last step follows from (2) and the fact that two circles intersect whenever one circle intersects the boundary of the other. By symmetry

$$(3) \quad \sigma(\mathfrak{F}_2)D_1 = \sigma(\mathfrak{F}_1)D_2 = 0.$$

With the aid of (ii) and the definitions of \mathfrak{F}_1 and \mathfrak{F}_2 we check that the sets

$$D_1 + \sigma(\mathfrak{F}_1) \quad \text{and} \quad D_2 + \sigma(\mathfrak{F}_2)$$

are closed and utilize (1), (2), and (3) to see that their intersection, namely

$$D_1 D_2 + \sigma(\mathfrak{F}_1)D_2 + \sigma(\mathfrak{F}_2)D_1 + \sigma(\mathfrak{F}_1)\sigma(\mathfrak{F}_2),$$

is vacuous. Consequently

$$C = D + \sigma(\mathfrak{F}) = [D_1 + \sigma(\mathfrak{F}_1)] + [D_2 + \sigma(\mathfrak{F}_2)]$$

and thus C is disconnected in contradiction to the hypothesis that C is a continuum.

5. Density theory. That the use of \mathfrak{R}'_x rather than \mathfrak{C}'_x in density definitions (see 2.11) is a convenience rather than a necessity is pointed out in the following

5.1. REMARK. If $\phi \in \mathcal{U}$, $A \subset E_2$, and $x \in E_2$, then

$$\mathfrak{D}_\phi^\nabla(A, x) = \lim_{r \rightarrow 0+} \frac{\phi(A\mathfrak{C}_x^r)}{2r}, \quad \mathfrak{D}_\phi^\Delta(A, x) = \overline{\lim}_{r \rightarrow 0+} \frac{\phi(A\mathfrak{C}_x^r)}{2r},$$

$$\mathfrak{D}_\phi^\Delta(A, x) = \overline{\lim}_{(\text{diam } \Gamma) \rightarrow 0, \Gamma \in \mathfrak{S}} \frac{\phi(A\Gamma)}{\text{diam } \Gamma}$$

where $\mathfrak{S} = E_\Gamma [x \in \Gamma = \mathfrak{C}_y^r \text{ for some } y \in E_2 \text{ and some } r > 0]$.

5.2. REMARK. Let $\phi \in \mathcal{M}$. For each positive integer n , let Δ_n^∇ , Δ_n^Δ and Δ_n^Δ be the functions on E_2 such that $x \in E_2$ implies

$$\Delta_n^\nabla(x) = \inf_{0 < r < 1/n} \frac{\phi(\mathfrak{R}_x^r)}{2r},$$

$$\Delta_n^\Delta(x) = \sup_{0 < r < 1/n} \frac{\phi(\mathfrak{R}_x^r)}{2r},$$

$$\Delta_n^\Delta(x) = \sup_{\Gamma \in \mathfrak{S}, \text{diam } \Gamma < 1/n} \frac{\phi(\Gamma)}{\text{diam } \Gamma}, \quad \text{where}$$

$$\mathfrak{S} = E_\Gamma [\Gamma \text{ is a circle and } x \in \Gamma],$$

and note that the first of these functions is upper semi-continuous while the other two are lower semi-continuous and thus all three are Borel functions.

Also let Δ^∇ , Δ^Δ , and Δ^Δ be the functions on E_2 such that $x \in E_2$ implies

$$\Delta^\nabla(x) = \lim_{n \rightarrow \infty} \Delta_n^\nabla(x), \quad \Delta^\Delta(x) = \lim_{n \rightarrow \infty} \Delta_n^\Delta(x), \quad \Delta^\Delta(x) = \lim_{n \rightarrow \infty} \Delta_n^\Delta(x).$$

We note that these functions are likewise Borel functions and moreover that for $x \in E_2$

$$\Delta^\nabla(x) = \mathfrak{D}_\phi^\nabla(x), \quad \Delta^\Delta(x) = \mathfrak{D}_\phi^\Delta(x) \quad \text{and} \quad \Delta^\Delta(x) = \mathfrak{D}_\phi^\Delta(x).$$

Thus Lemma 5.3 below is established.

The proofs of Lemmas 5.4 through 5.10 involve for the most part merely certain well known uniformization principles. The reader will presumably wish to make use of 3.1(i) in verifying Lemma 5.6. Lemmas 5.7 and 5.8 are corollaries of 5.6. Also leaving to the reader the proofs of Lemmas 5.5 and 5.10, we now suggest proofs for Lemmas 5.4 and 5.9.

If $\epsilon > 0$ and $\mathfrak{D}_\phi^\Delta(x) > 0$ for ϕ almost all x , then for n sufficiently large the bounded Borel set

$$E_x [|x| < n \text{ and } \mathfrak{D}_\phi^\Delta(x) \geq 1/n]$$

not only has the ϕ measure of its complement less than ϵ , but, by Theorem

3.6, contains a closed subset C , necessarily bounded, such that $\phi(\tilde{C}) < \epsilon$. Thus Lemma 5.4 which follows is proved.

We next indicate a proof of Lemma 5.9.

Letting R be the set of all positive rational numbers and checking that

$$E_x [\mathfrak{D}_\phi^\Delta(x) < 1.01 \mathfrak{D}_\phi^\nabla(x)] = \sum_{t \in R} E_x [t < \mathfrak{D}_\phi^\nabla(x) \leq \mathfrak{D}_\phi^\Delta(x) < 1.01t],$$

we choose a positive rational number α such that $\phi(B) > 0$ where B is the Borel set

$$E_x [\alpha < \mathfrak{D}_\phi^\nabla(x) \leq \mathfrak{D}_\phi^\Delta(x) < 1.01\alpha].$$

Accordingly, we select an integer n so large that $\phi(B') > 0$ where B' is the bounded Borel set

$$BE_x [|x| \leq n \text{ and } \alpha < \Delta_n^\nabla(x) \leq \Delta_n^\Delta(x) < 1.01\alpha].$$

Remembering Theorem 3.6, let C be a bounded closed subset of B' with $\phi(C) > 0$, let $\delta = n^{-1}$ and note that

$$\alpha \leq \phi(\mathfrak{R}_x^r)/2r \leq 1.01\alpha < 1.01 \mathfrak{D}_\phi^\nabla(x)$$

for $x \in C$ and $0 < r < \delta$.

With these remarks in mind the reader should have little difficulty in supplying proofs for Lemmas 5.5, 5.6 and 5.10.

5.3. LEMMA. *If $\alpha > 0$, $\phi \in \mathfrak{M}$, then the three sets*

$$E_x [\mathfrak{D}_\phi^\nabla(x) > \alpha], \quad E_x [\mathfrak{D}_\phi^\Delta(x) > \alpha], \quad E_x [\mathfrak{D}_\phi^\Delta(x) > \alpha]$$

are Borel sets.

5.4. LEMMA. *If $\phi \in \mathfrak{M}$, $\epsilon > 0$, and if $\mathfrak{D}_\phi^\Delta(x) > 0$ for ϕ almost all x , then there is a number $\alpha > 0$ and a bounded closed set C with $\phi(\tilde{C}) < \epsilon$ such that $\mathfrak{D}_\phi^\Delta(x) > \alpha$ for $x \in C$.*

5.5. LEMMA. *If $\phi \in \mathfrak{M}$, $\alpha > 0$, $\epsilon > 0$, and if $\mathfrak{D}_\phi^\nabla(x) > \alpha$ for ϕ almost all x , then there is a number $\delta > 0$ and a closed set C with $\phi(\tilde{C}) < \epsilon$ such that*

$$\alpha \leq \phi(\mathfrak{R}_x^r)/2r$$

for $x \in C$ and $0 < r < \delta$.

5.6. LEMMA. *If $\phi \in \mathfrak{M}$, $\mu > 0$, $\epsilon > 0$, $A \subset E_2$, and $\mathfrak{D}_\phi^\Delta(x) < \mu$ for ϕ almost all x in A , then there are positive numbers δ and λ and a bounded closed set C such that:*

$$\lambda < \mu, \quad \phi(A\tilde{C}) < \epsilon;$$

$$\phi(\mathfrak{R}_x^r) \leq 2\lambda r$$

whenever $x \in AC$ and $0 < r < \delta$.

5.7. LEMMA. If $\phi \in \mathfrak{M}$, $\epsilon > 0$, and if $\mathfrak{D}_\phi^\Delta(x) < \infty$ for ϕ almost all x , then there is a finite number $\lambda > 1$ and a bounded closed set C with $\phi(\bar{C}) < \epsilon$ such that

$$\phi(\mathfrak{R}_x^r) \leq 2\lambda r$$

for $x \in C$ and $0 < r < \lambda^{-1}$.

5.8. LEMMA. If $\phi \in \mathfrak{M}$, $\mu > 0$, $\epsilon > 0$, $A \subset E_2$, and $\mathfrak{D}_\phi^\Delta(x) < \mu$ for ϕ almost all $x \in A$, then there is a subset B of A and a $\delta > 0$ such that

$$\phi(B) > \phi(A) - \epsilon$$

and

$$\phi(\mathfrak{R}_x^r) \leq 2\mu r$$

for $x \in B$ and $0 < r < \delta$.

5.9. LEMMA. If $\phi \in \mathfrak{M}$ and $\mathfrak{D}_\phi^\Delta(x) < 1.01 \mathfrak{D}_\phi^\nabla(x)$ for ϕ almost all x , then there are positive numbers α and δ and a bounded closed set C with $\phi(C) > 0$ such that

$$\alpha \leq \frac{\phi(\mathfrak{R}_x^r)}{2r} \leq 1.01\alpha < 1.01 \mathfrak{D}_\phi^\nabla(x)$$

for $x \in C$ and $0 < r < \delta$.

5.10. LEMMA. If $\phi \in \mathfrak{M}$ and $3\mathfrak{D}_\phi^\Delta(x) < 4\mathfrak{D}_\phi^\nabla(x)$ for ϕ almost all x , then there are positive numbers α , δ and η , with $1 < \eta < 3^{-14}$, and a bounded closed set C , with $\phi(C) > 0$, such that

$$\alpha \leq \phi(\mathfrak{R}_x^r)/2r, \quad \phi(\Gamma)/\text{diam } \Gamma \leq \alpha\eta$$

whenever $x \in C$, Γ is a circle with $x \in \Gamma$, $0 < 2r < \delta$, $\text{diam } \Gamma < \delta$.

The next theorem, of which we make considerable use, is well known and follows immediately from Theorem 3.5 of [5]. Referring to 2.25 for the meaning of \frown , we state

5.11. THEOREM. If $M > 0$ and a set A is covered by a family \mathfrak{F} of circles each with diameter less than M , then there is a countable disjointed subfamily \mathfrak{G} of \mathfrak{F} such that

$$A \subset \sum_{\Gamma \in \mathfrak{G}} \hat{\Gamma}.$$

This Theorem 5.11 may also be proved directly, even though A is unbounded, by a method closely patterned after the procedure outlined on p. 109 of [7].

5.12. THEOREM. If $\phi \in \mathfrak{M}$ and A is a Borel set, then $\mathfrak{D}_\phi^\Delta(\bar{A}, x) = 0$ for L almost all x in A .

Proof (by contradiction). Supposing the set

$$AE_x [\mathfrak{D}_\phi^\Delta(\tilde{A}, x) > 0]$$

has positive L measure, we select a positive number η such that

$$(1) \quad L(B) > 0$$

where

$$B = E_x [\mathfrak{D}_\phi^\Delta(\tilde{A}, x) > \eta].$$

Take $0 < \epsilon < 1$, choose a closed subset C of \tilde{A} such that $\phi(\tilde{A}\tilde{C}) < \epsilon$, and let \mathfrak{F} be the family of those circles of the form \mathfrak{R}_x^r where $x \in B$,

$$0 < r < \frac{\epsilon}{10} \frac{\text{dist}(\{x\}, C)}{1 + \text{dist}(\{x\}, C)}, \quad \frac{\phi(\tilde{A}\mathfrak{R}_x^r)}{2r} > \eta.$$

With the help of Lemma 5.11 we pick a countable disjointed subfamily \mathfrak{G} of \mathfrak{F} such that

$$B \subset \sum_{\Gamma \in \mathfrak{G}} \hat{\Gamma}.$$

Noting $\text{diam } \Gamma < \epsilon$ and $\Gamma \subset \tilde{C}$ for $\Gamma \in \mathfrak{G}$ we are assured by 2.6 and the definition of \mathfrak{F} that

$$\begin{aligned} L_\epsilon(B) &\leq \sum_{\Gamma \in \mathfrak{G}} (\text{diam } \hat{\Gamma}) = 5 \sum_{\Gamma \in \mathfrak{G}} (\text{diam } \Gamma) < \frac{5}{\eta} \sum_{\Gamma \in \mathfrak{G}} \phi(\tilde{A}\Gamma) \\ &= \frac{5}{\eta} \phi[\tilde{A}\sigma(\mathfrak{G})] \leq \frac{5}{\eta} \phi(\tilde{A}\tilde{C}) < \frac{5}{\eta} \epsilon. \end{aligned}$$

From this last relation we have $L(B) = 0$ in contradiction to (1). The proof is complete.

If Γ is a circle of diameter d and $x \in \Gamma$ then $\Gamma \subset \mathfrak{R}_x^d$ and we have

5.13. THEOREM. *If $\phi \in \mathfrak{U}$, $A \in E_2$, and $x \in E$, then*

$$\mathfrak{D}_\phi^\Delta(A, x)/2 \leq \mathfrak{D}_\phi^\Delta(A, x) \leq \mathfrak{D}_\phi^\Delta(A, x).$$

5.14. THEOREM. *If $\phi \in \mathfrak{M}$ and A is a Borel set, then, for L almost all x in A ,*

$$\mathfrak{D}_\phi^\nabla(x) = \mathfrak{D}_\phi^\nabla(A, x), \quad \mathfrak{D}_\phi^\Delta(x) = \mathfrak{D}_\phi^\Delta(A, x), \quad \mathfrak{D}_\phi^\Delta(x) = \mathfrak{D}_\phi^\Delta(A, x).$$

Proof. We check that

$$\begin{aligned} \mathfrak{D}_\phi^\nabla(x) &\leq \mathfrak{D}_\phi^\nabla(A, x) + \mathfrak{D}_\phi^\Delta(\tilde{A}, x) \leq \mathfrak{D}_\phi^\nabla(x) + \mathfrak{D}_\phi^\Delta(\tilde{A}, x), \\ \mathfrak{D}_\phi^\Delta(x) &\leq \mathfrak{D}_\phi^\Delta(A, x) + \mathfrak{D}_\phi^\Delta(\tilde{A}, x) \leq \mathfrak{D}_\phi^\Delta(x) + \mathfrak{D}_\phi^\Delta(\tilde{A}, x), \end{aligned}$$

$$\mathfrak{D}_\phi^\Delta(x) \leq \mathfrak{D}_\phi^\Delta(A, x) + \mathfrak{D}_\phi^\Delta(\bar{A}, x) \leq \mathfrak{D}_\phi^\Delta(x) + 2\mathfrak{D}_\phi^\Delta(\bar{A}, x),$$

for $x \in E_2$.

Recourse to Theorem 5.12 completes the proof.

Using intersecting measure we obtain the

5.15. COROLLARY. *If $\phi \in \mathfrak{U}$, $B \subset A \subset E_2$, $\phi(A) < \infty$, and B is a Borel set, then, for L almost all $x \in B$,*

$$\mathfrak{D}_\phi^\nabla(A, x) = \mathfrak{D}_\phi^\nabla(B, x), \quad \mathfrak{D}_\phi^\Delta(A, x) = \mathfrak{D}_\phi^\Delta(B, x), \quad \mathfrak{D}_\phi^\Delta(A, x) = \mathfrak{D}_\phi^\Delta(B, x).$$

5.16. THEOREM. *If $A \subset E_2$ and $L(A) < \infty$, then $\mathfrak{D}_L^\Delta(A, x) \leq 1$ for L almost all $x \in A$.*

Proof (by contradiction). Assuming the set $AE_x[\mathfrak{D}_L^\Delta(A, x) > 1]$ has positive L measure, let $\eta > 1$ be such that

$$(1) \quad L(B) > 0$$

where

$$B = AE_x[\mathfrak{D}_L^\Delta(A, x) > \eta].$$

Let ϵ be a positive number and use Definition 2.6 to ascertain a positive number δ such that both

$$L(A) \leq L_\delta(A) + \epsilon, \quad L(B) \leq L_\delta(B) + \epsilon.$$

Select a family \mathfrak{F} of circles covering B such that $\Gamma \in \mathfrak{F}$ implies

$$(2) \quad \text{diam } \Gamma < \delta/5$$

and

$$L(A\Gamma) > \eta (\text{diam } \Gamma).$$

Relying upon Theorem 5.11 we let \mathfrak{G} be a countable disjointed subfamily of \mathfrak{F} such that

$$B \subset \sum_{\Gamma \in \mathfrak{G}} \hat{\Gamma}$$

and observe from (2) and Definition 2.6 not only that

$$L_\delta(B) \leq \sum_{\Gamma \in \mathfrak{G}} (\text{diam } \hat{\Gamma})$$

but also, upon setting $D = \sigma(\mathfrak{G})$, that

$$L_\delta(AD) \leq \sum_{\Gamma \in \mathfrak{G}} (\text{diam } \Gamma).$$

From Definitions 2.7 and 2.6 follow $0 \leq L(AD) - L_\delta(AD)$, $0 \leq L(A\bar{D}) - L_\delta(A\bar{D})$, and also

$$L_\delta(AD) + L_\delta(A\tilde{D}) \geq L_\delta(A);$$

whereas from the measurability of D follows

$$L(AD) + L(A\tilde{D}) = L(A).$$

Consequently

$$\begin{aligned} 0 \leq L(AD) - L_\delta(AD) &\leq [L(AD) - L_\delta(AD)] + [L(A\tilde{D}) - L_\delta(A\tilde{D})] \\ &\leq L(A) - L_\delta(A) < \epsilon \end{aligned}$$

and

$$L(AD) \leq L_\delta(AD) + \epsilon.$$

Remembering the disjointedness of \mathfrak{G} we have

$$\begin{aligned} \eta \sum_{\Gamma \in \mathfrak{G}} (\text{diam } \Gamma) &\leq \sum_{\Gamma \in \mathfrak{G}} L(A\Gamma) = L[A\sigma(\mathfrak{G})] = L(AD) \\ &\leq L_\delta(AD) + \epsilon \leq \sum_{\Gamma \in \mathfrak{G}} (\text{diam } \Gamma) + \epsilon, \end{aligned}$$

and therefore

$$\sum_{\Gamma \in \mathfrak{G}} (\text{diam } \Gamma) \leq \frac{\epsilon}{\eta - 1}.$$

Recapitulation now reveals

$$L(B) - \epsilon \leq L_\delta(B) \leq \sum_{\Gamma \in \mathfrak{G}} (\text{diam } \hat{\Gamma}) = 5 \sum_{\Gamma \in \mathfrak{G}} \text{diam } \Gamma \leq \frac{5\epsilon}{\eta - 1}$$

and we use the arbitrariness of ϵ to see that $L(B) = 0$ in contradiction to (1). The proof is complete.

5.17. THEOREM. *If $A \subset E_2$ and $L(A) < \infty$ then $\mathfrak{D}_L^\Delta(A, x) \leq 1$ for L almost all x .*

Proof. Relying on 3.4 let B be a Borel set containing A with $L(B)$ finite and note from 5.12, 5.13, and 5.16 that $\mathfrak{D}_L^\Delta(B, x) \leq 1$ for L almost all x . Thus the desired conclusion is reached.

5.18. LEMMA. *If $\phi \in \mathfrak{M}$, $A \subset E_2$, and $\mathfrak{D}_\phi^\Delta(x) > \mu \geq 0$ for L almost all x in A , then $\mu L(A) \leq 5\phi(E_2)$.*

Proof. Let $B = AE_2[\mathfrak{D}_\phi^\Delta(x) > \mu]$. Let δ be a positive number and \mathfrak{F} be a family of circles covering B such that

$$(\text{diam } \Gamma) < \delta/5, \quad \phi(\Gamma) \geq \mu \text{ diam } \Gamma \quad \text{for } \Gamma \in \mathfrak{F}.$$

Relying on Theorem 5.11, let \mathfrak{G} be a disjointed subfamily of \mathfrak{F} such that

$B \subset \sum_{\Gamma \in \mathfrak{G}} \hat{\Gamma}$. We thus have

$$\mu L_\delta(B) \leq \mu \sum_{\Gamma \in \mathfrak{G}} \text{diam } \hat{\Gamma} = 5\mu \sum_{\Gamma \in \mathfrak{G}} \text{diam } \Gamma \leq 5 \sum_{\Gamma \in \mathfrak{G}} \phi(\Gamma) = 5\phi[\sigma(\mathfrak{G})] \leq 5\phi(E_2).$$

Therefore $\mu L(A) = \mu L(B) \leq 5\phi(E_2)$.

5.19. COROLLARY. *If $\phi \in \mathfrak{M}$, then $E_x[\mathfrak{D}_\phi^\Delta(x) = \infty]$ has L measure zero.*

Using intersecting measure and applying Lemma 5.18 we have

5.20. LEMMA. *If $\phi \in \mathfrak{M}$, $A \subset E_2$, and $\mathfrak{D}_\phi^\Delta(A, x) > \mu \geq 0$ for L almost all x in A , then $\mu L(A) \leq 5\phi(A)$.*

5.21. LEMMA. *If $\phi \in \mathfrak{M}$, A is a Borel set, and $\mathfrak{D}_\phi^\Delta(x) > \mu \geq 0$ for L almost all x in A , then $\mu L(A) \leq 5\phi(A)$.*

Proof. Since A is a Borel set we note from 5.14 that $\mathfrak{D}_\phi^\Delta(x) = \mathfrak{D}_\phi^\Delta(A, x)$ for L almost all x in A , and use Lemma 5.20.

5.22. LEMMA. *If $\phi \in \mathfrak{M}$, A is a Borel set, and $\mathfrak{D}_\phi^\Delta(x) > 0$ for L almost all $x \in A$, then $L(B) = 0$ whenever B is a Borel subset of A with $\phi(B) = 0$.*

Proof. Let B be a Borel subset of A with $\phi(B) = 0$. Letting

$$B_n = BE_x[\mathfrak{D}_\phi^\Delta(x) > 1/n]$$

for each positive integer n , we see from Lemma 5.3 that B_n is a Borel set and thus from Lemma 5.21 that $L(B_n) = 0$. Furthermore $B_1 + B_2 + B_3 + \cdots$ is L almost all of B and thus $L(B) = 0$.

5.23. LEMMA. *If $\phi \in \mathfrak{M}$, $A \subset E_2$, and $\mathfrak{D}_\phi^\Delta(x) < \mu$ for ϕ almost all $x \in A$, then $\phi(A) \leq 2\mu L(A)$.*

Proof. Let $\epsilon > 0$. Relying on Lemma 5.8, determine a set $B \subset A$ and a number $\delta > 0$ such that

$$\phi(B) \geq \phi(A) - \epsilon$$

and

$$\phi(\mathfrak{R}_x^r)/2r \leq \mu$$

for $x \in B$ and $0 < r < \delta$.

Also determine a family \mathfrak{F} of open subsets of E_2 such that

$$B \subset \sigma(\mathfrak{F}), \quad L(B) + \epsilon \geq \sum_{s \in \mathfrak{F}} \text{diam } s;$$

$$sB \neq 0, \quad \text{diam } s < \delta/2$$

whenever $s \in \mathfrak{F}$.

For $s \in \mathfrak{F}$ let Γ_s be a circle with center at a point of sB and radius equal to

diam s . We then have the relation

$$\begin{aligned}\phi(A) - \epsilon &\leq \phi(B) \leq \sum_{s \in \mathfrak{F}} \phi(Bs) \leq \sum_{s \in \mathfrak{F}} \phi(\Gamma_s) \leq \mu \sum_{s \in \mathfrak{F}} \text{diam } \Gamma_s \\ &= 2\mu \sum_{s \in \mathfrak{F}} \text{diam } s \leq 2\mu[L(B) + \epsilon] \leq 2\mu[L(A) + \epsilon].\end{aligned}$$

The conclusion $\phi(A) \leq 2\mu L(A)$ follows.

5.24. COROLLARY. If $\phi \in \mathfrak{M}$ and $L(A) < \infty$, then $A E_x[\mathfrak{D}_\phi^\Delta(x) = 0]$ has ϕ measure zero.

5.25. LEMMA. If $\phi \in \mathfrak{M}$, and $\mathfrak{D}_\phi^\Delta(x) < \infty$ for ϕ almost all x , then corresponding to $\epsilon > 0$ there is a $\delta > 0$ such that $\phi(A) < \epsilon$ whenever $L(A) < \delta$.

Proof. Let $\epsilon > 0$. Take $\mu > 0$ so large that $\phi(B) < 2^{-1}\epsilon$ where

$$B = E[\mathfrak{D}_\phi^\Delta(x) \geq \mu].$$

Putting $\delta = \epsilon(4\mu)^{-1}$ we use 5.23 and see: $L(A) < \delta$ implies

$$\begin{aligned}\phi(A) &\leq \phi(AB) + \phi(A\tilde{B}) \leq \phi(B) + \phi(A\tilde{B}) \leq \epsilon/2 + 2\mu L(A\tilde{B}) \\ &\leq \epsilon/2 + 2\mu L(A) < \epsilon/2 + \epsilon/2 = \epsilon.\end{aligned}$$

5.26. COROLLARY. If $\phi \in \mathfrak{M}$, and $\mathfrak{D}_\phi^\Delta(x) < \infty$ for ϕ almost all x , then $\phi(A) = 0$ whenever $L(A) = 0$.

5.27. THEOREM. If $\phi \in \mathfrak{U}$, $\mu > 0$, A is rectifiable, and $\mathfrak{D}_\phi^\nabla(x) < \mu$ for ϕ almost all $x \in A$, then $\phi(A) \leq 10\mu L(A)$.

Proof. Let f be a function on the open interval $E_x[0 < t < 1]$ to E_2 which satisfies a Lipschitz condition and whose range R includes A .

We divide the remainder of the proof into three parts.

PART I. If $\alpha \subset A$ and $\text{diam } \alpha = 0$ then $\phi(\alpha) = 0$.

Proof (by contradiction). Suppose $\phi(\alpha) > 0$.

The set α is nonvacuous and there is a point p such that

$$|\alpha| = \{p\}, \quad \mathfrak{D}_\phi^\nabla(p) = \infty.$$

Consequently

$$\phi\left(A E_x[\mathfrak{D}_\phi^\nabla(x) = \infty]\right) \geq \phi(\alpha) > 0$$

in contradiction to the hypothesis that $\mathfrak{D}_\phi^\nabla(x) < \mu$ for ϕ almost all x in A .

PART II. If $\alpha \subset A$ and $\alpha \subset \beta \subset R$ with β connected, then $\phi(\alpha) \leq 10\mu L(\beta)$.

Proof. If $\text{diam } \beta = 0$ then $\text{diam } \alpha = 0$ and $\phi(\alpha) = 0$ by part I.

We therefore assume $\text{diam } \beta > 0$.

Let $\alpha' = \alpha E_x[\mathfrak{D}_\phi^\nabla(x) < \mu]$ and note that

$$(1) \quad \phi(\alpha') = \phi(\alpha).$$

Let \mathfrak{F} be the family of circles such that $\Gamma \in \mathfrak{F}$ if and only if there is an $x \in \alpha'$ and an r with

$$(2) \quad 0 < r \leq \text{diam } \beta/2$$

such that

$$(3) \quad \Gamma = \mathfrak{R}_x^r \quad \text{and} \quad \phi(\widehat{\Gamma}) < \mu \text{diam } (\widehat{\Gamma}).$$

Clearly \mathfrak{F} covers α' . Furthermore, since β is connected, we have from (2) and Theorem 4.2 that

$$(4) \quad \Gamma \in \mathfrak{F} \text{ implies } (\text{diam } \Gamma) \leq 2L(\beta\Gamma).$$

Now select, according to Theorem 5.11, a countable disjointed subfamily \mathfrak{G} of \mathfrak{F} such that

$$(5) \quad \alpha' \subset \sum_{\Gamma \in \mathfrak{G}} \widehat{\Gamma}.$$

From (1), (5), (3), Abbreviation 2.25, (4) and the disjointedness of \mathfrak{G} we have

$$\begin{aligned} \phi(\alpha) &= \phi(\alpha') \leq \phi\left(\sum_{\Gamma \in \mathfrak{G}} \widehat{\Gamma}\right) \leq \sum_{\Gamma \in \mathfrak{G}} \phi(\widehat{\Gamma}) \leq \sum_{\Gamma \in \mathfrak{G}} \mu(\text{diam } \widehat{\Gamma}) \\ &= 5\mu \sum_{\Gamma \in \mathfrak{G}} \text{diam } \Gamma \leq 10\mu \sum_{\Gamma \in \mathfrak{G}} L(\beta\Gamma) \\ &= 10\mu L[\beta\sigma(\mathfrak{G})] \leq 10\mu L(\beta). \end{aligned}$$

PART III. $\phi(A) \leq 10\mu L(A)$.

Proof. Let $\epsilon > 0$ and, by 3.4, let D be an open set containing A such that

$$(6) \quad L(DR) \leq L(A) + \epsilon.$$

Let \mathfrak{F} be the disjointed family of open intervals such that $\sigma(\mathfrak{F})$ is the open set $E_\epsilon[f(t) \in D]$, and, for $I \in \mathfrak{F}$, let

$$I^* = \sum_{t \in I} \{f(t)\}.$$

Now letting

$$\bar{\mathfrak{F}} = \sum_{I \in \mathfrak{F}} \{I^*\}$$

we note that $\bar{\mathfrak{F}}$ is a countable family of connected Borel subsets of E_2 with

$$DR = \sum_{I \in \mathfrak{F}} \sum_{t \in I} \{f(t)\} = \sum_{I \in \mathfrak{F}} I^* = \sigma(\bar{\mathfrak{F}}).$$

We next select a finite subfamily \mathfrak{G} of $\bar{\mathfrak{F}}$ such that

$$\phi(DR\tilde{C}) < \epsilon$$

where

$$C = \sigma(\mathfrak{G}).$$

Notice that

$$(7) \quad C \subset DR$$

and

$$(8) \quad \phi(A) = \phi(AC) + \phi(A\tilde{C}) = \phi(AC) + \phi(DR\tilde{C}) < \phi(AC) + \epsilon.$$

Let \mathfrak{F} be the family such that $\beta \in \mathfrak{F}$ if and only if β is of the form $\sigma(\mathfrak{Y})$ where: \mathfrak{Y} is a subfamily of \mathfrak{G} ; $\sigma(\mathfrak{Y})$ is connected; and $\sigma(\mathfrak{Y})\sigma(\mathfrak{G}-\mathfrak{Y})=0$. We note that \mathfrak{F} is a finite disjointed family of connected Borel sets.

If it were possible for s to be a member of \mathfrak{G} and to be a subset of no member of \mathfrak{F} we could be led by an iterative procedure to the false proposition that \mathfrak{G} is infinite. Accordingly $C \subset \sigma(\mathfrak{F})$ and, since it is evident that $\sigma(\mathfrak{F}) \subset C$, we conclude that

$$(9) \quad C = \sigma(\mathfrak{F}).$$

Recalling that \mathfrak{F} is a disjointed family of connected Borel sets we use (8), (9), Part II, (7), (6) to obtain

$$\begin{aligned} \phi(A) &\leq \phi(AC) + \epsilon = \phi[A\sigma(\mathfrak{F})] + \epsilon \leq \sum_{\beta \in \mathfrak{F}} \phi(A\beta) + \epsilon \\ &\leq 10\mu \sum_{\beta \in \mathfrak{F}} L(\beta) + \epsilon = 10\mu L[\sigma(\mathfrak{F})] + \epsilon \\ &= 10\mu L(C) + \epsilon \leq 10\mu L(DB) + \epsilon \leq 10\mu [L(A) + \epsilon] + \epsilon. \end{aligned}$$

Therefore $\phi(A) \leq 10\mu L(A)$.

The proof of 5.27 is complete.

5.28. COROLLARY. If $\phi \in \mathfrak{U}$, then $AE_x[\mathfrak{D}_\phi^\nabla(x)=0]$ has ϕ measure zero for each rectifiable set A .

5.29. COROLLARY. If $\phi \in \mathfrak{M}$, $\mathfrak{D}_\phi^\nabla(x) < \infty$ for ϕ almost all x , then $\phi(B)=0$ whenever B is a rectifiable set for which $L(B)=0$.

5.30. THEOREM. If $\phi \in \mathfrak{M}$, $\mu > 0$, $\phi(A) > 0$, $\mathfrak{D}_\phi^\Delta(x) > \mu$ for $x \in A$, and if ψ is the function on E_2 such that $s \subset E_2$ implies $\psi(s)$ is the infimum of numbers of the form $\phi(S)$ where S is an open set containing s , then $\psi \in \mathfrak{M}$ and there is a point $x_0 \in A$ such that

$$\mathfrak{D}_\psi^\Delta(A, x_0) > \mu/7.$$

Proof. A simple check shows that $\psi \in \mathfrak{M}$. In order to prove there is a point $x_0 \in A$ such that

$$\mathfrak{D}_\psi^\Delta(A, x_0) > \mu/7$$

we shall show that the relation,

$$(1) \quad \mathfrak{D}_{\psi}^{\Delta}(A, x) \leq \mu/7 \quad \text{for } x \in A,$$

leads to a contradiction.

Let $\epsilon > 0$, let H be an open set such that $H \subset A$ and

$$\phi(H) \leq \psi(A) + \epsilon,$$

and let \mathfrak{F} be the family of circles Γ such that

$$(2) \quad \Gamma \subset H, \quad 0 < \text{diam } \Gamma < 1, \quad \psi(A\hat{\Gamma}) < 5\phi(\Gamma)/6.$$

In the next paragraph we show that \mathfrak{F} covers A .

Supposing $x \in A$, choose positive numbers r_1, r_2, r_3, \dots such that $r_n \rightarrow 0$ as $n \rightarrow \infty$;

$$0 \leq \lim_{n \rightarrow \infty} 2r_n/\phi(\mathfrak{R}_x^{r_n}) < 1/\mu.$$

With the aid of (1) we find

$$\lim_{n \rightarrow \infty} \frac{\psi(A\mathfrak{R}_x^{r_n})}{\phi(\mathfrak{R}_x^{r_n})} = 5 \lim_{n \rightarrow \infty} \left[\frac{\psi(A\mathfrak{R}_x^{r_n})}{10r_n} \cdot \frac{2r_n}{\phi(\mathfrak{R}_x^{r_n})} \right] \leq 5\mathfrak{D}_{\psi}^{\Delta}(A, x) \cdot \frac{1}{\mu} \leq \frac{5}{7} < \frac{5}{6}.$$

Accordingly for n sufficiently large $\mathfrak{R}_x^{r_n} \in \mathfrak{F}$ and $x \in \mathfrak{R}_x^{r_n} \subset \sigma(\mathfrak{F})$.

Now use Theorem 5.11 to select a countable disjointed subfamily \mathfrak{G} of \mathfrak{F} for which $A \subset \sum_{\Gamma \in \mathfrak{G}} \hat{\Gamma}$. Therefore, remembering (2), we infer the relation

$$\begin{aligned} \psi(A) = \psi\left(\sum_{\Gamma \in \mathfrak{G}} A\hat{\Gamma}\right) &\leq \sum_{\Gamma \in \mathfrak{G}} \psi(A\hat{\Gamma}) < \frac{5}{6} \sum_{\Gamma \in \mathfrak{G}} \phi(\Gamma) = \frac{5}{6} \phi[\sigma(\mathfrak{G})] \\ &\leq \frac{5}{6} \phi(H) \leq \frac{5}{6} [\psi(A) + \epsilon] \end{aligned}$$

from which follows the false proposition

$$0 < \phi(A) \leq \psi(A) \leq \frac{5}{6} \psi(A) < \psi(A).$$

5.31. LEMMA. If $\phi \in \mathfrak{M}$, $A \subset E_2$,

$$\mathfrak{D}_{\phi}^{\Delta}(x) > 0 \quad \text{and} \quad \mathfrak{D}_{\phi}^{\Delta}(A, x) = 0 \quad \text{for } x \in A,$$

then $\phi(A) = 0$.

Proof. Let $\epsilon > 0$, $\eta = \epsilon[\phi(E_2) + 1]^{-1}$ and \mathfrak{F} be the family of circles Γ such that $0 < \text{diam } \Gamma < 1$ and $\phi(A\hat{\Gamma}) < \eta\phi(\Gamma)$. In the next paragraph we show that \mathfrak{F} covers A .

Letting $x \in A$, choose positive numbers r_1, r_2, r_3, \dots , such that: $r_n \rightarrow 0$ as $n \rightarrow \infty$;

$$0 \leq \lim_{n \rightarrow \infty} \frac{2r_n}{\phi(\mathfrak{R}_x^{r_n})} < \infty.$$

We have

$$\lim_{n \rightarrow \infty} \frac{\phi(A \mathfrak{R}_x^{5r_n})}{\phi(\mathfrak{R}_x^{r_n})} = 5 \lim_{n \rightarrow \infty} \left[\frac{\phi(A \mathfrak{R}_x^{5r_n})}{10r_n} \cdot \frac{2r_n}{\phi(\mathfrak{R}_x^{r_n})} \right] = 0.$$

Accordingly for n sufficiently large $\mathfrak{R}_x^{r_n} \in \mathfrak{F}$ and $x \in \mathfrak{R}_x^{r_n} \subset \sigma(\mathfrak{F})$.

Now use Theorem 5.11 to select a countable disjointed subfamily \mathfrak{G} of \mathfrak{F} for which $A \subset \sum_{\Gamma \in \mathfrak{G}} \Gamma$, and note that

$$\phi(A) = \phi\left(\sum_{\Gamma \in \mathfrak{G}} A\Gamma\right) \leq \sum_{\Gamma \in \mathfrak{G}} \phi(A\Gamma) \leq \eta \sum_{\Gamma \in \mathfrak{G}} \phi(\Gamma) = \eta \phi[\sigma(\mathfrak{G})] \leq \eta \phi(E_2) + \epsilon.$$

Thus $\phi(A) = 0$.

5.32. THEOREM. *If $\phi \in \mathfrak{M}$, $A \subset E_2$, and $\mathfrak{D}_\phi^\Delta(x) > 0$ for ϕ almost all $x \in A$, then $\mathfrak{D}_\phi^\Delta(A, x) > 0$ for ϕ almost all $x \in A$.*

Proof. Let

$$C = A \ E_x [\mathfrak{D}_\phi^\Delta(x) > 0] \quad \text{and} \quad B = A \ E_x [\mathfrak{D}_\phi^\Delta(A, x) = 0].$$

Consequently $\mathfrak{D}_\phi^\Delta(BC, x) = 0$ for $x \in BC$, and we use Lemma 5.31 to obtain

$$0 = \phi(BC) = \phi(B).$$

5.33. THEOREM. *If $\phi \in \mathfrak{U}$, $A \subset B$, $\phi(B) < \infty$, and $\mathfrak{D}_\phi^\Delta(B, x) > 0$ for ϕ almost all $x \in A$, then $\mathfrak{D}_\phi^\Delta(A, x) > 0$ for ϕ almost all x in A .*

Proof. Use intersecting measure on B .

6. Rectifiability and directionality. Theorems 6.1, 6.2, and 6.3 which follow are rather immediate consequences of 2.17.

6.1. THEOREM. *Sgn $(A, x) = 0$ if and only if $A \subset \{x\} \subset E_2$.*

6.2. THEOREM. *If $x \in E_2$, $A \subset E_2$, then*

$$\text{dir}(A, x) = \prod_{r \in \mathfrak{R}} \text{Sgn}(AK_r, x)$$

where \mathfrak{R} is any set of positive numbers with 0 as a cluster point.

6.3. THEOREM. *If $x \in E_2$, $A \subset E_2$, then*

$$\text{dir}_\phi(A, x) = \prod_{\beta \in \mathfrak{S}} \text{Sgn}(A\tilde{\beta}, x)$$

where

$$\mathfrak{S} = E_\beta [\beta \subset A, \mathfrak{D}_\phi^\Delta(\beta, x) = 0].$$

Recalling 2.8 we obtain

6.4. THEOREM. *If ϕ is counting measure, $x \in E_2$, and $A \subset E_2$, then*

$$\text{dir}_\phi(A, x) = \text{dir}(A, x).$$

6.5. THEOREM. *If $\phi \in \mathbb{U}$, $A \subset E_2$, $B \subset E_2$, and $x \in E_2$, then*

- (i) $\text{Sgn}(A+B, x) = \text{Sgn}(A, x) + \text{Sgn}(B, x)$,
- (ii) $\text{dir}(A+B, x) = \text{dir}(A, x) + \text{dir}(B, x)$,
- (iii) $\text{dir}_\phi(A+B, x) = \text{dir}_\phi(A, x) + \text{dir}_\phi(B, x)$.

Proof. Since the closure of the union of two sets is the union of the closures of these sets we see that (i) holds. If β_1 and β_2 are sets such that $\mathfrak{D}_\phi^\Delta(\beta_1, x) = \mathfrak{D}_\phi^\Delta(\beta_2, x) = 0$, then $\mathfrak{D}_\phi^\Delta(\beta_1 + \beta_2, x) = 0$; and we see that (iii) holds. Letting ϕ be the counting measure of 2.8 and 3.14, we see that (ii) follows from (iii).

6.6. THEOREM. *If $\phi \in \mathbb{U}$, $x \in A \subset E_2$, and C is a bounded closed set with $C \text{ dir}_\phi(A, x) = 0$, then there is a set $B \subset A$ such that $\mathfrak{D}_\phi^\Delta(B, x) = 0$ and $C \text{ Sgn}(A \tilde{B}, x) = 0$.*

Proof. For $z \in C$: let β_z be such that

$$\beta_z \subset A, \mathfrak{D}_\phi^\Delta(\beta_z, x) = 0; \{z\} \text{ Sgn}(A \tilde{\beta}_z, x) = 0;$$

recall that $\text{Sgn}(A \tilde{\beta}_z, x)$ is closed; take S_z to be an open set of which z is an element and for which $S_z \text{ Sgn}(A \tilde{\beta}_z, x) = 0$. The Heine-Borel Theorem allows us to select a finite subset C' of C such that

$$C \subset \sum_{z \in C'} S_z.$$

Letting

$$B = \sum_{z \in C'} \beta_z,$$

checking that

$$\mathfrak{D}_\phi^\Delta(B, x) \leq \sum_{z \in C'} \mathfrak{D}_\phi^\Delta(\beta_z, x) = 0,$$

we observe that $z \in C'$ implies

$$S_z \text{ Sgn}(A \tilde{B}, x) \subset S_z \text{ Sgn}(A \tilde{\beta}_z, x) = 0.$$

Thus

$$C \text{ Sgn}(A \tilde{B}, x) \subset \sum_{z \in C'} S_z \text{ Sgn}(A \tilde{B}, x) = 0$$

and the proof is complete.

By taking $C = \mathbb{G}_{(0,0)}^1$ in the above theorem we have the

6.7. THEOREM. If $\phi \in \mathcal{U}$ and $A \subset E_2$ then $\text{dir}_\phi(A, x) = 0$ if and only if $\mathfrak{D}_\phi^\Delta(A - \{x\}, x) = 0$.

Letting ϕ be counting measure in Theorem 6.7 we have

6.8. COROLLARY. If $A \subset E_2$ then $\text{dir}(A, x) \neq 0$ if and only if x is a cluster point of A .

Using 6.7 and Theorem 5.12 we have the following

6.9. COROLLARY. If $\phi \in \mathcal{M}$, A is a Borel set, then $\text{dir}_\phi(\tilde{A}, x) = 0$ for L almost all $x \in A$.

From 6.5(iii) and Corollary 6.9 we have the

6.10. THEOREM. If $\phi \in \mathcal{M}$ and A is a Borel subset of E_2 , then $\text{dir}_\phi(A, x) = \text{dir}_\phi(x)$ for L almost all $x \in A$.

Use of intersecting measure leads to the

6.11. COROLLARY. If $\phi \in \mathcal{U}$, $B \subset A \subset E_2$, $\phi(A) < \infty$, and B is a Borel set, then $\text{dir}_\phi(B, x) = \text{dir}_\phi(A, x)$ for L almost all $x \in B$.

6.12. THEOREM. If A is strictly rectifiable, then there is a set $W \subset A$ with $L(A \setminus W) = 0$ such that $x \in W$ implies:

- (i) A is directional at x ;
- (ii) $\mathfrak{D}_L^\nabla(A, x) = \mathfrak{D}_L^\Delta(A, x) = 1$;
- (iii) corresponding to $\xi \in \text{dir}(A, x)$ there is a continuum $\alpha \subset A$ such that $\mathfrak{D}_L^\nabla(\alpha, x) = 1/2$ and $\text{dir}(\alpha, x) = \xi$.

Proof. If $\text{diam } A = 0$ then $L(A) = 0$ and the theorem is trivial.

We assume henceforth that $\text{diam } A > 0$.

Let g be a function on $[0, 1]$ to E_2 which satisfies a Lipschitz condition and whose range is A . Using the process known as parametrizing according to arc length, we ascertain a number $l > 0$ and a function f on $[0, l]$ with range A for which

$$(1) \quad \begin{aligned} |f(t'') - f(t')| &< t'' - t' \quad \text{for } 0 \leq t' \leq t'' \leq l, \\ |f'(t)| &= 1 \quad \text{for Lebesgue almost all } t \text{ in } [0, l]. \end{aligned}$$

We now define

$$T_1 = [0, l] \setminus E_t[|f'(t)| \neq 1] + \{0\} + \{l\},$$

$$B_1 = \sum_{t \in T_1} \{f(t)\},$$

$$B_2 = A E_x[\mathfrak{D}_L^\Delta(A, x) > 1],$$

$$W = A \tilde{B}_1 \tilde{B}_2.$$

Thus $L(B_2)=0$ by Theorem 5.17. From (1) we conclude $L(B_1) \leq$ (Lebesgue measure of T_1) = 0. Therefore

$$L(A\tilde{W}) = 0.$$

We set about showing that W has the additional properties ascribed to it by the theorem. To this end let x be any point in W and select a number τ , with $0 < \tau < l$, for which $x = f(\tau)$. Define $z = f'(\tau)$, note that $|z| = 1$, take $\delta > 0$ such that

$$(2) \quad 0 < |t - \tau| < \delta \quad \text{implies} \quad \left| \frac{f(t) - f(\tau)}{t - \tau} - z \right| < 1/2.$$

Let

$$A_1 = \sum_{t \in [\tau - \delta, \tau]} \{f(t)\}, \quad A_2 = \sum_{t \in [\tau, \tau + \delta]} \{f(t)\}, \quad A_3 = A_1 + A_2.$$

Note that A_1 , A_2 and A_3 are closed connected sets (continua).

We divide the remainder of the proof into 6 parts.

PART I. If $t' < t''$, $\tau - \delta \leq t' \leq t'' \leq \tau + \delta$, then $0 < |f(t'') - f(t')|$.

Proof (by contradiction). Assuming $f(t') = f(t'')$ we infer from (2) that

$$\begin{aligned} 0 < t'' - t' &= |z| (t'' - t') = | (t'' - \tau)z + (\tau - t')z | \\ &\leq | [t'' - \tau]z - [f(t'') - f(\tau)] | + | [\tau - t']z - [f(\tau) - f(t')] | \\ &\leq (t'' - \tau)/2 + (\tau - t')/2 = (t'' - t')/2. \end{aligned}$$

As an immediate consequence of Part I we have

PART II. $A_1 A_2 = \{x\}$, $\text{diam } A_1 > 0$, $\text{diam } A_2 > 0$.

PART III. $\mathfrak{D}_L^\nabla(A_1, x) = 1/2$, $\mathfrak{D}_L^\nabla(A_2, x) = 1/2$,

$$\mathfrak{D}_L^\nabla(A, x) = \mathfrak{D}_L^\Delta(A, x) = \mathfrak{D}_L^\nabla(A_3, x) = \mathfrak{D}_L^\Delta(A_3, x) = 1, \quad \mathfrak{D}_L^\Delta(A\tilde{A}_3) = 0.$$

Proof Since A_1 and A_2 are continua with positive diameter, it follows from Theorem 4.2 that $\mathfrak{D}_L^\nabla(A_1, x) \geq 1/2$ and $\mathfrak{D}_L^\nabla(A_2, x) \geq 1/2$. Hence, since $A_1 + A_2 = A_3$, $L(A_1 A_2) = L(\{x\}) = 0$ and $\mathfrak{D}_L^\Delta(A, x) \leq 1$, we infer

$$\begin{aligned} 1 &\leq \mathfrak{D}_L^\nabla(A_1, x) + \mathfrak{D}_L^\nabla(A_2, x) \leq \mathfrak{D}_L^\nabla(A_1 + A_2, x) = \mathfrak{D}_L^\nabla(A_3, x) \\ &\leq \mathfrak{D}_L^\nabla(A, x) \leq \mathfrak{D}_L^\Delta(A, x) \leq 1. \end{aligned}$$

Furthermore, inasmuch as $\mathfrak{D}_L^\nabla(A_3, x) \leq \mathfrak{D}_L^\Delta(A_3, x) \leq \mathfrak{D}_L^\Delta(A, x)$, we also have $\mathfrak{D}_L^\Delta(A_3, x) = 1$. Using that A_3 is an L measurable (closed) subset of A , we obtain

$$0 \leq \mathfrak{D}_L^\Delta(A\tilde{A}_3, x) \leq \mathfrak{D}_L^\Delta(A, x) - \mathfrak{D}_L^\nabla(A_3, x) = 1 - 1 = 0$$

and thus Part III is proved.

PART IV. $\text{dir } (A_1, x) = \{-z\}$, $\text{dir } (A_2, x) = \{z\}$.

Proof. Let q_1, q_2, q_3, \dots be points distinct from x with $q_n \rightarrow x$ as $n \rightarrow \infty$ and such that either

$$q_n \in A_1 \text{ for } n = 1, 2, 3, \dots \quad \text{or} \quad q_n \in A_2 \text{ for } n = 1, 2, 3, \dots$$

For each positive integer n : let t_n be such that $f(t_n) = q_n$ and $|t_n - \tau| < \delta$; note that (2) helps us infer

$$|q_n - x| = |f(t_n) - f(\tau)| \geq |t_n - \tau|/2.$$

Accordingly $t_n \rightarrow \tau$ as $n \rightarrow \infty$. By letting $\nu = -1$ or $+1$ according as q_1 is or is not an element of A_1 , we see that $\nu(t_n - \tau) > 0$ for each positive integer n , and also see that

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{q_n - x}{|q_n - x|} &= \lim_{n \rightarrow \infty} \frac{f(t_n) - f(\tau)}{|f(t_n) - f(\tau)|} = \lim_{n \rightarrow \infty} \frac{f(t_n) - f(\tau)}{\nu(t_n - \tau)} \cdot \frac{\nu(t_n - \tau)}{|f(t_n) - f(\tau)|} \\ &= \nu z \lim_{n \rightarrow \infty} \left| \frac{t_n - \tau}{f(t_n) - f(\tau)} \right| = \nu \frac{z}{|z|} = \nu z. \end{aligned}$$

PART V. $\text{dir}(A, x) = \{-z\} + \{z\}$ and A is directional at x .

Proof. Let ξ be any point in $\text{dir}(A, x)$. Let p_1, p_2, p_3, \dots be points in A distinct from x with

$$(3) \quad \lim_{n \rightarrow \infty} p_n = x, \quad \lim_{n \rightarrow \infty} \frac{p_n - x}{|p_n - x|} = \xi.$$

For each positive integer n : let

$$(4) \quad \rho_n = \text{dist}(\{p_n\}, A_3)/2, \quad r_n = |p_n - x| + \rho_n;$$

check that

$$\begin{aligned} \rho_n &\leq 2\rho_n \leq |p_n - x|, \quad \mathfrak{R}_{p_n}^{\rho_n} \subset \mathfrak{R}_x^{r_n}, \quad A_3 \mathfrak{R}_{p_n}^{\rho_n} = 0, \\ A\tilde{A}_3 \mathfrak{R}_x^{r_n} &\supset A\tilde{A}_3 \mathfrak{R}_x^{r_n} \mathfrak{R}_{p_n}^{\rho_n} = A\tilde{A}_3 \mathfrak{R}_{p_n}^{\rho_n} = A \mathfrak{R}_{p_n}^{\rho_n}; \end{aligned}$$

observe that Theorem 4.2 implies

$$L(A \mathfrak{R}_{p_n}^{\rho_n}) \geq \rho_n;$$

infer

$$(5) \quad 0 < |p_n - x| \leq r_n \leq 2|p_n - x|;$$

$$(6) \quad \frac{L(A\tilde{A}_3 \mathfrak{R}_x^{r_n})}{2r_n} \geq \frac{L(A \mathfrak{R}_{p_n}^{\rho_n})}{2r_n} \geq \frac{\rho_n}{4|p_n - x|}.$$

Thus inasmuch as $\mathfrak{D}_L^\Delta(A\tilde{A}_3, x) = 0$ by Part III, we see from (5) and (6) that

$$(7) \quad \lim_{n \rightarrow \infty} \frac{\rho_n}{|p_n - x|} = 0.$$

Now, for each positive integer n : recall the definition (4) of ρ_n ; remember that A_3 is closed; choose q_n so that

$$(8) \quad q_n \in A_3, \quad |p_n - q_n| = 2\rho_n \leq |p_n - x|;$$

and note

$$(9) \quad |q_n - x| \geq |p_n - x| - |p_n - q_n| = |p_n - x| - 2\rho_n.$$

With (7) in mind select a positive number N so that $n > N$ implies $2\rho_n < |p_n - x|$ which in turn, by virtue of (9), implies $|q_n - x| > 0$. Consequently $n > N$ implies

$$\begin{aligned} \left| \frac{p_n - x}{|p_n - x|} - \frac{q_n - x}{|q_n - x|} \right| &= \left| \frac{p_n - q_n + q_n - x}{|p_n - x|} - \frac{q_n - x}{|q_n - x|} \right| \\ &\leq \frac{2\rho_n}{|p_n - x|} + \left| \frac{q_n - x}{|p_n - x|} - \frac{q_n - x}{|q_n - x|} \right| \\ &= \frac{2\rho_n}{|p_n - x|} + |q_n - x| \left| \frac{|q_n - x| - |p_n - x|}{|p_n - x||q_n - x|} \right| \\ &\leq \frac{2\rho_n}{|p_n - x|} + \frac{|q_n - p_n|}{|p_n - x|} = \frac{4\rho_n}{|p_n - x|}. \end{aligned}$$

This last relation, (7), and (3) combine to yield

$$\lim_{n \rightarrow \infty} \frac{q_n - x}{|q_n - x|} = \lim_{n \rightarrow \infty} \frac{p_n - x}{|p_n - x|} = \xi$$

and, since (3) and (8) assure us that $q_n \rightarrow x$ with q_n in A_3 as $n \rightarrow \infty$, it is immediate that

$$\xi \in \text{dir}(A_3, x).$$

By our choice of ξ we have now shown

$$\text{dir}(A, x) \subset \text{dir}(A_3, x) \subset \text{dir}(A, x),$$

so that use of 6.5(iii) and Part IV yields

$$\text{dir}(A, x) = \text{dir}(A_3, x) = \text{dir}(A_1, x) + \text{dir}(A_2, x) = \{-z\} + \{z\}.$$

Referring to 2.19 and 2.20 we conclude at once that A is directional at x .

PART VI. If $\xi \in \text{dir}(A, x)$, then there is a continuum $\alpha \subset A$ such that $\mathfrak{D}_L^\nabla(\alpha, x) = 1/2$ and $\text{dir}(\alpha, x) = \xi$.

Proof. From Part V we know $\xi = -z$ or $\xi = z$. Recalling Part III we take $\alpha = A_1$ in the event $\xi = -z$ and take $\alpha = A_2$ in the event $\xi = z$. Since the sets A_1 and A_2 are continua the proof is complete.

6.13. LEMMA. If $\phi \in \mathfrak{M}$, A is strictly rectifiable, $\mu > 0$, $\mathfrak{D}_\phi^\Delta(x) > \mu$ for L al-

most all x in A , and $\mathfrak{D}_\phi^\nabla(x) < \infty$ for ϕ almost all x , then A is ϕ directional at ϕ almost all of its points.

Proof. Identifying the notation of the theorem and that of 6.12 we let W be the set whose existence is assured by 6.12, and let y be any point in W . Since $L(A \tilde{W}) = 0$, Remark 2.15(ii) and Corollary 5.29 enables us to infer $\phi(A \tilde{W}) = 0$. Therefore, since A is directional at y according to 6.12, we know that $\text{dir}(A, x)$ is diametral and that completion of our proof is tantamount to the verification of

$$(1) \quad \text{dir}_\phi(A, y) = \text{dir}(A, y).$$

Select $\xi \in \text{dir}(A, x)$ and let, as in 6.12, α be a continuum contained in A for which

$$\mathfrak{D}_L^\nabla(\alpha, y) \geq 1/2, \quad \text{dir}(\alpha, y) = \{\xi\}.$$

Since $\mathfrak{D}_\phi^\Delta(x) > \mu$ for L almost all x in A , we are assured by 5.21 of the following relation,

$$\frac{\phi(\alpha \mathfrak{R}_y^r)}{2r} \geq \frac{\mu}{5} \frac{L(\alpha \mathfrak{R}_y^r)}{2r} \quad \text{for } r > 0,$$

which convinces us that

$$\mathfrak{D}_\phi^\nabla(\alpha, y) \geq \mu/10 > 0.$$

Thus, since $\phi(\{y\}) = 0$ by 5.29, we have

$$\mathfrak{D}_\phi^\Delta(\alpha - \{y\}, y) = \mathfrak{D}_\phi^\Delta(\alpha, y) \geq \mathfrak{D}_\phi^\nabla(\alpha, y) > 0.$$

Hence from 6.7 it follows that

$$0 \neq \text{dir}_\phi(\alpha, y) \subset \text{dir}(\alpha, y) = \{\xi\}.$$

Accordingly

$$\xi \in \text{dir}_\phi(\alpha, y) \subset \text{dir}_\phi(A, y),$$

and recalling the choice of ξ , it is apparent that

$$\text{dir}(A, y) \subset \text{dir}_\phi(A, y) \subset \text{dir}(A, y).$$

The validity of (1) is established and the proof is complete.

We later obtain as an incidental result that a strictly rectifiable set A is L directional at L almost all points of A . This special result may now be obtained by defining the function ϕ on \mathcal{E}_2 so that $\phi(S) = L(AS)$ for $S \subset E_2$ and applying Lemma 6.13.

6.14. LEMMA. If $\phi \in \mathfrak{M}$, A is a strictly rectifiable set, and $\mathfrak{D}_\phi^\nabla(x) < \infty$ for ϕ almost all x , then $\text{dir}_\phi(x)$ is diametral for ϕ almost all x in A .

Proof. Let $\epsilon > 0$. Since $AE_x[\mathfrak{D}_\phi^\Delta(x) = 0]$ has ϕ measure zero by 5.24, we select such a number μ that

$$(1) \quad 0 < \mu < 1, \quad \phi(A\bar{B}) < \epsilon,$$

where B is the Borel set $AE_x[\mathfrak{D}_\phi^\Delta(x) > \mu]$.

Let ψ be the auxiliary function on \mathcal{E}_2 such that

$$\psi(s) = \phi(Bs) + L(A\bar{B}s) \quad \text{for } s \subset E_2.$$

Check that $\psi \in \mathfrak{M}$.

From 2.15(ii) and 5.29 it is clear that $\phi(Bs) = 0$ whenever $L(s) = 0$. Thus

$$(2) \quad L(s) = 0 \text{ implies } \psi(s) = 0.$$

Since Corollary 5.19 implies the set

$$E_x[\mathfrak{D}_\psi^\Delta(x) = \infty]$$

has L measure zero, it follows from (2) that

$$(3) \quad \mathfrak{D}_\psi^\nabla(x) \leq \mathfrak{D}_\psi^\Delta(x) < \infty \quad \text{for } \psi \text{ almost all } x.$$

Since

$$(4) \quad \psi(\beta) = \phi(\beta) \quad \text{for } \beta \subset B,$$

we use 5.14 to infer

$$(5) \quad \mathfrak{D}_\psi^\Delta(x) = \mathfrak{D}_\psi^\Delta(B, x) = \mathfrak{D}_\phi^\Delta(B, x) = \mathfrak{D}_\phi^\Delta(x) > \mu$$

for L almost all x in B ; on the other hand; using 5.15 and 6.12, we also infer

$$(6) \quad \mathfrak{D}_\psi^\Delta(x) = \mathfrak{D}_\psi^\Delta(\bar{B}, x) = \mathfrak{D}_L^\Delta(A\bar{B}, x) = \mathfrak{D}_L^\Delta(A, x) = 1 > \mu$$

for L almost all $x \in A\bar{B}$. Relations (5) and (6) assure us that

$$(7) \quad \mathfrak{D}_\psi^\Delta(x) > \mu \quad \text{for } L \text{ almost all } x \text{ in } A.$$

We combine (3), (7) and 6.13 and discover that A is ψ directional at ψ almost all of its points. Accordingly

$$(8) \quad \text{dir}_\psi(A, x) \text{ is diametral}$$

for ψ almost all x in A .

From (4) and 6.3 it may be inferred that

$$\text{dir}_\psi(B, x) = \text{dir}_\phi(B, x) \quad \text{for } x \in B.$$

From this and 6.11 it follows that

$$(9) \quad \text{dir}_\psi(A, x) = \text{dir}_\psi(B, x) = \text{dir}_\phi(B, x) = \text{dir}_\phi(x)$$

for L , and hence ψ , almost all x in B . Consequently, in view of (8) and (9), we see that $\text{dir}_\phi(x)$ is diametral for ψ almost all x in B .

Hence $\text{dir}_\phi(x)$ is diametral for ϕ almost all x in B .

Reference to (1) completes the proof.

6.15. THEOREM. *If $\phi \in \mathcal{U}$, A is ϕ rectifiable, $\phi(A) < \infty$, and $\mathcal{D}_\phi^\nabla(A, x) < \infty$ for ϕ almost all x in A , then A is ϕ directional at ϕ almost all of its points.*

Proof (by contradiction). Choose B so that $B \subset A$, $\phi(B) > 0$, and $\text{dir}_\phi(A, x)$ is nondiametral for $x \in B$. Use the ϕ rectifiability of A to obtain a strictly rectifiable set C for which $\phi(BC) > 0$.

Let Φ be the function on \mathcal{E}_2 such that $\Phi(s) = \phi(As)$ for $s \subset E_2$ and note: $\Phi \in \mathcal{M}$; $\mathcal{D}_\Phi^\nabla(x) < \infty$ for Φ almost all x ;

$$(1) \quad \Phi(BC) = \phi(BC) > 0.$$

Since $\Phi(\beta) = \phi(\beta)$ for each $\beta \subset A$ it follows from 6.5(iii) and 6.3 that $x \in E_2$ implies

$$\text{dir}_\Phi(x) = \text{dir}_\phi(A, x) + \text{dir}_\phi(\tilde{A}, x) = \text{dir}_\phi(A, x) = \text{dir}_\phi(A, x).$$

Accordingly $\text{dir}_\phi(x)$ is nondiametral for each $x \in BC$, and 6.14 assures us that $\Phi(BC) = 0$ in contradiction to (1).

6.16. REMARK. In Theorem 6.15 the hypothesis that $\phi(A) < \infty$ may be dispensed with. We neither prove nor use this fact.

6.17 LEMMA. *If $\phi \in \mathcal{M}$, $\mathcal{D}_\phi^\nabla(x) < \infty$ for ϕ almost all x , and A is rectifiable, then there is a set $B \subset A$ such that: $\phi(A\tilde{B}) = 0$; B is directional at each of its points; and*

$$\text{dir}(B, x) = \text{dir}_\phi(B, x) = \text{dir}_\phi(A, x) = \text{dir}(A, x)$$

for each $x \in B$.

Proof. Let C be a strictly rectifiable set containing A and define

$$W_1 = CE_x [C \text{ is directional at } x],$$

$$W_2 = AE_x [A \text{ is } \phi \text{ directional at } x],$$

$$B = W_1 W_2.$$

Note that $B \subset A$ and use 6.12, 5.29, and 6.15 to infer

$$L(A\tilde{W}_1) = 0 = \phi(A\tilde{W}_1), \quad \phi(A\tilde{W}_2) = 0, \quad \phi(A\tilde{B}) = 0.$$

Thus, with the aid of 6.5(iii) and 6.7, verify that $x \in E_2$ implies

$$(1) \quad \begin{aligned} \text{dir}_\phi(A, x) &= \text{dir}_\phi(A\tilde{B}, x) + \text{dir}_\phi(B, x) \\ &= \text{dir}_\phi(B, x) \subset \text{dir}(B, x) \subset \text{dir}(A, x) \subset \text{dir}(C, x). \end{aligned}$$

Since both $\text{dir}_\phi(A, x)$ and $\text{dir}(C, x)$ are diametral for each $x \in B$, we conclude from (1) that

$$\text{dir}(B, x) = \text{dir}_\phi(B, x) = \text{dir}_\phi(A, x) = \text{dir}(A, x)$$

for each $x \in B$, and that B is directional at each of its points.

6.18. THEOREM. *If $\phi \in \mathcal{U}$, $\phi(A) < \infty$, $\mathcal{D}_\phi^\nabla(A, x) < \infty$ for ϕ almost all x in A , and A is ϕ rectifiable, then A contains a subset B such that $\phi(A \setminus B) = 0$ and such that B is directional and ϕ directional at each of its points.*

Proof. Use intersecting measure and 6.3, and apply Lemma 6.17.

6.19. THEOREM. *If $\phi \in \mathcal{U}$, $\phi(A) < \infty$, $\mathcal{D}_\phi^\nabla(A, x) < \infty$ for ϕ almost all x in A , and A is ϕ rectifiable, then corresponding to each positive number ϵ there is a set $B \subset A$ such that $\phi(A \setminus B) < \epsilon$ and such that B is directional at each of its points.*

Proof. Use 6.18 and the definition of ϕ rectifiability.

7. **Restrictedness and rectifiability.** We recall, before giving the next theorem and its proof, that i is the point $(0, 1)$ of E_2 and that for x in E_2 the numbers x_1 and x_2 are such that $x = (x_1, x_2)$.

7.1. THEOREM. *If $\eta > 0$, A is a bounded subset of E_2 , and $\mathcal{R}_i^\eta \text{Sgn}(A, x)$ is vacuous whenever $x \in A$, then A is rectifiable.*

Proof. The theorem being obvious in case there is a $p \in E_2$ such that $A \subset \{p\}$, we assume that A contains at least two points.

Let p' and p'' be distinct elements of A and set $P = p'' - p'$. Since $\mathcal{R}_i^\eta \text{Sgn}(A, x)$ is vacuous for x in A we know that

$$|i - P|/|P| \geq \eta,$$

from which follow successively the relations

$$\begin{aligned} 1 - P_2/|P| + |P_1|/|P| &\geq \eta, \\ P_2 &\leq |P_1| + (1 - \eta)|P| \leq |P_1| + (1 - \eta)(|P_1| + |P_2|), \\ P_2 - (1 - \eta)|P_2| &\leq 2|P_1|. \end{aligned}$$

Thus if $P_2 \geq 0$ we conclude from the last relation that

$$P_2 \leq 2\eta^{-1}|P_1|,$$

an inequality which is obvious if $P_2 < 0$. Accordingly

$$p_2'' - p_2' \leq 2\eta^{-1}|p_1'' - p_1'|$$

and we have shown

$$x_2'' - x_2' \leq 2\eta^{-1}|x_1'' - x_1'| \text{ for } x' \in A \text{ and } x'' \in A.$$

This implies

$$|x_2'' - x_2'| \leq 2\eta^{-1} |x_1'' - x_1'| \quad \text{for } x' \text{ and } x'' \text{ in } A$$

which in turn implies

$$(1) \quad |x_2'' - x_2'| \leq 2\eta^{-1} |x_1'' - x_1'| \quad \text{for } x' \text{ and } x'' \text{ in clsr } A.$$

Letting

$$\bar{A} = \text{clsr } A, \quad C = \sum_{x \in \bar{A}} \{x_1\},$$

we note that C is a bounded closed subset of E_1 and that (1) assures us that \bar{A} is the graph of a function (or, if we prefer, is a function) with domain C . Denoting this function by h we observe

$$(2) \quad \sum_{t \in C} \{(t, h(t))\} = \bar{A}.$$

Taking J as the closed interval containing C with end points in C we extend the function h to the open intervals comprising $J - C$ by linear interpolation to obtain a function g on J which, because of (1), satisfies the Lipschitz condition

$$|g(t'') - g(t')| \leq 2\eta^{-1} |t'' - t'| \quad \text{for } t' \in J \text{ and } t'' \in J.$$

Finally we define f on J by

$$f(t) = (t, g(t)) \quad \text{for } t \in J.$$

Clearly f satisfies the Lipschitz condition

$$|f(t'') - f(t')| \leq (1 + 2\eta^{-1}) |t'' - t'| \quad \text{for } t' \in J \text{ and } t'' \in J.$$

By (2) the range of f includes \bar{A} which in turn includes A .

7.2. THEOREM. *If $\eta > 0$, A is a bounded subset of E_2 , $z \in E_2$ with $|z| = 1$, and if $\mathcal{R}_z^* \text{Sgn}(A, x)$ is vacuous whenever $x \in A$, then A is rectifiable.*

The proof is obvious and may be effected by the rotational device of letting

$$\bar{S} = \sum_{x \in S} \{-izx\},$$

selecting A_0 so that $\bar{A}_0 = A$, and checking that A is congruent to the set A_0 which is rectifiable by 7.1.

This scheme is employed in more detail in the proof of 7.19 where the argument is slightly more complicated due to the presence of a measure.

7.3. THEOREM. *If $\eta > 0$, $r > 0$, A is a bounded subset of E_2 , $z \in E_2$ with $|z| = 1$, and if $\mathcal{R}_z^* \text{Sgn}(A \mathcal{R}_z', x)$ is vacuous whenever $x \in A$, then A is rectifiable.*

Proof. Let \mathfrak{F} be a finite family of sets such that $A \subset \sigma(\mathfrak{F}) \subset E_2$ and $\text{diam } s < 2^{-1r}$ whenever $s \in \mathfrak{F}$. Clearly if $s \in \mathfrak{F}$ and $x \in s$ then $As \subset A\mathfrak{R}_x^r$ and $\text{Sgn}(As, x) \subset \text{Sgn}(A\mathfrak{R}_x^r, x)$. Thus by 7.2, $s \in \mathfrak{F}$ implies As is rectifiable and therefore, \mathfrak{F} being finite, $\sum_{s \in \mathfrak{F}} (As)$ is rectifiable.

7.4. THEOREM. *If A is restricted at each of its points, then there is a countable family \mathfrak{F} of rectifiable sets for which*

$$A = \sigma(\mathfrak{F}).$$

Proof. Recall 2.22. Let $\Gamma = E_2 E_x [|z| = 1]$. Let B be such a countable subset of Γ that its closure is Γ . Let R be the set of positive rationals less than 1. Let N be the set of positive integers. Let $A_n = \mathfrak{R}_{(0,0)}^n A$.

Checking that

$$A = \sum_{s \in B} \sum_{r \in R} \sum_{r \in R} \sum_{n \in N} \omega(n, r, \eta, z)$$

where

$$\omega(n, r, \eta, z) = A_n E_x [\mathfrak{R}_x^r \text{Sgn}(A_n \mathfrak{R}_x^r, x) = 0],$$

we see in the light of 7.3 that \mathfrak{F} may be taken as the countable family

$$\sum_{s \in B} \sum_{r \in R} \sum_{r \in R} \sum_{n \in N} \{\omega(n, r, \eta, z)\}.$$

7.5. DEFINITION. For a and λ positive numbers and $x \in E_2$ we define

$$\langle x, a, \lambda \rangle = E_y [|y_1 - x_1| < a \text{ and } |y_2 - x_2| < \lambda a].$$

7.6. LEMMA. *If $x \in E_2$, $0 < \lambda$, $0 < \eta$, $0 < a$, $\lambda\eta = 40$, and if x' and x'' are elements of $\langle x, 5a, \lambda \rangle$ for which $x_2' - x_2'' = \lambda a$, then there are sets α' and α'' such that*

$$\begin{aligned} \alpha' \{x'\} + \alpha'' \{x''\} &= 0, & \alpha' + \alpha'' &= \langle x, 5a, \lambda \rangle, \\ \text{Sgn}(\alpha', x') &\subset \mathfrak{R}_i^r, & \text{Sgn}(\alpha'', x'') &\subset \mathfrak{R}_{-i}^r. \end{aligned}$$

Proof. Consider the sets

$$\alpha' = E_y \left[|y_1 - x_1| < 5a \text{ and } \frac{x_2' + x_2''}{2} \leq y_2 < x_2 + 5\lambda a \right],$$

$$\alpha'' = E_y \left[|y_1 - x_1| < 5a \text{ and } x_2 - 5\lambda a < y_2 \leq \frac{x_2' + x_2''}{2} \right].$$

Clearly $\alpha' \{x'\} = 0$, $\alpha'' \{x''\} = 0$ and $\alpha' + \alpha'' = \langle x, 5a, \lambda \rangle$. We shall now prove

$$(1) \quad \text{Sgn}(\alpha', x') \subset \mathfrak{R}_i^r.$$

Let $p' \in \alpha'$, denote $p' - x'$ by ξ' and note the relations

$$|\xi'| \geq \xi'_2 = p'_2 - x'_2 \geq \frac{x'_2 + x'_{2'}}{2} - x'_2 = \frac{x'_{2'} - x'_2}{2} = \frac{\lambda a}{2} > 0$$

and

$$|\xi'_1| < 10a,$$

which together imply

$$\begin{aligned} \left| i - \frac{p' - x'}{|p' - x'|} \right| &= \left| i - \frac{\xi'}{|\xi'|} \right| \leq 1 - \frac{\xi'_2}{|\xi'|} + \frac{|\xi'_1|}{|\xi'|} \\ (2) \quad &= \frac{|\xi'| - \xi'_2 + |\xi'_1|}{|\xi'|} \leq \frac{|\xi_1| + \xi'_2 - \xi'_2 + |\xi'_1|}{|\xi'|} \\ &= \frac{2|\xi'_1|}{|\xi'|} \leq \frac{4|\xi'_1|}{\lambda a} < \frac{40a}{\lambda a} = \frac{40}{\lambda} = \eta. \end{aligned}$$

Thus (1) is established.

Similarly taking $p'' \in \alpha''$ and denoting $x'' - p''$ by ξ'' we note the relations

$$|\xi''| \geq \lambda a/2 > 0 \quad \text{and} \quad |\xi''_1| < 10a$$

which together imply

$$\left| -i - \frac{p'' - x''}{|p'' - x''|} \right| = \left| i - \frac{\xi''}{|\xi''|} \right| < \eta.$$

We conclude

$$\text{Sgn}(\alpha'', x'') \subset \mathfrak{R}_{-i}^{\eta}.$$

As a corollary we have

7.7. LEMMA. If $x \in E_2$, $0 < \lambda$, $0 < \eta$, $0 < a$, $\lambda\eta = 40$, $y \in \langle x, 5a, \lambda \rangle$, $|y_2 - x_2| = \lambda a$, then there are sets β' and β'' such that

$$\beta'\{x\} + \beta''\{y\} = 0, \quad \beta' + \beta'' = \langle x, 5a, \lambda \rangle,$$

$$\text{Sgn}(\beta', x) + \text{Sgn}(\beta'', y) \subset \mathfrak{R}_{-i}^{\eta} + \mathfrak{R}_i^{\eta}.$$

7.8. LEMMA. If $x \in A \subset E_2$, $0 < \eta < 1$, $\lambda\eta = 40$, $\omega = 10^{-2}$, and $\text{Sgn}(A, x) \mathfrak{R}_i^{\omega\eta}$ is nonvacuous, then there is a number $b > 0$ such that

$$0 \neq A E_y [(|y_1 - x_1| < b) \text{ and } (|y_2 - x_2| = \lambda b)].$$

Proof. Since $\text{Sgn}(A, x) \mathfrak{R}_i^{\omega\eta}$ is nonvacuous we select a point $p \in A$ such that

$$\left| \frac{p - x}{|p - x|} - i \right| < \omega\eta < 10^{-2},$$

let $\xi = p - x$, and check in order:

$$\begin{aligned} \left| \frac{\xi}{|\xi|} - i \right| &= \left| \frac{\xi_1}{|\xi|} + i \left(\frac{\xi_2}{|\xi|} - 1 \right) \right| < \omega\eta < 10^{-2}, \\ \left| \frac{\xi_2}{|\xi|} - 1 \right| &< 10^{-2}, \quad |\xi_2| > 0, \\ |\xi_1| < \omega\eta |\xi| &\leq \omega\eta |\xi_1| + \omega\eta |\xi_2| \leq \frac{|\xi_1|}{2} + \omega\eta |\xi_2|, \\ |\xi_1| &\leq 2\omega\eta |\xi_2| = \frac{4}{5} \frac{|\xi_2|}{\lambda} < \frac{|\xi_2|}{\lambda}. \end{aligned}$$

Accordingly

$$|p_1 - x_1| < \frac{|p_2 - x_2|}{\lambda}.$$

We now define b so that $\lambda b = |p_2 - x_2|$, note that $|p_1 - x_1| < b$, and conclude

$$AE_y [(|y_1 - x_1| < b) \text{ and } (|y_2 - x_2| = \lambda b)]$$

is nonvacuous (in fact p is in it).

7.9. DEFINITION. We say A is *countably ϕ rectifiable* if and only if $\phi \in \mathcal{U}$ and there is a countable family \mathfrak{F} of rectifiable sets such that

$$\phi[A - \sigma(\mathfrak{F})] = 0.$$

It should be noted that A is countably ϕ rectifiable whenever it is ϕ rectifiable. Furthermore if $\phi(A) < \infty$ then A is countably ϕ rectifiable when and only when A is ϕ rectifiable. It is likewise obvious that $\sigma(\mathcal{U})$ is countably ϕ rectifiable whenever \mathcal{U} is a countable family of countably ϕ rectifiable sets.

7.10. NOTATION. Whenever $\phi \in \mathcal{U}$ the notation ϕ^0 will be used to denote that function ψ on \mathcal{E}_2 such that $s \subset E_2$ implies $\psi(s)$ is the infimum of numbers of the form $\phi(S)$ where S is an open set containing s .

Thus from 3.12 it follows that $\phi^0 \in \mathcal{U}$ or $\phi^0 \in \mathcal{M}$ according as $\phi \in \mathcal{U}$ or $\phi \in \mathcal{M}$.

7.11. THEOREM. If $\phi \in \mathcal{U}$, $\epsilon > 0$, $0 < \eta < 1$, $\lambda\eta = 40$, $\delta > 0$, $0 < \rho \leq 40^{-1}\delta$, $\xi \in A \subset E_2$, and if corresponding to each $x \in A$ there is a set B such that

- (i) $(\mathcal{R}_x + \mathcal{R}_x^r) \text{Sgn}(\bar{B}, x) = 0$,
- (ii) $\phi(B\mathcal{R}_x^r) \leq 2r\epsilon$ for $0 < r \leq \delta$,

then there is a rectifiable set D such that

$$\phi^0(A\tilde{D}\mathcal{R}_\xi^0) \leq 240\lambda\rho\epsilon.$$

Proof. Let

$$(1) \quad K = A\mathcal{R}_\xi^0,$$

- (2) $\omega = 10^{-2},$
 (3) $D = KE \left[\mathfrak{R}_i^{\omega} \text{Sgn} (K, x) = 0 \right],$
 (4) $C = K\tilde{D}.$

From 7.1 it follows that D is rectifiable.

Notice that

$$(5) \quad D + C = K,$$

and that verification (in Part IV below) of the relation

$$\phi^0(C) \leq 240\lambda\rho\epsilon$$

will complete the proof.

For $x \in C$ we define a numerical set T_x as follows: $b \in T_x$ if and only if $b > 0$ and

$$(6) \quad 0 \neq KE \left[(|y_1 - x_1| < b) \text{ and } (|y_2 - x_2| = \lambda b) \right].$$

Accordingly for $x \in C$: we use (2), (3), (4) and Lemma 7.8 to check that $T_x \neq 0$; with the aid of (6) and (1) we infer

$$(7) \quad \lambda \sup T_x \leq \text{diam } K \leq 2\rho < \infty;$$

we select a number $a_x \in T_x$ such that

$$(8) \quad 0 < a_x \leq \sup T_x \leq 2a_x < \infty.$$

Letting

$$\mathfrak{F} = \sum_{x \in C} \{E_i[|t - x_1| < a_x]\}, \quad X = \sum_{x \in C} \{x_1\},$$

observing that $X \subset E_1$ and that X is covered by the family \mathfrak{F} of open intervals, we use the one-dimensional analogue of 5.11 to obtain a countable subset C_0 of C for which

$$(9) \quad \sum_{x \in C_0} \{E_i[|t - x_1| < a_x]\} \text{ is disjointed}$$

and

$$(10) \quad X \subset \sum_{x \in C_0} E_i[|t - x_1| < 5a_x].$$

We divide the remainder of the proof into four parts.

PART I.

$$\sum_{x \in C_0} 2a_x \leq 6\rho.$$

Proof. With the aid of (4) and (1) we obtain

$$\text{diam } X \leq \text{diam } C \leq \text{diam } K \leq 2\rho$$

so that we may use (8) and (7) to infer

$$\text{diam} \sum_{x \in C_0} E_t [|t - x_1| < a_x] \leq 2\rho + \text{diam } X + 2\rho \leq 6\rho.$$

Recourse to (9) completes this proof.

PART II.

$$C \subset \sum_{x \in C_0} \langle x, 5a_x, \lambda \rangle.$$

Proof (by contradiction). Suppose y is in C and

$$(11) \quad y \notin \sum_{x \in C_0} \langle x, 5a_x, \lambda \rangle,$$

note that $y_1 \in X$ and use (10) to secure an x' in C for which

$$(12) \quad |y_1 - x'_1| < 5a_{x'}.$$

Now the statement, $y \in \langle x, 5a_{x'}, \lambda \rangle$, which is a consequence of (11), combines with (12) to yield

$$5\lambda a_{x'} \leq |y_2 - x'_2|.$$

Determining b so that $\lambda b = |y_2 - x'_2|$ we have

$$(13) \quad 5\lambda a_{x'} \leq |y_2 - x'_2| = \lambda b.$$

Thus (13) implies the relation

$$5a_{x'} \leq b < 5b$$

which combines with (12) to tell us

$$(14) \quad |y_1 - x'_1| < b.$$

From the relations $y \in C \subset K$, (14), (13), (6) follows the proposition

$$b \in T_{x'}$$

which, viewed in the light of (8) and (13), implies the contradiction

$$b \leq \sup T_{x'} \leq 2a_{x'} < 5a_{x'} \leq b.$$

PART III. If $x \in C_0$ then

$$\phi(\langle x, 5a_x, \lambda \rangle) \leq 80\lambda a_x \epsilon.$$

Proof. Since $a_x \in T_x$ we use (6) to ascertain a point $y \in K$ with

$$|y_1 - x_1| < a_x, \quad |y_2 - x_2| = \lambda a_x.$$

Noting that x and y are in $\langle x, 5a_x, \lambda \rangle$ with $|y_2 - x_2| = \lambda a_x$, and invoking Lemma 7.7, we choose sets β' and β'' for which

$$\beta'\{x\} + \beta''\{y\} = 0, \quad \beta' + \beta'' = \langle x, 5a_x, \lambda \rangle,$$

$$\text{Sgn}(\beta', x) + \text{Sgn}(\beta'', y) \subset \mathfrak{R}_{-i}^\eta + \mathfrak{R}_i^\eta.$$

Putting $d = \text{diam } \langle x, 5a_x, \lambda \rangle$, and checking that $1 \leq \lambda < \infty$, we use (8) and (7) to obtain

$$(16) \quad d \leq 10a_x + 10\lambda a_x \leq 20\lambda a_x \leq 40\rho \leq \delta.$$

Because of (16), (ii), (i), and the fact that x and y are in $K \subset A$, we can and do select sets B_1 and B_2 such that

$$(17) \quad \phi(B_1 \mathfrak{R}_x^d) \leq 2d\epsilon, \quad \phi(B_2 \mathfrak{R}_y^d) \leq 2d\epsilon,$$

and

$$(\mathfrak{R}_{-i}^\eta + \mathfrak{R}_i^\eta)[\text{Sgn}(\bar{B}_1, x) + \text{Sgn}(\bar{B}_2, y)] = 0.$$

Referring to (15) and the last relation we deduce

$$\text{Sgn}(\beta' \bar{B}_1, x) + \text{Sgn}(\beta'' \bar{B}_2, y) \subset (\mathfrak{R}_{-i}^\eta + \mathfrak{R}_i^\eta)[\text{Sgn}(\bar{B}_1, x) + \text{Sgn}(\bar{B}_2, y)] = 0,$$

and thus, from 6.1 and (15), conclude $\beta' \bar{B}_1 + \beta'' \bar{B}_2 \subset \beta'\{x\} + \beta''\{y\} = 0$. Accordingly the relation

$$\begin{aligned} \langle x, 5a_x, \lambda \rangle &= \beta' + \beta'' = \beta' \bar{B}_1 + \beta' B_1 + \beta'' \bar{B}_2 + \beta'' B_2 \\ &= \beta' B_1 + \beta'' B_2 \subset B_1 \langle x, 5a_x, \lambda \rangle + B_2 \langle x, 5a_x, \lambda \rangle \\ &\subset B_1 \mathfrak{R}_x^d + B_2 \mathfrak{R}_y^d \end{aligned}$$

may be combined with (17) and (16) to give

$$\phi(\langle x, 5a_x, \lambda \rangle) \leq \phi(B_1 \mathfrak{R}_x^d) + \phi(B_2 \mathfrak{R}_y^d) \leq 4d\epsilon \leq 80\lambda a_x \epsilon.$$

PART IV. $\phi^0(C) \leq 240\lambda\rho\epsilon$.

Proof. Using the fact that the set

$$\sum_{x \in C_0} \langle x, 5a_x, \lambda \rangle$$

is open and also 7.10, Parts II, III, and I, we discover

$$\begin{aligned} \phi^0(C) &\leq \phi\left(\sum_{x \in C_0} \langle x, 5a_x, \lambda \rangle\right) \leq \sum_{x \in C_0} \phi(\langle x, 5a_x, \lambda \rangle) \\ &\leq \sum_{x \in C_0} 80\lambda a_x \epsilon = 40\lambda \epsilon \sum_{x \in C_0} 2a_x \leq 240\lambda\rho\epsilon. \end{aligned}$$

7.12. THEOREM. If $\phi \in \mathfrak{U}$, $0 < \eta < 1$, $\delta > 0$, A is a bounded subset of E_2 , and if corresponding to each $x \in A$ there is a set B such that

$$(i) (\mathfrak{R}_{-i}^{\eta} + \mathfrak{R}_i^{\eta}) \operatorname{Sgn} (\bar{B}, x) = 0,$$

$$(ii) \phi(B\mathfrak{R}_x^r) \leq 2r \text{ for } 0 < r \leq \delta,$$

then there is a rectifiable set C such that $\phi^0(A\bar{C}) < \infty$.

Proof. Let \mathfrak{F} be a finite family of nonvacuous sets each with diameter less than $40^{-1}\delta$ such that $A = \sigma(\mathfrak{F})$. Letting $\lambda\eta = 40$, $\rho = 40^{-1}\delta$, $\epsilon = 1$, we use 7.11 to associate with each $s \in \mathfrak{F}$ a point $x_s \in s$ and a rectifiable set D_s such that

$$\phi^0(s\bar{D}_s) = \phi^0(s\bar{D}_s\mathfrak{R}_{x_s}^{\rho}) \leq 240\lambda\rho\epsilon < \infty.$$

Accordingly the rectifiable set

$$\sum_{s \in \mathfrak{F}} D_s$$

has the properties demanded of C by the theorem.

7.13. THEOREM. If $\phi \in \mathfrak{M}$, $\phi(A) > 0$,

$$\mathfrak{D}_{\phi}^{\Delta}(x) > 0 \quad \text{and} \quad (\{-i\} + \{i\}) \operatorname{dir}_{\phi}(x) = 0$$

for ϕ almost all $x \in A$, then there is a rectifiable set with positive ϕ measure.

Proof. Ascertain $\mu > 0$ so that $\phi(A_1) > 0$ where

$$A_1 = AE_x [\mathfrak{D}_{\phi}^{\Delta}(x) > \mu].$$

Remember that $\operatorname{Sgn}(s, x)$ is closed for $s \subset E_2$ and $x \in E_2$, and select a positive number $\eta < 1$ such that $\phi(A_2) > 0$ where A_2 is the set for which $x \in A_2$ if and only if there is a set B such that

$$x \in A_1, \quad (\mathfrak{R}_{-i}^{\eta} + \mathfrak{R}_i^{\eta}) \operatorname{Sgn} (\bar{B}, x) = 0, \quad \mathfrak{D}_{\phi}^{\Delta}(B, x) = 0.$$

Choose λ and ϵ so that

$$\lambda\eta = 40,$$

$$(1) \quad 240\lambda\epsilon = \mu/10.$$

Take $\delta > 0$ so that $\phi(A_3) > 0$ where A_3 is the set for which $x \in A_3$ if and only if there is a set B such that

$$x \in A_2, \quad (\mathfrak{R}_{-i}^{\eta} + \mathfrak{R}_i^{\eta}) \operatorname{Sgn} (\bar{B}, x) = 0,$$

$$\phi(B\mathfrak{R}_x^r) \leq 2r\epsilon \quad \text{for } 0 < r \leq \delta.$$

Noting $A_3 \subset A_1$ with $\phi(A_3) > 0$, we infer that $\mathfrak{D}_{\phi}^{\Delta}(x) > \mu$ for $x \in A_3$, use 7.10 and 5.30 to obtain a point ξ in A_3 for which $\mathfrak{D}_{\phi}^{\Delta}(A_3, \xi) > 7^{-1}\mu$ and fix a number ρ such that

$$0 < \rho \leq \delta/40,$$

$$(2) \quad \phi^0(A_3\mathfrak{R}_{\xi}^{\rho}) > 2\mu\rho/7 > \mu\rho/7.$$

With the aid of 7.11 we so secure a rectifiable set D that

$$\phi^0(A_3 \tilde{D} \mathfrak{R}_\epsilon^0) \leq 240\lambda\rho\epsilon.$$

Viewing this last relation in the light of (2) and (1) we discover

$$(3) \quad \begin{aligned} \phi^0(D) &\geq \phi^0(A_3 D \mathfrak{R}_\epsilon^0) = \phi^0(A_3 \mathfrak{R}_\epsilon^0) - \phi^0(A_3 \tilde{D} \mathfrak{R}_\epsilon^0) \\ &\geq \mu\rho/7 - 240\lambda\rho\epsilon > \mu\rho/10 - \mu\rho/10 = 0. \end{aligned}$$

Now let $K = \text{clsr } D$. Since D is rectifiable, K is likewise. From 3.12 and (3) follows the relation

$$\phi(K) = \phi^0(K) \geq \phi^0(D) > 0.$$

Consequently there is a rectifiable set, namely K , with positive ϕ measure. The proof is complete.

7.14. THEOREM. If $\phi \in \mathfrak{M}$, $\phi(A) > 0$,

$$\mathfrak{D}_\phi^\Delta(x) > 0 \quad \text{and} \quad (\{-i\} + \{i\}) \text{dir}_\phi(x) = 0$$

for ϕ almost all $x \in A$, then there is a rectifiable subset of A with positive ϕ measure.

Proof. Let $\Phi \in \mathfrak{M}$ be the function on E_2 such that $s \subset E_2$ implies $\Phi(s) = \phi(As)$. Since $\mathfrak{D}_\Phi^\Delta(x) = \mathfrak{D}_\phi^\Delta(A, x)$ for $x \in E_2$ we infer from 5.32 that $\mathfrak{D}_\Phi^\Delta(x) > 0$ for ϕ almost all x in A . Inasmuch as the relation

$$\text{dir}_\Phi(x) \subset \text{dir}_\phi(x) \quad \text{for } x \in E_2$$

follows immediately from Definition 2.17, it is evident that

$$(\{-i\} + \{i\}) \text{dir}_\Phi(x) = 0$$

for ϕ almost all $x \in A$. Consequently

$$\mathfrak{D}_\Phi^\Delta(x) > 0, \quad (\{-i\} + \{i\}) \text{dir}_\Phi(x) = 0$$

not only for ϕ almost all x in A but for Φ almost all x .

Thus 7.13 guarantees us a rectifiable set C with $\Phi(C) > 0$ and we obtain the rectifiable set AC with $\phi(AC) = \Phi(C) > 0$.

7.15. LEMMA. If $\phi \in \mathfrak{M}$ and if every Borel set with positive ϕ measure contains a rectifiable set with positive ϕ measure, then E_2 is ϕ rectifiable.

Proof. Let $\epsilon > 0$ and \mathfrak{F} be the family of all rectifiable sets. Denote

$$\sup_{\alpha \in \mathfrak{F}} \phi(\alpha)$$

by M and let $C_1 \subset C_2 \subset C_3 \subset \dots$ be closed rectifiable sets for which $\phi(C_n) \rightarrow M$ as $n \rightarrow \infty$. Setting

$$C = C_1 + C_2 + C_3 + \dots$$

we observe that C is a Borel set and that, whether $\phi(\tilde{C}) > 0$ or $\phi(\tilde{C}) = 0$, our hypotheses enable us to select a number $\mu > 0$ and a rectifiable subset A_0 of \tilde{C} such that $\phi(A_0) = \mu\phi(\tilde{C})$. Since $C_n + A_0$ is then rectifiable with $C_n A_0 \subset C A_0 \subset C\tilde{C} = 0$ for each positive integer n , we have

$$M + \mu\phi(\tilde{C}) = M + \phi(A_0) = \lim_{n \rightarrow \infty} [\phi(C_n) + \phi(A_0)] = \lim_{n \rightarrow \infty} \phi(C_n + A_0) \leq M,$$

from which it follows that

$$\mu\phi(\tilde{C}) = 0, \quad \phi(\tilde{C}) = 0.$$

Inasmuch as $\phi(\tilde{C}_n) \rightarrow \phi(\tilde{C}) = 0$ as $n \rightarrow \infty$ we know that $\phi(\tilde{C}_n) < \epsilon$ for n sufficiently large. Accordingly E_2 is ϕ rectifiable.

7.16. THEOREM. If $\phi \in \mathfrak{M}$, $A \subset E_2$,

$$\mathfrak{D}_\phi^\Delta(x) > 0 \quad \text{and} \quad (\{-i\} + \{i\}) \operatorname{dir}_\phi(x) = 0$$

for ϕ almost all $x \in A$, then A is ϕ rectifiable.

Proof. Let $\Phi \in \mathfrak{M}$ be the function on E_2 such that $s \subset E_2$ implies $\Phi(s) = \phi(As)$. A glance at the *proof* of 7.14 convinces us that

$$\mathfrak{D}_\Phi^\Delta(x) > 0, \quad (\{-i\} + \{i\}) \operatorname{dir}_\Phi(x) = 0$$

for Φ almost all x . Thus, corresponding to each Borel set B with positive Φ measure there is, by 7.14, a rectifiable subset of B with positive Φ measure. Accordingly 7.15 assures us E_2 is Φ rectifiable. Hence A is ϕ rectifiable.

7.17. THEOREM. If $\phi \in \mathfrak{U}$, $\phi^0(A) < \infty$,

$$\mathfrak{D}_\phi^\Delta(x) > 0 \quad \text{and} \quad (\{-i\} + \{i\}) \operatorname{dir}_\phi(x) = 0$$

for ϕ almost all $x \in A$, then A is ϕ rectifiable.

Proof. Since $\phi^0(A) < \infty$ we can and do select an open set G containing A for which $\phi(G) < \infty$. Letting Φ be the function on E_2 such that $s \subset E_2$ implies $\Phi(s) = \phi(Gs)$ we note: $\Phi \in \mathfrak{M}$;

$$\mathfrak{D}_\Phi^\Delta(x) = \mathfrak{D}_\phi^\Delta(x), \quad \operatorname{dir}_\Phi(x) = \operatorname{dir}_\phi(x)$$

for each x in the *open* set G . Consequently

$$\mathfrak{D}_\Phi^\Delta(x) > 0, \quad (\{-i\} + \{i\}) \operatorname{dir}_\Phi(x) = 0$$

for ϕ and Φ almost all $x \in A$. Theorem 7.16 now assures us that A is Φ rectifiable. Thus A , being a subset of G , is ϕ rectifiable.

7.18. THEOREM. If $\phi \in \mathfrak{U}$, $A \subset E_2$,

$$\mathfrak{D}_\phi^\Delta(x) > 0 \quad \text{and} \quad (\{-i\} + \{i\}) \operatorname{dir}_\phi(x) = 0$$

for ϕ almost all $x \in A$, then A is countably ϕ rectifiable.

Proof. Let R be the set of positive rationals less than 1. Let N be the set of positive integers. For $n \in N$, $\eta \in R$, $\delta \in R$ we let $\omega(n, \eta, \delta)$ be the set defined by: $x \in \omega(n, \eta, \delta)$ if and only if there is a set B such that

$$(0) \quad x \in A, \quad |x| \leq n,$$

$$(1) \quad (\mathfrak{R}_{-i}^n + \mathfrak{R}_i^n) \text{ Sgn } (\tilde{B}, x) = 0,$$

$$(2) \quad \phi(B\mathfrak{R}_x^r) \leq 2r \quad \text{for } 0 < r \leq \delta.$$

Setting

$$(3) \quad H = \sum_{\delta \in R} \sum_{\eta \in R} \sum_{n \in N} \omega(n, \eta, \delta)$$

we check that

$$A \supset H \supset A \int_{\tilde{x}} [(\{-i\} + \{i\}) \text{dir}_{\phi}(x) = 0],$$

infer from our hypotheses that

$$(4) \quad \phi(A\tilde{H}) = 0,$$

and divide the remainder of the proof into two parts.

PART I. If $n \in N$, $\eta \in R$, $\delta \in R$, then $\omega(n, \eta, \delta)$ is countably rectifiable.

Proof. Let $D = \omega(n, \eta, \delta)$. Since (0) implies D is a bounded subset of A , we use (1), (2), and 7.12 to obtain a rectifiable set C such that $\phi^0(D\tilde{C}) < \infty$. From 7.17 it follows that $D\tilde{C}$ is ϕ rectifiable. Thus the set D , which equals $DC + D\tilde{C}$, is ϕ rectifiable and hence, a fortiori, countably ϕ rectifiable.

PART II. A is countably ϕ rectifiable.

Proof. From (3) and Part I we know H is countably ϕ rectifiable. In view of (4) it is now evident that A is countably ϕ rectifiable.

7.19. THEOREM. If $\phi \in \mathfrak{U}$, $A \subset E_2$, Z is a diametral set,

$$\mathfrak{D}_{\phi}^{\Delta}(x) > 0 \quad \text{and} \quad Z \text{dir}_{\phi}(x) = 0$$

for ϕ almost all $x \in A$, then A is countably ϕ rectifiable.

Proof. Let $\{-z\} + \{z\} = Z$; $x' = -zix$, for $x \in E_2$;

$$\bar{s} = \sum_{x \in s} \{x'\}$$

for $s \subset E_2$. Let ψ be the function on E_2 such that

$$\psi(s) = \phi(\bar{s}) \quad \text{for } s \subset E_2.$$

Observe that $|x' - y'| = |x - y|$ for x and y in E_2 , check that $\psi \in \mathfrak{U}$, and note that

$$\overline{\sum_{s \in \mathfrak{F}} s} = \sum_{s \in \mathfrak{F}} \bar{s}, \quad \overline{\prod_{s \in \mathfrak{F}} s} = \prod_{s \in \mathfrak{F}} \bar{s}$$

for each family \mathfrak{F} of subsets of E_2 . Next infer

$$(1) \quad \overline{\text{Sgn}(s, x)} = \text{Sgn}(\bar{s}, x'), \quad \mathfrak{D}_\psi^\Delta(s, x) = \mathfrak{D}_\phi^\Delta(\bar{s}, x'), \\ \overline{\text{dir}_\psi(x)} = \text{dir}_\phi(x')$$

whenever $s \subset E_2$ and $x \in E_2$.

Setting

$$(2) \quad A_1 = A \ E_\psi [\mathfrak{D}_\phi^\Delta(y) > 0 \text{ and } Z \text{ dir}_\phi(y) = 0]$$

we select A_0 so that $\bar{A}_0 = A_1$ and verify that

$$(3) \quad \overline{(\{-i\} + \{i\}) \text{ dir}_\psi(x)} = (\{-z\} + \{z\}) \text{ dir}_\phi(x') = Z \text{ dir}_\phi(x') = 0$$

for $x \in A_0$. With the help of (2), (1), (3), 7.18 and 7.9 we secure a countable family \mathfrak{G} of rectifiable sets such that

$$\psi[A_0 - \sigma(\mathfrak{G})] = 0.$$

Consequently

$$0 = \phi[\bar{A}_0 - \overline{\sigma(\mathfrak{G})}] = \phi\left[A_1 - \sum_{s \in \mathfrak{G}} \bar{s}\right].$$

Since \bar{s} is congruent to the rectifiable set s for each $s \in \mathfrak{G}$ it is now clear that A_1 is countably ϕ rectifiable and, inasmuch as our hypotheses and (2) together imply $\phi(A\bar{A}_1) = 0$, it is evident that A is likewise countably ϕ rectifiable.

7.20. THEOREM. *If $\phi \in \mathfrak{U}$ and E_2 is ϕ restricted at ϕ almost all of its points then E_2 is countably ϕ rectifiable.*

Proof. Letting

$$\Gamma = E_2 E_x [|z| = 1], \quad \alpha(z) = E_x [(\{-z\} + \{z\}) \text{ dir}_\phi(x) = 0],$$

we select a countable subset β of Γ whose closure is Γ . Since $\text{dir}_\phi(x)$ is closed for $x \in E_2$ it is clear that $\phi(\bar{B}) = 0$ where

$$B = \sum_{z \in \beta} \alpha(z).$$

Now for $z \in \beta$ we gather from 2.24 that $\mathfrak{D}_\phi^\Delta(x) > 0$ for ϕ almost all x in $\alpha(z)$ and hence conclude from 7.19 that $\alpha(z)$ is countably ϕ rectifiable.

Accordingly B , and hence E_2 , is countably ϕ rectifiable.

7.21. THEOREM. *If $\phi \in \mathfrak{U}$, and A is ϕ restricted at ϕ almost all of its points then A is countably ϕ rectifiable.*

Proof. Let $\Phi \in \mathcal{U}$ be the function on \mathcal{C}_2 such that $\Phi(s) = \phi(As)$ for $s \in E_2$. With the help of 6.5 (iii), 6.7, 6.3, we deduce

$$\text{dir}_\Phi(x) = \text{dir}_\Phi(A, x) + \text{dir}_\Phi(\tilde{A}, x) = \text{dir}_\Phi(A, x) = \text{dir}_\Phi(A, x)$$

for $x \in A$. Checking that $\mathcal{D}_\Phi^\Delta(x) = \mathcal{D}_\Phi^\Delta(A, x)$ for $x \in E_2$ it follows from Definition 2.24 and our hypotheses that E_2 is Φ restricted not only for ϕ almost all $x \in A$ but for Φ almost all x . Accordingly Theorem 7.20 implies E_2 is countably Φ rectifiable. That A is countably ϕ rectifiable is now evident.

7.22. THEOREM. *If $\phi \in \mathcal{U}$, $\phi(A) < \infty$, and A is ϕ restricted at ϕ almost all of its points then A is ϕ rectifiable.*

Proof. Use 7.21 and the fact that $\phi(A) < \infty$.

7.23. THEOREM. *If $\phi \in \mathcal{U}$, and A is countably ϕ rectifiable then $\mathcal{D}_\phi^\nabla(A, x) > 0$ for ϕ almost all $x \in A$.*

Proof. Introduce intersecting measure and use 5.28.

7.24. THEOREM. *If $\phi \in \mathcal{U}$, B is ϕ restricted at ϕ almost all of its points, $A \subset B$, then A is ϕ restricted at ϕ almost all of its points.*

Proof. Inasmuch as $\text{dir}_\phi(A, x) \subset \text{dir}_\phi(B, x)$ for $x \in E_2$, the only difficulty arises in verifying that $\mathcal{D}_\phi^\Delta(A, x) > 0$ for ϕ almost all $x \in A$.

Since B is countably ϕ rectifiable by 7.21, we see at once that A is likewise countably ϕ rectifiable. From 7.23 it is apparent that

$$\mathcal{D}_\phi^\Delta(A, x) \geq \mathcal{D}_\phi^\nabla(A, x) > 0$$

for ϕ almost all $x \in A$.

7.25. THEOREM. *If $\phi \in \mathcal{U}$, and A is restricted at each of its points then A is countably ϕ rectifiable.*

Proof. Use 7.4.

7.26. THEOREM. *If $\phi \in \mathcal{U}$, $\phi(A) < \infty$, and A is restricted at each of its points, then A is ϕ rectifiable.*

Proof. Use 7.25.

7.27. THEOREM. *If A is ϕ directional at x , then A is ϕ restricted at x .*

Proof. Letting $\{-z\} + \{z\} = \text{dir}_\phi(A, x)$, we note that $\{-iz\} + \{iz\}$ is a diametral set whose intersection with $\text{dir}_\phi(A, x)$ is vacuous. Moreover inasmuch as $\text{dir}_\phi(A, x) \neq 0$ it follows from 6.7 that $\mathcal{D}_\phi^\Delta(A, x) > 0$.

Obviously we also have

7.28. THEOREM. *If A is directional at x , then A is restricted at x .*

7.29. THEOREM. *If $\phi \in \mathcal{U}$, $\phi(A) < \infty$, and if corresponding to each positive*

number ϵ there is a set $B \subset A$ such that $\phi(A \setminus B) < \epsilon$ and such that B is restricted at each of its points, then A is ϕ rectifiable.

Proof. See 7.26.

7.30. REMARK. If in Theorem 7.21 we replace ϕ by counting measure, the theorem which emerges is noticeably weaker than 7.4. Thus in view of the meaning (see 2.22) of the phrase " A is restricted at x " it is natural to regard the phrase " A is ϕ restricted at x " as short (compare with 2.24) for:

$$\mathfrak{D}_{\phi}^{\Delta}(A, x) > 0$$

and there is a point $z \in E_2$ with $|z| = 1$ for which

$$\{z\} \operatorname{dir}_{\phi}(A, x) = 0.$$

The futility of this viewpoint will now be shown.

Throughout this remark we shall call a set a *triangle* if and only if it is a closed set consisting of the interior and boundary of an equilateral triangle with its base horizontal and below the opposite vertex. Thus the length of each of the three sides of a triangle is equal to the diameter of the triangle.

For T a triangle and n a positive integer: construct $2n+1$ triangles, each with diameter equal to

$$\frac{\operatorname{diam} T}{2n+1},$$

so placed that the union of their bases is the base of T ; next take the middle triangle and vertically translate it upward until its top vertex coincides with the top vertex of T ; denote the family of $2n+1$ triangles thus obtained by $O_n(T)$.

Let T_0 be a triangle with diameter 1. Let $\mathfrak{F}_0 = \{T_0\}$. Let $\mathfrak{F}_1, \mathfrak{F}_2, \mathfrak{F}_3, \dots$ be so determined that

$$\mathfrak{F}_n = \sum_{T \in \mathfrak{F}_{n-1}} O_n(T)$$

for each positive integer n .

Noting that $\sigma(\mathfrak{F}_1) \supset \sigma(\mathfrak{F}_2) \supset \sigma(\mathfrak{F}_3) \supset \dots$ we define

$$\mathfrak{F} = \mathfrak{F}_1 + \mathfrak{F}_2 + \mathfrak{F}_3 + \dots$$

and⁽²⁾

$$\mathcal{A} = \prod_{n=1}^{\infty} \sigma(\mathfrak{F}_n).$$

(²) This set \mathcal{A} seems to have many properties in common with the set P described on pages 327 and 328 of [1].

Since $T \in \mathfrak{F}$ implies the projection of $\mathcal{A}T$ on the base of T is the base of T we have little difficulty in verifying that

$$L(\mathcal{A}T) = \text{diam } T \quad \text{for } T \in \mathfrak{F}.$$

In particular

$$L(\mathcal{A}) = 1.$$

An appraisal of our geometric construction convinces us that

$$\{e^{i\theta}\} \text{dir}_L(\mathcal{A}, x) = 0$$

for $x \in \mathcal{A}$ and $\pi/3 < \theta < 2\pi/3$. In fact a more careful analysis reveals that

$$\{e^{i\theta}\} \text{dir}_L(\mathcal{A}, x) = 0$$

for $x \in \mathcal{A}$ and $0 < \theta < \pi$.

We shall term a triangle in \mathfrak{F} an *apex* triangle if and only if its top vertex coincides with the top vertex of some larger triangle in \mathfrak{F} .

Let W be the set so defined that $x \in W$ if and only if $x \in \mathcal{A}$ and the number of apex triangles of which x is an element is finite. Inasmuch as

$$\prod_{n=1}^{\infty} \left(1 - \frac{1}{2n+1}\right) = 0,$$

we conclude $L(W) = 0$.

A check shows that

$$\mathfrak{D}_L^{\nabla}(\mathcal{A}, x) = 0 \quad \text{for } x \in \mathcal{A}\tilde{W};$$

consequently $\mathfrak{D}_L^{\nabla}(\mathcal{A}, x) = 0$ for L almost all x in \mathcal{A} .

Since $L(\mathcal{A}) = 1 > 0$ it follows from 11.1 (1) and 11.1 (4) that \mathcal{A} is not L rectifiable.

A further examination of \mathcal{A} makes clear that

$$\mathfrak{D}_L^{\Delta}(\mathcal{A}, x) \geq 1/2 \quad \text{for } x \in \mathcal{A};$$

and a closer scrutiny discloses that

$$\mathfrak{D}_L^{\Delta}(\mathcal{A}, x) = 1$$

for L almost all x in \mathcal{A} .

Now consider the two statements which follow.

STATEMENT I. If $\phi \in \mathfrak{U}$, $\phi(\mathcal{A}) < \infty$,

$$\mathfrak{D}_{\phi}^{\Delta}(\mathcal{A}, x) > 0 \quad \text{and} \quad (\{-i\} + \{i\}) \text{dir}_{\phi}(\mathcal{A}, x) = 0$$

for ϕ almost all x in \mathcal{A} , then \mathcal{A} is ϕ rectifiable.

STATEMENT II. If $\phi \in \mathfrak{U}$, $\phi(\mathcal{A}) < \infty$,

$$\mathfrak{D}_{\phi}^{\Delta}(A, x) > 0 \quad \text{and} \quad \{i\} \operatorname{dir}_{\phi}(A, x) = 0$$

for ϕ almost all x in A , then A is ϕ rectifiable.

Since $e^{i\pi/2} = i$, it follows from 7.22 and the properties of \mathcal{A} that Statement I is true and Statement II is false.

8. Density ratios. Recalling abbreviation 2.27 we state

8.1. THEOREM. If $\psi \in \mathfrak{M}$, $P = (0, 0)$, $Q = (1, 0)$, $A \subset B \subset E_2$, and if further

(i) $P \in A$ and $Q \in A$,

(ii) $1 \leq \psi(\mathfrak{R}_x^r)/2r \leq 1.01$ for $x \in B$ and $0 < r \leq 2$,

(iii) $1 \leq \psi(B\mathfrak{R}_x^r)/2r \leq 1.01$ for $x \in A$ and $0 < r \leq 2$,

then $\psi(P * Q) > .06$.

Proof. Let

$$\begin{aligned} \Gamma &= \mathfrak{R}_P^1, & \rho &= (2.4)^{1/2}, & \Gamma'_1 &= \mathfrak{R}_P^{\rho}, \\ \Gamma_2 &= \mathfrak{R}_Q^1, & \Gamma'_2 &= \mathfrak{R}_Q^{.6}, & \Gamma'_2 &= \mathfrak{R}_Q^2, \\ R &= (1.6, 0), & \Gamma_3 &= \mathfrak{R}_R^{1.4}, & H &= \Gamma'_2 - \Gamma'_1. \end{aligned}$$

It is possible that the reader may find it instructive to draw a judicious number of the above circles. However the reasoning here given is purely arithmetic in form.

In order to prove $\psi(P * Q) > .06$ we assume hereinafter that $\psi(P * Q) \leq .06$ and divide the remainder of the argument into parts culminating in Part VI which is easily recognized as false.

PART I. $\psi(\Gamma_1\Gamma_2) \leq .06$.

Proof. $\Gamma_1\Gamma_2 = P * Q$.

PART II. $\Gamma_1 + \Gamma_2 + \Gamma_3 \subset \Gamma'_2$.

Proof. $x \in \Gamma_1$ implies

$$|x - Q| \leq |x - P| + |P - Q| < 1 + 1 = 2$$

which in turn implies $x \in \Gamma'_2$.

Obviously $\Gamma_2 \subset \Gamma'_2$.

Finally $x \in \Gamma_3$ implies

$$|x - Q| \leq |x - R| + |R - Q| < 1.4 + .6 = 2, \quad \text{and} \quad x \in \Gamma'_2.$$

PART III. $\Gamma_3\Gamma_1 \subset \Gamma_2\Gamma_1$.

Proof. Suppose $x \in \Gamma_3\Gamma_1$ and let $r_1 = |x - P|$, $r_2 = |x - Q|$, $r_3 = |x - R|$. Since $x \in \Gamma_3\Gamma_1$ we know

$$(1) \quad r_1^2 < 1, \quad r_3^2 < (1.4)^2 = 49/25.$$

We also have

$$(2) \quad x_1^2 + x_2^2 = r_1^2,$$

$$(3) \quad (x_1 - 8/5)^2 + x_2^2 = (x_1 - 1.6)^2 + x_2^2 = r_3^2,$$

$$(4) \quad r_2^2 = (x_1 - 1)^2 + x_2^2.$$

From (2) and (3) we find

$$80x_1 = 25r_1^2 - 25r_3^2 + 64.$$

From (4) and (2) we have

$$40r_2^2 = 40 + 40r_1^2 - 80x_1.$$

Thus

$$\begin{aligned} 40r_2^2 &= 40 + 40r_1^2 - 25r_1^2 + 25r_3^2 - 64 \\ &= -24 + 15r_1^2 + 25r_3^2. \end{aligned}$$

Use of (1) yields

$$40r_2^2 < -24 + 15 + 25(49/25) = 40.$$

Accordingly $r_2 < 1$ and $x \in \Gamma_2$, and since $x \in \Gamma_1$, we have shown that $x \in \Gamma_2 \cap \Gamma_1$.

The proof of Part III is complete.

PART IV. If $x \in H$, then $|x - R| < .31$.

Proof. Let $r_1 = |x - P|$, $r_2 = |x - Q|$, $r_3 = |x - R|$. Since $x \in H = \Gamma'_2 - \Gamma'_1$ we know

$$(5) \quad r_2^2 < .36, \quad r_1^2 \geq 2.4.$$

We also know

$$(6) \quad x_1^2 + x_2^2 = r_1^2,$$

$$(7) \quad (x_1 - 1)^2 + x_2^2 = r_2^2,$$

$$(8) \quad r_3^2 = (x_1 - 1.6)^2 + x_2^2.$$

From (6) and (7) we find

$$2x_1 = r_1^2 - r_2^2 + 1.$$

From (8) and (6) we have

$$r_3^2 = 2.56 + r_1^2 - 3.2x_1.$$

Thus

$$\begin{aligned} r_3^2 &= 2.56 + r_1^2 - 1.6r_1^2 + 1.6r_2^2 - 1.6 \\ &= .96 + 1.6r_2^2 - .6r_1^2. \end{aligned}$$

Use of (5) yields

$$r_3^2 < .96 + (1.6)(.36) - (.6)(2.4) = .096 < (.31)^2.$$

The proof of Part IV is complete.

PART V. *The set BH is vacuous.*

Proof (by contradiction). Suppose $x \in BH$ and let $r_3 = |x - R|$. By Part IV we know

$$(9) \quad r_3 < .31.$$

Now let

$$\Gamma = \mathbb{R}_x^{1.4-r_3}$$

and note that $y \in \Gamma$ implies

$$|y - R| \leq |y - x| + |x - R| < 1.4 - r_3 + r_3 = 1.4,$$

which in turn implies $y \in \Gamma_3$. Thus $\Gamma \subset \Gamma_3$ and, using Part III, we infer that

$$(10) \quad \Gamma\Gamma_1 \subset \Gamma_3\Gamma_1 \subset \Gamma_2\Gamma_1$$

and also, making use of Part II, that

$$\Gamma_2'' \supset \Gamma_1 + \Gamma_3 \supset \Gamma_1 + \Gamma.$$

Hence from (ii), (10), and Part I we obtain

$$\begin{aligned} 4.04 &\geq \psi(\Gamma_2'') \geq \psi(\Gamma_1 + \Gamma) = \psi(\Gamma_1) + \psi(\Gamma) - \psi(\Gamma\Gamma_1) \\ &\geq 2 + 2.8 - 2r_3 - \psi(\Gamma_2\Gamma_1) \geq 4.8 - 2r_3 - .06. \end{aligned}$$

Thus $4.04 \geq 4.74 - 2r_3$. That is,

$$r_3 \geq 2.37 - 2.02 = .35$$

in contradiction to (9).

PART VI. $3.131 > 3.14$.

Proof. Since $H = \Gamma_2' - \Gamma_1'$ we have

$$\Gamma_2' = \Gamma_2'\Gamma_1' + (\Gamma_2' - \Gamma_1') = \Gamma_2'\Gamma_1' + H,$$

and since BH is vacuous by Part V, we see that

$$B\Gamma_2' = B\Gamma_2'\Gamma_1'.$$

Therefore $B\Gamma_1' \supset B\Gamma_2'$ and hence

$$B\Gamma_1' \supset B\Gamma_1' + B\Gamma_2' \supset B\Gamma_1 + B\Gamma_2',$$

inasmuch as $\Gamma_1 \subset \Gamma_1'$. Using (iii), the fact that $\Gamma_2' \subset \Gamma_2$, and Part I, we infer

$$\begin{aligned} 2.02\rho &\geq \psi(B\Gamma_1') \geq \psi(B\Gamma_1 + B\Gamma_2') \\ &= \psi(B\Gamma_1) + \psi(B\Gamma_2') - \psi(B\Gamma_1\Gamma_2') \\ &\geq \psi(B\Gamma_1) + \psi(B\Gamma_2') - \psi(\Gamma_1\Gamma_2) \\ &\geq 2 + 1.2 - .06. \end{aligned}$$

Consequently $2.02\rho \geq 3.14$. Remembering that $\rho = (2.4)^{1/2} < 1.55$ we get

$$3.131 = (2.02)(1.55) > 2.02\rho \geq 3.14.$$

The proof of Part VI is complete.

8.2. THEOREM. If $\phi \in \mathfrak{M}$, $\phi(E_2) > 0$, and $\mathfrak{D}_\phi^\Delta(x) < 1.01 \mathfrak{D}_\phi^\nabla(x)$ for ϕ almost all x , then there is a closed set C with $\phi(C) > 0$ and a number $\mu > 0$ such that

$$\phi(p * q) \geq \mu |q - p|$$

for $p \in C$ and $q \in C$.

Proof. In accordance with 5.9 we select positive numbers α and δ_1 and a bounded closed set D with $\phi(D) > 0$ such that

$$\alpha \leq \frac{\phi(\mathfrak{R}_\xi^\rho)}{2\rho} \leq 1.01\alpha < 1.01\mathfrak{D}_\phi^\nabla(\xi)$$

for $0 < \rho < \delta_1$ and $\xi \in D$. Letting $\Phi \in \mathfrak{M}$ be the function on \mathcal{E}_2 such that

$$\Phi(s) = \phi(Ds) \quad \text{for } s \subset E_2,$$

and checking that $\mathfrak{D}_\Phi^\Delta(\xi) \leq \mathfrak{D}_\phi^\Delta(\xi) < \infty$ for ϕ and Φ almost all ξ we are assured by 5.14 and 5.26 that

$$\alpha < \mathfrak{D}_\phi^\nabla(\xi) = \mathfrak{D}_\Phi^\nabla(D, \xi) = \mathfrak{D}_\Phi^\nabla(\xi)$$

for L and Φ almost all ξ in D . Inasmuch as $\Phi(D) = \phi(D) > 0$, Lemma 5.5 yields us a number $\delta_2 > 0$ and a bounded closed set D_0 such that

$$\Phi(D_0) > 0, \quad \alpha \leq \frac{\Phi(\mathfrak{R}_\xi^\rho)}{2\rho},$$

for $\xi \in D_0$ and $0 < \rho < \delta_2$. Now take

$$\delta = \frac{\delta_1 \delta_2}{\delta_1 + \delta_2}$$

so that

$$0 < \delta < \delta_1, \quad 0 < \delta < \delta_2.$$

Set $D_1 = D_0 D$. Observe that $\phi(D_1) = \phi(D_0 D) = \Phi(D_0) > 0$ and that

$$\alpha \leq \frac{\Phi(\mathfrak{R}_\xi^\rho)}{2\rho} = \frac{\phi(D\mathfrak{R}_\xi^\rho)}{2\rho} \leq \frac{\phi(\mathfrak{R}_\xi^\rho)}{2\rho} \leq 1.01\alpha$$

for $\xi \in D_1$ and $0 < \rho < \delta$. Since the bounded closed set D_1 is expressible as a finite sum of closed sets each with diameter less than $2^{-1}\delta$ we now can and do select a closed subset C of D_1 for which

$$\text{diam } C < \delta/2, \quad \phi(C) > 0, \quad C \subset D.$$

Clearly

$$(1) \quad \alpha \leq \frac{\phi(\mathcal{R}_\xi^\rho)}{2\rho} \leq 1.01\alpha \quad \text{for } \xi \in D \text{ and } 0 < \rho < \delta,$$

$$(2) \quad \alpha \leq \frac{\phi(D\mathcal{R}_\xi^\rho)}{2\rho} \leq 1.01\alpha \quad \text{for } \xi \in C \text{ and } 0 < \rho < \delta.$$

Having effected this uniformization we put

$$(3) \quad \mu = .06\alpha$$

and suppose $p \in C$, $q \in C$. We shall complete the proof by showing that

$$(4) \quad \phi(p * q) \geq \mu |q - p|.$$

To this end let $d = |q - p|$, note that (4) is trivially true if $d = 0$, assume henceforth that $d > 0$, let $P = (0, 0)$, $Q = (1, 0)$, and T be the function on E_2 (that is the finite complex plane) such that

$$T(x) = (q - p)x + p \quad \text{for } x \in E_2.$$

Clearly T is a continuous, univalent function whose domain and whose range is E_2 , indeed for $x \in E_2$ and $y \in E_2$ we have

$$|T(y) - T(x)| = d |y - x|.$$

Introduce the abbreviations

$$x' = T(x), \quad \bar{s} = \sum_{x \in s} \{x'\};$$

check that

$$\overline{sS} = \bar{s}\bar{S}, \quad \overline{s\mathcal{R}_x^r} = \bar{s}\mathcal{R}_{x'}^{r^d}$$

whenever $x \in E_2$, $r > 0$, $s \subset E_2$, $S \subset E_2$; define ψ as that function on \mathcal{E}_2 such that

$$\psi(s) = \frac{1}{\alpha d} \phi(\bar{s}) \quad \text{for } s \subset E_2.$$

Verify that $\psi \in \mathfrak{M}$ and that

$$(5) \quad \frac{\psi(s\mathcal{R}_x^r)}{2r} = \frac{1}{\alpha} \frac{\phi(\bar{s}\mathcal{R}_{x'}^{r^d})}{2r^d}$$

whenever $x \in E_2$ and $0 < r < \infty$. Determine sets A and B so that

$$\bar{A} = C, \quad \bar{B} = D.$$

Since $x \in B$ implies $x' \in D$, since $x \in A$ implies $x' \in C$, and since $0 < r \leq 2$

implies $0 < rd \leq 2 \text{ diam } C < \delta$, it follows from (5), (1), and (2) that

$$1 \leq \frac{\psi(\mathbb{R}_x^r)}{2r} \leq 1.01 \quad \text{for } x \in B \text{ and } 0 < r \leq 2,$$

$$1 \leq \frac{\psi(B\mathbb{R}_x^r)}{2r} \leq 1.01 \quad \text{for } x \in A \text{ and } 0 < r \leq 2.$$

Recalling that $P = (0, 0)$, $Q = (1, 0)$ we have

$$A \subset B, \quad p = P', \quad q = Q', \quad P \in A, \quad Q \in A,$$

so that Theorem 8.1 implies

$$\psi(P * Q) > .06.$$

A simple check shows

$$\overline{P * Q} = p * q,$$

from which we deduce

$$\frac{1}{\alpha d} \phi(p * q) = \frac{1}{\alpha d} \phi(\overline{P * Q}) = \psi(P * Q) > .06.$$

With the help of (3) we thus conclude that

$$\phi(p * q) > .06\alpha d = .06\alpha |q - p| = \mu |q - p|$$

and that (4) is true.

8.3. THEOREM. *If $\phi \in \mathcal{M}$, $3\mathcal{D}_\phi^\Delta(x) < 4\mathcal{D}_\phi^\nabla(x)$ for ϕ almost all x , then there is a closed set C with $\phi(C) > 0$ and a number $\mu > 0$ such that*

$$\phi(p * q) \geq \mu |q - p|$$

for $p \in C$ and $q \in C$.

Proof. By Lemma 5.10 there are positive numbers α, δ, η with $1 < \eta < 4/3$ and a bounded closed set C' with $\phi(C') > 0$ such that the relations $x \in C'$, Γ is a circle with $x \in \Gamma$, $0 < 2r < \delta$, $\text{diam } \Gamma < \delta$ together imply the relations

$$(1) \quad \alpha \leq \frac{\phi(\mathbb{R}_x^r)}{2r}, \quad \frac{\phi(\Gamma)}{\text{diam } \Gamma} \leq \alpha\eta.$$

Since C' is expressible as a finite sum of closed sets each with diameter less than $3^{-1}\delta$, we choose a closed subset C of C' with $\phi(C) > 0$ and diameter less than $3^{-1}\delta$.

Putting $\mu = \alpha(4 - 3\eta)$, observing the positivity of μ , and supposing $p \in C$ and $q \in C$ we propose to complete the proof by showing

$$(2) \quad \phi(p * q) \geq \mu |q - p|.$$

To this end let $d = |q - p|$, note that (2) is trivially true if $d = 0$, assume henceforth that $d > 0$, and define

$$p' = \frac{p + q}{2}, \quad \Gamma_1 = \mathfrak{R}_p^d, \quad \Gamma_2 = \mathfrak{R}_q^d, \quad \Gamma_3 = \mathfrak{R}_{p'}^{1.5d}.$$

Since $x \in \Gamma_1$ implies $|x - p'| \leq |x - p| + |p - p'| < d + .5d$ and since $x \in \Gamma_2$ implies $|x - p'| \leq |x - q| + |q - p'| < d + .5d$ we see that $\Gamma_1 + \Gamma_2 \subset \Gamma_3$. Accordingly

$$\phi(\Gamma_3) \geq \phi(\Gamma_1) + \phi(\Gamma_2) - \phi(\Gamma_1 \Gamma_2).$$

From this, (1), and the fact that $2d \leq 3d \leq 3 \text{ diam } C < \delta$ it follows that

$$3d\eta\alpha \geq 4d\alpha - \phi(\Gamma_1 \Gamma_2) = 4d\alpha - \phi(p * q)$$

and hence that

$$\phi(p * q) \geq \alpha(4 - 3\eta)d = \mu |q - p|.$$

With the verification of (2) the proof is complete.

9. Density ratios and rectifiability.

9.1. THEOREM. If

(i) $\phi \in \mathfrak{M}$ and $\mu > 0$,

(ii) $0 < \mathfrak{D}_\phi^\Delta(x) < \infty$ for ϕ almost all x ,

(iii) C is a bounded closed set with $\phi(C) > 0$,

(iv) $\phi(p * q) \geq \mu |q - p|$ for $p \in C$ and $q \in C$,

then there is a rectifiable set of positive ϕ measure.

Proof. We determine: according to Lemma 5.4, a number $\alpha > 0$ and a closed set A such that

$$\phi(\tilde{A}) < \phi(C)/4,$$

and

$$\mathfrak{D}_\phi^\Delta(x) > \alpha \quad \text{for } x \in A;$$

and next, according to Lemma 5.7, a number $\lambda > 1$ and a closed set B such that

$$\phi(\tilde{B}) < \phi(C)/4$$

and

$$\phi(\mathfrak{R}_x^r) \leq 2r\lambda$$

for $x \in B$ and $0 < r < \lambda^{-1}$. Let $C_1 = ABC$, note that C_1 is closed and that

$$\phi(C_1) > \phi(C)/2 > 0,$$

- (1) $\mathfrak{D}_\phi^\Delta(x) > \alpha > 0$ for $x \in C_1$,
 (2) $\phi(\mathfrak{R}_x^r) \leq 2r\lambda$ for $x \in C_1$ and $0 < r < 1/\lambda$.

Let ϵ be a number such that both

$$(3) \quad 0 < \epsilon < \frac{\alpha\mu}{2 + 240\lambda}$$

and

$$(4) \quad 0 < \epsilon < \alpha/2.$$

We use 5.12 and 5.26 to see that $\mathfrak{D}_\phi^\Delta(\tilde{C}_1, x) = 0$ for ϕ almost all x in C_1 and now ascertain $x_0 \in C_1$ and a number δ such that

$$(5) \quad 0 < \delta < 1/\lambda$$

and

$$(6) \quad \frac{\phi(C_1 \mathfrak{E}_{x_0}^r)}{2r} < \epsilon \quad \text{for } 0 < r < \delta.$$

Next, relying on (1), we let r_0 be a number such that

$$(7) \quad \begin{aligned} 0 < r_0 < \delta/12, \\ \phi(\mathfrak{R}_{x_0}^{r_0}) > \alpha r_0 \end{aligned}$$

and note, using (4) and recalling 2.9,

$$\phi(C_1 \mathfrak{E}_{x_0}^{r_0}) \geq \phi(C_1 \mathfrak{R}_{x_0}^{r_0}) = \phi(\mathfrak{R}_{x_0}^{r_0}) - \phi(\tilde{C}_1 \mathfrak{R}_{x_0}^{r_0}) > \alpha r_0 - 2r_0\epsilon > 0.$$

Let

$$C_2 = C_1 \mathfrak{E}_{x_0}^{r_1} \quad \text{where} \quad r_1 = \inf_i E_i [\mathfrak{E}_{x_0}^i \supset C_1 \mathfrak{E}_{x_0}^{r_0}],$$

and check that

$$C_1 \mathfrak{E}_{x_0}^{r_1} = C_1 \mathfrak{E}_x^{r_0}$$

and

$$(8) \quad C_1 \text{ bndr } \mathfrak{E}_{x_0}^{r_1} \neq 0.$$

It is clear that $r_1 \leq r_0$, but we observe furthermore that

$$0 < r_1 \leq r_0;$$

for if it were possible that $r_1 = 0$, it would follow that $C_2 = \{x_0\}$ and that

$$\phi(\{x_0\}) = \phi(C_2) = \phi(C_1 \mathfrak{E}_{x_0}^{r_1}) = \phi(C_1 \mathfrak{E}_{x_0}^{r_0}) > 0 \quad \text{and} \quad \mathfrak{D}_\phi^\Delta(x_0) = \infty,$$

contrary to (2). Therefore

$$(9) \quad \phi(C_2) > r_0(\alpha - 2\epsilon) \geq r_1(\alpha - 2\epsilon).$$

We define

$$\gamma = \text{bndr } \mathcal{R}_{x_0}^1.$$

Since $\text{bndr } \mathcal{R}_{x_0}^1 = \text{bndr } C_{x_0}^1$ we have from (8)

$$(10) \quad 0 \neq \gamma C_1 \mathcal{C}_{x_0}^{r_0} \subset \gamma C_2.$$

Let \mathfrak{F} be the family of circles such that $\Gamma \in \mathfrak{F}$ if and only if there is an x and an r such that $\Gamma = \mathcal{R}_x^r$ where

$$(11) \quad x \in C_2, \quad 0 < r < 2r_1, \quad \text{and} \quad \phi(\tilde{C}_1 \mathcal{R}_x^r) \geq \mu r.$$

Now, using the covering Theorem 5.11, we let \mathfrak{G} be a countable disjointed subfamily of \mathfrak{F} such that

$$\sum_{s \in \mathfrak{G}} s \supset \sigma(\mathfrak{F}).$$

Clearly for each $d > 0$

$$(12) \quad \mathfrak{G} \cap E[(\text{diam } s) > d] \text{ is finite.}$$

Therefore considering the family $\sum_{s \in \mathfrak{G}} \{s\}$ in relation to Theorem 4.1, we select a subfamily \mathfrak{S} of \mathfrak{G} such that the family

$$\sum_{s \in \mathfrak{S}} \{s\}$$

is semi-disjointed and

$$\sum_{s \in \mathfrak{S}} s = \sum_{s \in \mathfrak{G}} s.$$

We now let

$$C_3 = C_2 + \gamma + \sum_{s \in \mathfrak{S}} (\text{clsr } s)$$

and

$$D = \left(C_3 - \sum_{s \in \mathfrak{S}} s \right) + \sum_{s \in \mathfrak{S}} (\text{bndr } s),$$

and divide the remainder of the proof into six parts.

PART I. C_3 is closed.

Proof (by contradiction). Suppose $p_0 \in \tilde{C}_3(\text{clsr } C_3)$. Since $(C_2 + \gamma)$ is closed and $\text{clsr } s$ is closed for each $s \in \mathfrak{S}$, we select distinct members s_1, s_2, s_3, \dots of \mathfrak{S} with

$$\lim_{n \rightarrow \infty} \text{dist} [\{p_0\}, (\text{clsr } s_n)] = 0.$$

From (12) we infer

$$\lim_{n \rightarrow \infty} (\text{diam } s_n) = 0.$$

Thus, since the center of each of these circles is an element of C_2 , it follows that p_0 is a cluster point of the closed set C_2 , and $p_0 \in C_2 \subset C_3$.

PART II. C_3 is connected.

Proof (by contradiction). Suppose C' and C'' are disjoint nonvacuous closed sets with $C_3 = C' + C''$, $\gamma \subset C'$. Evidently

$$(\text{clsr } s) \subset C' \quad \text{or} \quad (\text{clsr } s) \subset C'' \quad \text{for } s \in \mathfrak{S}.$$

To show that $C'' \subset \mathfrak{C}_{x_0}^{\gamma_1}$ we assume the contrary: there is a circle $s \in \mathfrak{S}$ with $(\text{clsr } s) \subset C''$, $(\text{clsr } s) \cap \mathfrak{C}_{x_0}^{\gamma_1} \neq \emptyset$. Since the center of s is in $C_2 \subset \mathfrak{C}_{x_0}^{\gamma_1}$ we have $0 \neq (\text{clsr } s) \cap \gamma \subset C'' \cap C' = \emptyset$.

Hence $C'' \subset \mathfrak{C}_{x_0}^{\gamma_1} \cap \tilde{C}' \subset \mathfrak{C}_{x_0}^{\gamma_1} \cap \tilde{\gamma} = \mathfrak{R}_{x_0}^{\gamma_1}$.

Let $C''' = C' + \mathfrak{R}_{x_0}^{\gamma_1}$. Clearly C''' is closed with $C''' \cap C'' = \emptyset$. It is evident that $0 \neq C_1 C''$, $0 \neq C_1 C' \subset C_1 C'''$. Choose points $p \in C_1 C''$ and $q \in C_1 C'''$ such that

$$|p - q| = \text{dist}(C_1 C'', C_1 C''') > 0.$$

Since $\gamma C_1 \neq \emptyset$ by (10) and since $p \in \mathfrak{R}_{x_0}^{\gamma_1}$ we have $|p - q| < 2r_1$. Obviously $(p * q) \subset \tilde{C}_1$, $p \in C_2$ and (iv) implies

$$\phi(\mathfrak{R}_p^{|p-q|} \tilde{C}_1) \geq \phi(p * q) \geq \mu |p - q|.$$

Hence $\mathfrak{R}_p^{|p-q|} \in \mathfrak{S}$, $\mathfrak{C}_p^{|p-q|}$ is contained in C_3 , is a continuum, and is split into two disjoint closed sets $C'' \cap \mathfrak{C}_p^{|p-q|}$ and $C''' \cap \mathfrak{C}_p^{|p-q|}$ with p in the first and q in the second.

PART III. D is a continuum.

Proof. By Parts I and II the set C_3 is a continuum. From Lemma 4.7 it follows that D is a continuum.

PART IV. $L(D) < \infty$.

Proof. Evidently

$$(13) \quad D \subset C_2 + \gamma + \sum_{s \in \mathfrak{S}} (\text{bndr } s).$$

Noting that \mathfrak{S} is disjointed and $\phi \in \mathfrak{M}$ we use (11) to obtain

$$(14) \quad \begin{aligned} \sum_{s \in \mathfrak{S}} \text{diam } s &\leq \frac{2}{\mu} \sum_{s \in \mathfrak{S}} \phi(\tilde{C}_1 s) \leq \frac{2}{\mu} \sum_{s \in \mathfrak{S}} \phi(s) \leq \frac{2}{\mu} \phi[\sigma(\mathfrak{S})] < \infty, \\ L \left[\sum_{s \in \mathfrak{S}} (\text{bndr } s) \right] &\leq 5\pi \sum_{s \in \mathfrak{S}} (\text{diam } s) < \infty. \end{aligned}$$

Remembering (1) and Lemma 5.20 we have

$$(15) \quad L(C_2) = L(C_1 \mathfrak{C}_{x_0}^{r_1}) \leq (5/\alpha) \phi(C_1 \mathfrak{C}_{x_0}^{r_1}) < \infty.$$

Combining (13), (14), (15) and the relation $L(\gamma) = 2\pi r_1 < \infty$ we reach the conclusion $L(D) < \infty$.

PART V. D is rectifiable.

Proof. D is a continuum of finite Carathéodory linear measure. Hence Theorem 4.6 implies that D is rectifiable.

PART VI. $\phi(D) > 0$.

Proof. Since $\sum_{s \in \mathfrak{F}} s$ is a Borel subset of C_3 and $D \supset C_3 - \sum_{s \in \mathfrak{F}} s$ we have

$$\phi(D) \geq \phi\left(C_3 - \sum_{s \in \mathfrak{F}} s\right) = \phi(C_3) - \phi\left(\sum_{s \in \mathfrak{F}} s\right).$$

Since $C_3 \supset C_2$ we have, using (9),

$$\phi(C_3) \geq \phi(C_2) \geq r_1(\alpha - 2\epsilon).$$

Next, since $\mathfrak{F} \subset \mathfrak{F}$, we use (11), (7), (5), (2) to obtain the relation

$$(16) \quad \begin{aligned} \phi\left(\sum_{s \in \mathfrak{F}} s\right) &\leq \sum_{s \in \mathfrak{F}} \phi(s) \leq \lambda \sum_{s \in \mathfrak{F}} \text{diam } s = 5\lambda \sum_{s \in \mathfrak{F}} \text{diam } s \\ &\leq \frac{10\lambda}{\mu} \sum_{s \in \mathfrak{F}} \phi(\tilde{C}_1 s). \end{aligned}$$

From the facts that \mathfrak{F} is disjointed, $\sigma(\mathfrak{F}) \subset \mathcal{R}_{x_0}^{12r_1}$ and from (6) we have that

$$\sum_{s \in \mathfrak{F}} \phi(\tilde{C}_1 s) = \phi[\tilde{C}_1 \sigma(\mathfrak{F})] \leq \phi(\tilde{C}_1 \mathcal{R}_{x_0}^{12r_1}) < 24r_1\epsilon$$

and thus, referring back to (16), that

$$\phi\left(\sum_{s \in \mathfrak{F}} s\right) \leq \frac{240\lambda r_1 \epsilon}{\mu}.$$

Finally, remembering the bounds for ϵ given in (3), we have

$$\phi(D) \geq r_1(\alpha - 2\epsilon) - \frac{240\lambda r_1 \epsilon}{\mu} = r_1 \left[\alpha - \epsilon \left(\frac{(2 + 240\lambda)}{\mu} \right) \right] > 0.$$

We now state three theorems and prove the third; the first two follow from the third.

9.2. THEOREM. If $\phi \in \mathfrak{M}$ and $\mathfrak{D}_\phi^\Delta(x) < 1.01 \mathfrak{D}_\phi^\nabla(x)$ for almost all x , then E_2 is ϕ rectifiable.

9.3. THEOREM. If $\phi \in \mathfrak{M}$ and $3\mathfrak{D}_\phi^\Delta(x) < 4\mathfrak{D}_\phi^\nabla(x)$ for almost all x , then E_2 is ϕ rectifiable.

9.4. THEOREM. If $\phi \in \mathfrak{M}$,

$$A_1 = E_x [\mathfrak{D}_\phi^\Delta(x) < 1.01 \mathfrak{D}_\phi^\nabla(x)], \quad A_2 = E_x [3\mathfrak{D}_\phi^\Delta(x) < 4\mathfrak{D}_\phi^\nabla(x)]$$

and $\phi(\tilde{A}_1)\phi(\tilde{A}_2) = 0$, then E_2 is ϕ rectifiable.

Proof. Let $\nu = 1$ or 2 according as $\phi(\tilde{A}_1)$ is or is not zero and note that $\phi(\tilde{A}_\nu) = 0$.

Letting B be any Borel set with positive ϕ measure, we introduce the intersecting measure $\Phi \in \mathfrak{M}$ for which

$$\Phi(s) = \phi(Bs) \quad \text{for } s \subset E_2,$$

and note, with the aid of 5.14, that for L almost all $x \in B$

$$\mathfrak{D}_\phi^\nabla(x) = \mathfrak{D}_\phi^\nabla(B, x) = \mathfrak{D}_\Phi^\nabla(x), \quad \mathfrak{D}_\phi^\Delta(x) = \mathfrak{D}_\Phi^\Delta(x), \quad \mathfrak{D}_\phi^\Delta(x) = \mathfrak{D}_\Phi^\Delta(x).$$

Thus defining

$$D_1 = E_x [\mathfrak{D}_\Phi^\Delta(x) < 1.01 \mathfrak{D}_\Phi^\nabla(x)], \quad D_2 = E_x [3\mathfrak{D}_\Phi^\Delta(x) < 4\mathfrak{D}_\Phi^\nabla(x)],$$

noting that $\mathfrak{D}_\Phi^\Delta(x) < \infty$ for ϕ almost all x and using 5.26 we obtain

$$0 = L(BA, \tilde{D}_\nu) = \phi(BA, \tilde{D}_\nu) = \phi(BA, \tilde{D}_\nu) + \phi(B\tilde{A}_\nu, \tilde{D}_\nu) = \phi(B\tilde{D}_\nu) = \Phi(\tilde{D}_\nu).$$

Relying upon 8.2 or 8.3 according as $\nu = 1$ or $\nu = 2$, we select a number $\mu > 0$ and a bounded closed set C such that $\Phi(C) > 0$ and

$$\Phi(p * q) \geq \mu |q - p|$$

for $p \in C$ and $q \in C$. Clearly

$$0 < \mathfrak{D}_\Phi^\Delta(x) < \infty$$

for Φ almost all x so that we may now invoke 9.1 to ascertain a rectifiable set β of positive Φ measure. From this it follows at once that $B\beta$ is a rectifiable subset of B with

$$\phi(B\beta) = \Phi(\beta) > 0.$$

Recourse to Lemma 7.14 completes the proof.

By using intersecting measure we obtain the following two theorems from 9.2 and 9.3 respectively.

9.5. THEOREM. If $\phi \in \mathfrak{U}$, $\phi(A) < \infty$, $\mathfrak{D}_\phi^\Delta(A, x) < 1.01 \mathfrak{D}_\phi^\nabla(A, x)$ for ϕ almost all x in A , then A is ϕ rectifiable.

9.6. THEOREM. If $\phi \in \mathfrak{U}$, $\phi(A) < \infty$, $3\mathfrak{D}_\phi^\Delta(A, x) < 4\mathfrak{D}_\phi^\nabla(A, x)$ for ϕ almost all x in A , then A is ϕ rectifiable.

10. Rectifiability and density ratios.

10.1. THEOREM. If $\phi \in \mathfrak{M}$, $\mathfrak{D}_\phi^\nabla(x) < \infty$ for ϕ almost all x , and A is strictly rectifiable, then

$$0 < \mathfrak{D}_\phi^\nabla(x) = \mathfrak{D}_\phi^\Delta(x) < \infty$$

for ϕ almost all x in A .

Proof. Letting

$$C' = A - E_x [\mathfrak{D}_L^\nabla(A, x) = \mathfrak{D}_L^\Delta(A, x) = 1],$$

$$C'' = AE_x [\mathfrak{D}_\phi^\Delta(x) = \infty],$$

$$C''' = AE_x [\mathfrak{D}_\phi^\nabla(x) = 0],$$

we see in the light of 5.29 that any subset of A which has L measure zero has ϕ measure zero, and use 6.12, 5.19 and 5.13, 5.28 to obtain

$$0 = L(C') = \phi(C'), \quad 0 = L(C'') = \phi(C''), \quad 0 = \phi(C''').$$

Thus

$$(1) \quad \phi(A - A') = 0$$

where

$$A' = A - (C' + C'' + C''').$$

Let \mathcal{L} be the function on \mathcal{E}_2 such that $\mathcal{L}(s) = L(As)$ for $s \in \mathcal{E}_2$, note $\mathcal{L} \in \mathfrak{M}$ and that

$$(2) \quad 0 < \mathfrak{D}_\phi^\nabla(x) \leq \mathfrak{D}_\phi^\Delta(x) < \infty, \quad \mathfrak{D}_{\mathcal{L}}^\nabla(x) = \mathfrak{D}_{\mathcal{L}}^\Delta(x) = 1$$

for each $x \in A'$.

Let V be the function on A' such that $x \in A'$ implies $V(x)$ is the family of circles of the form \mathfrak{R}_x^r where

$$\phi(\mathfrak{R}_x^{5r}) < 6 \frac{\mathfrak{D}_\phi^\Delta(x)}{\mathfrak{D}_\phi^\nabla(x)} \phi(\mathfrak{R}_x^r).$$

Let F be the function on A' such that $x \in A'$ implies $F(x)$ is the family of those circles Γ for which $x \in \Gamma$ and

$$\phi(\widehat{\Gamma}) < 6 \frac{\mathfrak{D}_\phi^\Delta(x)}{\mathfrak{D}_\phi^\nabla(x)} \phi(\Gamma).$$

Note $V(x) \subset F(x)$ whenever $x \in A'$.

Notation 2.9 of [5] is used in the next three of the four parts into which the remainder of the proof is divided.

PART I. If $x \in A'$, then

$$\lim_{V(x) \ni \Gamma \rightarrow x} \frac{\phi(\Gamma)}{\text{diam } \Gamma} = \mathfrak{D}_\phi^\nabla(x).$$

Proof. Let r_1, r_2, r_3, \dots be positive numbers such that: $r_n \rightarrow 0$ as $n \rightarrow \infty$; and

$$(3) \quad \mathfrak{D}_\phi^\nabla(x) = \lim_{n \rightarrow \infty} \phi(\mathfrak{R}_x^{r_n}).$$

We have

$$\overline{\lim}_{n \rightarrow \infty} \frac{\phi(\mathfrak{R}_x^{5r_n})}{\phi(\mathfrak{R}_x^{r_n})} = 5 \overline{\lim}_{n \rightarrow \infty} \frac{\phi(\mathfrak{R}_x^{5r_n})}{10r_n} \cdot \frac{2r_n}{\phi(\mathfrak{R}_x^{r_n})} \leq 5 \frac{\mathfrak{D}_\phi^\Delta(x)}{\mathfrak{D}_\phi^\nabla(x)} < 6 \frac{\mathfrak{D}_\phi^\Delta(x)}{\mathfrak{D}_\phi^\nabla(x)}.$$

Accordingly for n sufficiently large

$$\phi(\mathfrak{R}_x^{5r_n}) < 6 \frac{\mathfrak{D}_\phi^\Delta(x)}{\mathfrak{D}_\phi^\nabla(x)} \phi(\mathfrak{R}_x^{r_n})$$

and $\mathfrak{R}_x^{r_n} \in V(x)$. From this and (3) we conclude

$$\mathfrak{D}_\phi^\nabla(x) \leq \lim_{V(x) \ni \Gamma \rightarrow x} \frac{\phi(\Gamma)}{\text{diam } \Gamma} \leq \mathfrak{D}_\phi^\nabla(x).$$

PART II. If $x \in A'$, then

$$\mathfrak{D}_\phi^\Delta(x) = \lim_{F(x) \ni \Gamma \rightarrow x} \frac{\phi(\Gamma)}{\text{diam } \Gamma}.$$

Proof. Let $\Gamma_1, \Gamma_2, \Gamma_3, \dots$ be circles, each of which has x in it, with

$$(4) \quad \text{diam } \Gamma_n \rightarrow 0 \quad \text{and} \quad \frac{\phi(\Gamma_n)}{\text{diam } \Gamma_n} \rightarrow \mathfrak{D}_\phi^\Delta(x)$$

as $n \rightarrow \infty$. For each positive integer n we set $d_n = \text{diam } \Gamma_n$ and obtain

$$\overline{\lim}_{n \rightarrow \infty} \frac{\phi(\widehat{\Gamma}_n)}{\phi(\Gamma_n)} = 5 \overline{\lim}_{n \rightarrow \infty} \frac{\phi(\widehat{\Gamma}_n)}{5d_n} \cdot \frac{d_n}{\phi(\Gamma_n)} \leq 5 \frac{\mathfrak{D}_\phi^\Delta(x)}{\mathfrak{D}_\phi^\Delta(x)} < 6 \leq 6 \frac{\mathfrak{D}_\phi^\Delta(x)}{\mathfrak{D}_\phi^\nabla(x)}.$$

Thus for n sufficiently large $\Gamma_n \in F(x)$. From this and (4) we conclude

$$\mathfrak{D}_\phi^\Delta(x) \leq \lim_{F(x) \ni \Gamma \rightarrow x} \frac{\phi(\Gamma)}{\text{diam } \Gamma} \leq \mathfrak{D}_\phi^\Delta(x).$$

PART III. For ϕ almost all x in A' we have

$$0 < \mathfrak{D}_{\phi}^{\nabla}(x) = \mathfrak{D}_{\phi}^{\Delta}(x) < \infty.$$

Proof. Remembering (2), the definition of F and Parts I and II we see that F is a diametrically regular Borelian blanket. (In [5] see Definitions 6.1, 6.5, 6.6.) Denoting

$$A' \overset{x}{E} \left[0 \leq \liminf_{F(x) \ni \Gamma \rightarrow x} \frac{\mathcal{L}(\Gamma)}{\phi(\Gamma)} < \infty \right]$$

by A'' we use Definition 2.3 and Theorem 11.4 of [5] to gain the information that $\phi(A' - A'') = 0$.

For $x \in A''$ we denote the number

$$\liminf_{F(x) \ni \Gamma \rightarrow x} \frac{\mathcal{L}(\Gamma)}{\phi(\Gamma)}$$

by $h(x)$.

Now

$$(5) \quad \frac{\mathcal{L}(\Gamma)}{\phi(\Gamma)} \cdot \frac{\phi(\Gamma)}{\text{diam } \Gamma} = \frac{\mathcal{L}(\Gamma)}{\text{diam } \Gamma}$$

whenever $\Gamma \in F(x)$ and $x \in A''$. Thus using Part I, the inclusion of $V(x)$ in $F(x)$, (5), the definition of V , and (2) we deduce that $x \in A''$ implies

$$\begin{aligned} h(x) \mathfrak{D}_{\phi}^{\nabla}(x) &= h(x) \liminf_{V(x) \ni \Gamma \rightarrow x} \frac{\phi(\Gamma)}{\text{diam } \Gamma} = \liminf_{V(x) \ni \Gamma \rightarrow x} \frac{\mathcal{L}(\Gamma)}{\phi(\Gamma)} \liminf_{V(x) \ni \Gamma \rightarrow x} \frac{\phi(\Gamma)}{\text{diam } \Gamma} \\ &\geq \liminf_{V(x) \ni \Gamma \rightarrow x} \frac{\mathcal{L}(\Gamma)}{\text{diam } \Gamma} \geq \mathfrak{D}_{\mathcal{L}}^{\nabla}(x) = 1. \end{aligned}$$

On the other hand using Part II, (5), the definition of F , and (2) we deduce that $x \in A''$ implies

$$\begin{aligned} h(x) \mathfrak{D}_{\phi}^{\Delta}(x) &= h(x) \overline{\liminf}_{F(x) \ni \Gamma \rightarrow x} \frac{\phi(\Gamma)}{\text{diam } \Gamma} = \liminf_{F(x) \ni \Gamma \rightarrow x} \frac{\mathcal{L}(\Gamma)}{\phi(\Gamma)} \overline{\liminf}_{F(x) \ni \Gamma \rightarrow x} \frac{\phi(\Gamma)}{\text{diam } \Gamma} \\ &\leq \overline{\liminf}_{F(x) \ni \Gamma \rightarrow x} \frac{\mathcal{L}(\Gamma)}{\text{diam } \Gamma} \leq \mathfrak{D}_{\mathcal{L}}^{\Delta}(x) = 1. \end{aligned}$$

Accordingly $x \in A''$ implies that

$$1 \leq h(x) \mathfrak{D}_{\phi}^{\nabla}(x) \leq h(x) \mathfrak{D}_{\phi}^{\Delta}(x) \leq 1$$

and hence that

$$0 < \mathfrak{D}_{\phi}^{\nabla}(x) = \mathfrak{D}_{\phi}^{\Delta}(x) < \infty.$$

Since $\phi(A' - A'') = 0$ the proof of Part III is complete.

PART IV. For ϕ almost all x in A we have

$$0 < \mathfrak{D}_{\phi}^{\nabla}(x) = \mathfrak{D}_{\phi}^{\Delta}(x) < \infty.$$

Proof. Use (1) and Part III.

10.2. THEOREM. If $\phi \in \mathfrak{M}$, $\mathfrak{D}_{\phi}^{\nabla}(x) < \infty$ for ϕ almost all x and E_2 is ϕ rectifiable, then

$$0 < \mathfrak{D}_{\phi}^{\nabla}(x) = \mathfrak{D}_{\phi}^{\Delta}(x) < \infty$$

for ϕ almost all x .

Proof. Let $\epsilon > 0$, and choose a strictly rectifiable set C such that $\phi(\tilde{C}) < \epsilon$. Putting

$$B = E_2 - \underset{x}{E} [0 < \mathfrak{D}_{\phi}^{\nabla}(x) = \mathfrak{D}_{\phi}^{\Delta}(x) < \infty]$$

we are assured by 10.1 that

$$\phi(BC) = 0.$$

Accordingly

$$\phi(B) \leq \phi(BC) + \phi(B\tilde{C}) \leq 0 + \phi(\tilde{C}) < \epsilon.$$

10.3. THEOREM. If $\phi \in \mathfrak{U}$, $\phi(A) < \infty$, $\mathfrak{D}_{\phi}^{\nabla}(A, x) < \infty$ for ϕ almost all x in A , and A is ϕ rectifiable, then

$$0 < \mathfrak{D}_{\phi}^{\nabla}(A, x) = \mathfrak{D}_{\phi}^{\Delta}(A, x) < \infty$$

for ϕ almost all x in A .

Proof. Use intersecting measure and 10.2.

11. **Interrelations.** A unified and fairly complete picture of our previous results may be obtained from

11.1. THEOREM. If $\phi \in \mathfrak{U}$, $\phi(A) < \infty$, $\mathfrak{D}_{\phi}^{\nabla}(A, x) < \infty$ for ϕ almost all x in A , then the following nine propositions are equivalent:

- (1) A is ϕ rectifiable;
- (2) $\mathfrak{D}_{\phi}^{\Delta}(A, x) < 1.01 \mathfrak{D}_{\phi}^{\nabla}(A, x)$ for ϕ almost all x in A ;
- (3) $3 \mathfrak{D}_{\phi}^{\Delta}(A, x) < 4 \mathfrak{D}_{\phi}^{\nabla}(A, x)$ for ϕ almost all x in A ;
- (4) $0 < \mathfrak{D}_{\phi}^{\nabla}(A, x) = \mathfrak{D}_{\phi}^{\Delta}(A, x) < \infty$ for ϕ almost all x in A ;
- (5) $0 < \mathfrak{D}_{\phi}^{\nabla}(A, x) = \mathfrak{D}_{\phi}^{\Delta}(A, x) < \infty$ for ϕ almost all x in A ;
- (6) A is ϕ directional at ϕ almost all of its points;
- (7) A is ϕ restricted at ϕ almost all of its points;
- (8) Corresponding to each positive number ϵ there is a set $B \subset A$ such that $\phi(A \setminus B) < \epsilon$ and such that B is directional at each of its points;
- (9) Corresponding to each positive number ϵ there is a set $B \subset A$ such that $\phi(A \setminus B) < \epsilon$ and such that B is restricted at each of its points.

Proof. Our argument consists of 13 steps each of which is given in the left column of the table below with its corresponding supporting reference or comment juxtaposed in the right column.

| | |
|-----------------|---------|
| (1) implies (6) | 6.15 |
| (6) implies (7) | 7.27 |
| (7) implies (1) | 7.22 |
| (1) implies (8) | 6.19 |
| (8) implies (9) | 7.28 |
| (9) implies (1) | 7.29 |
| (1) implies (5) | 10.3 |
| (5) implies (3) | obvious |
| (3) implies (1) | 9.6 |
| (1) implies (5) | 10.3 |
| (5) implies (4) | obvious |
| (4) implies (2) | obvious |
| (2) implies (1) | 9.5 |

The proof is complete.

In the following three theorems we investigate the connection between ϕ and L rectifiability.

11.2. THEOREM. *If $\phi \in \mathfrak{U}$, $\phi(A) < \infty$, A is L rectifiable, and if $\mathfrak{D}_\phi^\Delta(A, x) < \infty$ for ϕ almost all x in A , then A is ϕ rectifiable.*

Proof. Letting $\Phi \in \mathfrak{M}$ be the function on \mathcal{E}_2 such that

$$\Phi(s) = \phi(As) \qquad \qquad \qquad \text{for } s \subset E_2$$

the truth of the theorem is evident from 5.25 and the definitions of ϕ and L rectifiability.

11.3. REMARK. It is possible to construct a ϕ in \mathfrak{U} and a closed set A such that

$$\begin{aligned} \phi(A) &= 1, & L(A) &= 0, \\ \mathfrak{D}_\phi^\nabla(A, x) &= 0, & \mathfrak{D}_\phi^\Delta(A, x) &= \infty \end{aligned}$$

for $x \in A$.

The set A is thus obviously L rectifiable but by 11.1 (1) and 11.1 (4) it is not ϕ rectifiable.

11.4. THEOREM. *If $\phi \in \mathfrak{U}$, $L(A) < \infty$, A is ϕ rectifiable, and $\mathfrak{D}_\phi^\Delta(A, x) > 0$ for L almost all x in A , then A is L rectifiable.*

Proof. Let Φ be as in 11.2, let $\epsilon > 0$, choose a positive number μ so small (see 3.1(i)) that $L(A\bar{B}) < 2^{-1}\epsilon$ where B is the Borel set

$$E_\# [\mathfrak{D}_\Phi^\Delta(x) > \mu].$$

Letting C be a closed rectifiable set with $\Phi(\tilde{C}) < 10^{-1}\mu\epsilon$ we are helped by 5.21 to obtain

$$\begin{aligned} L(A\tilde{C}) &= L(AB\tilde{C}) + L(A\tilde{B}\tilde{C}) \leq L(B\tilde{C}) + L(A\tilde{B}) \\ &\leq L(B\tilde{C}) + \frac{\epsilon}{2} \leq \frac{5}{\mu} \Phi(B\tilde{C}) + \frac{\epsilon}{2} \leq \frac{5}{\mu} \Phi(\tilde{C}) + \frac{\epsilon}{2} < \epsilon. \end{aligned}$$

Thus AC is a rectifiable subset of A with $L(A\tilde{C}) < \epsilon$.

The L rectifiability of A is established.

11.5. REMARK. Let A be a set for which $L(A) < \infty$ and which is not L rectifiable; the set \mathcal{A} of 7.30 is such a set. Let ϕ be the function on \mathcal{E}_2 such that $\phi(s) = 0$ for $s \in E_2$. Clearly A is ϕ and not L rectifiable.

As an immediate consequence of the two preceding theorems we have

11.6. THEOREM. If $\phi \in \mathcal{U}$, $\phi(A) < \infty$, $L(A) < \infty$, $\mathfrak{D}_\phi^\Delta(A, x) > 0$ for L almost all x in A , $\mathfrak{D}_\phi^\Delta(A, x) < \infty$ for ϕ almost all x in A , then A is ϕ rectifiable if and only if A is L rectifiable.

BIBLIOGRAPHY

1. A. S. Besicovitch, *On the fundamental geometrical properties of linearly measurable plane sets of points* (II). Math. Ann. vol. 115 (1938) pp. 296–329.
2. C. Carathéodory, *Über das lineare Mass von Punktmengen—eine Verallgemeinerung des Längenbegriffs*, Nachr. Ges. Wiss. Göttingen, 1914, pp. 404–426.
3. H. Hahn, *Theorie der reellen Funktionen*, vol. 1, Berlin, 1921.
4. C. Kuratowski, *Topologie I*, Warsaw, 1933.
5. A. P. Morse, *A theory of covering and differentiation*, Trans. Amer. Math. Soc. vol. 55 (1944).
6. F. Roger, *Sur quelques applications métriques de la notion de contingent bilatéral*, C. R. Acad. Sci. Paris vol. 201 (1935) pp. 28–30.
7. S. Saks, *Theory of the integral*, Warsaw, 1937.

SUPPLEMENTARY BIBLIOGRAPHY

1. T. Estermann, *Über Carathéodory und Minkowski Verallgemeinerungen des Längenbegriffs*, Abh. Math. Sem. Hamburgischen Univ., 1925, pp. 73–116.
2. J. Favard, *Une définition de la longueur et de l'aire*, C. R. Acad. Sci. Paris vol. 194 (1932) pp. 344–346.
3. W. Gross, *Über des Flächenmass von Punktmengen*, Monatshefte für Mathematik und Physik vol. 27 (1918) pp. 145–176.
4. ———, *Über des lineare Mass von Punktmengen*, Monatshefte für Mathematik und Physik vol. 28 (1918) pp. 177–193.
5. O. Jansen, *Über einige stetige Kurven, über Bogenlängen, linearen Inhalt und Flächeninhalt*, Inaugural-Dissertation, Königsberg, 1907.
6. R. L. Jeffery, *Sets of k -extent in n -dimensional space*, Trans. Amer. Math. Soc. vol. 35 (1933) pp. 629–647.
7. H. Minkowski, *Über die Begriffe Länge, Oberfläche und Volumen*, Jber. Deutschen Math. Verein. vol. 9 (1900) pp. 115–121.
8. A. P. Morse and J. F. Randolph, *Gillespie measure*, Duke Math. J. vol. 6 (1940) pp. 408–419.

9. J. P. Schauder, *The theory of surface measure*, Fund. Math. vol. 8 (1926) pp. 1-48.
10. H. Steinhaus, *Zur Praxis der Rektifikation und zum Längenbegriff*, Sächsische Akademie zu Leipzig, Berichte vol. 82 (1930) pp. 120-130.
11. ———, *Sur la portée puratique et théorique de quelques théorèmes sur la mesure des ensembles de droites*, Comptes rendus du premier congress des pays Slaves, 1929, pp. 348-354.
12. W. H. Young and Grace Chisholm, *The theory of sets of points*, 1906.

UNIVERSITY OF CALIFORNIA,
BERKELEY, CALIF.
CORNELL UNIVERSITY,
ITHACA, N. Y.