

HOMOLOGICAL ALGEBRA IN LOCALLY COMPACT ABELIAN GROUPS

BY
MARTIN MOSKOWITZ⁽¹⁾

I. Introduction. This paper is concerned with homological algebra in the category \mathcal{L} of locally compact abelian topological groups. The morphisms of \mathcal{L} are the continuous homomorphisms. However, only certain exact sequences, resolutions etc., are admissible. These are sequences whose continuous homomorphisms are open onto their respective ranges. Such maps are called proper and the corresponding sequences, proper exact. Concomitant with this is the fact that although \mathcal{L} is an additive category it is not abelian.

The material in §II may be regarded as preparatory, although there may be some independent interest here. Various structural facts are proven (some of which are well known) and basic properties of important dual subcategories of \mathcal{L} are investigated.

In §III the projectives and injectives of \mathcal{L} are computed. It turns out that subgroups of projectives are projective and quotient groups of injectives are injective. Vector groups are characterized by the fact that they are both projective and injective. Finally, necessary and sufficient conditions are given for the existence of proper resolutions.

In §§IV and V continuous versions of the functors Hom and \otimes (via dualization) are defined on certain subcategories of \mathcal{L} and their functorial properties, including exactness, are investigated. These extend the usual notions for discrete groups. In order to do this it is necessary to study topological groups of continuous multilinear functions. Sufficient, and in a sense, necessary conditions are given for the various groups to be locally compact. Under these conditions the appropriate functors are shown to be isomorphic. Finally, more or less explicit computations are made for Hom , \otimes , etc., sharpening some of the earlier results, and their geometric and structural significance is investigated.

In §VI Tor and Ext are defined by resolutions as derived functors of \otimes and Hom , and their functorial properties are studied. It turns out that Tor_n and Ext_n vanish for $n \geq 2$, and that Tor_1 and Ext_1 are computable. Knowledge of Ext_1 in turn gives information about certain group extensions in \mathcal{L} .

Throughout this paper, complete duality of all concepts and theorems is obtained. In order to accomplish this some concessions have to be made to the topology at various stages. For this reason the functors Hom , \otimes , Tor , and Ext are not defined on \mathcal{L} but only on certain subcategories.

Received by the editors September 7, 1965.

⁽¹⁾ Research partially supported by the National Science Foundation.

This paper relies heavily on the Pontrjagin Duality Theorem and related structural and character theoretic facts about locally compact abelian groups. These facts are to be found for the most part in [1] or [2]. Material on discrete abelian group theory, and homological algebra are to be found in [3] and [6] respectively. Whenever possible we have deliberately chosen to refer to textbooks rather than the original papers since there the material is systematically organized. One theorem of some importance in the theory of locally compact groups is the so-called Open Mapping Theorem due to Pontrjagin. (See [2, Vol. I].) Since it seems to be less well known than the above material and is used so frequently here, it is worthwhile stating.

Let G and H be arbitrary locally compact groups and $f: G \rightarrow H$ a continuous surjective homomorphism. If G is the countable union of compact sets, then f is open.

Now a locally compact group G is called compactly generated if there exists a compact symmetric neighborhood U of 1 so that $G = \bigcup_{n=1}^{\infty} U^n$. Evidently a compactly generated group is the countable union of compact sets, and it is the class of compactly generated abelian groups to which we will apply the Open Mapping Theorem.

Before proceeding we make the following notational conventions: By G in \mathcal{L} we mean that G is a locally compact abelian group with the group operation written additively. We denote by R, Z, T the group of real numbers, integers and reals modulo 1 respectively. If G and H are in \mathcal{L} and $f: G \rightarrow H$ is a continuous homomorphism we denote by G^\wedge and H^\wedge the character groups of G and H and by $f^\wedge: H^\wedge \rightarrow G^\wedge$ the continuous homomorphism dual to f . If G is in \mathcal{L} and H is a closed subgroup of G , then (G^\wedge, H) stands for the annihilator of H in G^\wedge . The symbol \cong always means an isomorphism of topological groups. If $\{G_i\}_{i \in I}$ is a family of groups in \mathcal{L} we denote their direct product by $G_1 \oplus \cdots \oplus G_n$ if the family is finite and by $\prod_{i \in I} G_i$ in general. If all the G_i are equal we shall write G^I . The group G always denotes a locally compact abelian group, and G_0 its identity component. The symbols S° , S^- , and id_S denote the interior, closure, and identity map $S \rightarrow S$ respectively.

Finally, I would like to express my gratitude and appreciation to Professor G. P. Hochschild who, as thesis adviser, gave without limit his time, energy, and advice. In addition to many valuable suggestions Professor Hochschild gave me much encouragement during the entire period of the preparation of this paper.

II. Structural results.

PROPOSITION 2.1. *Let G_1 and G_2 be in \mathcal{L} , $f: G_1 \rightarrow G_2$ be a continuous homomorphism, and G_1^\wedge, G_2^\wedge and f^\wedge be their duals. Then,*

- (1) $\text{Ker } f^\wedge = (G_2^\wedge, f(G_1)^\wedge)$,
- (2) $f^\wedge(G_2^\wedge)^\wedge = (G_1^\wedge, \text{Ker } f)$,
- (3) f^\wedge is a monomorphism if and only if $f(G_1)$ is dense in G_2 .

Proof. (1) It is easy to see that for any closed subgroup H_1 of G_1 ,

$$(f^\wedge)^{-1}(G_1^\wedge, H_1) = (G_2^\wedge, f(H_1)^\wedge).$$

The result follows by taking $H_1 = G_1$.

(2) Apply (1) to f^\wedge . Then $\text{Ker } f^{\wedge\wedge} = (G_1^\wedge, f^\wedge(G_2^\wedge)^-)$ and therefore, by the duality theorem, $\text{Ker } f = (G_1, f^\wedge(G_2^\wedge)^-)$. Taking annihilators we have

$$f^\wedge(G_2^\wedge)^- = (G_1^\wedge, \text{Ker } f).$$

(3) f^\wedge is a monomorphism if and only if $(0) = \text{Ker } f^\wedge = (G_2^\wedge, f(G_1)^-)$, i.e., if and only if $f(G_1)^- = G_2$.

PROPOSITION 2.2. *Let G_1 and G_2 be in \mathcal{L} . If $f: G_1 \rightarrow G_2$ is a proper homomorphism then*

- (1) $f(G_1)$ is a closed subgroup of G_2 ,
- (2) $\text{Ker } f^\wedge = (G_2^\wedge, f(G_1))$,
- (3) f^\wedge is a proper homomorphism.

Proof. (1) Since $f(G_1)$ is a continuous open image of a locally compact space, it is locally compact. A locally compact subgroup of a topological group is closed.

(2) This follows immediately from (1) and Proposition 2.1, Part 1.

(3) f factors into $g\pi$ where $\pi: G_1 \rightarrow G_1/\text{Ker } f$ is the canonical epimorphism and $g: G_1/\text{Ker } f \rightarrow G_2$ is the continuous monomorphism induced by f . Since $f = g\pi$ where π is continuous, and f is a proper homomorphism, g is a proper monomorphism.

Now, as is known (see [2]), the dual π^\wedge of the proper epimorphism π is a proper monomorphism; and the dual g^\wedge of the *proper* monomorphism g is a proper epimorphism. So $f^\wedge = \pi^\wedge g^\wedge$ and is therefore an open map followed by an open map onto its range.

DEFINITION. A sequence

$$(S) \quad \cdots \longrightarrow G_i \xrightarrow{f_i} G_{i+1} \xrightarrow{f_{i+1}} G_{i+2} \longrightarrow \cdots$$

where each G_i is in \mathcal{L} and each f_i is a continuous homomorphism is called a *complex* if $f_i(G_i) \subset \text{Ker } f_{i+1}$ for all i . The sequence is called *exact* if for each i , $f_i(G_i) = \text{Ker } f_{i+1}$. A complex or an exact sequence is called *proper* if each f_i is a proper homomorphism. Note that if the G_i are compactly generated then an exact sequence is automatically proper exact by the Open Mapping Theorem.

THEOREM 2.1. *If the sequence*

$$(S) \quad \cdots \longrightarrow G_i \xrightarrow{f_i} G_{i+1} \xrightarrow{f_{i+1}} G_{i+2} \longrightarrow \cdots$$

is proper exact then its dual

$$\cdots \longrightarrow G_{i+2}^\wedge \xrightarrow{f_{i+1}^\wedge} G_{i+1}^\wedge \xrightarrow{f_i^\wedge} G_i^\wedge \longrightarrow \cdots$$

is also proper exact. In particular, the dual of a short proper exact sequence $0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$ is short proper exact.

Proof. By Proposition 2.2, f_i^\wedge is a proper homomorphism and

$$\text{Ker } f_i^\wedge = (G_{i+1}^\wedge, f_i(G_i)).$$

But $(G_{i+1}^\wedge, f_i(G_i)) = (G_{i+1}^\wedge, \text{Ker } f_{i+1})$ since $f_i(G_i) = \text{Ker } f_{i+1}$. Proposition 2.2 implies that $(G_{i+1}^\wedge, \text{Ker } f_{i+1}) = f_{i+1}^\wedge(G_{i+2}^\wedge)$. Thus $\text{Ker } f_i^\wedge = f_{i+1}^\wedge(G_{i+2}^\wedge)$.

REMARK. It is clear that the dual of a complex is a complex and that of a proper complex is a proper complex.

PROPOSITION 2.3. Let G be in \mathcal{L} and H_1 and H_2 closed subgroups of G with $H_1 \subset H_2$. Then $(H_2^\wedge, H_1) \cong (G^\wedge, H_1)/(G^\wedge, H_2)$.

Proof. Consider the short proper exact sequence

$$0 \rightarrow H_2/H_1 \rightarrow G/H_1 \rightarrow G/H_2 \rightarrow 0.$$

As is known (see [2]), the dual sequence is

$$0 \rightarrow (G^\wedge, H_2) \rightarrow (G^\wedge, H_1) \rightarrow (H_2^\wedge, H_1) \rightarrow 0$$

and is proper exact by Theorem 2.1.

THEOREM 2.2. Let the sequence (S) be a proper complex and

$$\cdots \longleftarrow G_i^\wedge \xleftarrow{f_i^\wedge} G_{i+1}^\wedge \xleftarrow{f_{i+1}^\wedge} G_{i+2}^\wedge \longleftarrow \cdots$$

be its dual. Denote by H^i the i th cohomology group of the former sequence and H_i the i th homology group of the latter. Then H^i and H_i are in \mathcal{L} and $H^i = H_i^\wedge$.

Proof. The fact that both complexes are proper implies that H^i and H_i are in \mathcal{L} by Proposition 2.2. The proof will be complete if we can show that for each i , $(H^{i+1})^\wedge = H_{i+1}$. By Proposition 2.2, $\text{Ker } f_i^\wedge = (G_{i+1}^\wedge, f_i(G_i))$ and $f_{i+1}^\wedge(G_{i+2}^\wedge) = (G_{i+1}^\wedge, \text{Ker } f_{i+1})$. Hence $\text{Ker } f_i^\wedge / f_{i+1}^\wedge(G_{i+2}^\wedge) = (G_{i+1}^\wedge, f_i(G_i)) / (G_{i+1}^\wedge, \text{Ker } f_{i+1})$. However, the latter term is isomorphic with $\text{Ker } (f_{i+1}^\wedge, f_i(G_i))$ which as is known (see [2]) is isomorphic with $(\text{Ker } f_{i+1} / f_i(G_i))^\wedge$.

Next we investigate the structure of the class of groups without small subgroups.

DEFINITION. A group G in \mathcal{L} has no small subgroups if there exists a neighborhood U of 0 which does not contain any nontrivial subgroups.

PROPOSITION 2.4. A locally compact group without small subgroups has a countable fundamental system of neighborhoods of (0).

Proof. Let U_0 be a compact symmetric neighborhood of (0) containing no nontrivial subgroups of G . Construct a sequence U_0, U_1, \dots of compact symmetric neighborhoods of (0) such that $2U_i \subset U_{i-1}$ for all $i > 0$. Then $\bigcap_{i=0}^\infty U_i$ is evidently a subgroup of G contained in U_0 . Hence $\bigcap_{i=0}^\infty U_i = (0)$. Now let U be any neighborhood of (0) in G . U contains an open neighborhood U' of (0) such that $U' \subset U_0$. Consider the family of closed subsets $U_i \cap \tilde{U}'$ of the compact space $U_0 \cap \tilde{U}'$ where \tilde{U}' is the complement of U' in G . The intersection of any finite subfamily

of these is again a $U_j \cap \tilde{U}'$, and the intersection of all of these is empty. Hence there exists an i so that $U_i \cap \tilde{U}' = \emptyset$, so that $U_i \subset U' \subset U$. Therefore, the U_i 's are a countable fundamental system of neighborhoods of $(0)^{(2)}$.

DEFINITION. A *test neighborhood*⁽³⁾ of a group G in \mathcal{L} is a neighborhood Γ_* of (0) in G such that for each neighborhood Γ of (0) in G there is an integer $k(\Gamma)$ which has the property that for every x in G , if $x, 2x, \dots, k(\Gamma)x \in \Gamma_*$, then $x \in \Gamma$.

THEOREM 2.3. *A group G has a test neighborhood if and only if it has no small subgroups.*

Proof. Suppose G is a locally compact abelian group without small subgroups. Let Γ_* be a compact symmetric neighborhood of (0) which contains no subgroups, and Γ be any neighborhood of (0) in G . By looking at $\Gamma \cap \Gamma_*$ we may assume that $\Gamma \subset \Gamma_*$. If there is no integer $k(\Gamma)$ with the above property, then for each integer k there is an $x(k)$ in G such that $x(k), 2x(k), \dots, kx(k) \in \Gamma_*$, but $x(k) \notin \Gamma$. Γ_* is compact, and by Proposition 2.4, G obeys the first axiom of countability. Hence there exists a subsequence $x(k_i) \rightarrow_i y$, where $y \in \Gamma_*$. Let j be any fixed positive integer. Choose i so that $k_i > j$. Then $jx(k_i) \in \Gamma_*$. But $jx(k_i) \rightarrow_i jy \in \Gamma_*$. Since Γ_* is symmetric, it contains the subgroup generated by y . Thus $y = 0$. Since $x(k_i) \rightarrow_i 0$, there exists an i after which $x(k_i) \in \Gamma$. This contradicts the assumption.

Conversely, let Γ_* be a test neighborhood of (0) . Suppose Γ_* contains a subgroup H . Let $x \in H$. Then x belongs to every neighborhood of (0) , so that $x = 0$. Thus Γ_* contains no nontrivial subgroups.

PROPOSITION 2.5. *Let G be in \mathcal{L} , and let H be a closed subgroup of G .*

- (1) *If G has no small subgroups, then H has no small subgroups.*
- (2) *If H is open and has no small subgroups, then G has no small subgroups.*

PROPOSITION 2.6. *Let $G = G_1 \oplus G_2 \oplus \dots \oplus G_n$. G has no small subgroups if and only if G_i has no small subgroups for $i = 1, \dots, n$.*

The proofs of Propositions 2.5 and 2.6 are clear.

THEOREM 2.4. *Let G be in \mathcal{L} . Then G has no small subgroups if and only if $G \cong R^n \oplus T^m \oplus D$ where R^n is an n -dimensional vector group, T^m is an m -dimensional torus, and D is a discrete group.*

Proof. In any case, the general structure theorem for groups in \mathcal{L} (see [1]) yields $G = R^n \oplus H$ where H contains a compact open subgroup C . Now suppose that G has no small subgroups. Then by Propositions 2.6 and 2.5, C has no small subgroups. Because the characters separate the points of C , and C is compact without small subgroups, C is isomorphic to a closed subgroup of T^k , a finite-dimensional

(2) This proof is modeled after one of Yamabe's.

(3) The existence of a test neighborhood in T was utilized by A. Weil in [1].

torus. This means that C^\wedge is a quotient group of Z^k and is therefore finitely generated. Hence $C^\wedge \cong Z^m \oplus F$ where F is a finite group. Thus $C \cong T^m \oplus F$. Now $H/T^m/C/T^m \cong H/C$. But since C/T^m is finite, and H/C is discrete, it follows that H/T^m is discrete, that is, T^m is open in H . Since T^m is divisible, it is a direct summand. So $H = T^m \oplus D$ where $D \cong H/T^m$, a discrete group. Therefore, $G \cong R^n \oplus T^m \oplus D$.

Conversely, since R , T , and D clearly have no small subgroups, Proposition 2.6 implies that a group of the form $R^n \oplus T^m \oplus D$ has no small subgroups. This completes the proof. An immediate consequence is

COROLLARY 1. *If G has no small subgroups then G_0 is open in G , and $G_0 \cong R^n \oplus T^m$. Furthermore, n , m , and D form a complete set of invariants for G .*

COROLLARY 2. *The following conditions are equivalent.*

- (1) G is compact without small subgroups.
- (2) $G \cong F \oplus T^m$ where F is a finite group.
- (3) G^\wedge is discrete and finitely generated.

Proof. (1) and (2) are equivalent since if G has no small subgroups and is compact, then $n=0$ and D is finite, and conversely, (2) and (3) are equivalent by the Fundamental Theorem of Abelian Groups applied to G^\wedge .

Now we consider the class of compactly generated groups. The structure of these groups can be derived from Theorem 2.1 together with the general structure theorem and the fact that toral subgroups are direct summands (see 2). However, we make use of Theorem 2.4 which, in addition, yields the fact that they are dual to the class of groups without small subgroups.

THEOREM 2.5. *Let G be in \mathcal{L} . Then G is compactly generated if and only if $G \cong R^n \oplus Z^m \oplus C$ where n and m are integers and C is a compact group.*

Proof. Suppose G is compactly generated. Let U be a compact neighborhood of (0) , which generates G . Let Γ_* be a neighborhood of (0) in T which contains no nontrivial subgroups. Denote by $W(U, \Gamma_*)$ the neighborhood of (0) in G^\wedge made up of the characters ξ of G which take U into Γ_* . $W(U, \Gamma_*)$ contains no nontrivial subgroups. Assume $n\xi \in W(U, \Gamma_*)$ for all integers n . Then $\xi(nU) = (n\xi)(U) \subset \Gamma_*$ for all n , and so $\xi(\bigcup_{n=1}^\infty nU) = \xi(G) \subset \Gamma_*$. $\xi(G)$ is evidently a subgroup of Γ_* . Hence $\xi=0$. Thus, if G is compactly generated then G^\wedge has no small subgroups and is therefore isomorphic to $R^n \oplus T^m \oplus D$, by Theorem 2.4. Hence, $G \cong R^n \oplus Z^m \oplus G$.

Conversely, a group G of the form $R^n \oplus Z^m \oplus C$ where C is compact, is evidently compactly generated. It is clear that n , m , and C are invariants and hence a complete set of invariants. C is characterized by the fact that it contains every compact subgroup of G . C is called the maximum compact subgroup of G .

COROLLARY 1. *G is compactly generated if and only if G^\wedge has no small subgroups. This follows immediately from Theorems 2.4 and 2.5.*

COROLLARY 2. *A compactly generated group G has no nontrivial compact subgroups if and only if G^\wedge is connected. G is connected if and only if G^\wedge is torsion free.*

THEOREM 2.6. *Let G be in \mathcal{L} and H be a closed subgroup. Then*

- (1) *G has no small subgroups if and only if H and G/H have no small subgroups;*
- (2) *G is compactly generated if and only if H and G/H are compactly generated.*

Proof. It suffices to prove (1) since (2) follows by dualization. Suppose that H and G/H have no small subgroups. Then there exists a neighborhood U of (0) in G such that $U \cap H$ contains no nontrivial subgroups, and also $\pi(U)$ contains no nontrivial subgroups, where π is the canonical epimorphism $G \rightarrow G/H$. Now let L be a subgroup of G , and suppose $L \subset U$. Then $\pi(L) \subset \pi(U)$, and therefore $L \subset H$. Since $L \subset U \cap H$, we know $L = (0)$.

Conversely, suppose G has no small subgroups. By Proposition 2.5, H has no small subgroups. We show that G/H has no small subgroups. First we observe that it is sufficient to show that $G_0 + H/H$ has no small subgroups. In fact, as we noted in Theorem 2.4, Corollary 1, G_0 and hence, $G_0 + H$, is open in G . Now, $G/H/G_0 + H/H \cong G/G_0 + H$, which is discrete, and hence has no small subgroups. It then follows from the part of the theorem already proven that G/H has no small subgroups.

Because G_0 is open, $G_0 + H/H \cong G_0/G_0 \cap H$. Thus it is sufficient to prove the theorem in the case that G is connected, so that $G \cong R^n \oplus T^m$.

Consider the simply connected covering group V of G with ϕ the covering map. Let π be the canonical map of G onto G/H . $\phi^{-1}(H)$ is a closed subgroup of V . Since ϕ and π are continuous open epimorphisms, so is $\pi\phi$. Therefore, $G/H \cong V/\text{Ker } \pi\phi$ which clearly has no small subgroups.

We give an application of our methods in proving the following known approximation theorem.

THEOREM 2.7. *Let G be in \mathcal{L} and U be a neighborhood of 0 in G . Then there exists a compact subgroup H of G , so that $H \subset U$ and G/H has no small subgroups; that is, any group in \mathcal{L} is a projective limit of groups without small subgroups.*

Proof. By taking U sufficiently small and identifying G with $G^{\wedge\wedge}$ we may assume $U = W(V, \Gamma)$ where V is a compact symmetric neighborhood of 0 in G^\wedge , and Γ is a neighborhood of 0 in T . Let $L = \bigcup_{n=1}^{\infty} nV$. Then

$$0 \rightarrow L \rightarrow G^\wedge \rightarrow G^\wedge/L \rightarrow 0$$

is a short proper exact sequence, where L is compactly generated and G^\wedge/L is discrete. By Theorem 2.1, the dual sequence is the short proper exact sequence $0 \rightarrow (G^\wedge/L)^\wedge \rightarrow G \rightarrow L^\wedge \rightarrow 0$. Let $H = (G^\wedge/L)^\wedge$. Then H is compact and $G/H \cong L^\wedge$, which, by Corollary 1 of Theorem 2.5, has no small subgroups, since L is compactly generated. If ξ is a character of G^\wedge/L , then $\xi \in G^{\wedge\wedge} = G$, and $\xi(L) = (0)$, so that $\xi(V) = (0) \subset \Gamma$. Hence $\xi \in U$. Thus $H \subset U$.

Let G and H be in \mathcal{L} . We denote by $\text{Hom}(G, H)$ the set of continuous homomorphisms from G to H ; by $\sum f^\wedge(H^\wedge)$ the subgroup of G^\wedge generated by $f^\wedge(H^\wedge)$ for $f \in \text{Hom}(G, H)$; and by $\bigcap \text{Ker } f$ the intersection of $\text{Ker } f$ for $f \in \text{Hom}(G, H)$.

THEOREM 2.8. *Let G and H be in \mathcal{L} . Then $\text{Hom}(G, H)$ separates the points of G if and only if $\sum f^\wedge(H^\wedge)$ is dense in G^\wedge .*

Proof. $\text{Hom}(G, H)$ separates points of G if and only if $\bigcap \text{Ker } f = (0)$, that is, if and only if $(G^\wedge, \bigcap \text{Ker } f) = (G^\wedge, (0)) = G^\wedge$. But $(G^\wedge, \bigcap \text{Ker } f) = (\sum (G^\wedge, \text{Ker } f))^\perp$ (see [2]). Hence $\text{Hom}(G, H)$ separates the points of G if and only if $\sum (G^\wedge, \text{Ker } f)$ is dense in G^\wedge . However, $(G^\wedge, \text{Ker } f) = (f^\wedge(H^\wedge))^\perp$, by Proposition 2.1. The result follows if we can show that if $\sum (f^\wedge(H^\wedge))^\perp$ is dense in G^\wedge , then $\sum f^\wedge(H^\wedge)$ is dense in G^\wedge . Now let $\xi \in G^\wedge$ and U be a neighborhood of 0 in G^\wedge . Choose U_1 so that $2U_1 \subset U$. There exists ξ_1, \dots, ξ_n so that $\xi_i \in f_i^\wedge(H^\wedge)^\perp$ and $(\sum_{i=1}^n \xi_i) - \xi \in U_1$. Let U_2 be a neighborhood of 0 in G^\wedge satisfying $nU_2 \subset U_1$, and choose $\eta_i \in H^\wedge$ so that $f_i^\wedge(\eta_i) - \xi_i \in U_2$, $i = 1, \dots, n$. Then $\sum_i f_i^\wedge(\eta_i) - \xi \in 2U_1 \subset U$.

COROLLARY. *$\text{Hom}(G, R)$ separates points of G if and only if the set of points of G^\wedge lying on one-parameter subgroups is dense in G^\wedge .*

Proof. In particular, if we take $H = R$, then $R = R^\wedge$. It follows that $\sum f^\wedge(R^\wedge)$ equals the subgroup of G^\wedge generated by the points which lie on one-parameter subgroups. Now, for any abelian topological group A ,

$$\sum \{f(R) : f \in \text{Hom}(R, A)\} = \bigcup \{f(R) : f \in \text{Hom}(R, A)\}.$$

For suppose $x = \sum_{i=1}^n x_i$, where $x_i \in f_i(R)$; that is $x_i = f_i(t_i)$. We can assume $x_i \neq 0$ for all i . Choose a new parametrization, say g_i of f_i so that $x_i = g_i(1)$. Then $g(t) = \sum_{i=1}^n g_i(t)$ is a one-parameter subgroup of A , and $g(1) = x$.

We now study dual properties of compactly generated groups and groups without small subgroups, which generalize the usual facts about compact-discrete duality.

THEOREM 2.9. *Let G be a compactly generated group in \mathcal{L} and G^\wedge be its character group. Then (1) G is torsion free if and only if G^\wedge is divisible. (2) G is divisible if and only if G^\wedge is torsion free.*

Proof. For each positive integer n , we define $f_n: G \rightarrow G$ by $f_n(x) = nx$. Clearly, f_n is a continuous homomorphism. By Theorem 2.5, $G \cong R^k \oplus T^m \oplus C$, and evidently f_n operates componentwise. Now $f_n(R^k) = R^k$, since vector groups are divisible. $f_n(Z^m)$ is closed in Z^m since Z^m is discrete; finally $f_n(C)$ is compact since f_n is continuous. The product of closed sets is closed in the product topology; hence $f_n(G) = f_n(R^k) \oplus f_n(Z^m) \oplus f_n(C)$ is closed in G , and is therefore locally compact. The Open Mapping Theorem guarantees that f_n is proper. By Proposition 2.2, f_n^\wedge is proper, and $f_n^\wedge(G^\wedge) = (G^\wedge, \text{Ker } f_n)$. One computes easily that $f_n^\wedge(\xi) = n\xi$ for $\xi \in G^\wedge$. Hence $(G^\wedge, \text{Ker } f_n) = nG^\wedge$. Now $n(G^\wedge) = G^\wedge$ if and only if $\text{Ker } f_n = (0)$.

Thus G^\wedge is divisible if and only if G is torsion free. Part 2 of this theorem follows in a similar way from $n(G) = (G, \text{Ker } f_n^\wedge)$.

COROLLARY. *Let G be a connected group in \mathcal{L} and H a closed pure subgroup of G . Then H is connected.*

Proof. By Corollary 2 of Theorem 2.5, G^\wedge is torsion free. Hence by Theorem 2.9, G is divisible. Now by definition, see [3], $nH = H \cap nG = H$ for each integer n . Thus H is divisible. Since H is compactly generated, by Theorem 2.6, it follows from Theorem 2.9 that H^\wedge is torsion free and hence from Corollary 2 of Theorem 2.5 that H is connected.

III. Projectives and injectives.

DEFINITION. I is an injective of \mathcal{L} if I is in \mathcal{L} , and for all short proper exact sequences and continuous homomorphisms $g: G_1 \rightarrow I$ there is a continuous homomorphism $g^-: G_2 \rightarrow I$ so that $g^-f_1 = g$.

$$\begin{array}{ccccccc} 0 & \longrightarrow & G_1 & \xrightarrow{f_1} & G_2 & \xrightarrow{f_2} & G_3 \longrightarrow 0 \\ & & \searrow g & & \swarrow g^- & & \\ & & & & I & & \end{array}$$

P is a projective if P is in \mathcal{L} , and for all short proper exact sequences and continuous homomorphisms $g: P \rightarrow G_3$ there is a continuous homomorphism $g^-: P \rightarrow G_2$ so that $f_2g^- = g$.

$$\begin{array}{ccccccc} 0 & \longrightarrow & G_1 & \xrightarrow{f_1} & G_2 & \xrightarrow{f_2} & G_3 \longrightarrow 0 \\ & & & & \swarrow g^- & & \nearrow g \\ & & & & P & & \end{array}$$

THEOREM 3.1. I is injective if and only if I^\wedge is projective.

Proof. Suppose we are given

$$\begin{array}{ccccccc} 0 & \longrightarrow & G_1 & \xrightarrow{f_1} & G_2 & \xrightarrow{f_2} & G_3 \longrightarrow 0 \\ & & & & & & \nearrow g \\ & & & & & & I^\wedge \end{array}$$

We dualize.

$$\begin{array}{ccccccc} 0 & \longrightarrow & G_3^\wedge & \xrightarrow{f_2^\wedge} & G_2^\wedge & \xrightarrow{f_1^\wedge} & G_1^\wedge \longrightarrow 0 \\ & & \searrow g^\wedge & & \swarrow h & & \\ & & & & I^{\wedge\wedge} & & \end{array}$$

is exact.

Since $I^{\wedge\wedge}$ is injective there is a continuous homomorphism $h: G_2^\wedge \rightarrow I^{\wedge\wedge}$ such that $hf_2^\wedge = g^\wedge$. Thus $g = f_2h^\wedge$. So h^\wedge is the required lift of g . The converse is proven similarly.

PROPOSITION 3.1. *If $\{I_a : a \in A\}$ is a family of injectives of \mathcal{L} , such that $\prod_{a \in A} I_a$ is in \mathcal{L} then $\prod_{a \in A} I_a$ is an injective of \mathcal{L} .*

PROPOSITION 3.2. *Conversely, if $I = \prod_{a \in A} I_a$ and is injective, then I_a is injective for all $a \in A$.*

Propositions 3.1 and 3.2 follow immediately from the definitions.

PROPOSITION 3.3. *If I is injective for \mathcal{L} then I is connected.*

Proof. Consider the proper exact sequence $0 \rightarrow Z \rightarrow R \rightarrow T \rightarrow 0$. Let $y \in I$. Define a continuous homomorphism $f: Z \rightarrow I$ by $f(n) = ny$. Since I is injective, f extends to $f^-: R \rightarrow I$. Now $f^-(R)$ is connected and hence is contained in I_0 . In particular $y \in I_0$. Thus $I \subset I_0$. In fact, each point of I lies on a 1 parameter subgroup.

PROPOSITION 3.4. *If I is in \mathcal{L} and is injective for the class of compactly generated groups, then I is injective for \mathcal{L} .*

Proof. By considering the class of extensions $0 \rightarrow nZ \rightarrow Z \rightarrow Z_n \rightarrow 0$ whose groups are compactly generated, we conclude that I is divisible. Let G be in \mathcal{L} , H be a closed subgroup of G , and $f: H \rightarrow I$ be a continuous homomorphism. We show f extends to a continuous homomorphism $G \rightarrow I$. G has an open, compactly generated subgroup, say L . By assumption, $f|_{L \cap H}$ extends to a continuous homomorphism $f_1: L \rightarrow I$. Let f_2 be the homomorphism of the external direct sum $L \oplus H$ into I , defined by $f_2((l, h)) = f_1(l) + f(h)$. The kernel of the canonical epimorphism of $L \oplus H$ onto the subgroup $L + H$ of G consists of the elements $(x, -x)$ with x in $L \cap H$. Evidently these elements lie in the kernel of f_2 . Hence f_2 induces a homomorphism f_3 of $L + H$ into I such that $f_3(l + h) = f_1(l) + f(h)$. Moreover, f_3 extends f_1 , since f is a homomorphism. Since L is open and f_1 is continuous on L , f_3 is continuous, because any homomorphic extension of a continuous homomorphism defined on an open subgroup is itself continuous. Since I is divisible and $L + H$ is open, f_3 extends to a continuous homomorphism $f_4: G \rightarrow I$, and evidently f_4 extends f .

THEOREM 3.2. *The following conditions on the locally compact abelian group I are equivalent.*

- (1) *I is injective for the class of compactly generated groups,*
- (2) *I is injective for \mathcal{L} ,*
- (3) *$I \cong R^n \oplus T^r$, the direct product of a vector group and a (possibly infinite-dimensional) torus⁽⁴⁾.*

Proof. If I is injective, then by Proposition 3.3, $I = R^n \oplus C$ where C is compact and connected. As a direct summand, C is also injective, by Proposition 3.2. Hence, C^\wedge is a discrete projective of \mathcal{L} , and is therefore a projective of \mathcal{D} , the

(⁴) It was recently pointed out to me by Professor G. P. Hochschild that the fact that R is injective was proven by J. Dixmier [7], for a different purpose.

category of discrete abelian groups. As a projective of \mathcal{D} , C^\wedge is a free abelian group, i.e., a direct sum of copies of \mathbb{Z} . Hence C is a direct product of copies of T .

Conversely, $R^n \oplus T^\sigma$ is injective for \mathcal{L} . In fact, by Proposition 3.1, it is sufficient to prove that R and T are injective. As is known (see [2]) T is injective for \mathcal{L} . We now show that R is injective for \mathcal{L} . It is sufficient to show that R is injective for the class of compactly generated groups. Let G_2 be compactly generated and let G_1 be a closed subgroup of G_2 . Since G_1 is compactly generated by Theorem 2.6, Theorem 2.5 shows that $G_i \cong W_i \oplus C_i$ where W_i is the direct sum of a vector group and a discrete finitely generated free abelian group, and C_i is a compact group for $i=1, 2$. Since W_1 admits a topological embedding as a closed subgroup of a vector group, it follows that W_1 has no nontrivial compact subgroups because vector groups have this property. Let $\pi: G_2 \rightarrow W_2$ be the canonical epimorphism with kernel C_2 . Let $\alpha: W_1 \rightarrow W_2$ be the restriction of π to W_1 . Then $\text{Ker } \alpha = W_1 \cap C_2 = (0)$ so that α is a continuous monomorphism. Since C_2 is compact, π is a closed map (see [1]). Now W_1 is a closed subgroup of G_1 , which is a closed subgroup of G_2 . Hence $\pi(W_1)$ is closed in W_2 and therefore is locally compact. Since W_1 is compactly generated, α is a proper monomorphism, by the Open Mapping Theorem. In addition, since C_2 is the maximum compact subgroup of G_2 , it follows that $C_1 \subset C_2$.

Now let $f \in \text{Hom}(G_1, R)$, and write A for $\pi(W_1)$. Clearly, $f(C_1) = (0)$. Since $\alpha^{-1}: A \rightarrow W_1$ is a topological group isomorphism, it follows that

$$(f|_{W_1}) \circ \alpha^{-1} \in \text{Hom}(A, R).$$

Suppose there exists $g \in \text{Hom}(W_2, R)$ so that $g|_A = (f|_{W_1}) \circ \alpha^{-1}$. Then define $f^-: G_2 \rightarrow R$ by $f^-(x+y) = g(x)$ where $x \in W_2$ and $y \in C_2$. Evidently $f^- \in \text{Hom}(G_2, R)$. If $s+t \in G_1$ where $s \in W_1$ and $t \in C_1$, then $f(s+t) = f(s) + f(t) = f(s)$, since $f(C_1) = (0)$. But $g(\alpha(s)) = f(s)$, by the choice of g . Hence $f(s+t) = g(\alpha(s))$. On the other hand, $f^-(s+t) = f^-(s) + f^-(t) = f(s)$, since, clearly, $f^-(C_1) = (0)$. Now $s = x+y$, where $x \in W_2$ and $y \in C_2$. Therefore $f^-(s) = g(x)$, by the definition of f^- . But $x = \alpha(s)$. So $f^-(s) = g(\alpha(s))$. Therefore $f^-(s+t) = g(\alpha(s)) = f(s+t)$ and $f^-|_{G_1} = f$.

Thus we have reduced the problem to showing that if W_1 is a subgroup of W_2 (W_i as above), and $f \in \text{Hom}(W_1, R)$ then there exists $g \in \text{Hom}(W_2, R)$ so that $g|_{W_1} = f$. Since one can embed W_2 as a closed subgroup of a vector group, say V , if f can be extended to $g \in \text{Hom}(V, R)$, then $g|_{W_2}$ will be the desired extension of f . Thus we may assume that W_2 is a vector group. We have $W_1 = V_1 \oplus Z^{m_1}$, where V_1 is a vector group. Let $\{x_1, \dots, x_n\}$ be a basis of V_1 as a vector space, and $\{y_1, \dots, y_k\}$ a basis of Z^{m_1} as a finitely generated free abelian group. Clearly $\{x_1, \dots, x_n, y_1, \dots, y_k\}$ are linearly independent in V and can be extended to a basis of V , say $\{x_1, \dots, x_n, y_1, \dots, y_k, z_1, \dots, z_j\}$. If

$$x = \sum_{i=1}^n a_i x_i + \sum_{i=1}^k b_i y_i + \sum_{i=1}^j c_i z_i$$

is in V , let g be the linear (and hence continuous) functional defined by

$$g(x) = \sum_{i=1}^n a_i f(x_i) + \sum_{i=1}^k b_i f(y_i).$$

Since f is a continuous homomorphism on V_1 , f is linear on V_1 and therefore g extends f . This completes the proof that R is injective. We have shown that (2) and (3) are equivalent. Proposition 3.4 completes the proof of Theorem 3.2.

COROLLARY 1. *Let G be in \mathcal{L} and H be a closed subgroup of G . Any one-parameter subgroup f of G/H lifts to a one-parameter subgroup f^- of G .*

Proof. This follows directly from Theorem 3.2 and Theorem 3.1. The above corollary and our previous results yield an easy proof of the following known result.

COROLLARY 2. *Let G be in \mathcal{L} . G is connected if and only if the set of points of G which lie on one-parameter subgroups is dense in G .*

Proof. Suppose that G is connected. Let U be any neighborhood of 0 and let $x \in G$. By Theorem 2.7, choose a compact subgroup H of G so that $H \subset U$ and G/H has no small subgroups. Denote the canonical epimorphism $G \rightarrow G/H$ by π . G/H is connected and therefore is of the form $R^n \oplus T^m$. Clearly, each point of G/H lies on a one-parameter subgroup. Hence there is an $f \in \text{Hom}(R, G/H)$ with the property that $f(1) = \pi(x)$. By Corollary 1 of Theorem 3.2, f lifts to $f^- \in \text{Hom}(R, G)$: that is, $\pi f^- = f$. Hence, $\pi f^-(1) = f(1) = \pi(x)$, so that $f^-(1) - x \in H \subset U$. Thus, the set of points of G that lie on one-parameter subgroups is dense in G . The converse is clear.

COROLLARY 3. *Let G be a locally compact abelian group. $\text{Hom}(G, R)$ separates the points of G if and only if G^\wedge is connected, i.e., if and only if $G = V \oplus D$, where V is a vector group and D is a torsion-free discrete group.*

Proof. This follows directly from Corollary 2 of Theorem 3.2 and the Corollary to Theorem 2.8.

THEOREM 3.3. *The following conditions on a locally compact abelian group P are equivalent.*

- (1) P is projective for the class of groups without small subgroups.
- (2) P is projective for \mathcal{L} .
- (3) $P \cong R^n \oplus \sum_{\sigma} Z$, the direct product of a vector group and a discrete free abelian group.

Proof. Theorem 3.3 is dual to Theorem 3.2.

COROLLARY 1. *If G is a projective of \mathcal{L} and H is a closed subgroup of G , then H is a projective of \mathcal{L} .*

Proof. By Theorem 3.3, it is sufficient to show H is isomorphic to the direct product of a vector group and a discrete free abelian group. From Theorem 3.3 we know that $G = V \oplus D$, where V is a vector group and D is a discrete free abelian group. Then $H_0 \subset G_0 = V$, so H_0 is a closed connected subgroup of V . This implies that H_0 is itself a vector group and hence an injective for \mathcal{L} , so that H_0 is a direct factor of H ; $H \cong H_0 \oplus H/H_0$. We may therefore assume, for the purpose of proving the corollary, that $H_0 = (0)$. Now G has no small subgroups. Hence, by Theorem 2.6, H has no small subgroups. By Corollary 1 of Theorem 2.4, H_0 is open in H . Hence H is discrete. Let π be the canonical epimorphism $G \rightarrow D$ with kernel V . Then $\text{Ker}(\pi|_H) = H \cap V$, a discrete subgroup of a vector group, which is therefore free. Moreover, $\pi|_H$ induces a monomorphism $H/H \cap V \rightarrow D$. Hence $H/H \cap V$ is isomorphic with a subgroup of a free group and is itself free. Let

$$\gamma: H \rightarrow H/H \cap V$$

be the canonical epimorphism, and consider $\text{id}(H/H \cap V)$. Since $H/H \cap V$ is a discrete free abelian group it is a projective for \mathcal{L} , so that $\text{id}(H/H \cap V)$ lifts to a (continuous) homomorphism $f: H/H \cap V \rightarrow H$ with the property that $\gamma f = \text{id}(H/H \cap V)$. Hence $H \cong (H \cap V) \oplus (H/H \cap V)$. Since both $H \cap V$ and $H/H \cap V$ are free, H is free.

COROLLARY 2. *Dually, proper homomorphic images of injectives are injectives, and in particular, the image under a continuous homomorphism of a torus is a torus. (Proper and continuous are equivalent here.)*

The following characterization of vector groups illustrates their particular importance.

COROLLARY 3. *A group G is both projective and injective for \mathcal{L} if and only if G is a vector group.*

Proof. If G is projective then $G \cong V \oplus \sum_{\sigma} Z$, the direct product of a vector group and a discrete free abelian group. Since G is injective it must be connected, hence $G = V$ a vector group. Conversely a vector group is both projective and injective.

THEOREM 3.4. *G_1 is injective if and only if G_1 is compactly generated and every proper short exact sequence beginning with G_1 splits.*

Proof. If G_1 is injective, then evidently every proper exact sequence

$$0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$$

splits and, by Theorem 3.2, G_1 is compactly generated.

Now assume G is compactly generated, and all short exact sequences beginning with G split. We have $G \cong R^n \oplus Z^m \oplus C$, where C is compact. Because the characters of C separate points, C can be embedded in $T^{\sigma(G)}$, via

$$0 \longrightarrow C \xrightarrow{f_1} T^{\sigma(G)},$$

say. Since f_1 is a continuous monomorphism and C is compact, f_1 is proper. If we denote by f_2 the canonical epimorphism of $T^{\sigma(G)} \rightarrow T^{\sigma(G)}/f_1(C)$, then

$$0 \longrightarrow C \xrightarrow{f_1} T^{\sigma(G)} \xrightarrow{f_2} T^{\sigma(G)}/f_1(C) \longrightarrow 0$$

is a proper short exact sequence. By Corollary 2 of the previous theorem, $T^{\sigma(G)}/f_1(C) \cong T^{\sigma'(G)}$, a torus. Since $0 \rightarrow Z^m \rightarrow R^m \rightarrow T^m \rightarrow 0$ is a proper short exact sequence beginning with Z^m , and $0 \rightarrow R^n \rightarrow R^n \rightarrow 0 \rightarrow 0$ is a proper short exact sequence beginning with R^n , it follows directly that

$$0 \rightarrow G \rightarrow R^{n+m} \oplus T^{\sigma(G)} \rightarrow T^{\sigma'(G)+m} \rightarrow 0$$

is a proper short exact sequence. This sequence splits, so G is a direct summand of $R^{n+m} \oplus T^{\sigma(G)}$ which is injective. Hence G is injective by Corollary 2 of Theorem 3.3.

COROLLARY. *If G is in \mathcal{L} then vector subgroups and toral subgroups of G are direct summands.*

THEOREM 3.5. *A group G_3 is projective if and only if G_3 has no small subgroups, and every proper short exact sequence $0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$ ending with G_3 splits.*

Proof. Theorem 3.5 is dual to Theorem 3.4.

DEFINITION. For a locally compact abelian group G we say that a sequence of homomorphisms

$$(0) \longrightarrow G \xrightarrow{e} I_0 \xrightarrow{d_0} I_1 \xrightarrow{d_1} I_2 \xrightarrow{d_2} \cdots$$

is an injective resolution of G if each I_n is injective and the sequence is proper exact. We say that a sequence of homomorphisms

$$\cdots \longrightarrow P_3 \xrightarrow{d_2} P_2 \xrightarrow{d_1} P_1 \xrightarrow{d_0} P_0 \xrightarrow{\pi} G \longrightarrow (0)$$

is a projective resolution of G if each P_n is projective and the sequence is proper exact.

DEFINITION. If $I_n = (0)$, $n \geq 2$ or $P_n = (0)$, $n \geq 2$, we call the resolution short.

THEOREM 3.6. (1) *G has an injective resolution if and only if G is compactly generated, and in that case there is a short injective resolution.*

(2) *G has a projective resolution if and only if G has no small subgroups, and in that case there is a short projective resolution.*

Proof. Let G be compactly generated. In the course of proving Theorem 3.4, we constructed the proper short exact sequence $0 \rightarrow G \rightarrow R^{n+m} \oplus T^\sigma \rightarrow T^{\sigma'} \rightarrow 0$. This is a short injective resolution.

Conversely, if $G \in \mathcal{L}$ and has an injective resolution

$$0 \longrightarrow G \xrightarrow{e} I_0 \xrightarrow{d_0} I_1 \xrightarrow{d_1} I_2 \xrightarrow{d_2} \cdots,$$

then $e(G) \cong G$. But $e(G)$ is locally compact and therefore a closed subgroup of I_0 . Since I_0 is injective it is compactly generated. Hence $e(G)$, and therefore G , is compactly generated.

(2) follows from (1) by dualization. The above injective resolution and its dual are clearly the ones to compute with.

IV. Multilinear functions and tensor products in commutative topological groups.

Let G_1, \dots, G_r and G be commutative topological groups. We call a map $\mu: \prod_{i=1}^r G_i \rightarrow G$ r -linear if for all x_i and y_i in G_i ($i = 1, \dots, r$), $\mu(x_1, \dots, x_i + y_i, \dots, x_r) = \mu(x_1, \dots, x_i, \dots, x_r) + \mu(x_1, \dots, y_i, \dots, x_r)$. We denote by $M_G(G_1, \dots, G_r)$ the set of all r linear continuous G valued maps on $\prod_{i=1}^r G_i$ ⁽⁵⁾. $M_G(G_1, \dots, G_r)$ becomes a commutative group under pointwise addition.

We topologize $M_G(G_1, \dots, G_r)$ by uniform convergence on compact subsets. More precisely, for each compact subset F of $\prod_{i=1}^r G_i$, and for each neighborhood Γ of 0 in G we define $W(F, \Gamma)$ to be the set $\{\mu: \mu \in M_G(G_1, \dots, G_r) \text{ and } \mu(F) \subset \Gamma\}$. It is easily verified that, with this family of sets as a fundamental system of neighborhoods of 0, $M_G(G_1, \dots, G_r)$ is a topological group.

THEOREM 4.1. (1) *If G_1, \dots, G_r are compactly generated and G is compact without small subgroups, then $M_G(G_1, \dots, G_r)$ is locally compact.*

(2) *If G_1, \dots, G_r are compact and G is without small subgroups, then $M_G(G_1, \dots, G_r)$ is discrete.*

(3) *If G_1, \dots, G_r are discrete and G is compact, then $M_G(G_1, \dots, G_r)$ is compact.*

Proof. (1) $\prod_{i=1}^r G_i$ is clearly locally compact. The topology on the function space $M_G(G_1, \dots, G_r)$ is the compact-open topology. In this setting the Ascoli Theorem states that a subset F of the space of all continuous maps $\prod_{i=1}^r G_i \rightarrow G$, with the compact-open topology, is compact if and only if (a) F is closed, (b) for each $x \in \prod_{i=1}^r G_i$, the set $\{f(x): f \in F\}$ has compact closure in G , and (c) F is equicontinuous at each point x of $\prod_{i=1}^r G_i$.

For each $i = 1, \dots, r$ let U_i be a compact neighborhood of 0 generating G_i . We denote $\prod_{i=1}^r U_i$ by U . Since G has no small subgroups, it possesses a closed test neighborhood of 0, say Γ_* . Thus $W(U, \Gamma_*)$ is a neighborhood of 0 in $M_G(G_1, \dots, G_r)$. We show that $W(U, \Gamma_*)$ is compact.

Since Γ_* is a closed neighborhood, it is clear that $W(U, \Gamma_*)$ is closed. $W(U, \Gamma_*)$ satisfies condition (b) automatically, since G is compact.

Now we show that $W(U, \Gamma_*)$ is equicontinuous at x , where $x = (x_1, \dots, x_r)$. For each $i = 1, \dots, r$, $x_i = \sum_{l=1}^{n_i} x_{il}$ where n_i is an integer and $x_{il} \in U_i$. We denote $\max \{n_i: i = 1, \dots, r\}$ by $n(x)$. If Γ is any neighborhood of 0 in G , let Γ' be a neighborhood of 0 in G with the property that $(2^r - 1)\Gamma' \subset \Gamma$. Choose Γ'' small enough so that $(r - 1)n(x)\Gamma'' \subset \Gamma'$, and $k(\Gamma'')$ as usual. Finally, for each $i = 1, \dots, r$, choose a neighborhood of 0, V_i , in G_i with the property that $k(\Gamma'')V_i \subset U_i$. Let $V(x, \Gamma)$

⁽⁵⁾ We note that for $r > 1$ continuity at 0 does not guarantee continuity everywhere.

$= \prod_{i=1}^r V_i$. This is a neighborhood of 0 in $\prod_{i=1}^r G_i$. We show that $\mu(x + V(x, \Gamma)) \subset \mu(x) + \Gamma$ for every $\mu \in W(U, \Gamma_*)$.

For each $(v_1, \dots, v_r) \in V$,

$$\mu(x_1 + v_1, \dots, x_r + v_r) = \mu(x_1, \dots, x_r) + \sum_{2^r-2} + \mu(v_1, \dots, v_r).$$

Since $(2^r - 1)\Gamma' \subset \Gamma$, it is sufficient to show that $\mu(v_1, \dots, v_r)$ and each of the terms in \sum_{2^r-2} belong to Γ' for every $\mu \in W(U, \Gamma_*)$. First we observe that

$$\mu(U_1 \times \dots \times V_i \times \dots \times U_r) \subset \Gamma''$$

for each $\mu \in W(U, \Gamma_*)$ and each $i = 1, \dots, r$, since

$$j\mu(u_1, \dots, v_i, \dots, u_r) = \mu(u_1, \dots, jv_i, \dots, u_r) \in \mu(U) \subset \Gamma_*,$$

for all $j = 1, \dots, k(\Gamma'')$. In particular, $\mu(V_1 \times \dots \times V_r) \subset \Gamma'$. Also, for each $q = 1, \dots, r-1$,

$$\begin{aligned} \mu(v_1, \dots, v_q, x_{q+1}, \dots, x_r) &= \mu\left(v_1, \dots, v_q, \sum_{l=1}^{n(x)} x_{q+1} l_{q+1}, \dots, \sum_{l=1}^{n(x)} x_r l_r\right) \\ &= \sum_{l_{q+1}, \dots, l_r=1}^{n(x)} \mu(v_1, \dots, v_q, x_{q+1} l_{q+1}, \dots, x_r l_r) \in (r-1)n(x)\Gamma'' \subset \Gamma' \end{aligned}$$

and we may deal similarly with the other terms of \sum_{2^r-2} . This completes the proof of part 1.

(2) Let Γ_* be a neighborhood of 0 in G which contains no nontrivial subgroups. Since $\prod_{i=1}^r G_i$ is compact, $W(\prod_{i=1}^r G_i, \Gamma_*)$ is a neighborhood of 0 in $M_G(G_1, \dots, G_r)$. Let $\mu \in W(\prod_{i=1}^r G_i, \Gamma_*)$ and suppose that $\mu(x_1, \dots, x_r) \neq 0$ for some $(x_1, \dots, x_r) \in \prod_{i=1}^r G_i$. Then $\mu(G_1 \times \{x_2\} \times \dots \times \{x_r\})$ is a nontrivial homomorphic image of G_1 and is contained in Γ_* ; a contradiction. Thus

$$W\left(\prod_{i=1}^r G_i, \Gamma_*\right) = (0),$$

and $M_G(G_1, \dots, G_r)$ is discrete.

(3) $\prod_{i=1}^r G_i$ is clearly discrete and hence the compact subsets F of $\prod_{i=1}^r G_i$ are finite. This means that $M_G(G_1, \dots, G_r)$ is a subspace of $G^{\prod_{i=1}^r G_i}$ with the product topology. G is compact, so by the Tychonoff Theorem, $G^{\prod_{i=1}^r G_i}$ is also compact. We show that $M_G(G_1, \dots, G_r)$ is closed in $G^{\prod_{i=1}^r G_i}$. Let $\{\mu_a : a \in A\}$ be a net in $M_G(G_1, \dots, G_r)$ which converges to a map $f: \prod_{i=1}^r G_i \rightarrow G$. If x_i and y_i belong to G_i , where i is any integer between 1 and r , then

$$\mu_a(x_1, \dots, x_i + y_i, \dots, x_r) \xrightarrow{a} f(x_1, \dots, x_i + y_i, \dots, x_r),$$

$$\mu_a(x_1, \dots, x_i, \dots, x_r) \xrightarrow{a} f(x_1, \dots, x_i, \dots, x_r),$$

$$\mu_a(x_1, \dots, y_i, \dots, x_r) \xrightarrow{a} f(x_1, \dots, y_i, \dots, x_r),$$

by definition of the product topology. Because of the continuity of the group operation in G and the fact that each μ_a is r linear, it is clear that f is r linear. Since $\prod_{i=1}^r G_i$ is discrete, $f \in M_G(G_1, \dots, G_r)$. This completes the proof.

If $r=1$ we write $\text{Hom}(G_1, G)$ for $M_G(G_1, \dots, G_r)$. In this case, we observe that equicontinuity at 0 implies equicontinuity at every point. Let U_1 be any compact neighborhood of 0 in G_1 and Γ_* as above. We show that $W(U_1, \Gamma_*)$ is equicontinuous at 0 and is therefore compact. If Γ is any neighborhood of 0, choose V so that $k(\Gamma)V \subset U_1$. Then for $v \in V$ and $h \in W(U_1, \Gamma_*)$,

$$jh(v) = h(jv) \in h(U_1) \subset \Gamma_*,$$

for $j=1, \dots, k(\Gamma)$. Therefore, $h(v) \in \Gamma$. Thus we have the following result.

COROLLARY. *If G_1 is locally compact, and G is compact without small subgroups, then $\text{Hom}(G_1, G)$ is locally compact.*

THEOREM 4.2. *Let G_1, \dots, G_r be locally compact abelian groups, and G be an arbitrary abelian topological group. For $r > 1$, $M_G(G_1, \dots, G_r)$ is isomorphic with $\text{Hom}(G_1, M_G(G_2, \dots, G_r))$ as a topological group.*

Proof. If S and T are topological spaces we denote the set of all continuous maps $S \rightarrow T$ by $\mathcal{M}(S, T)$. If $\mathcal{M}(S, T)$ is regarded as a topological space with the compact open topology we write $\mathcal{M}^*(S, T)$ for this. A well known property of the compact open topology is the following: if S and T are locally compact spaces and X an arbitrary space, then the map $\tau: \mathcal{M}(S \times T, X) \rightarrow \mathcal{M}(S, \mathcal{M}^*(T, X))$ defined by $\tau(f)(s)(t) = f(s, t)$ for $f \in \mathcal{M}(S \times T, X)$ and $(s, t) \in S \times T$, is a homeomorphism of $\mathcal{M}^*(S \times T, X)$ onto $\mathcal{M}^*(S, \mathcal{M}^*(T, X))$. Now, $M_G(G_1, \dots, G_r)$ and

$$\text{Hom}(G_1, M_G(G_2, \dots, G_r))$$

are subspaces of $\mathcal{M}^*(\prod_{i=1}^r G_i, G)$ and $\mathcal{M}^*(G_1, M_G(G_2, \dots, G_r))$ respectively. The latter can be identified with a subspace of $\mathcal{M}^*(G_1, \mathcal{M}^*(\prod_{i=2}^r G_i, G))$. The map τ sends the first homeomorphically onto the second and is evidently a group homomorphism.

COROLLARY 1. *If G_1 is compactly generated and G_2 has no small subgroups, then $\text{Hom}(G_1, G_2)$ is locally compact.*

Proof. By Theorem 4.2, $M_G(G_1, G_2) \cong \text{Hom}(G_1, \text{Hom}(G_2, G))$. Specializing to $G=T$ we find that $M_T(G_1, G_2^\wedge) \cong \text{Hom}(G_1, G_2)$. Since G_2 has no small subgroups, G_2^\wedge is compactly generated and, since T is compact without small subgroups, Theorem 4.1 applies.

COROLLARY 2. *If G_1 and G_2 are any locally compact abelian groups then $\text{Hom}(G_1, G_2)$ is isomorphic with $\text{Hom}(G_2^\wedge, G_1^\wedge)$.*

Proof. As we have already seen, $M_T(G_1, G_2^\wedge) \cong \text{Hom}(G_1, G_2)$. Now $M_T(G_1, G_2^\wedge) \cong M_T(G_2^\wedge, G_1)$, and the result follows by applying the same isomorphism with (G_2^\wedge, G_1^\wedge) in the place of (G_1, G_2) .

The following classes of groups are dual to one another: locally compact groups are self dual; compactly generated groups are dual to the groups without small subgroups; discrete finitely generated groups are dual to compact groups without small subgroups. As a result of these facts, and of Corollary 2, theorems of the sort considered below can be immediately dualized. For example, we have shown that

(1) If G is compactly generated and H has no small subgroups then $\text{Hom}(G, H)$ is locally compact.

(2) If G is any locally compact group and H is compact without small subgroups then $\text{Hom}(G, H)$ is locally compact.

Statement (1) is self dual whereas statement (2) dualizes as follows. If G is a finitely generated discrete group and H is any locally compact group then $\text{Hom}(G, H)$ is locally compact. We show that, in a sense, these are the best possible results.

THEOREM 4.3. (1') *If $\text{Hom}(G, H)$ is locally compact for all H without small subgroups then G is compactly generated.*

(2') *If $\text{Hom}(G, H)$ is locally compact for all H then G is a finitely generated discrete group.*

Dually, (1'') If $\text{Hom}(G, H)$ is locally compact for all compactly generated groups G then H is without small subgroups.

(2'') *If $\text{Hom}(G, H)$ is locally compact for all G then H is compact without small subgroups.*

Proof. It is sufficient to prove (1') and (2').

(1') We actually prove a stronger result; namely, that if $\text{Hom}(G, D)$ is locally compact for a certain fixed discrete group then G is compactly generated.

For the moment, let D denote an arbitrary discrete group. We may assume that $\text{Hom}(G, D)$ has a compact neighborhood of 0 of the form $W(F, \Gamma)$ where F is a compact subset of G and Γ is a closed neighborhood of 0 in D . Since D is discrete we may assume $\Gamma = (0)$. By a compactness argument, F is contained in a compact neighborhood U of 0 in G , which we may assume is symmetric. $W(U, (0))$ is a closed neighborhood of 0 in $\text{Hom}(G, D)$ which is contained in $W(F, (0))$. Hence $W(U, (0))$ is compact. By the Ascoli Theorem, for each $x \in G$, $\{f(x) : f \in W(U, (0))\}$ has compact closure in D , that is $\{f(x) : f(U) = (0)\}$ is finite. Let $L = \bigcup_{n=1}^{\infty} nU$. Then L is an open compactly generated subgroup of G . It is clear that for each $x \in G$, $\{f(x) : f(L) = (0)\}$ is finite. Now take D to be the direct sum of infinitely many copies of the additive group of the rationals mod 1. We show that $G = L$.

Suppose $x \in G$ but $x \notin L$. Since D is divisible, the 0-map, $L \rightarrow (0)$, can be extended to a homomorphism $f: G \rightarrow D$ with the property that $f(x) \neq 0$. Since L is open in G , f is continuous. Furthermore, because of the choice of D , there are infinitely many such extensions all differing in value at x . For if $mx \in L$ for some

integer m , any choice of $f(x)$ subject to the condition $\text{order } (f(x)) \mid m$ yields an extension of the sort described above, and there are infinitely many summands. On the other hand if mx is never in L , $f(x)$ is completely arbitrary. Hence, $\{f(x) : f(L) = (0)\}$ is infinite. This contradiction completes the proof of (1').

(2') If H ranges over all the groups without small subgroups then by (1') G is compactly generated. Now take $H = T^\sigma$, any infinite-dimensional torus. We make use of the evident result that if G is any abelian topological group and $\{H_a : a \in A\}$ is a family of abelian topological groups, then

$$\text{Hom}\left(G, \prod_{a \in A} H_a\right) \cong \prod_{a \in A} \text{Hom}(G, H_a).$$

We have therefore, $\text{Hom}(G, H) \cong \prod_\sigma \text{Hom}(G, T) = \prod_\sigma G^\wedge$. Thus $\prod_\sigma G^\wedge$ is locally compact. Hence all but a finite number of the factors are compact, so that G^\wedge is compact and G is discrete. Since G is compactly generated and discrete, it is finitely generated.

DEFINITION. If G_1, \dots, G_r is a family of locally compact abelian groups we define $\bigotimes_{i=1}^r G_i = M_T(G_1, \dots, G_r)^\wedge$.

Notice that for any locally compact abelian group G since $M_T(Z, G)$ is naturally isomorphic with $\text{Hom}(G, T)$ by Theorem 4.2, it follows that $Z \otimes G$ is naturally isomorphic with G .

THEOREM 4.4. $\bigotimes_{i=1}^r G_i$ is an abelian topological group. If G_1, \dots, G_r are compactly generated, compact, or discrete, then $\bigotimes_{i=1}^r G_i$ is locally compact, compact, or discrete, respectively⁽⁶⁾.

Proof. This follows directly from Theorem 4.1.

DEFINITION. We define the tensor map $\phi: \prod_{i=1}^r G_i \rightarrow \bigotimes_{i=1}^r G_i$ by

$$\phi(x_1, \dots, x_r)(\mu) = \mu(x_1, \dots, x_r)$$

for all $\mu \in M_T(G_1, \dots, G_r)$. It is clear that $\phi(x_1, \dots, x_r)$ is a continuous linear map $M_T(G_1, \dots, G_r) \rightarrow T$, and thus a character of $M_T(G_1, \dots, G_r)$.

Now let D be the subgroup of $\bigotimes_{i=1}^r G_i$ that is generated by $\phi(G_1 \times \dots \times G_r)$. Obviously, D is the set of all finite sums of elements of the form $\phi(x_1, \dots, x_r)$.

THEOREM 4.5. (1) ϕ is a continuous r -linear map.

(2) If $M_T(G_1, \dots, G_r)$ is locally compact⁽⁷⁾ then D is dense in $\bigotimes_{i=1}^r G_i$. In particular, if the G_i are discrete then $D = \bigotimes_{i=1}^r G_i$.

Proof. ϕ is r -linear, because each of the $\mu \in M_T(G_1, \dots, G_r)$ is r -linear. To show ϕ is continuous at x it is sufficient to take a neighborhood $\phi(x) + W(F, \Gamma)$ of $\phi(x)$

⁽⁶⁾ Actually, in §V we shall see that if G_1, \dots, G_r are compactly generated then $\bigotimes_{i=1}^r G_i$ is also compactly generated.

⁽⁷⁾ Whenever we make the convenient hypothesis that $M_T(G_1, \dots, G_r)$ is locally compact we really have in mind more accessible conditions; namely, that either the G_i are compactly generated (Theorem 4.1, Part 1) or they are discrete (Theorem 4.1, Part 3).

with F a compact subset of $M_T(G_1, \dots, G_r)$ and Γ a neighborhood of 0 in T , and find a neighborhood $U(F, \Gamma)$ of x in $\prod_{i=1}^r G_i$ so that $\phi(U) \subset \phi(x) + W(F, \Gamma)$, that is, $\mu(U) \subset \mu(x) + \Gamma$ for all $\mu \in F$. Thus the continuity of ϕ at x is equivalent to the equicontinuity of F at x . However, since F is compact, the Ascoli Theorem guarantees that F is equicontinuous at every point. Thus ϕ is continuous.

It is clear that $\mu^\wedge(\phi(x_1, \dots, x_r)) = \phi(x_1, \dots, x_r)(\mu) = \mu(x_1, \dots, x_r)$. Hence if μ^\wedge annihilates $\phi(\prod_{i=1}^r G_i)$ then $\mu = 0$, and so μ^\wedge is 0 because \wedge is an isomorphism $M_T(G_1, \dots, G_r) \rightarrow M_T(G_1, \dots, G_r)^\wedge$. Consequently the annihilator of D in $M_T(G_1, \dots, G_r)^\wedge$ is (0). Since $M_T(G_1, \dots, G_r)$ is a locally compact abelian group, the result follows.

The following lemma is a known fact about the compact open topology.

LEMMA. *If T is a locally compact space, then the composition map*

$$\mathcal{M}^*(T, X) \times \mathcal{M}^*(S, T) \rightarrow \mathcal{M}^*(S, X)$$

is continuous.

Now let G and H be in \mathcal{L} . It follows from the above lemma that the composition map $\text{Hom}(G, H) \times M_G(G_1, \dots, G_r) \rightarrow M_H(G_1, \dots, G_r)$ is continuous. In particular for each $\mu \in M_G(G_1, \dots, G_r)$, we define the map $\rho_\mu: G^\wedge \rightarrow M_T(G_1, \dots, G_r)$ by $\rho_\mu(\xi) = \xi \circ \mu$ for $\xi \in G^\wedge$. Clearly, ρ_μ is a continuous homomorphism. Furthermore, it follows from the definitions of the maps involved that for each μ in $M_G(G_1, \dots, G_r)$ we have $\rho_\mu^\wedge \circ \phi = \omega_G \circ \mu$, where ω_G is the natural isomorphism of G onto G^\wedge .

THEOREM 4.6. *Let G and G_1, \dots, G_r be in \mathcal{L} . To each $\mu \in M_G(G_1, \dots, G_r)$ there corresponds a $\chi_\mu \in \text{Hom}(\bigotimes_{i=1}^r G_i, G)$ satisfying $\chi_\mu \phi = \mu$. If $M_T(G_1, \dots, G_r)$ is locally compact, then the condition $\chi_\mu \phi = \mu$ characterizes χ_μ uniquely.*

Proof. Define $\chi_\mu = \omega_G^{-1} \rho_\mu^\wedge$ where ω_G and ρ_μ are as above. Then χ_μ is a continuous homomorphism $\bigotimes_{i=1}^r G_i \rightarrow G$, since ω_G^{-1} and ρ_μ^\wedge are continuous homomorphisms. Thus $\chi_\mu \in \text{Hom}(\bigotimes_{i=1}^r G_i, G)$. Moreover, we have $\chi_\mu \phi = \omega_G^{-1} \rho_\mu^\wedge \phi = \omega_G^{-1} \omega_G \mu = \mu$.

If f_μ is another continuous homomorphism satisfying $f_\mu \phi = \mu$, then, by linearity, $f_\mu|_D = \chi_\mu|_D$. Since f_μ and χ_μ are both continuous and D is dense, it follows that $f_\mu = \chi_\mu$. This completes the proof.

Denote by \mathcal{A} the category of all abelian topological groups. We define two functors $\mathcal{F}, \mathcal{G}: \mathcal{L} \rightarrow \mathcal{A}$ by $\mathcal{F}(G) = M_G(G_1, \dots, G_r)$, $\mathcal{G}(G) = \text{Hom}(\bigotimes_{i=1}^r G_i, G)$. If G and H are in \mathcal{L} and h is a continuous homomorphism then we define $\mathcal{F}(h)(\mu) = h \circ \mu$ for $\mu \in M_G(G_1, \dots, G_r)$ and $\mathcal{G}(h)(f) = h \circ f$, for $f \in \text{Hom}(\bigotimes_{i=1}^r G_i, G)$. It is clear that \mathcal{F} and \mathcal{G} are additive covariant functors.

THEOREM 4.7. *If $M_T(G_1, \dots, G_r)$ is locally compact then the functors \mathcal{F} and \mathcal{G} are isomorphic.*

Proof. We denote the map $\mu \rightarrow \chi_\mu$ defined above by

$$\chi: M_G(G_1, \dots, G_r) \rightarrow \text{Hom}\left(\bigotimes_{i=1}^r G_i, G\right).$$

If $f \in \text{Hom}(\bigotimes_{i=1}^r G_i, G)$ then $f \circ \phi \in M_G(G_1, \dots, G_r)$. Since f and $\chi_{f \circ \phi}$ are continuous homomorphisms that coincide on D , we have $f = \chi_{f \circ \phi}$. On the other hand, if $\chi_\mu = \chi_\nu$ then $\chi_\mu \phi = \chi_\nu \phi$, i.e., $\mu = \nu$. Thus χ is a bijection.

We denote by ρ the map $\mu \rightarrow \rho_\mu$. Thus

$$\rho: M_G(G_1, \dots, G_r) \rightarrow \text{Hom}(G^\wedge, M_T(G_1, \dots, G_r)).$$

One checks easily that ρ is a continuous homomorphism.

Now consider the dualization map

$$\lambda: \text{Hom}(G^\wedge, M_T(G_1, \dots, G_r)) \rightarrow \text{Hom}\left(\bigotimes_{i=1}^r G_i, G\right)$$

defined by $\lambda(h) = \omega_G^{-1} h^\wedge$, for $h \in \text{Hom}(G^\wedge, M_T(G_1, \dots, G_r))$, where ω_G is the natural isomorphism of $G \rightarrow G^{\wedge\wedge}$. By Corollary 2, Theorem 4.2, the dualization map $\text{Hom}(A, B) \rightarrow \text{Hom}(B^\wedge, A^\wedge)$ is continuous. Hence λ is continuous. As is easily seen, $\chi = \lambda\rho$, so that χ is a continuous homomorphism. If $W(F, \Gamma)$ is a neighborhood of 0 in $M_G(G_1, \dots, G_r)$, since the tensor map ϕ is continuous, $\phi(F)$ is compact. Hence $W(\phi(F), \Gamma)$ is a neighborhood of 0 in $\text{Hom}(\bigotimes_{i=1}^r G_i, G)$ which is contained in $\chi(W(F, \Gamma))$. Thus χ is an open map. We have shown that χ is a topological group isomorphism.

If h is a continuous homomorphism $h: G \rightarrow H$ we show that the following diagram is commutative:

$$\begin{array}{ccc} \mathcal{F}(G) & \xrightarrow{\chi} & \mathcal{G}(G) \\ \downarrow \mathcal{F}(h) & & \downarrow \mathcal{G}(h) \\ \mathcal{F}(H) & \xrightarrow{\chi} & \mathcal{G}(H) \end{array}$$

First, for each $\mu \in M_G(G_1, \dots, G_r)$ and $d \in D$, it is clear that $\chi_{h \circ \mu}(d) = h(\chi_\mu(d))$. Since D is dense and $\chi_{h \circ \mu}$ and $h \circ \chi_\mu$ are continuous, it follows that $\chi_{h \circ \mu} = h \circ \chi_\mu$. Hence $\chi \mathcal{F}(h)(\mu) = \chi_{h \circ \mu} = h \circ \chi_\mu = \mathcal{G}(h)\chi_\mu$. Thus $\chi \mathcal{F}(h) = \mathcal{G}(h)\chi$. This completes the proof.

COROLLARY 1. *If $M_T(G_1, \dots, G_r)$ is locally compact the property of equivalence of r -linear maps and linear maps characterizes $\bigotimes_{i=1}^r G_i$ up to natural topological group isomorphisms. In particular, if all the G_i are discrete then $\bigotimes_{i=1}^r G_i$ is the usual tensor product for abstract abelian groups.*

COROLLARY 2. *If G , G_1 , G_2 and $M_T(G_1, G_2)$ are locally compact then $\text{Hom}(G_1 \otimes G_2, G)$ is isomorphic with $\text{Hom}(G_1, \text{Hom}(G_2, G))$.*

This follows directly from Theorems 4.7 and 4.2.

THEOREM 4.8. *Let G and G_1, \dots, G_r be abelian topological groups, and let H and H_1, \dots, H_r be subgroups of the respective G 's. Denote the natural epimorphisms $G \rightarrow G/H$ by π , and $\prod_{i=1}^r G_i \rightarrow \prod_{i=1}^r G_i/H_i$ by f . Suppose $\mu \in M_G(G_1, \dots, G_r)$, and $\mu(G_1 \times \dots \times H_i \times \dots \times G_r) \subset H$ for $i=1, \dots, r$. Then there exists a unique $\nu \in M_{G/H}(G_1/H_1, \dots, G_r/H_r)$ with the property that $\pi\mu = \nu f$.*

Proof. It is easy to see that for (x_1, \dots, x_r) and $(y_1, \dots, y_r) \in \prod_{i=1}^r G_i$,

$$\mu(x_1, \dots, x_r) - \mu(y_1, \dots, y_r) = \sum_{i=1}^r \mu(y_1, \dots, y_{i-1}, x_i - y_i, x_{i+1}, \dots, x_r).$$

Hence if $x_i + H_i = y_i + H_i$ for $i = 1, \dots, r$, then

$$\mu(y_1, \dots, y_{i-1}, x_i - y_i, x_{i+1}, \dots, x_r) \in H$$

for each i , and therefore $\sum_{i=1}^r \mu(y_1, \dots, y_{i-1}, x_i - y_i, x_{i+1}, \dots, x_r) \in H$, that is, $\mu(x_1, \dots, x_r) + H = \mu(y_1, \dots, y_r) + H$. Thus the equation $\pi\mu = \nu f$ yields a well defined ν . Because μ is r -linear, ν is also r -linear. Since π and μ are continuous and f is open, it follows that ν is continuous. Clearly ν is unique.

COROLLARY. *In particular, if $\mu(G_1 \times \dots \times H_i \times \dots \times G_r) = (0)$ for all i then there is a unique $\nu \in M_G(G_1/H_1, \dots, G_r/H_r)$ such that $\nu f = \mu$.*

DEFINITION. Let G_1, \dots, G_r be in \mathcal{L} and for each i let H_i be a closed subgroup of G_i . Denote by $[H_1, \dots, H_r]$ the closure of the subgroup of $\bigotimes_{i=1}^r G_i$ generated by $\{\phi(x_1, \dots, x_r) : x_i \in H_i \text{ for some } i = 1, \dots, r\}$, where ϕ is the tensor map on $\prod_{i=1}^r G_i$.

THEOREM 4.9. *If G_1, \dots, G_r are compactly generated then $\bigotimes_{i=1}^r (G_i/H_i)$ is isomorphic with $\bigotimes_{i=1}^r G_i/[H_1, \dots, H_r]$.*

Proof. Since $\phi(G_1 \times \dots \times H_i \times \dots \times G_r) \subset [H_1, \dots, H_r]$ for $i = 1, \dots, r$ and ϕ is a continuous r -linear map, Theorem 4.8 shows that there is a continuous r -linear map $\nu: \prod_{i=1}^r G_i/H_i \rightarrow \bigotimes_{i=1}^r G_i/[H_1, \dots, H_r]$ with the property that $\pi\phi = \nu f$ where $f: \prod_{i=1}^r G_i \rightarrow \prod_{i=1}^r G_i/H_i$ and $\pi: \bigotimes_{i=1}^r G_i \rightarrow \bigotimes_{i=1}^r G_i/[H_1, \dots, H_r]$ are the canonical epimorphisms. Let ϕ' be the tensor map on $\prod_{i=1}^r G_i/H_i$. By Theorem 4.6 there exists a continuous linear map $\chi_\nu: \bigotimes_{i=1}^r G_i/H_i \rightarrow \bigotimes_{i=1}^r G_i/[H_1, \dots, H_r]$ satisfying $\chi_\nu\phi' = \nu$. Similarly, since $\phi'f$ is continuous r -linear there is a continuous linear map $\xi: \bigotimes_{i=1}^r G_i \rightarrow \bigotimes_{i=1}^r G_i/H_i$ with the property that $\xi\phi = \phi'f$ so that $\chi_\nu\xi\phi = \chi_\nu\phi'f = \nu f = \pi\phi$. Since $M_T(G_1, \dots, G_r)$ is locally compact, by Theorem 4.1, it follows from Theorem 4.6 that $\chi_\nu\xi = \pi$. Because ξ is continuous and π is open, χ_ν is open. Since π is surjective, so is χ_ν . It remains to show that χ_ν is a monomorphism.

Let $\sum_j \phi(x_1^{(j)}, \dots, x_r^{(j)}) \in \bigotimes_{i=1}^r G_i$ where for each j one of the $x_i^{(j)} \in H_i$ for some $i = 1, \dots, r$. Then

$$\xi\left(\sum_j \phi(x_1^{(j)}, \dots, x_r^{(j)})\right) = \sum_j \xi\phi(x_1^{(j)}, \dots, x_r^{(j)}) = \sum_j \phi'f(x_1^{(j)}, \dots, x_r^{(j)}) = \sum_j 0 = 0$$

since for each j , $\phi'f(x_1^{(j)}, \dots, x_r^{(j)}) = \phi'(\dots, 0, \dots)$ and ϕ' is r -linear. Since ξ is continuous it follows that $\xi[H_1, \dots, H_r] = (0)$. Hence there is a continuous linear map $\xi^-: \bigotimes_{i=1}^r G_i/[H_1, \dots, H_r] \rightarrow \bigotimes_{i=1}^r G_i/H_i$ satisfying $\xi^-\pi = \xi$, so that $\xi^-\chi_\nu\xi\phi = \xi^-\pi\phi = \xi\phi$. Hence $\xi^-\chi_\nu\xi = \xi$ by Theorem 4.6, and therefore $\xi^-\chi_\nu$ is the identity on $\xi(\bigotimes_{i=1}^r G_i)$. Now, $\xi(\bigotimes_{i=1}^r G_i) \supset \xi\phi(\prod_{i=1}^r G_i) = \phi'f(\prod_{i=1}^r G_i) = \phi'(\prod_{i=1}^r G_i/H_i)$

since f is surjective. Consequently $\xi(\bigotimes_{i=1}^r G_i)$ contains the subgroup D' of $\bigotimes_{i=1}^r G_i/H_i$ generated by $\phi'(\prod_{i=1}^r G_i/H_i)$. Since G_i/H_i is compactly generated, by Theorem 2.6, $M_T(G_1/H_1, \dots, G_r/H_r)$ is locally compact, by Theorem 4.1, and hence D' is dense, by Theorem 4.5. Thus $\xi(\bigotimes_{i=1}^r G_i)$ is dense. Since $\xi^{-1}\chi_v$ is the identity on a dense subgroup it equals the identity everywhere, and therefore χ_v is a monomorphism. This completes the proof.

COROLLARY. *Under the hypothesis of Theorem 4.9, if the H_i are all open then $[H_1, \dots, H_r]$ is open.*

Proof. If the H_i are open then each G_i/H_i is discrete and hence $\bigotimes_{i=1}^r G_i/H_i$ is discrete. By Theorem 4.9, $\bigotimes_{i=1}^r G_i/[H_1, \dots, H_r] \cong \bigotimes_{i=1}^r G_i/H_i$. Therefore, $[H_1, \dots, H_r]$ is open.

PROPOSITION 4.1. *Let G be compactly generated, D discrete and R the additive group of real numbers. Then $\text{Hom}(G, R) \cong R^{n+m}$ and $\text{Hom}(G, D) \cong D^m \oplus \text{Hom}(C, D)$ where $G = R^n \oplus Z^m \oplus C$ is the canonical representation of G in terms of its maximum compact subgroup C .*

Proof. By Corollary 2 of Theorem 4.2, $\text{Hom}(G, R) \cong \text{Hom}(R^\wedge, G^\wedge)$. However, $G^\wedge \cong R^n \oplus T^m \oplus C^\wedge$ and $R^\wedge \cong R$. A previous remark yields $\text{Hom}(G, R) \cong \text{Hom}(R, R)^n \oplus \text{Hom}(R, T)^m \oplus \text{Hom}(R, C^\wedge)$. It is clear that $\text{Hom}(R, R) \cong R$. Since R is connected and C^\wedge is discrete, $\text{Hom}(R, C^\wedge) = (0)$. Thus $\text{Hom}(G, R) \cong R^n \oplus (R^\wedge)^m \cong R^{n+m}$.

Similarly, $\text{Hom}(G, D) \cong \text{Hom}(D^\wedge, G^\wedge)$, which is isomorphic to $\text{Hom}(D^\wedge, R)^n \oplus \text{Hom}(D^\wedge, T)^m \oplus \text{Hom}(D^\wedge, C^\wedge)$. It is clear that $\text{Hom}(D^\wedge, R) = (0)$, $\text{Hom}(D^\wedge, T) \cong D$ and $\text{Hom}(D^\wedge, C^\wedge) \cong \text{Hom}(C, D)$. Thus $\text{Hom}(G, D) \cong D^m \oplus \text{Hom}(C, D)$. This completes the proof.

THEOREM 4.10. *If G_1 is a closed subgroup of a compactly generated group G_2 and H has no small subgroups then the continuous restriction homomorphism*

$$f: \text{Hom}(G_2, H) \rightarrow \text{Hom}(G_1, H)$$

is open.

Proof. Since H has no small subgroups, $H \cong R^n \oplus T^m \oplus D$ where D is a discrete group. As we have observed before, $\text{Hom}(G_i, H)$ is naturally isomorphic with $\text{Hom}(G_i, R)^n \oplus \text{Hom}(G_i, T)^m \oplus \text{Hom}(G_i, D)$. An easy direct sum argument shows that it suffices to prove that the following partial maps are open

$$f_1: \text{Hom}(G_2, R) \rightarrow \text{Hom}(G_1, R),$$

$$f_2: \text{Hom}(G_2, T) \rightarrow \text{Hom}(G_1, T),$$

$$f_3: \text{Hom}(G_2, D) \rightarrow \text{Hom}(G_1, D).$$

(1) Since R is injective, f_1 is an epimorphism. Both $\text{Hom}(G_1, R)$ and $\text{Hom}(G_2, R)$ are locally compact, by a corollary to Theorem 4.1. In fact, Proposition 4.1 shows

that $\text{Hom}(G_2, R)$ is a vector group. The open mapping theorem guarantees that f_1 is open.

(2) It is easy to see that $f_2 = i^\wedge$ where $i: G_1 \rightarrow G_2$ is the inclusion of G_1 into G_2 . Hence f_2 is a proper epimorphism.

(3) $\text{Hom}(G_1, D) \cong D^m \oplus \text{Hom}(C, D)$, by Proposition 4.1. But the latter is isomorphic to $D^m \oplus M_T(C, D^\wedge)$. By Theorem 4.1, $M_T(C, D^\wedge)$ is discrete, because C and D^\wedge are compact. Thus $\text{Hom}(G_1, D)$ is a finite direct product of discrete groups. Since $\text{Hom}(G_1, D)$ is discrete, f_3 is an open map.

V. Tensor products of homomorphisms, exactness, and structural properties.

Let G_1, \dots, G_r and H_1, \dots, H_r be in \mathcal{L} and assume $M_T(G_1, \dots, G_r)$ and $M_T(H_1, \dots, H_r)$ are locally compact. Denote the respective tensor maps by $\phi: \prod_{i=1}^r G_i \rightarrow \bigotimes_{i=1}^r G_i$ and $\phi': \prod_{i=1}^r H_i \rightarrow \bigotimes_{i=1}^r H_i$. If for each $i=1, \dots, r$ we are given a continuous homomorphism $f_i: G_i \rightarrow H_i$, define the map

$$[f_1, \dots, f_r]: \prod_{i=1}^r G_i \rightarrow \prod_{i=1}^r H_i$$

by $[f_1, \dots, f_r](x_1, \dots, x_r) = (f_1(x_1), \dots, f_r(x_r))$. It is clear that $\phi'[f_1, \dots, f_r]$ is a continuous r -linear map $\prod_{i=1}^r G_i \rightarrow \bigotimes_{i=1}^r H_i$. Since $M_T(H_1, \dots, H_r)$ is in \mathcal{L} , so is $\bigotimes_{i=1}^r H_i$. Hence since $M_T(G_1, \dots, G_r)$ is in \mathcal{L} there exists, by Theorem 4.6, a unique continuous linear map denoted by $f_1 \otimes \dots \otimes f_r: \bigotimes_{i=1}^r G_i \rightarrow \bigotimes_{i=1}^r H_i$ satisfying $(f_1 \otimes \dots \otimes f_r) \cdot \phi = \phi'[f_1, \dots, f_r]$.

If, for $i=1, \dots, r$, $g_i: G_i \rightarrow H_i$ is another family of continuous homomorphisms then for each i , $f_1 \otimes \dots \otimes f_i + g_i \otimes \dots \otimes f_r = f_1 \otimes \dots \otimes f_i \otimes \dots \otimes f_r + f_1 \otimes \dots \otimes g_i \otimes \dots \otimes f_r$. These equations are derived by observing that the functions on either side agree on $\phi(\prod_{i=1}^r G_i)$ and hence, by Theorem 4.6, on $\bigotimes_{i=1}^r G_i$. In particular, if $f_i = 0$ for some i then $f_1 \otimes \dots \otimes f_r = 0$. It is clear that if id_{G_i} is the identity map on G_i then $\text{id}_{G_1} \otimes \dots \otimes \text{id}_{G_r} = \text{id}_{\bigotimes_{i=1}^r G_i}$. Given continuous homomorphisms

$$G_i \xrightarrow{f_i} H_i \xrightarrow{g_i} L_i$$

where $i=1, \dots, r$, a similar application of Theorem 4.6 shows that $g_1 f_1 \otimes \dots \otimes g_r f_r = (g_1 \otimes \dots \otimes g_r) \cdot (f_1 \otimes \dots \otimes f_r)$. If $f_i: G_i \rightarrow H_i$ is an isomorphism then $f_1 \otimes \dots \otimes f_r$ is an isomorphism and $(f_1 \otimes \dots \otimes f_r)^{-1} = f_1^{-1} \otimes \dots \otimes f_r^{-1}$.

Dually, we define $[f_1, \dots, f_r]^*: M_T(H_1, \dots, H_r) \rightarrow M_T(G_1, \dots, G_r)$ by

$$[f_1, \dots, f_r]^*(\mu) = \mu \cdot [f_1, \dots, f_r]$$

for $\mu \in M_T(H_1, \dots, H_r)$. One sees easily that $\omega[f_1, \dots, f_r]^* = (f_1 \otimes \dots \otimes f_r)^\wedge \omega'$ where ω denotes the natural isomorphism $M_T(G_1, \dots, G_r) \rightarrow M_T(G_1, \dots, G_r)^\wedge$ and ω' the natural isomorphism $M_T(H_1, \dots, H_r) \rightarrow M_T(H_1, \dots, H_r)^\wedge$. Hence $[f_1, \dots, f_r]^*$ is the continuous homomorphism dual to $f_1 \otimes \dots \otimes f_r$. Whenever it is clear which maps f_1, \dots, f_r are being considered we will abbreviate $[f_1, \dots, f_r]^*$ by \bullet , and the image of a subset S of $M_T(H_1, \dots, H_r)$ in $M_T(G_1, \dots, G_r)$ by S^\bullet .

THEOREM 5.1. *Let $M_T(G_1, \dots, G_r)$ and $M_T(H_1, \dots, H_r)$ be locally compact. If f_1, \dots, f_r are proper epimorphisms then $f_1 \otimes \dots \otimes f_r$ is a proper epimorphism.*

Proof. Clearly $[f_1, \dots, f_r]$ is an epimorphism. Consequently,

$$(f_1 \otimes \dots \otimes f_r) \phi \left(\prod_{i=1}^r G_i \right) = \phi' [f_1, \dots, f_r] \left(\prod_{i=1}^r G_i \right) = \phi' \left(\prod_{i=1}^r H_i \right).$$

It follows by linearity that $f_1 \otimes \dots \otimes f_r(D) = D'$ where D and D' are the subgroups of $\bigotimes_{i=1}^r G_i$ and $\bigotimes_{i=1}^r H_i$ generated by $\phi(\prod_{i=1}^r G_i)$ and $\phi'(\prod_{i=1}^r H_i)$ respectively. Hence $f_1 \otimes \dots \otimes f_r(\bigotimes_{i=1}^r G_i)$ is dense in $\bigotimes_{i=1}^r H_i$. This implies that \bullet is a (continuous) monomorphism. We show that \bullet is proper. Let $W(F, \Gamma)$ be an open neighborhood of 0 in $M_T(H_1, \dots, H_r)$ where F is compact. Clearly F is contained in $\prod_{i=1}^r F_{H_i}$, a compact box. Hence $W(F, \Gamma)^\bullet \supset W(\prod_{i=1}^r F_{H_i}, \Gamma)^\bullet$. It is sufficient to show that $W(\prod_{i=1}^r F_{H_i}, \Gamma)^\bullet$ contains the intersection of $M_T(H_1, \dots, H_r)^\bullet$ with an open neighborhood of 0 in $M_T(G_1, \dots, G_r)$. Thus we may assume $F = \prod_{i=1}^r F_{H_i}$. Let U_i be a compact neighborhood of 0 in G_i . Since f_i is a proper epimorphism, $\{f_i(x_i + U_i^\circ) : x_i \in G_i\}$ is an open covering of H_i and hence of F_{H_i} . By compactness $F_{H_i} \subset \bigcup_{j=1}^{n_i} f_i(x_i^{(j)} + U_i^\circ)$ where $x_i^{(j)} \in G_i$. Let

$$F_{G_i} = \bigcup_{j=1}^{n_i} x_i^{(j)} + U_i.$$

It is clear that F_{G_i} is compact and $f_i(F_{G_i}) \supset F_{H_i}$. Evidently, $W(\prod_{i=1}^r F_{H_i}, \Gamma)^\bullet \supset M_T(H_1, \dots, H_r)^\bullet \cap W(\prod_{i=1}^r F_{G_i}, \Gamma)$. Thus \bullet is a proper monomorphism and hence $f_1 \otimes \dots \otimes f_r$ is a proper epimorphism, by Theorem 2.1. This completes the proof.

Henceforth we restrict ourselves to the case $r=2$ and slightly change the notation.

PROPOSITION 5.1. *If $g: G_1 \rightarrow G_2$ and $h: H_1 \rightarrow H_2$ are proper epimorphisms where G_i and $H_i \in \mathcal{L}$ for $i=1, 2$, then $M_T(G_2, H_2)^\bullet$ is the set of $\nu \in M_T(G_1, H_1)$ satisfying $\nu(G_1 \times \text{Ker } h) = (0) = \nu(\text{Ker } g \times H_1)$.*

Proof. Suppose $\nu \in M_T(G_2, H_2)^\bullet$. If $x \in \text{Ker } g$, then $\nu(x, y) = \mu(0, h(y)) = (0)$, since μ is bilinear. Hence $\nu(\text{Ker } g \times H_1) = (0)$. Similarly, $\nu(G_1 \times \text{Ker } h) = (0)$. On the other hand, if $\nu \in M_T(G_1, H_1)$ and $\nu(G_1 \times \text{Ker } h) = (0) = \nu(\text{Ker } g \times H_1)$, then by Theorem 4.8, ν induces a map μ in $M_T(G_1/\text{Ker } g, H_1/\text{Ker } h)$ with the property that $\mu f = \nu$, where f is the map from $G_1 \times H_1 \rightarrow (G_1/\text{Ker } g) \times (H_1/\text{Ker } h)$, defined by $f(x, y) = (x + \text{Ker } g, y + \text{Ker } h)$. Since $G_2 \cong G_1/\text{Ker } g$, and $H_2 \cong H_1/\text{Ker } h$, after identification via these isomorphisms, $f = [g, h]$ and therefore, $\nu = \mu[g, h]$. This completes the proof.

THEOREM 5.2. *Let*

$$0 \longrightarrow G_1 \xrightarrow{g_1} G_2 \xrightarrow{g_2} G_3 \longrightarrow 0$$

be a proper exact sequence with G_1 and G_3 , or equivalently G_2 , compactly generated. Let H be compactly generated. Then

$$G_1 \otimes H \xrightarrow{g_1 \otimes \text{id}_H} G_2 \otimes H \xrightarrow{g_2 \otimes \text{id}_H} G_3 \otimes H \longrightarrow (0)$$

is a proper exact sequence.

Proof. By Theorem 2.1, it is sufficient to show that

$$0 \longrightarrow M_T(G_3, H) \xrightarrow{\bullet_2} M_T(G_2, H) \xrightarrow{\bullet_1} M_T(G_1, H)$$

is proper exact. By Theorem 5.1, \bullet_2 is a proper monomorphism. Since $\text{Ker } \bullet_1$ is the set of all $\mu \in M_T(G_2, H)$ that annihilate $g_1(G_1) \times H = \text{Ker } g_2 \times H$, it follows from Proposition 5.1 that $\text{Ker } \bullet_1 = M_T(G_3, H)^{\bullet_2}$. Because of the isomorphism of functors established in §IV, it follows from Theorem 4.10 that \bullet_1 is proper. This completes the proof.

COROLLARY. Let G be compactly generated and Z_n be the cyclic group of order n . Then $Z_n \otimes G \cong G/nG$.

Proof. Consider the proper exact sequence $0 \rightarrow nZ \rightarrow Z \rightarrow Z_n \rightarrow 0$. By Theorem 5.2, $nZ \otimes G \rightarrow Z \otimes G \rightarrow Z_n \otimes G \rightarrow 0$ is proper exact. Hence $Z_n \otimes G \cong Z \otimes G / \text{image of } nZ \otimes G$. However as we noted earlier this may be identified with G/nG . Note that nG is a closed subgroup of G since G is compactly generated. This completes the proof.

Now let \mathcal{C} and \mathcal{N} , respectively, denote the categories of groups in \mathcal{L} that are compactly generated and have no small subgroups. We define four families of functors of G . These are defined only on the categories and only for those H 's indicated below. The mappings are those already used several times.

- (1) $* \otimes H: \mathcal{C} \rightarrow \mathcal{L}$, where H is in \mathcal{C} . This is a covariant additive functor.
- (2) $M_T(*, H): \mathcal{C} \rightarrow \mathcal{L}$, where H is in \mathcal{C} . This is a contravariant additive functor.
- (3) $\text{Hom}(*, H): \mathcal{C} \rightarrow \mathcal{L}$, where H is in \mathcal{N} . This is a contravariant additive functor.
- (4) $\text{Hom}(H, *): \mathcal{N} \rightarrow \mathcal{L}$, where H is in \mathcal{C} . This is a covariant additive functor.

DEFINITION. A covariant additive functor \mathcal{F} is called proper terminally exact if for all proper exact sequences $0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$ in the domain of \mathcal{F} , $\mathcal{F}(G_1) \rightarrow \mathcal{F}(G_2) \rightarrow \mathcal{F}(G_3) \rightarrow 0$ is a proper exact sequence, and is called proper initially exact if $0 \rightarrow \mathcal{F}(G_1) \rightarrow \mathcal{F}(G_2) \rightarrow \mathcal{F}(G_3)$ is a proper exact sequence.

Similarly a contravariant additive functor \mathcal{F} is called proper terminally exact if for all proper exact sequences $0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$ in the domain of \mathcal{F} , $\mathcal{F}(G_3) \rightarrow \mathcal{F}(G_2) \rightarrow \mathcal{F}(G_1) \rightarrow 0$ is a proper exact sequence, and is called proper initially exact if $0 \rightarrow \mathcal{F}(G_3) \rightarrow \mathcal{F}(G_2) \rightarrow \mathcal{F}(G_1)$ is a proper exact sequence. These definitions are continuous analogues of those given in [6].

REMARK. Throughout §V, we will utilize fully the fact that all notions and theorems dualize; in particular, that \mathcal{C} and \mathcal{N} and certain of their subclasses are

dual categories, that the dual of a short proper exact sequence is a short proper exact sequence, and that for G and H in \mathcal{C} , $(G \otimes H)^\wedge$, $M_T(G, H)$, $\text{Hom}(G, H^\wedge)$ and $\text{Hom}(H, G^\wedge)$ are isomorphic functors. Thus one proof replaces four.

In proving Theorem 5.2, we have actually proven

THEOREM 5.3. *The functor $* \otimes H$ is a proper terminally exact functor, and $M_T(*, H)$, $\text{Hom}(*, H)$, and $\text{Hom}(H, *)$ are proper initially exact functors.*

COROLLARY. *Let G_1 and G_2 be in \mathcal{C} and H in \mathcal{N} . If $g: G_1 \rightarrow G_2$ is a proper homomorphism then the induced map $g^*: \text{Hom}(G_2, H) \rightarrow \text{Hom}(G_1, H)$ obtained by composition is a proper homomorphism.*

Let G_1, G_2 and H be in \mathcal{C} . If $g: G_1 \rightarrow G_2$ is a proper homomorphism then the induced maps $G_1 \otimes H \rightarrow G_2 \otimes H$ and $M_T(G_2, H) \rightarrow M_T(G_1, H)$ are proper homomorphisms.

Let G_1 and G_2 be in \mathcal{N} and H in \mathcal{C} . If $g: G_1 \rightarrow G_2$ is a proper homomorphism then the induced map $\text{Hom}(H, G_1) \rightarrow \text{Hom}(H, G_2)$ is a proper homomorphism.

The above Corollary generalizes Theorem 4.10.

Proof. Since $g(G_1)$ is a closed subgroup of G_2 , we can write $g = i\pi$ where $i: g(G_1) \rightarrow G_2$ and $\pi: G_1 \rightarrow g(G_1)$ are proper homomorphisms; i is injective and π surjective. Hence $g^* = \pi^* i^*$. It follows from Theorem 4.10 that i^* is an open map. Thus, to show that g^* is proper, it is sufficient to show that π^* is proper, that is, we are reduced to proving the theorem in the case where g is surjective. In this case, consider the proper exact sequence $0 \rightarrow \text{Ker } g \rightarrow G_1 \rightarrow G_2 \rightarrow 0$. It follows from Theorem 5.3 that

$$0 \longrightarrow \text{Hom}(G_2, H) \xrightarrow{g^*} \text{Hom}(G_1, H) \longrightarrow \text{Hom}(\text{Ker } g, H)$$

is a proper exact sequence. In particular, g^* is a proper homomorphism. The proof is completed by dualization.

DEFINITION. A covariant or contravariant additive functor \mathcal{F} is called proper exact if it is both proper terminally exact and proper initially exact (see [6]).

THEOREM 5.4. *$* \otimes H$, $M_T(*, H)$ and $\text{Hom}(H, *)$ are proper exact functors if and only if the maximum compact subgroup of H is (0) . $\text{Hom}(*, H)$ is a proper exact functor if and only if H is connected.*

Proof. $\text{Hom}(*, H)$ is proper exact if and only if H is injective for the class of compactly generated groups. As we have already proven, the injectives for \mathcal{L} are the same as those for compactly generated groups, namely, $R^n \oplus T^\sigma$, direct sums of vector groups and tori. H is a group without small subgroups; hence, σ is finite. So $\text{Hom}(*, H)$ is a proper exact functor if and only if $H = R^n \oplus T^m$, i.e., if and only if H is a connected group without small subgroups. Consequently, because of the isomorphism of various functors, and Theorems 2.1 and 2.5 Corollary 2, the result follows.

COROLLARY. *We remark that $\text{Hom}(*, H)$ is proper exact if and only if it is proper exact relative to the sequence $0 \rightarrow Z \rightarrow R \rightarrow T \rightarrow 0$, for, as we saw earlier, a group which is injective for this sequence must be connected, and hence is injective for all sequences, by Theorem 5.4. Since the sequence $0 \rightarrow Z \rightarrow R \rightarrow T \rightarrow 0$ is self dual, the same statements are true for $* \otimes H$, $M_T(*, H)$ and $\text{Hom}(H, *)$.*

DEFINITION. A functor preserving proper exactness for a class of extensions is called a proper exact functor relative to that class.

THEOREM 5.5. *Relative to sequences $0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$, where G_1 is open in G_2 , $* \otimes H$ and $M_T(*, H)$ are proper exact if and only if H is torsion free. $\text{Hom}(*, H)$ is proper exact if and only if H is divisible. Relative to sequences $0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$, where G_1 is compact, $\text{Hom}(H, *)$ is proper exact if and only if H is torsion free.*

Proof. By dualization, it is sufficient to prove that $\text{Hom}(*, H)$ is proper exact if and only if H is divisible. If $\text{Hom}(*, H)$ is proper exact relative to the sequences $0 \rightarrow nZ \rightarrow Z \rightarrow Z_n \rightarrow 0$ then evidently H is divisible. The converse follows from the fact that G_1 is open in G_2 .

THEOREM 5.6. *Relative to sequences $0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$ with each G_i compact, $* \otimes H$ and $M_T(*, H)$ are proper exact if and only if the maximum compact subgroup of H is connected. $\text{Hom}(*, H)$ is proper exact if and only if H/H_0 is torsion free. Relative to sequences $0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$, of discrete groups, $\text{Hom}(H, *)$ is proper exact if and only if the maximum compact subgroup of H is connected.*

Proof. If H is connected, $\text{Hom}(*, H)$ is proper exact, by Theorem 5.4. Thus we are led to the case of $H = D$, a discrete group. D is torsion free; for suppose $y \in D$, and the order of $y = n$. Consider the sequence $0 \rightarrow Z_n \rightarrow T \rightarrow T \rightarrow 0$. The natural map of Z_n into D cannot extend to T since T is connected. Conversely, if D is a torsion free discrete group, and $\xi: G_1 \rightarrow D$ is a continuous homomorphism, then $\xi(G_1)$ is compact and hence finite, since D is discrete. Because D is torsion free, ξ is trivial and extends trivially. Dualizing, Theorem 5.6 is completely proven.

THEOREM 5.7. *Relative to sequences $0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$, where G_2 is connected and G_1 is a pure subgroup of G_2 , $* \otimes H$ and $M_T(*, H)$ are proper exact functors for all H in \mathcal{C} . $\text{Hom}(*, H)$ is a proper exact functor for all H in \mathcal{N} . Relative to sequences $0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$ where G_2 and G_3 are torsion free, $\text{Hom}(H, *)$ is a proper exact functor for all H in \mathcal{C} .*

Proof. If H is connected, $\text{Hom}(*, H)$ is proper exact, by Theorem 5.4. Thus, we are led to the case $H = D$, a discrete group. Now, by Corollary 1 of Theorem 2.9, G_1 is connected. Hence any continuous homomorphism $G_1 \rightarrow D$ is trivial and extends trivially. The proof is completed by dualization.

Next we give an explicit computation of $\text{Hom}(G_1, G_2)$, $G_1 \otimes G_2$, and $M_T(G_1, G_2)$ in terms of known invariants of G_1 and G_2 .

THEOREM 5.8. *Let G be compactly generated and H have no small subgroups. Let their canonical decompositions be $G = R^n \oplus Z^m \oplus C$ and $H = R^k \oplus T^l \oplus D$. Then $\text{Hom}(G, H) \cong R^{(n+m)k} \oplus (G^\wedge)^l \oplus D^m \oplus \text{Hom}(C, D)$.*

Dually, if G_1 and G_2 are compactly generated groups whose decompositions for $i=1, 2$ are $G_i = R^{n_i} \oplus Z^{m_i} \oplus C_i$, where C_i is the maximal compact subgroup, then

$$G_1 \otimes G_2 \cong R^{(n_1+m_1)n_2} \oplus G_1^{m_2} \oplus C_2^{m_1} \oplus C_1 \otimes C_2,$$

and

$$M_T(G_1, G_2) \cong R^{(n_1+m_1)n_2} \oplus (G_1^\wedge)^{m_2} \oplus (C_2^\wedge)^{m_1} \oplus M_T(C_1, C_2).$$

Proof. The first part follows directly from the additive properties of Hom , and Proposition 4.1. The second part now follows by dualization.

We use Theorem 5.8 to compute some specific examples and give the significance of tensor products of various groups. The first such result gives a sharpening of Theorem 4.4.

COROLLARY 1. *If G_1 and G_2 are in \mathcal{C} then so is $G_1 \otimes G_2$.*

Proof. The result follows from Theorem 5.8, since $C_1 \otimes C_2$ is compact by Theorem 4.4.

COROLLARY 2. *If G_1 is connected and G_2 is compactly generated then*

$$G_1 \otimes G_2 \cong R^{n_1 n_2} \oplus G_1^{m_2}.$$

Proof. G_1 is connected if and only if $m_1=0$ and C_1 is connected. In this case $C_2^{m_1}=(0)$ and $(C_1 \otimes C_2)^\wedge \cong \text{Hom}(C_1, C_2^\wedge)=(0)$. Hence $C_1 \otimes C_2=(0)$. Thus $G_1 \otimes G_2 \cong R^{n_1 n_2} \oplus G_1^{m_2}$. From this we conclude several facts:

(1) If G_1 is connected and G_2 is compactly generated then $G_1 \otimes G_2$ is connected. This is clear.

(2) If G_1 is connected and G_2 is compact then $G_1 \otimes G_2=(0)$. G_2 is compact if and only if $n_2=m_2=0$. The result follows.

(3) If G_1 is connected and G_2 is compact then $M_G(G_1, G_2)=(0)$ for all locally compact commutative groups G .

For $G=T$, we see that $M_T(G_1, G_2) \cong (G_1 \otimes G_2)^\wedge=(0)$. Now let G be any locally compact group and ξ a character of G . If $\mu \in M_G(G_1, G_2)$ then $\xi\mu \in M_T(G_1, G_2)$, so that $\xi\mu(x_1, x_2)=0$ for all characters ξ . Since the characters of G separate points, $\mu=0$.

(4) If G_1 and G_2 are connected then $G_1 \otimes G_2 \cong R^{n_1 n_2}$. The tensor product of connected groups of vector dimensions n_1 and n_2 is isomorphic to the vector group of dimension $n_1 n_2$.

(5) If G has no small subgroups then $T \otimes G^\wedge$ is (naturally \cong with) the maximal toral subgroup of G .

We take $G_1=T$ and $G_2=G^\wedge$, a compactly generated group. Since $n_1=m_1=0$ and $C_1=T$, it follows that $T \otimes G^\wedge = T^{m_2}$, which is the maximal toral subgroup of G .

COROLLARY 3. *If G is a group without small subgroups then $\text{Hom}(R, G)$, the group of one parameter subgroups of G , together with $e: \text{Hom}(R, G) \rightarrow G$, the evaluation at 1, is the simply connected covering group of G_0 .*

Proof. $G = R^n \oplus T^m \oplus D$. Since $\text{Hom}(R, G) \cong M_T(R, G^\wedge)$, taking $n_2 = 1$, $m_2 = 0$ and $C_2 = (0)$, we see that $\text{Hom}(R, G) \cong R^{n+m} = R^{\dim G}$. Since uniform convergence on compact subsets implies pointwise convergence, the evaluation map e is a continuous homomorphism. Because $\text{Hom}(R, G)$ is a vector group, it is connected. This means that $e(\text{Hom}(R, G))$ is contained in G_0 . Since G_0 is connected, the corollary to Theorem 5.4 shows that $\text{Hom}(*, G_0)$ is proper exact for the sequence $0 \rightarrow Z \rightarrow R \rightarrow T \rightarrow 0$. Hence each point of G_0 lies on a one parameter subgroup, and e is surjective, as a map to G_0 . Since both groups are compactly generated the Open Mapping Theorem shows that e is proper. A comparison of the dimensions of $\text{Hom}(R, G)$ and G_0 shows that $\text{Ker } e$ is discrete, and so e is a covering map. Since $\text{Hom}(R, G)$ is a vector group the result follows.

Now we turn to $G \otimes H$ where G and H are compact.

PROPOSITION 5.1. *Let G be in \mathcal{C} and H be compact. Then $G \otimes H \cong (G/G_0) \otimes H$.*

Proof. Consider the proper exact sequence $0 \rightarrow G_0 \rightarrow G \rightarrow G/G_0 \rightarrow 0$. It follows that $G_0 \otimes H \rightarrow G \otimes H \rightarrow (G/G_0) \otimes H \rightarrow 0$ is proper exact. However, $(G_0 \otimes H)^\wedge \cong \text{Hom}(G_0, H^\wedge)$. Since G_0 is connected and H^\wedge is discrete, $\text{Hom}(G_0, H^\wedge) = (0)$, and therefore $G_0 \otimes H = (0)$. This means that $0 \rightarrow G \otimes H \rightarrow (G/G_0) \otimes H \rightarrow 0$ is proper exact, and consequently $G \otimes H \cong (G/G_0) \otimes H$.

COROLLARY. *If G and H are compact then $G \otimes H \cong (G/G_0) \otimes (H/H_0)$.*

Proof. By repeating the above argument we get $(G/G_0) \otimes H \cong (G/G_0) \otimes (H/H_0)$, and hence $G \otimes H \cong (G/G_0) \otimes (H/H_0)$, the tensor product of compact totally disconnected groups.

This suggests utilization of results of Braconnier [5], some of which, however, go back to Krull and others.

We denote by I_p the additive group of the p -adic integers, by (Z, \parallel_p) the subgroup of integers with the p -adic topology and by $Z(p^n)$ the cyclic group of order p^n . I_p is a compact commutative totally disconnected group.

Now we give the definition of a p -group in [5], and summarize some of the basic properties we shall utilize. Let G be in \mathcal{L} and define $f: Z \times G \rightarrow G$ by $f(n, x) = nx$.

DEFINITION. G is a p -group if for each fixed $x \in G$, f is continuous in n , where Z takes the p -adic topology. If G is discrete, this notion coincides with the usual one of a p -group in abstract abelian groups (see [3]). If G is a p -group then f extends to a continuous bilinear map, say $\mu: I_p \times G \rightarrow G$. G is a p -group if and only if G^\wedge is a p -group. I_p is a p -group.

DEFINITION. Let $G \in \mathcal{L}$. G_p is the set of those x 's in G for which $f(*, x)$ is a continuous function of n , where Z takes the p -adic topology. G_p is called the p -primary component of G .

The G_p 's are disjoint p subgroups of G . If G is totally disconnected then G_p is closed. If G is compact and totally disconnected then $G \cong \prod_p G_p$, the direct product of the G_p 's (product topology). This is called the primary decomposition of a compact totally disconnected group [5]. It is dual to the primary decomposition of a discrete torsion group [3].

PROPOSITION 5.2. *If G is a compact p -group, then $I_p \otimes G \cong G$.*

Proof. Let μ be the G -valued bilinear map on $I_p \times G$, derived from the fact that G is a locally compact p -group. Let χ_μ be the induced continuous homomorphism satisfying $\chi_\mu \phi = \mu$. χ_μ is surjective, since $\chi_\mu(I_p \otimes G) \supset \chi_\mu \phi(I_p \times G) = \mu(I_p \times G) \supset \mu((Z, \parallel_p) \times G)$. But $\mu(1, g) = g$, so that $\mu((Z, \parallel_p) \times G) \supset G$. Thus $\chi_\mu(I_p \otimes G) = G$. Since I_p and G are compact; so is $I_p \otimes G$. Thus the Open Mapping Theorem yields that χ_μ is open. We consider $\chi_\mu^\wedge: G^\wedge \rightarrow M_T(I_p, G)$, defined by $\chi_\mu^\wedge(\xi) = \xi \circ \mu$ for $\xi \in G^\wedge$, and show that χ_μ^\wedge is surjective. Suppose $\nu \in M_T(I_p, G)$. We consider the restriction of ν to $(Z, \parallel_p) \times G$. Let $\xi(x) = \nu(1, x)$. Clearly ξ is a character of G . Now $\xi\mu(n, x) = \xi(n, x) = n\xi(x) = n\nu(1, x) = \nu(n, x)$. So $\xi\mu = \nu$ on $(Z, \parallel_p) \times G$ which is dense in $I_p \times G$, because (Z, \parallel_p) is dense in I_p . Since $\xi\mu$ and ν are continuous, they are equal. Thus χ_μ^\wedge is surjective. Since G and $M_T(I_p, G)$ are locally compact, this implies that χ_μ is injective and hence an isomorphism.

PROPOSITION 5.3. *If G is any compact totally disconnected group then $I_p \otimes G \cong G_p$.*

Proof. Let $G \cong \prod_q G_q$ be the primary decomposition of G . $I_p \otimes G \cong \prod_q (I_p \otimes G_q)$. Now $I_p \otimes G_q = (0)$ for $q \neq p$. In fact, $(I_p \otimes G_q)^\wedge \cong \text{Hom}(I_p, G_q^\wedge)$ where G_q^\wedge is a discrete q group. If $\xi: I_p \rightarrow G_q^\wedge$ is a continuous homomorphism we know [5] that either (1) $\xi(I_p) \cong I_p$, or (2) $\xi(I_p) \cong Z(p^n)$ depending on whether ξ is injective or not. In any case, I_p is torsion free, and G_q^\wedge is a discrete q group. Hence (1) is impossible. Similarly, since $Z(p^n)$ has all its elements of order p^k , with $p \neq q$, (2) is impossible. Thus $\text{Hom}(I_p, G_q^\wedge) = (0)$, and therefore $I_p \otimes G_q = (0)$. Hence $I_p \otimes G \cong \prod_q I_p \otimes G_q \cong I_p \otimes G_p \cong G_p$, by Proposition 5.2.

THEOREM 5.9. *If G is any compact group, then*

- (1) $I_p \otimes G \cong (G/G_0)_p$, the p -primary component of G/G_0 .
- (2) $I_p \otimes I_q = (0)$, for $q \neq p$, and $I_p \otimes I_p \cong I_p$.
- (3) $G/G_0 \cong \prod_p (I_p \otimes G)$.

Proof. $I_p \otimes G \cong I_p \otimes (G/G_0)$ by Proposition 5.1. By Proposition 5.3, $I_p \otimes (G/G_0) \cong (G/G_0)_p$. Hence, $I_p \otimes G \cong (G/G_0)_p$. In particular, $I_p \otimes I_q = (0)$, for $q \neq p$, and $I_p \otimes I_p \cong I_p$. It follows that, for any compact group G , $G/G_0 \cong \prod_p I_p \otimes G$. This completes the proof.

COROLLARY. *Let G be in \mathcal{C} . Then $G \otimes H = (0)$ for all compact groups H if and only if G is connected.*

Proof. If H is compact and G is connected, then $G \otimes H = (0)$ by Corollary 3 of Theorem 5.8. Conversely, suppose $G \otimes H = (0)$ for all compact groups H .

We have $G \cong R^n \oplus Z^m \oplus C$, and hence $G \otimes H \cong (R \otimes H)^n \oplus (Z \otimes H)^m \oplus C \otimes H$. As was just noted, $R \otimes H = (0)$. Also, $Z \otimes H \cong H$. Since $G \otimes H = (0)$ for all H , it follows that $m = 0$. Because R^n is connected and $G \otimes H \cong C \otimes H$, we may assume that G is compact. By hypothesis, $I_p \otimes G = (0)$ for each prime p . Hence, by Theorem 5.9, $G/G_0 \cong \prod_p I_p \otimes G = (0)$. Thus $G = G_0$.

THEOREM 5.10. (1) *The tensor product of compact totally disconnected groups is a compact totally disconnected group whose p -primary component is $G_p \otimes H_p$.*

(2) *The tensor product of compact p -groups is a compact p -group.*

(3) *The tensor product of a compact p -group with a compact q -group is (0) for $p \neq q$.*

Proof. Let G and H be arbitrary compact groups. Then $G \otimes H$ is compact by Theorem 4.4. It is totally disconnected if and only if $(G \otimes H)^\wedge \cong \text{Hom}(G, H^\wedge)$ is a torsion group. Now if $f \in \text{Hom}(G, H^\wedge)$ then $f(G)$ is a finite subgroup of H^\wedge , since G is compact and H^\wedge is discrete. Let $n(f) = \text{order of } f(G)$. Then $n(f)f = 0$. Thus $(G \otimes H)^\wedge$ is a torsion group, whence $G \otimes H$ is compact and totally disconnected.

Now suppose, in addition, that G and H are totally disconnected. Let $G = \prod_p G_p$ and $H = \prod_q H_q$ be the primary decompositions of G and H , respectively. Then $G \otimes H \cong \prod_{p,q} G_p \otimes H_q$. Now if $p \neq q$ then

$$G_p \otimes H_q \cong I_p \otimes G \otimes I_q \otimes H \cong I_p \otimes I_q \otimes G \otimes H \cong (0) \otimes G \otimes H = (0).$$

Hence, $G \otimes H \cong \prod_p G_p \otimes H_p$. Moreover, $G_p \otimes H_p \cong I_p \otimes G \otimes I_p \otimes H \cong I_p \otimes I_p \otimes G \otimes H \cong I_p \otimes G \otimes H$. Since $G \otimes H$ is compact and totally disconnected, Proposition 5.3 shows that $I_p \otimes G \otimes H \cong (G \otimes H)_p$. Hence $(G \otimes H)_p \cong G_p \otimes H_p$, and $\prod_p G_p \otimes H_p$ is the primary decomposition of $G \otimes H$. (2) follows immediately from (1), and (3) was proven in the course of proving (1). This completes the proof.

Finally, Theorem 5.10 yields, in combination with the corollary to Proposition 5.1, the following result.

THEOREM 5.11. *If G and H are arbitrary compact groups then $G \otimes H$ is a compact totally disconnected group whose p -primary component is $(G/G_0)_p \otimes (H/H_0)_p$.*

VI. Ext and Tor. The standard homological algebra [6] applies in its entirety only to abelian categories. As we have noted in the Introduction, \mathcal{L} is not an abelian category. These difficulties have already been encountered in §§IV and V, where the functors Hom , \otimes and related functors were discussed. They were resolved by restricting the homomorphisms and the corresponding sequences to be proper, and by restricting the domains of definition of the various functors. In this section, we define the functors Ext and Tor as derived functors of Hom and \otimes by projective and injective resolutions. Similarly here, the resolutions employed must be proper. The domains of the functors Ext and Tor must be included in those of Hom and \otimes respectively and, in fact, will have to be restricted even further.

Even so, all the difficulties have not been systematically isolated. It is necessary to make ad hoc arguments at certain places within a proof in order to show that various subgroups are closed and, more particularly, that various homomorphisms are proper. Moreover, this procedure is not always possible in certain of the standard methods of proof of homological algebra, and the results must be gotten from others by dualization. However, with the restrictions on the domains, the admissible homomorphisms, and the resolutions indicated above, all the homological results that one would want are true. Although we do not work out every detail of the theory completely but merely assert that this or that standard homological method works, in one case the work is done in complete detail; namely in the proof of Theorem 6.6.

THEOREM 6.1. *Let G and H be in \mathcal{L} and suppose that*

$$\begin{array}{ccccccc} 0 & \longrightarrow & G & \xrightarrow{e} & I_0 & \xrightarrow{d_0} & I_1 \xrightarrow{d_1} I_2 \xrightarrow{d_2} \cdots \\ & & \downarrow f & & \downarrow f_0 & & \downarrow f_1 \downarrow f_2 \\ 0 & \longrightarrow & H & \xrightarrow{\varepsilon} & J_0 & \xrightarrow{\delta_0} & J_1 \xrightarrow{\delta_1} J_2 \xrightarrow{\delta_2} \cdots \end{array}$$

is a proper exact sequence and
is a complex,

where J_n is injective for $n=1, 2, \dots$. Then given $f \in \text{Hom}(G, H)$, there exists, for each integer n , an $f_n \in \text{Hom}(I_n, J_n)$ such that the above diagram is commutative. Moreover, any two such sequences of maps are homotopic in the sense that, if g_n is another such sequence of continuous homomorphisms, there exist $h_n \in \text{Hom}(I_n, J_{n-1})$, for $n=1, 2, \dots$, and $h_0 \in \text{Hom}(I_0, H)$ such that $f_n - g_n = \delta_{n-1}h_n + h_{n+1}d_n$, for $n=1, 2, \dots$, and $f_0 - g_0 = h_1d_0 + \varepsilon h_0$. (Actually, h_0 may be taken to be 0.)

Dually, let G and $H \in \mathcal{L}$ and suppose that

$$\begin{array}{ccccccc} \cdots & \xrightarrow{d_2} & P_2 & \xrightarrow{d_1} & P_1 & \xrightarrow{d_0} & P_0 \xrightarrow{p} G \longrightarrow 0 \\ & & \downarrow f_2 & & \downarrow f_1 & & \downarrow f_0 \downarrow f \\ \cdots & \xrightarrow{\delta_2} & Q_2 & \xrightarrow{\delta_1} & Q_1 & \xrightarrow{\delta_0} & Q_0 \xrightarrow{\pi} H \longrightarrow 0 \end{array}$$

is a complex where each P_n is projective and
is a proper exact sequence.

Then, given $f \in \text{Hom}(G, H)$, there exists for each integer n , an $f_n \in \text{Hom}(P_n, Q_n)$ such that the above diagram is commutative. Moreover, any two such sequences of maps are homotopic.

Proof. The proof proceeds along the usual lines (see [6]), except that the assumption that the top row is proper takes care of the presence of topology. The second statement follows from the first by dualization.

COROLLARY. *If* $0 \longrightarrow G \xrightarrow{e} I_0 \xrightarrow{d_0} I_1 \xrightarrow{d_1} \cdots$ *and*

$$\begin{array}{c} \downarrow \text{id}_G \\ 0 \longrightarrow G \xrightarrow{\varepsilon} J_0 \xrightarrow{\delta_0} J_1 \xrightarrow{\delta_1} \cdots \end{array}$$

are two injective resolutions of G then, for each n , there exists $f_n \in \text{Hom}(I_n, J_n)$ and $g_n \in \text{Hom}(J_n, I_n)$ such that $g_n f_n$ and $f_n g_n$ are homotopic to id_{I_n} and id_{J_n} , respectively.

Dually, any two projective resolutions of a group G are homotopic.

The proofs are obvious from Theorem 6.1.

In order to define $\text{Ext}^l(G, H)$ as a derived functor of $\text{Hom}(G, H)$ by injective resolution of H , it is necessary that G in \mathcal{C} and H in \mathcal{N} (see §V). However, since H is to have an injective resolution it is necessary, by Theorem 3.6, that H is in \mathcal{C} . Thus we must assume that G in \mathcal{C} and H in $\mathcal{C} \cap \mathcal{N}$.

DEFINITION. A group in $\mathcal{C} \cap \mathcal{N}$ is called elementary.

It follows directly from Theorems 2.4 and 2.5 that E is elementary if and only if $E \cong R^n \oplus T^m \oplus Z^s \oplus F$ where F is a finite group.

Let E be an elementary group and consider injective resolutions of

$$E: 0 \longrightarrow E \xrightarrow{e} I_0 \xrightarrow{d_0} I_1 \xrightarrow{d_1} I_2 \xrightarrow{d_2} \dots$$

where I_n is in \mathcal{N} , that is, I_n is elementary (Theorem 3.2). We prove the existence of such resolutions shortly. Let $G \in \mathcal{C}$. By the corollary to Theorem 5.3, the induced sequence

$$0 \longrightarrow \text{Hom}(G, E) \xrightarrow{e^*} \text{Hom}(G, I_0) \xrightarrow{d_0^*} \text{Hom}(G, I_1) \xrightarrow{d_1^*} \text{Hom}(G, I_2) \xrightarrow{d_2^*} \dots$$

which we denote by $\text{Hom}(G, I(E))$ is proper. Moreover, it is clear that $\text{Hom}(G, I(E))$ is a complex. The sequence obtained from $\text{Hom}(G, I(E))$ by deleting all terms after $\text{Hom}(G, I_1)$ is proper exact by Theorem 5.3. Finally, since $\text{Hom}(G, I(E))$ is a complex and the homomorphisms are proper, it follows that $d_{n-1}^*(\text{Hom}(G, I_{n-1}))$ is a closed subgroup of $\text{Ker } d_n^*$.

DEFINITION. We denote the n th homology group of $\text{Hom}(G, I(E))$ by $\text{Ext}_n^l(G, E)$. The corollary to Theorem 6.1 shows that the topological groups so defined are independent (to within natural isomorphisms) of the resolution. As usual $\text{Ext}_n^l(G, E)$ is an additive functor contravariant in G and covariant in E .

Since E is elementary, as was remarked above, $E \cong R^n \oplus T^m \oplus Z^s \oplus F$, where F is a finite group and hence is the direct product of a finite number of cyclic groups Z_{n_i} . As in §III, an injective resolution of E may be found by finding an injective resolution of the factors of E and then taking their direct product.

Injective resolutions of R , T , Z , and Z_n are the following:

$$0 \rightarrow R \rightarrow R \rightarrow 0 \rightarrow 0$$

$$0 \rightarrow T \rightarrow T \rightarrow 0 \rightarrow 0$$

$$0 \rightarrow Z \rightarrow R \rightarrow T \rightarrow 0$$

$$0 \rightarrow Z_n \rightarrow T \rightarrow T \rightarrow 0.$$

Thus, if E is elementary, there exists an injective resolution

$$0 \longrightarrow E \xrightarrow{e} I_0 \xrightarrow{d_0} I_1 \xrightarrow{d_1} I_2 \xrightarrow{d_2} \dots$$

where $I_n = (0)$ for $n \geq 2$ and I_n is elementary for $n = 1, 2$. Hence $d_n = 0$ for $n \geq 1$ so that $d_n^* = 0$ for $n \geq 1$.

PROPOSITION 6.1. *Let G be in \mathcal{C} and E be elementary. Then $\text{Ext}_0^l(G, E) \cong \text{Hom}(G, E)$, $\text{Ext}_n^l(G, E) = (0)$ for $n \geq 2$, and $\text{Ext}_1^l(G, E) \cong \text{Hom}(G, I_1)/d_0^*(\text{Hom}(G, I_0))$, where d_0^* is the map induced from the standard short resolution of E . In particular, $\text{Ext}_1^l(G, E)$ is a discrete group.*

Proof. For $n \geq 2$, $\text{Ker } d_n^* = \text{Hom}(G, I_n) = (0)$, since $I_n = (0)$. Thus $\text{Ext}_n^l(G, E) = (0)$. $\text{Ext}_1^l(G, E) = \text{Ker } d_1^*/d_0^*(\text{Hom}(G, I_0)) = \text{Hom}(G, I_1)/d_0^*(\text{Hom}(G, I_0))$. $\text{Ext}_0^l(G, E) = \text{Ker } d_0^* = e^*(\text{Hom}(G, E)) \cong \text{Hom}(G, E)$, since $0 \rightarrow \text{Hom}(G, E) \rightarrow \text{Hom}(G, I_0) \rightarrow \text{Hom}(G, I_1)$ is a proper exact sequence. Now because G is in \mathcal{C} and I_0 and I_1 are in \mathcal{N} , a dual version of Theorem 4.10 yields that $d_0^*: \text{Hom}(G, I_0) \rightarrow \text{Hom}(G, I_1)$ is an open map. Hence $d_0^*(\text{Hom}(G, I_0))$ is open in $\text{Hom}(G, I_1)$ and therefore $\text{Ext}_1^l(G, E)$ is discrete. This completes the proof of Proposition 6.1.

THEOREM 6.2. *If $0 \rightarrow E_1 \rightarrow E_2 \rightarrow E_3 \rightarrow 0$ is a proper exact sequence of elementary groups, and G is in \mathcal{C} , then*

$$(6.1) \quad \begin{aligned} 0 \rightarrow \text{Hom}(G, E_1) \rightarrow \text{Hom}(G, E_2) \rightarrow \text{Hom}(G, E_3) \\ \rightarrow \text{Ext}_1^l(G, E_1) \rightarrow \text{Ext}_1^l(G, E_2) \rightarrow \text{Ext}_1^l(G, E_3) \rightarrow 0 \end{aligned}$$

is a proper exact sequence. In particular, if $0 \rightarrow E \rightarrow I_0 \rightarrow I_1 \rightarrow I_2 \rightarrow \cdots$ is any injective resolution of E by elementary groups, then

$$0 \rightarrow \text{Hom}(G, E) \rightarrow \text{Hom}(G, I_0) \rightarrow \text{Hom}(G, I_1) \rightarrow \text{Ext}_1^l(G, E) \rightarrow 0$$

is a proper exact sequence. Similarly, if $0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$ is a proper exact sequence of groups in \mathcal{C} and E is elementary, then

$$\begin{aligned} 0 \rightarrow \text{Hom}(G_3, E) \rightarrow \text{Hom}(G_2, E) \rightarrow \text{Hom}(G_1, E) \\ \rightarrow \text{Ext}_1^l(G_3, E) \rightarrow \text{Ext}_1^l(G_2, E) \rightarrow \text{Ext}_1^l(G_1, E) \rightarrow 0 \end{aligned}$$

is a proper exact sequence.

Proof. The sequence obtained from 6.1 by deleting all terms after $\text{Hom}(G, E_3)$ is proper exact by Theorem 5.3. Clearly, all maps in 6.1 are continuous. They are trivially proper since $\text{Ext}_1^l(G, E_i)$ is discrete for $i = 1, 2, 3$. The proof that the sequence is exact follows the usual one in homological algebra [6]. It terminates because $\text{Ext}_n^l(G, E_i) = (0)$ for $i = 1, 2, 3$ and $n \geq 2$. This proves that 6.1 is proper exact. The second statement follows from this because $\text{Ext}_1^l(G, I_0) = (0)$ since I_0 is injective. The final statement follows from the standard exact sequence for a complex and a subcomplex.

We utilize Proposition 6.1 to compute Ext_1^l explicitly.

THEOREM 6.3. *Let G be in \mathcal{C} and E be elementary. Then*

$$\text{Ext}_1^l(G, E) \cong (C^\wedge)^s \oplus C^\wedge/n_1 C^\wedge \oplus \cdots \oplus C^\wedge/n_k C^\wedge$$

where C is the maximum compact subgroup of G and s, n_1, \dots, n_k are respectively the rank and torsion numbers of E/E_0 . In particular, $\text{Ext}_1^l(G, E) \cong \text{Ext}_1^l(C, E/E_0)$.

Proof. First we compute some special cases. If $E=Z$, the injective resolution

$$0 \longrightarrow Z \longrightarrow R \xrightarrow{\pi} T \longrightarrow 0$$

of Z show that $\text{Ext}_1^I(R, Z) = (0)$, since (R, π) is a covering group of T and R is simply connected. Also $\text{Ext}_1^I(Z, Z) = (0)$, clearly. If C is compact then $\text{Ext}_1^I(C, Z) = C^\wedge$, since R has no nontrivial compact subgroups.

Now $G \cong R^n \oplus Z^m \oplus C$, because G is compactly generated. Since Ext_1^I is additive, it follows that $\text{Ext}_1^I(G, Z) = C^\wedge$ where C is the maximum compact subgroup of G . If $E = Z_n$, the cyclic group of order n , the injective resolution

$$0 \longrightarrow Z_n \longrightarrow T \xrightarrow{\pi} T \longrightarrow 0$$

of Z_n shows that $\text{Ext}_1^I(R, Z_n) = (0)$, since (T, π) is a covering group of T and R is simply connected. Also $\text{Ext}_1^I(Z, Z_n) = (0)$, since T is divisible. If C is compact then, obviously, $\text{Ext}_1^I(C, Z_n) \cong C^\wedge / nC^\wedge$. Consequently, $\text{Ext}_1^I(G, Z_n) \cong C^\wedge / nC^\wedge$, where C is the maximum compact subgroup of G .

Finally, take $E = R^n \oplus T^m \oplus Z^s \oplus F$. Then $\text{Ext}_1^I(G, E_0) = (0)$ since E_0 is injective. Now, by additivity, $\text{Ext}_1^I(G, E) \cong \text{Ext}_1^I(G, E_0) \oplus \text{Ext}_1^I(G, Z)^s \oplus \text{Ext}_1^I(G, F)$. The result follows.

COROLLARY. If (1) the maximum compact subgroup of G is (0) , or

(2) E is connected, or

(3) G is torsion free and E/E_0 is finite then $\text{Ext}_1^I(G, E) = (0)$.

Proof. (1) is clear from Theorem 6.3. (2) has already been seen in the proof of Theorem 6.3. To prove (3), suppose E/E_0 is finite, so that $s=0$. Since G is torsion free, so is C . Hence C^\wedge is divisible, by Theorem 2.9, and $C^\wedge / n_i C^\wedge = (0)$ for $i=1, \dots, k$. The result follows.

Let G and H be in \mathcal{L} . We define $\text{Ext}^P(G, H)$ as a derived functor of $\text{Hom}(G, H)$ by taking projective resolutions of G . As above, it is clear that G must be an elementary group which we denote by E , and H in \mathcal{N} . Consider projective resolutions

$$\xrightarrow{d_2} P_2 \xrightarrow{d_1} P_1 \xrightarrow{d_0} P_0 \xrightarrow{\pi} E \longrightarrow 0$$

of E where $P_n \in \mathcal{C}$, i.e., is elementary (Theorem 3.3). Since a group is in \mathcal{C} if and only if its dual is in \mathcal{N} , by Theorem 2.5, Corollary 1, it follows that E is elementary if and only if E^\wedge is elementary. Hence the existence of such resolutions follows by dualizing the resolution constructed on page 394. By the Corollary to Theorem 5.3, the induced sequence

$$0 \longrightarrow \text{Hom}(E, H) \xrightarrow{\pi^*} \text{Hom}(P_0, H) \xrightarrow{d_0^*} \text{Hom}(P_1, H) \xrightarrow{d_1^*} \text{Hom}(P_2, H) \xrightarrow{d_2^*} \dots$$

which we denote by $\text{Hom}(P(E), H)$ is proper. Obviously, $\text{Hom}(P(E), H)$ is a complex. The sequence obtained from $\text{Hom}(P(E), H)$ by deleting all terms after $\text{Hom}(P_1, H)$ is proper exact, by Theorem 5.3. Finally, since $\text{Hom}(P(E), H)$ is a

complex and the homomorphisms are proper, it follows that $d_{n-1}^*(\text{Hom}(P_{n-1}, H))$ is a closed subgroup of $\text{Ker } d_n^*$.

DEFINITION. We denote the n th homology group of $\text{Hom}(P(E), H)$ by $\text{Ext}_n^P(E, H)$. The corollary to Theorem 6.1 shows that the topological groups so defined are independent (to within natural isomorphisms) of the resolution. As usual, $\text{Ext}_n^P(E, H)$ is an additive functor contravariant in E and covariant in H .

A standard argument [6] utilizing the appropriate double complex shows that on their common domain, i.e., pairs (G, H) where both G and H are elementary, $\text{Ext}_n^I(G, H)$ and $\text{Ext}_n^P(G, H)$ are naturally isomorphic.

REMARK. For G in \mathcal{C} and E elementary, the fact that $\text{Ext}_1^I(G, E)$ and $\text{Ext}_1^I(C, E/E_0)$ are naturally isomorphic follows from the exactness of the two homology sequences. In fact, consider the proper exact sequence $(0) \rightarrow E_0 \rightarrow E \rightarrow E/E_0 \rightarrow (0)$. It follows that $\rightarrow \text{Ext}_1^I(G, E_0) \rightarrow \text{Ext}_1^I(G, E) \rightarrow \text{Ext}_1^I(G, E/E_0) \rightarrow (0)$ is proper exact. But since E_0 is injective, $\text{Ext}_1^I(G, E_0) = (0)$. Therefore $\text{Ext}_1^I(G, E) \cong \text{Ext}_1^I(G, E/E_0)$. On the other hand, consider the sequence $(0) \rightarrow C \rightarrow G \rightarrow G/C \rightarrow (0)$. Then $\rightarrow \text{Ext}_1^I(G/C, E/E_0) \rightarrow \text{Ext}_1^I(G, E/E_0) \rightarrow \text{Ext}_1^I(C, E/E_0) \rightarrow (0)$ is proper exact. Since both G/C and E/E_0 are elementary, $\text{Ext}_1^I(G/C, E/E_0) \cong \text{Ext}_1^I(G/C, E/E_0)$. But since G/C is a projective it follows that $\text{Ext}_1^I(G/C, E/E_0) = (0)$. Hence $\text{Ext}_1^I(G, E/E_0) \cong \text{Ext}_1^I(C, E/E_0)$ so that $\text{Ext}_1^I(G, E) \cong \text{Ext}_1^I(C, E/E_0)$. The corresponding facts about the other functors can be proven similarly.

THEOREM 6.4. *If E is elementary and H is in \mathcal{N} then $\text{Ext}_n^P(E, H)$ and $\text{Ext}_n^I(H^\wedge, E^\wedge)$ are naturally isomorphic.*

Proof. Clearly, $\text{Ext}_0^P(E, H) \cong \text{Hom}(E, H)$. So if $n=0$, the theorem follows from the fact that $\text{Hom}(E, H)$ is naturally isomorphic with $\text{Hom}(H^\wedge, E^\wedge)$. In general, $\text{Ext}_n^P(E, H) = \text{Ker } d_n^*/d_{n-1}^*(\text{Hom}(P_{n-1}, H))$. Moreover, E^\wedge is elementary, H^\wedge is in \mathcal{C} , and

$$0 \longrightarrow E^\wedge \xrightarrow{\pi^\wedge} P_0^\wedge \xrightarrow{d_0^\wedge} P_1^\wedge \xrightarrow{d_1^\wedge} P_2^\wedge \xrightarrow{d_2^\wedge} \dots$$

is an injective resolution of E^\wedge , which is admissible in defining $\text{Ext}_n^I(H^\wedge, E^\wedge)$. Thus $\text{Ext}_n^I(H^\wedge, E^\wedge) \cong \text{Ker } (d_n^\wedge)^*/(d_{n-1}^\wedge)^*(\text{Hom}(H^\wedge, P_{n-1}^\wedge))$, where homology is computed from the complex

$$0 \longrightarrow \text{Hom}(H^\wedge, E^\wedge) \xrightarrow{(\pi^\wedge)^*} \text{Hom}(H^\wedge, P_0^\wedge) \xrightarrow{(d_0^\wedge)^*} \text{Hom}(H^\wedge, P_1^\wedge) \xrightarrow{(d_1^\wedge)^*} \text{Hom}(H^\wedge, P_2^\wedge) \xrightarrow{(d_2^\wedge)^*} \dots$$

Now for each n , $\text{Hom}(P_n, H)$ and $\text{Hom}(H^\wedge, P_n^\wedge)$ are isomorphic in a functorial way. Under this identification, one sees easily that $\text{Ker } (d_n^\wedge)^* = \text{Ker } d_n^*$ and $d_{n-1}^*(\text{Hom}(P_{n-1}, H)) = (d_{n-1}^\wedge)^*(\text{Hom}(H^\wedge, P_{n-1}^\wedge))$. This completes the proof.

COROLLARY 1. *If $0 \rightarrow E_1 \rightarrow E_2 \rightarrow E_3 \rightarrow 0$ is a proper exact sequence of elementary groups and H is in \mathcal{N} , then the following is a proper exact sequence.*

$$0 \rightarrow \text{Hom}(E_3, H) \rightarrow \text{Hom}(E_2, H) \rightarrow \text{Hom}(E_1, H) \rightarrow \text{Ext}_1^P(E_3, H) \rightarrow \text{Ext}_1^P(E_2, H) \rightarrow \text{Ext}_1^P(E_1, H) \rightarrow 0.$$

In particular, if $\cdots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow E \rightarrow 0$ is any projective resolution of E by elementary groups, then the following is a proper exact sequence.

$$0 \rightarrow \text{Hom}(E, H) \rightarrow \text{Hom}(P_0, H) \rightarrow \text{Hom}(P_1, H) \rightarrow \text{Ext}_1(E, H) \rightarrow 0.$$

Similarly, if $0 \rightarrow H_1 \rightarrow H_2 \rightarrow H_3 \rightarrow 0$ is a proper exact sequence of groups in \mathcal{N} and E is elementary, then

$$0 \rightarrow \text{Hom}(E, H_1) \rightarrow \text{Hom}(E, H_2) \rightarrow \text{Hom}(E, H_3) \\ \rightarrow \text{Ext}_1^P(E, H_1) \rightarrow \text{Ext}_1^P(E, H_2) \rightarrow \text{Ext}_1^P(E, H_3) \rightarrow 0$$

is a proper exact sequence.

COROLLARY 2. If E is elementary and H is in \mathcal{N} then $\text{Ext}_0^P(E, H) \cong \text{Hom}(E, H)$.

$$\text{Ext}_1^P(E, H) \cong (H/H_0)^m \oplus (H/H_0)/n_1(H/H_0) \oplus \cdots \oplus (H/H_0)/n_k(H/H_0)$$

where m is the dimension of the maximal toral subgroup of E , and n_1, \dots, n_k are torsion numbers of E/E_0 . In particular, $\text{Ext}_1^P(E, H)$ is a discrete group and is isomorphic with Ext_1^P (maximum compact subgroup of $E, H/H_0$). Finally $\text{Ext}_n^P(E, H) = (0)$ for $n \geq 2$.

The proofs of Corollaries 1 and 2 follow from Theorem 6.4 and the corresponding facts about Ext_n^t by dualization.

COROLLARY 3. If (1) H is connected, or

(2) E is torsion free, or

(3) E has no toral subgroup and H is divisible, then $\text{Ext}_1^P(E, H) = (0)$.

Proof. This follows directly from Corollary 2.

Let G and H be in \mathcal{L} . We define $\text{Tor}(G, H)$ as a derived functor of $G \otimes H$ by taking projective resolutions of H . As above, we must assume that G is in \mathcal{C} and H is elementary. We denote H by E . As above, consider projective resolutions

$$\xrightarrow{d_2} P_2 \xrightarrow{d_1} P_1 \xrightarrow{d_0} P_0 \xrightarrow{\pi} E \longrightarrow 0$$

of E , where P_n is elementary. It follows from the corollary to Theorem 5.3 that the induced sequence

$$\xrightarrow{d_2^*} G \otimes P_2 \xrightarrow{d_1^*} G \otimes P_1 \xrightarrow{d_0^*} G \otimes P_0 \xrightarrow{\pi^*} G \otimes E \longrightarrow 0,$$

which we denote by $G \otimes P(E)$, is proper. Obviously, $G \otimes P(E)$ is a complex. The sequence obtained from $G \otimes P(E)$ by deleting all terms before $G \otimes P_1$ is proper exact, by Theorem 5.3. Finally, since $G \otimes P(E)$ is a complex and the homomorphisms are proper, it follows that $d_n^*(G \otimes P_{n+1})$ is a closed subgroup of $\text{Ker } d_{n-1}^*$.

DEFINITION. We denote the n th homology group of $G \otimes P(E)$ by $\text{Tor}_n(G, E)$. The corollary to Theorem 6.1 shows that the topological groups so defined are

independent (to within natural isomorphisms) of the resolution. As usual, $\text{Tor}_n(G, E)$ is an additive functor covariant in both variables.

THEOREM 6.5. *If G is in \mathcal{C} and E is elementary, then $\text{Tor}_n(G, E)$ is naturally isomorphic with $(\text{Ext}_n^I(G, E^\wedge))^\wedge$.*

Proof. $\text{Tor}_0(G, E) = (G \otimes P_0)/d_0^*(G \otimes P_1) \cong (G \otimes P_0)/\text{Ker } \pi^* \cong G \otimes E$, since $G \otimes P_1 \rightarrow G \otimes P_0 \rightarrow G \otimes E \rightarrow 0$ is a proper exact sequence. Since E^\wedge is elementary and G is in \mathcal{C} , $\text{Ext}_0^I(G, E^\wedge)$ is naturally isomorphic with $\text{Hom}(G, E^\wedge)$, by Proposition 6.1. As was shown in §IV, $G \otimes E$ is naturally isomorphic with $\text{Hom}((G, E^\wedge)^\wedge)$. This proves the theorem in the case $n=0$. Now, E^\wedge is elementary, and

$$0 \longrightarrow E^\wedge \xrightarrow{\pi^\wedge} P_0^\wedge \xrightarrow{d_0^\wedge} P_1^\wedge \xrightarrow{d_1^\wedge} P_2^\wedge \xrightarrow{d_2^\wedge} \dots$$

is an injective resolution of E^\wedge , which is admissible in defining $\text{Ext}_n^I(G, E^\wedge)$. Hence, $\text{Ext}_n^I(G, E^\wedge) \cong \text{Ker } (d_n^\wedge)^*/(d_{n-1}^\wedge)^*(\text{Hom}(G, P_{n-1}^\wedge))$ for $n \geq 1$, where homology is computed from the complex

$$0 \longrightarrow \text{Hom}(G, E^\wedge) \xrightarrow{(\pi^\wedge)^*} \text{Hom}(G, P_0^\wedge) \xrightarrow{(d_0^\wedge)^*} \text{Hom}(G, P_1^\wedge) \xrightarrow{(d_1^\wedge)^*} \dots$$

However, the results of §V show that this sequence is precisely

$$0 \longrightarrow (G \otimes E)^\wedge \xrightarrow{(\pi^\wedge)^\wedge} (G \otimes P_0)^\wedge \xrightarrow{(d_0^\wedge)^\wedge} (G \otimes P_1)^\wedge \xrightarrow{(d_1^\wedge)^\wedge} \dots$$

The theorem now follows from Theorem 2.2.

COROLLARY 1. *If $0 \rightarrow E_1 \rightarrow E_2 \rightarrow E_3 \rightarrow 0$ is a proper exact sequence of elementary groups and G is in \mathcal{C} then the following is a proper exact sequence.*

$$0 \rightarrow \text{Tor}_1(G, E_1) \rightarrow \text{Tor}_1(G, E_2) \rightarrow \text{Tor}_1(G, E_3) \\ \rightarrow G \otimes E_1 \rightarrow G \otimes E_2 \rightarrow G \otimes E_3 \rightarrow 0.$$

If $\dots \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow E \rightarrow 0$ is any projective resolution of E by elementary groups, then $0 \rightarrow \text{Tor}_1(G, E) \rightarrow G \otimes P_1 \rightarrow G \otimes P_0 \rightarrow G \otimes E \rightarrow 0$ is a proper exact sequence. Similarly, if $0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$ is a proper exact sequence of groups in \mathcal{C} and E is elementary, then

$$0 \rightarrow \text{Tor}_1(G_1, E) \rightarrow \text{Tor}_1(G_2, E) \rightarrow \text{Tor}_1(G_3, E) \\ \rightarrow G_1 \otimes E \rightarrow G_2 \otimes E \rightarrow G_3 \otimes E \rightarrow 0$$

is a proper exact sequence.

COROLLARY 2. *Let G be in \mathcal{C} and E be elementary. Then $\text{Tor}_0(G, E) \cong G \otimes E$. $\text{Tor}_1(G, E) \cong (C)^m \oplus C[n_1] \oplus \dots \oplus C[n_k]$, where C is the maximum compact subgroup of G , m is the toral dimension of E , and n_1, \dots, n_k are the torsion numbers of E/E_0 . Here $C[n]$ denotes the elements of C whose order divides n . In particular, $\text{Tor}_1(G, E)$ is a compact group isomorphic with $\text{Tor}_1(C, \text{maximum compact subgroup of } E)$. Finally $\text{Tor}_n(G, E) = (0)$ for $n \geq 2$.*

The proofs of Corollary 1 and 2 follow from Theorem 6.5 and the corresponding facts about Ext_n^I by dualization.

COROLLARY 3. *Let G be in \mathcal{C} and E be elementary. If*

- (1) *the maximum compact subgroup of $G=(0)$, or*
- (2) *E is torsion free, or*
- (3) *G is torsion free and E has no toral subgroup, then $\text{Tor}_1(G, E)=(0)$.*

Proof. This follows immediately from Corollary 2.

DEFINITION. Let G and H be in \mathcal{L} . An extension of H by G is a short proper exact sequence

$$0 \longrightarrow H \xrightarrow{f} Y \xrightarrow{g} G \longrightarrow 0$$

where Y is an abelian topological group. By a theorem of Gleason, [9] or [4], Y is in \mathcal{L} .

DEFINITION. Two extensions

$$0 \longrightarrow H \xrightarrow{f} Y \xrightarrow{g} G \longrightarrow 0$$

and

$$0 \longrightarrow H \xrightarrow{f'} Y' \xrightarrow{g'} G \longrightarrow 0$$

of H by G are equivalent if there exists a proper homomorphism h , making the following diagram commutative.

$$\begin{array}{ccccccc} & & & Y & & & \\ & & f \nearrow & \downarrow h & \searrow g & & \\ 0 & \longrightarrow & H & & G & \longrightarrow & 0 \\ & & f' \searrow & \downarrow & \nearrow g' & & \\ & & & Y' & & & \end{array}$$

Since h is automatically bijective, h is a topological group isomorphism, so that this is an equivalence relation.

DEFINITION. Denote by $\mathcal{E}(G, H)$ the set of equivalence classes of extensions of H by G .

THEOREM 6.6. (1) *If G is in \mathcal{C} and E is elementary, then there is a natural bijective correspondence between $\mathcal{E}(G, E)$ and $\text{Ext}_1^I(G, E)$.*

(2) *If E is elementary and H is in \mathcal{N} , then there is a natural bijective correspondence between $\mathcal{E}(E, H)$ and $\text{Ext}_1^P(E, H)$.*

Moreover, these correspondences transform the Baer composition of extensions to the group operation in Ext_1 so that they are actually group isomorphisms.

Whereas in the case of abstract groups, the usual homological proofs of both (1) and its dual (2) work, in the continuous case there are difficulties connected with (2) and it must be gotten from (1) by dualization. Since the \wedge gives a bijective correspondence between group extensions $0 \rightarrow E \rightarrow Y \rightarrow G \rightarrow 0$, where E is elementary and G is in \mathcal{C} and $0 \rightarrow G^\wedge \rightarrow Y^\wedge \rightarrow E^\wedge \rightarrow 0$ where G^\wedge is in \mathcal{N} and E^\wedge is elementary, and since if h is an equivalence between extensions of E by G ,

then h^\wedge is an equivalence between their duals, it follows that there is a natural bijective correspondence between $\mathcal{E}(G, E)$ and $\mathcal{E}(E^\wedge, G^\wedge)$ for G in \mathcal{C} and E elementary. On the other hand, by Theorem 6.4, $\text{Ext}_1^p(G, E)$ and $\text{Ext}_1^p(E^\wedge, G^\wedge)$ are naturally isomorphic under the same assumptions. Thus, (2) follows from (1) by dualization. Hence it suffices to prove Part 1 of Theorem 6.6.

Let $0 \longrightarrow E \xrightarrow{f} Y \xrightarrow{g} G \longrightarrow 0$ be an extension where E is elementary and G is in \mathcal{C} .

$$\begin{array}{ccccccc} 0 & \longrightarrow & E & \xrightarrow{f} & Y & \xrightarrow{g} & G \longrightarrow 0 \\ & & \downarrow \text{id}_E & & \downarrow l_0 & & \downarrow l_1 \end{array}$$

Let $0 \longrightarrow E \xrightarrow{e} I_0 \xrightarrow{d} I_1 \longrightarrow 0$ be a short injective resolution of E where I_0 and I_1 are elementary. By Theorem 6.1, choose $l_0 \in \text{Hom}(Y, I_0)$ and $l_1 \in \text{Hom}(G, I_1)$ so that the above diagram is commutative. By Theorem 6.2, the sequence

$$0 \longrightarrow \text{Hom}(G, E) \longrightarrow \text{Hom}(G, I_0) \xrightarrow{\delta} \text{Hom}(G, I_1) \xrightarrow{\gamma} \text{Ext}_1^t(G, E) \longrightarrow 0$$

is proper exact. Define the continuous homomorphism $\lambda_1: \text{Hom}(G, G) \rightarrow \text{Hom}(G, I_1)$ by $\lambda_1(\xi) = l_1 \circ \xi$. Then $\gamma\lambda_1(\text{id}_G) \in \text{Ext}_1^t(G, E)$ and is clearly independent of the representative of $\mathcal{E}(G, E)$, thus giving a map from $\mathcal{E}(G, E) \rightarrow \text{Ext}_1^t(G, E)$.

Now, given any $l_1 \in \text{Hom}(G, I_1)$ we construct an extension

$$0 \longrightarrow E \xrightarrow{f_{l_1}} Y_{l_1} \xrightarrow{g_{l_1}} G \longrightarrow 0$$

as follows. Form the direct sum $G \oplus I_0$ and define the continuous homomorphism $\theta_{l_1}: G \oplus I_0 \rightarrow I_1$ by $\theta_{l_1}(x, x_0) = -l_1(x) + d(x_0)$. Let $Y_{l_1} = \text{Ker } \theta_{l_1}$. It follows from Corollary 1 of Theorem 2.5 and Proposition 2.6 that $G \oplus I_0$ is in \mathcal{C} . Since θ_{l_1} is continuous, Y_{l_1} is a closed subgroup of $G \oplus I_0$ and therefore Y_{l_1} is in \mathcal{C} by Theorem 2.6. Define $f_{l_1}: E \rightarrow Y_{l_1}$ by $f_{l_1}(u) = (0, e(u))$ for $u \in E$. Now, $(0, e(u)) \in Y_{l_1}$ since $\theta_{l_1}(0, e(u)) = -l_1(0) + d(e(u)) = 0 + 0$. Since e is a proper monomorphism, so is f_{l_1} . Define $g_{l_1}: Y_{l_1} \rightarrow G$ by $g_{l_1}(x, x_0) = x$, i.e., g_{l_1} is the restriction to Y_{l_1} of the projection $G \oplus I_0 \rightarrow G$. Hence, $g_{l_1} \in \text{Hom}(Y_{l_1}, G)$. Moreover, $\text{Ker } g_{l_1} = Y_{l_1} \cap I_0 = (0) + \text{Ker } d = (0) + e(E) = f_{l_1}(E)$. Furthermore,

$$g_{l_1}(Y_{l_1}) = \{x: x \in G \text{ and } l_1(x) \in d(I_0) = I_1\} = G,$$

so that g_{l_1} is a continuous epimorphism. Since Y_{l_1} is in \mathcal{C} , g_{l_1} is an open map. Thus,

$$0 \longrightarrow E \xrightarrow{f_{l_1}} Y_{l_1} \xrightarrow{g_{l_1}} G \longrightarrow 0$$

is an extension. Now define $l_0: Y_{l_1} \rightarrow I_0$ by $l_0(x, x_0) = x_0$. Since l_0 is the restriction of a continuous homomorphism, $l_0 \in \text{Hom}(Y_{l_1}, I_0)$. It is immediate that $l_0 \circ f_{l_1} = e \circ \text{id}_E$ and $l_1 \circ g_{l_1} = d \circ l_0$, so that the following diagram is commutative.

$$\begin{array}{ccccccc} 0 & \longrightarrow & E & \xrightarrow{f_{l_1}} & Y_{l_1} & \xrightarrow{g_{l_1}} & G \longrightarrow 0 \\ & & \downarrow \text{id}_E & & \downarrow l_0 & & \downarrow l_1 \\ 0 & \longrightarrow & E & \xrightarrow{e} & I_0 & \xrightarrow{d} & I_1 \longrightarrow 0 \end{array}$$

Since γ is surjective, this implies that the map $\mathcal{E}(G, E) \rightarrow \text{Ext}_1^l(G, E)$ is surjective. Now, let

$$\begin{array}{ccccccc} 0 & \longrightarrow & E & \xrightarrow{f} & Y & \xrightarrow{g} & G \longrightarrow 0 \\ & & \downarrow \text{id}_E & & \downarrow l_0 & & \downarrow l_1 \\ & & & & & & \end{array} \quad \begin{array}{l} \text{be an extension, where} \\ G \text{ is in } \mathcal{C} \text{ and } E \text{ is elementary.} \end{array}$$

Let $0 \longrightarrow E \xrightarrow{e} I_0 \xrightarrow{d} I_1 \longrightarrow 0$ be the given resolution of E

and let l_0, l_1 be extensions of id_E . Form

$$0 \longrightarrow E \xrightarrow{f_{l_1}} Y_{l_1} \xrightarrow{g_{l_1}} G \longrightarrow 0$$

as above. Then

$$0 \longrightarrow E \xrightarrow{f_{l_1}} Y_{l_1} \xrightarrow{g_{l_1}} G \longrightarrow 0$$

and

$$0 \longrightarrow E \xrightarrow{f} Y \xrightarrow{g} G \longrightarrow 0$$

are equivalent. To see this, define $h: Y \rightarrow G \oplus I_0$ by $h(y) = (g(y), l_0(y))$. Then h is a continuous homomorphism since g and l_0 are. If $y \in Y$, then $\theta_{l_1}(h(y)) = -l_1(g(y)) + d(l_0(y)) = 0$ since $dl_0 = l_1g$. Thus, $h(Y) \subset Y_{l_1}$. One sees easily that $hf = f_{l_1}$ and $g_{l_1}h = g$. Clearly, h is bijective. Since the given extension is proper exact, Y is in \mathcal{C} by Theorem 2.6. Hence h is open, by the Open Mapping Theorem, and the extensions are equivalent.

Let

$$0 \longrightarrow E \xrightarrow{f} Y \xrightarrow{g} G \longrightarrow 0 \quad \text{and} \quad 0 \longrightarrow E \xrightarrow{f'} Y' \xrightarrow{g'} G \longrightarrow 0$$

be two extensions. Form the respective l_1 and l'_1 . Suppose they map into the same cohomology class of $\text{Ext}_1^l(G, E)$, namely that $\gamma l_1(\text{id}_G) = \gamma l'_1(\text{id}_G)$. Then $\gamma l_1 = \gamma l'_1$ so that $l_1 - l'_1 \in \text{Ker } \gamma$. Since $\text{Ker } \gamma = \delta(\text{Hom}(G, I_0))$, it follows that $l_1 - l'_1 = d\xi$ for some $\xi \in \text{Hom}(G, I_0)$.

Define $\alpha: G \oplus I_0 \rightarrow G \oplus I_0$ by $\alpha(x, x_0) = (x, \xi(x) + x_0)$. Because ξ is a continuous homomorphism, so is α . Obviously, α is bijective. Since $G \oplus I_0$ is in \mathcal{C} , α is a topological group automorphism, by the Open Mapping Theorem.

Now form Y_{l_1} and $Y_{l'_1}$ as above. We show that $\alpha(Y_{l_1}) = Y_{l'_1}$. Now $\alpha(Y_{l_1}) = \{\alpha(x, x_0) : l_1(x) = d(x_0)\}$. But $l_1(x) = l'_1(x) + d\xi(x)$ for each $x \in G$. Thus,

$$\alpha(Y_{l_1}) = \{(x, \xi(x) + x_0) : l'_1(x) + d\xi(x) = d(x_0)\}$$

which in turn equals $\{(x, x_0 - \xi(x)) : l'_1(x) = d(x_0 - \xi(x))\}$. Since as x_0 ranges over I_0 , so does $x_0 - \xi(x)$, the result follows. Hence $\alpha|_{Y_{l_1}}$ is a topological group isomorphism $Y_{l_1} \rightarrow Y_{l'_1}$. Also, for $u \in E$, $\alpha f_{l_1}(u) = \alpha(0, e(u)) = (0, \xi(0) + e(u)) = f_{l'_1}(u)$ and $g_{l'_1}\alpha(x, x_0) = g_{l'_1}(x, \xi(x) + x_0) = x = g_{l_1}(x, x_0)$. Thus $\alpha f_{l_1} = f_{l'_1}$ and $g_{l_1} = g_{l'_1}\alpha$ so that α is an equivalence. However, since

$$0 \longrightarrow E \xrightarrow{f} Y \xrightarrow{g} G \longrightarrow 0 \quad \text{and} \quad 0 \longrightarrow E \xrightarrow{f_{l_1}} Y_{l_1} \xrightarrow{g_{l_1}} G \longrightarrow 0$$

are equivalent, and

$$0 \longrightarrow E \xrightarrow{f'} Y' \xrightarrow{g'} G' \longrightarrow 0 \quad \text{and} \quad 0 \longrightarrow E \xrightarrow{f_{i_1}} Y_{i_1} \xrightarrow{g_{i_1}} G \longrightarrow 0$$

are equivalent, it follows that

$$0 \rightarrow E \rightarrow Y \rightarrow G \rightarrow 0 \quad \text{and} \quad 0 \rightarrow E \rightarrow Y' \rightarrow G \rightarrow 0$$

are equivalent. Thus the map $\mathcal{E}(G, E) \rightarrow \text{Ext}_1^I(G, E)$ is bijective.

If

$$0 \longrightarrow E \xrightarrow{f} Y \xrightarrow{g} G \longrightarrow 0 \quad \text{and} \quad 0 \longrightarrow E \xrightarrow{f'} Y' \xrightarrow{g'} G \longrightarrow 0$$

are two extensions of E by G , one defines their Baer product as follows: Form $Y \oplus Y'$, and let $A = \{(y, y') : (y, y') \in Y \oplus Y', g(y) = g'(y')\}$. A is a closed subgroup of $Y \oplus Y'$ since g and g' are continuous homomorphisms. Let

$$B = \{(-f(u), f'(u)) : u \in E\}.$$

B is a closed subgroup of $Y \oplus Y'$ because f and f' are proper homomorphisms. Since $gf = g'f' = 0$, $B \subset A$. Hence B is a closed subgroup of A . Let $Y'' = A/B$ and $\pi: A \rightarrow Y''$ be the canonical epimorphism. If $u \in E$, then $(f(u), 0)$ and $(0, f'(u)) \in A$. Define $f''(u) = \pi(f(u), 0) = \pi(0, f'(u))$. Then $f'': E \rightarrow Y''$ is a continuous homomorphism since f and π are. Let $\alpha: A \rightarrow G$ be defined by $\alpha(y, y') = g(y) = g'(y')$. Since α is the composition of g with the restriction to A of a projection, it is a continuous homomorphism. Also, $\alpha(-f(u), f'(u)) = g'f'(u) = 0$. Therefore α induces a continuous homomorphism $g'': Y'' \rightarrow G$. Since G and E are in \mathcal{C} , it follows from Theorem 2.6 that Y and Y' are in \mathcal{C} . Hence $Y \oplus Y'$ is in \mathcal{C} , by Proposition 2.6 and Corollary 1 of Theorem 2.5. Consequently A and therefore Y'' is in \mathcal{C} by Theorem 2.6. It is a strictly formal fact that

$$0 \longrightarrow E \xrightarrow{f''} Y'' \xrightarrow{g''} G \longrightarrow 0$$

is an exact sequence. Since E, Y, G are in \mathcal{C} it is proper. The equivalence class of this extension depends only on the equivalence classes of the given extensions. Thus $\mathcal{E}(G, E)$ is closed under Baer multiplication. It is also a strictly formal fact that the correspondence between $\mathcal{E}(G, E)$ and $\text{Ext}_1^I(G, E)$ is a homomorphism. Hence $\mathcal{E}(G, E) \rightarrow \text{Ext}_1^I(G, E)$ is an isomorphism. This completes the proof of Theorem 6.6⁽⁸⁾.

COROLLARY. Let $0 \rightarrow E \rightarrow Y \rightarrow G \rightarrow 0$ be an extension with G in \mathcal{C} and E elementary. If (1) The maximum compact subgroup of $G = (0)$, or

(2) E is connected, or

(3) G is torsion free and E/E_0 is finite, then $Y \cong G \oplus E$.

Let $0 \rightarrow H \rightarrow Y \rightarrow E \rightarrow 0$ be an extension with H in \mathcal{N} and E elementary.

If (1) H is connected, or

(2) the maximum compact subgroup of $E = (0)$, or

(3) H is divisible and the toral dimension of $E = (0)$, then $Y \cong H \oplus E$.

⁽⁸⁾ The proof of Theorem 6.6 is modeled after one in [6].

Proof. These corollaries follow directly from Theorem 6.6, the Corollary of Theorem 6.3 and Corollary 3 of Theorem 6.4.

REFERENCES

1. A. Weil, *L'integration dans les groupes topologiques et ses applications*, Hermann, Paris, 1953.
2. L. Pontrjagin, *Topologische Gruppen*, Teil 1, 2, Teubner, Leipzig, 1957.
3. I. Kaplansky, *Infinite Abelian groups*, The University of Michigan Press, Ann Arbor, 1954.
4. D. Montgomery and L. Zippin, *Topological transformation groups*, Interscience, New York, 1955.
5. J. Braconnier, *Sur les groupes topologiques localement compacts*, J. Math. Pures Appl. **27** (1948), 1–85.
6. H. Cartan and S. Eilenberg, *Homological algebra*, Princeton Univ. Press, Princeton, N. J., 1956.
7. J. Dixmier, *Quelques propriétés des groupes abéliens localement compacts*, Bull. Sci. Math. **81** (1957), 38–48.
8. H. Yamabe, *On the conjecture of Iwasawa and Gleason*, Ann. of Math. **58** (1953), 48–54.
9. A. Gleason, *Groups without small subgroups*, Ann. of Math. **56** (1952), 193–212.

UNIVERSITY OF CALIFORNIA,
BERKELEY, CALIFORNIA