

SYSTEMS OF TOEPLITZ OPERATORS ON H^2 . II

BY

H. R. POUSSON⁽¹⁾

1. Introduction. Let L^p ($0 < p \leq \infty$) be the usual Lebesgue space with respect to normalized Lebesgue measure on the unit circle. The space H^p ($0 < p \leq \infty$) will consist of analytic functions f on the unit disc such that $\lim_{r \rightarrow 1-} \|f(re^{i\theta})\|_p < \infty$. If $f \in H^p$, then the function defined a.e. by $f(e^{i\theta}) = \lim_{r \rightarrow 1-} f(re^{i\theta})$ is in L^p . Should $p \geq 1$, such functions in L^p are precisely those which have vanishing negative Fourier coefficients. The subspace of functions $f \in H^p$ such that $f(0) = 0$ is denoted by H_0^p .

The space of n by m dimensional, matrix-valued functions with terms in L^p is denoted by $L_{n \times m}^p$. The spaces $H_{n \times m}^p$ and $H_{0, n \times m}^p$ are defined analogously.

Let $\phi \in L_{n \times n}^\infty$. Define the operator $T_\phi: H_{n \times 1}^2 \rightarrow H_{n \times 1}^2$ by setting $T_\phi f = P\phi f$ for all $f \in H_{n \times 1}^2$ where $P: L_{n \times m}^2 \rightarrow H_{n \times m}^2$ is the projection operator. The operator T_ϕ can be considered as a system of Toeplitz operators on H^2 , and T_ϕ will be called the *Toeplitz operator* associated with the matrix-valued function ϕ .

This paper concerns conditions on $\phi \in L_{n \times n}^\infty$ which give an invertible T_ϕ . After defining suitable norms for matrix functions, we consider the problem of angle between manifolds of matrix functions. This gives results which are applicable to systems of Toeplitz operators. The approach is similar to the method used by Devinatz [2] for ϕ scalar.

2. Definitions and general results. Most of the preliminary results are stated without proof. The proofs are available in the literature or require only a simple generalization of proofs in the literature.

Denote normalized Lebesgue measure on the unit circle by $d\mu$.

For F a matrix function, let $F_k = \int F(e^{i\theta})e^{-ik\theta} d\mu$; $k = 0, \pm 1, \pm 2, \dots$. Denote the conjugate transpose, transpose, and complex conjugate of F by F^* , F' , and \bar{F} respectively. A matrix J will be called a *projection* matrix provided $J = J^2 = J^*$.

DEFINITION 2.1. A function $F \in H_{n \times n}^p$ is *outer* provided $\int \log |\det F| d\mu = \log |\det F_0| > -\infty$. A function $F \in H_{n \times n}^\infty$ is *inner* provided $F = JU$ where J is a constant projection matrix and U is unitary a.e.

Thus $F \in H_{n \times n}^p$ is outer if and only if $\det F \in H^{p/n}$ is outer.

THEOREM 2.2. If $F \in H_{n \times n}^2$ is outer, and if $F^{-1} \in L_{n \times n}^2$, then $F^{-1} \in H_{n \times n}^2$.

Received by the editors April 28, 1967.

(¹) The results of this paper are contained in the author's doctoral dissertation, University of Virginia, 1967. This work was partially supported by the National Science Foundation under grant NSF GP-3445.

A proof of the following is contained in [8].

THEOREM 2.3. *If $F, F^{-1} \in H_{n \times n}^2$, then F is an outer function.*

THEOREM 2.4. *If $W \in L_{n \times n}^1$ is hermitian positive semidefinite, then W has a factorization $W = BB^*$ where $B \in H_{n \times n}^2$ is outer provided $\int \log \det W d\mu > -\infty$.*

For a proof of the above, see [5, p. 193].

For $F \in H_{n \times n}^2$, $S(F)$ will denote the subspace of $H_{n \times n}^2$ spanned by $\{Fe^{ik\theta}\}_{k=0}^\infty$.

The following is proved by Masani [7, p. 286].

THEOREM 2.5. *Let $K \in H_{n \times n}^2$; $K \neq 0$. Then $K = FG$ where $F \in H_{n \times n}^2$ is outer and $G \in H_{n \times n}^\infty$ is inner. Also $S(G) = S(K)$.*

It can be shown that the Hilbert space adjoint T_ϕ^* of T_ϕ is T_{ϕ^*} ; hence T_ϕ is invertible if and only if T_{ϕ^*} is invertible.

THEOREM 2.6. *Let $\phi \in L_{n \times n}^\infty$. In order for T_ϕ to be invertible, it is necessary and sufficient that*

- (i) $\phi = G^*H$ where $G, G^{-1}, H, H^{-1} \in H_{n \times n}^2$; $S(G) = S(H) = H_{n \times n}^2$; and
- (ii) $K: f \rightarrow H^{-1}P(G^{-1})^*f$ defines a bounded operator from $L_{n \times 1}^2$ to $H_{n \times 1}^2$.

The proof of the above theorem and the proofs of the remaining results in this section are contained in [8].

COROLLARY 2.7. *If T_ϕ is invertible, then $\phi^{-1} \in L_{n \times n}^\infty$.*

THEOREM 2.8. *Suppose $\phi \in L_{n \times n}^\infty$ is positive definite. The following are equivalent.*

- (i) T_ϕ is invertible.
- (ii) $\text{ess inf det } \phi > 0$.
- (iii) $T_{\det \phi}$ is invertible.

REMARK 2.9. If $F \in H_{n \times n}^\infty$, then $T_\phi T_F = T_{\phi F}$ and $T_F T_\phi = T_{F^* \phi}$.

THEOREM 2.10. *Let $\phi \in H_{n \times n}^\infty$, then T_ϕ is invertible if and only if $\phi^{-1} \in H_{n \times n}^\infty$ in which case $T_\phi^{-1} = T_{\phi^{-1}}$.*

THEOREM 2.11. *If $\phi \in L_{n \times n}^\infty$ and T_ϕ is invertible, then there exists a factorization $\phi = UK$ where U is unitary, $K \in H_{n \times n}^\infty$ is outer, and both T_K and T_U are invertible.*

REMARK 2.12. If $\phi \in L_{n \times n}^\infty$ is unitary, then T_ϕ invertible implies $\phi = G^*H$ where $G, G^{-1}, H, H^{-1} \in H_{n \times n}^2$ and $HH^* = (GG^*)^{-1}$.

Proof. Consider Theorem 2.6 and use the fact that $\phi^* = \phi^{-1}$.

3. Matrix norms. Unless otherwise stated, all matrix functions will be n by n dimensional.

It follows from basic matrix theory that a matrix function F has a polar factorization $F = UH$ where U is unitary and H is positive semidefinite. Should $\det F \neq 0$ a.e., the factorization is unique and H is positive definite. It will be desirable to consider

polar factorizations where the hermitian factor is on the left of the unitary factor. When necessary to distinguish the two factorizations, subscripts will be used; i.e., $F = U_R H_R = H_L U_L$.

Two matrix functions F and G are unitarily equivalent, denoted $F \sim G$, provided there exists a unitary matrix function V such that $V^* G V = F$.

LEMMA 3.1. *If U is a unitary matrix function and if H is a positive definite hermitian matrix function, then $|\operatorname{tr} UH| \leq \operatorname{tr} H$.*

Proof. There exists a unitary function V such that $V^* U V = E$ is diagonal and $|e_{i,i}| = 1$ a.e. for $i = 1, 2, \dots, n$. Let $V^* H V = K$. Then $|\operatorname{tr} UH| = |\operatorname{tr} EK| = |\sum_{i=1}^n e_{i,i} k_{i,i}| \leq \operatorname{tr} K = \operatorname{tr} H$.

LEMMA 3.2. *If $F = U_R H_R = H_L U_L$ are the polar factorizations of F , then $\operatorname{tr} H_R = \operatorname{tr} H_L$.*

Proof. Applying 3.1 we have $\operatorname{tr} H_L = \operatorname{tr} U_L^* U_R H_R \leq \operatorname{tr} H_R$ and $\operatorname{tr} H_R = \operatorname{tr} U_L U_R^* H_L \leq \operatorname{tr} H_L$.

DEFINITION 3.3. For all $F \in L_{n \times n}^1$, define the norm of F , denoted $\|F\|_1$, as $\|F\|_1 = \int \operatorname{tr} (FF^*)^{1/2} d\mu$.

The properties of this norm, including the necessary triangle inequality, will be given by the next two lemmas.

LEMMA 3.4. *Let $F \in L_{n \times n}^1$ and let $F = U_R H_R = H_L U_L$ be polar factorizations of F . Then $\|F\|_1 = \int \operatorname{tr} H_R d\mu = \int \operatorname{tr} H_L d\mu$.*

Proof. We have

$$\|F\|_1 = \int \operatorname{tr} (FF^*)^{1/2} d\mu = \int \operatorname{tr} (U_R H_R^2 U_R^*)^{1/2} d\mu = \int \operatorname{tr} U_R H_R U_R^* d\mu = \int \operatorname{tr} H_R d\mu.$$

The conclusion now follows from 3.2.

If $F \in L_{n \times n}^1$ and U is unitary, then as an immediate consequence of 3.4, $\|UF\|_1 = \|F\|_1 = \|FU\|_1$ and $\|F^*\|_1 = \|F\|_1$.

LEMMA 3.5. *If $F, G \in L_{n \times n}^1$, then $\|F+G\|_1 \leq \|F\|_1 + \|G\|_1$.*

Proof. Consider polar factorizations $U_{F+G} H_{F+G}$, $U_F H_F$ and $U_G H_G$ of $F+G$, F and G respectively. Since $U_{F+G} H_{F+G} = U_F H_F + U_G H_G$, it follows from 3.4 and 3.1 that

$$\begin{aligned} \|F+G\|_1 &= \int \operatorname{tr} H_{F+G} d\mu \leq \int |\operatorname{tr} U_{F+G}^* U_F H_F| d\mu + \int |\operatorname{tr} U_{F+G}^* U_G H_G| d\mu \\ &\leq \int \operatorname{tr} H_F d\mu + \int \operatorname{tr} H_G d\mu = \|F\|_1 + \|G\|_1. \end{aligned}$$

The *Euclidean norm*, denoted $|A|$, of a constant matrix A is defined as $|A|$

$=(\operatorname{tr} AA^*)^{1/2}$. For proof that this is actually a norm, see [9, p. 125]. It is easily established that for constant matrices A and B , $|AB| \leq |A| |B|$.

DEFINITION 3.6. For all $F \in L_{n \times n}^2$, define the norm of F , denoted $\|F\|_2$, as $\|F\|_2 = (\int \operatorname{tr} FF^* d\mu)^{1/2} = (\int |F|^2 d\mu)^{1/2}$.

LEMMA 3.7. If $F, G \in L_{n \times n}^2$, then $|\int \operatorname{tr} FG d\mu| \leq \|F\|_2 \|G\|_2$.

Proof. It follows from Schwarz's inequality that

$$\begin{aligned} \left| \int \operatorname{tr} FG d\mu \right| &= \left| \sum \int f_{i,j} g_{j,i} d\mu \right| \leq \int \left(\sum |f_{i,j}| |g_{j,i}| \right) d\mu \\ &\leq \int \left(\sum |f_{i,j}|^2 \right)^{1/2} \left(\sum |g_{j,i}|^2 \right)^{1/2} d\mu \\ &\leq \left(\int \operatorname{tr} FF^* d\mu \right)^{1/2} \left(\int \operatorname{tr} GG^* d\mu \right)^{1/2} = \|F\|_2 \|G\|_2; \quad 1 \leq i, j \leq n. \end{aligned}$$

LEMMA 3.8. If $F, G \in L_{n \times n}^2$, then $\|FG\|_1 \leq \|F\|_2 \|G\|_2$.

Proof. Let $F \in L_{n \times n}^2$. Assume, without loss of generality, that $F^{-1} \in L_{n \times n}^2$. Applying 3.7, we have

$$\|FG\|_1 = \int \operatorname{tr} (FF^*)^{1/2} (FGG^*F^*)^{1/2} (FF^*)^{-1/2} d\mu \leq \|F\|_2 \|G\|_2.$$

THEOREM 3.9. Products of the form MN where M is in the unit ball of $H_{n \times n}^2$ and N is in the unit ball of $H_{0; n \times n}^2$ are dense in the open unit ball of $H_{0; n \times n}^1$.

Proof. Products of the form stated are contained in the unit ball of $H_{0; n \times n}^1$, see 3.8.

Now consider $F \in H_{0; n \times n}^1$ such that $\|F\|_1 < 1$ and $\int \log |\det F| d\mu > -\infty$. From 2.4, $(FF^*)^{1/2} = BB^*$ where $B \in H_{n \times n}^2$ is outer. Also, $\|B^{-1}F\|_2^2 = \|B\|_2^2 = \|F\|_1 < 1$. Hence B is contained in the unit ball of $H_{n \times n}^2$ and $B^{-1}F$ is contained in the unit ball of $L_{n \times n}^2$. By definition, $B \in H_{n \times n}^2$ is outer if and only if $\det B$ is outer. Hence $\det B$, considered as an analytic function in the open disc, does not vanish inside the disc. From this it follows that B^{-1} can be considered as an analytic function in the open disc. Clearly $B^{-1}F$ has an analytic extension inside the disc which vanishes at zero, since $F_0 = 0$. Since $B^{-1}F \in L_{n \times n}^2$ on the unit circle, $B^{-1}F \in H_{0; n \times n}^2$. Let $M = B$ and $N = B^{-1}F$.

Consider $F \in H_{0; n \times n}^1$ such that $\|F\|_1 < 1$ and $\int \log |\det F| d\mu = -\infty$. Let $K = e^{-i\theta}F$, then $|\det K| = |\det F|$ and $F_1 = K_0$. Using a form of Jensen's inequality, see [11, p. 123], $\log |\det K_0| \leq \int \log |\det K| d\mu = -\infty$. Therefore F_1 is singular.

Choose $\alpha > 0$ such that $F_1 + \alpha$ is nonsingular and $\|F_\alpha\|_1 < 1$ where $F_\alpha = F + \alpha e^{i\theta}$. Jensen's inequality implies $\int \log |\det F_\alpha| d\mu > -\infty$. It follows that functions F in the unit ball of $H_{0; n \times n}^1$ for which $\int \log |\det F| d\mu > -\infty$ are dense in the open unit ball of $H_{0; n \times n}^1$.

The *Banach norm*, denoted $|A|_B$, of a constant matrix A is defined as $|A|_B = \sup_{|X| \neq 0} |AX|/|X|$.

For an n by n matrix A , the Banach norm and the Euclidean norm are related by $|A|_B \leq |A| \leq n|A|_B$.

LEMMA 3.10. *If A is a constant matrix and if $A = UH$ is its polar factorization then $|A|_B = d_{1,1}$ where D is diagonal, $d_{1,1} \geq d_{2,2} \geq \cdots \geq d_{n,n}$ and $H \sim D$.*

Proof. Let X be any nonzero constant matrix and let V denote the unitary matrix such that $V^*HV = D$. Then $|AX| = (\text{tr } H^2 XX^*)^{1/2} = (\text{tr } D^2 V^* XX^* V)^{1/2} \leq d_{1,1}(\text{tr } XX^*)^{1/2} = d_{1,1}|X|$ and $|A|_B \leq d_{1,1}$.

Let E be the projection matrix which has every term zero with the exception of $e_{1,1} = 1$. Then

$$|AVE| = (\text{tr } UHVEV^*HU^*)^{1/2} = (\text{tr } V^*H^2VE)^{1/2} = (\text{tr } D^2E)^{1/2} = d_{1,1} = d_{1,1}|VE|$$

and $|A|_B \geq d_{1,1}$.

DEFINITION 3.11. For all $F \in L_{n \times n}^\infty$, define the norm of F , denoted $\|F\|_\infty$, as $\|F\|_\infty = \text{ess sup}_\theta |F(e^{i\theta})|_B$.

As an immediate consequence of 3.10 we have the following.

Note 3.12. Let $F \in L_{n \times n}^\infty$, let $F = UH$ be its polar factorization and let $H \sim D$ where D is diagonal with $d_{1,1} \geq d_{2,2} \geq \cdots \geq d_{n,n}$. Then $\|F\|_\infty = \|d_{1,1}\|_\infty$.

THEOREM 3.13. *Suppose $G \in L_{n \times n}^\infty$. The linear functional $I(\cdot)$ on $L_{n \times n}^1$ defined for all $F \in L_{n \times n}^1$ by $I(F) = \int \text{tr } FG \, d\mu$ is bounded and $\|I\| = \|G\|_\infty$. All bounded linear functionals on $L_{n \times n}^1$ have this form.*

Proof. Let $U_G H_G$ be the polar factorization of $G \in L_{n \times n}^\infty$ and let V denote the unitary matrix function such that $V^* H_G V = D$ where D is diagonal, $d_{1,1} \geq d_{2,2} \geq \cdots \geq d_{n,n}$. Suppose $U_F H_F$ is the polar factorization of an arbitrary $F \in L_{n \times n}^1$. Set $K = V^* U_F H_F U_G V$ and let J be the following diagonal (unitary) matrix. If $k_{i,i} \neq 0$, set $j_{i,i} = \bar{k}_{i,i} |k_{i,i}|^{-1}$, and if $k_{i,i} = 0$, set $j_{i,i} = 1$; $i = 1, 2, \dots, n$. Then

$$\begin{aligned} |I(F)| &= \left| \int \text{tr } U_F H_F U_G H_G \, d\mu \right| = \left| \int \text{tr } KD \, d\mu \right| \leq \int \left(\sum_{i=1}^n |k_{i,i}| |d_{i,i}| \right) d\mu \\ &\leq \|d_{1,1}\|_\infty \int \left(\sum_{i=1}^n |k_{i,i}| \right) d\mu = \|G\|_\infty \int \text{tr } JK \, d\mu \\ &= \|G\|_\infty \int \text{tr } U_G V J V^* U_F H_F \, d\mu \leq \|G\|_\infty \int \text{tr } H_F \, d\mu = \|G\|_\infty \|F\|_1. \end{aligned}$$

Therefore $I(\cdot)$ is bounded and $\|I\| \leq \|G\|_\infty$.

Now for $\alpha > 0$, take $E = [e^{i\theta} \mid |G(e^{i\theta})|_B \geq \|G\| - \alpha]$. Then $\mu(E) = \int \chi_E \, d\mu \neq 0$ where χ_E is the characteristic function of E .

Let Q denote the projection matrix which has every term equal to zero with the

exception of $q_{1,1}=1$. Set $F_\alpha = Q\chi_E/u(E)$, then $\|F_\alpha\|_1=1$. If $K_\alpha = V^*F_\alpha VU_G^*$, then $\|K_\alpha\|_1 = \|F_\alpha\|_1 = 1$ and

$$\begin{aligned} I(K_\alpha) &= \int_E \operatorname{tr} \left[\frac{V^* Q V U_G^*}{u(E)} \cdot G \right] d\mu = \frac{1}{u(E)} \int_E \operatorname{tr} V^* Q V H_G d\mu \\ &= \frac{1}{u(E)} \int_E \operatorname{tr} J D d\mu = \frac{1}{u(E)} \int_E |G|_B d\mu. \end{aligned}$$

Hence $\|G\|_\infty - \alpha \leq |I(K_\alpha)| \leq \|I\|$. Thus $\|G\|_\infty = \|I\|$.

It can be deduced from the analogous result for the scalar case that all bounded linear functionals on $L_{n \times n}^1$ have the form asserted.

4. Manifolds of matrix functions. Let $W \in L_{n \times n}^1$ be positive definite hermitian. Denote by $L_{n \times n}^2(W)$ the space of matrix functions F for which $\|F\|_W^2 = \int \operatorname{tr} F F^* W d\mu < \infty$. This is a Hilbert space with inner product defined by $(F, G)_W = \int \operatorname{tr} F G^* W d\mu$ for all $F, G \in L_{n \times n}^2(W)$.

Consider $F \in L_{n \times n}^2(W)$ such that F is orthogonal to the trigonometric polynomials, all of which are clearly contained in $L_{n \times n}^2(W)$. Then for all constant matrices C , and $k=0, \pm 1, \pm 2, \dots$, $(e^{ik\theta} C, F)_W = \int \operatorname{tr} e^{ik\theta} C F^* W d\mu = 0$. Since $\|F^* W^{1/2}\|_2^2 = \|F\|_W^2$, $F^* W^{1/2} \in L_{n \times n}^2$, so $F^* W \in L_{n \times n}^1$ and has a Fourier series expansion. Thus $F^* W = 0$. If $\det W \neq 0$ a.e., then $F^* = 0$ a.e. Hence the trigonometric polynomials are dense in $L_{n \times n}^2(W)$ provided $\det W \neq 0$ a.e.

Denote by $\Delta_{n \times n}$ the space of all analytic trigonometric polynomials with n by n constant matrix coefficients. Let $\Delta_{0; n \times n}$ be the subspace of all $A \in \Delta_{n \times n}$ for which $A_0 = 0$. Denote by $H_{n \times n}^2(W)$ and $H_{0; n \times n}^2(W)$ the subspaces of $L_{n \times n}^2(W)$ generated by $\Delta_{n \times n}$ and $\Delta_{0; n \times n}$ respectively. Let $\bar{H}_{0; n \times n}^2(W)$ be the subspace of $L_{n \times n}^2(W)$ generated by all conjugate transposes of elements of $\Delta_{0; n \times n}$.

DEFINITION 4.1. Let

$$\delta = \sup \{ |(F, K^*)_W| \mid F \in H_{n \times n}^2(W); K^* \in \bar{H}_{0; n \times n}^2(W); \|F\|_W \leq 1; \|K^*\|_W \leq 1 \}.$$

The manifolds $H_{n \times n}^2(W)$ and $\bar{H}_{0; n \times n}^2(W)$ are at positive angle provided $\delta < 1$.

Consider a scalar function $w \in L^1$. If $\log w$ is summable, then there exists an outer function $h \in H^1$ such that $|h|=w$. Helson and Szego [6] have proved the following.

THEOREM 4.2. *The manifolds $H^2(w)$ and $\bar{H}_0^2(w)$ are at positive angle if and only if there exists $g \in H^\infty$ and $\alpha > 0$ such that $|g(e^{i\theta})| \geq \alpha$ a.e. and $|\arg hg| \leq \pi/2 - \alpha$.*

A matrix form of this result will be shown to hold.

If A and B are hermitian, then $A \leq B$ provided $B - A$ is positive semidefinite.

Let $W \in L_{n \times n}^1$ be positive definite hermitian and suppose $\log \det W$ is summable. Then there exists two factorizations of W ; that is, $W = A^* A = B B^*$ where $A, B \in H_{n \times n}^2$; A and B are both outer functions. Let $U = B^{-1} A^* = B^* (A^{-1})$. Then U is unitary and $W = B U A$. In the theorem and proof which follow W, U, A and B will be as just described.

THEOREM 4.3. *The manifolds $H_{n \times n}^2(W)$ and $\bar{H}_{0; n \times n}^2(W)$ are at positive angle if and only if there exists $G \in H_{n \times n}^\infty$ and $\alpha > 0$ such that*

$$(4.4) \quad \alpha + G^*G \leq U^*G + G^*U.$$

Proof. Since $W = BUA$, we have from 4.1 that $\delta = \sup_{F, K} \left| \int \operatorname{tr} FKW d\mu \right| = \sup_{F, K} \left| \int \operatorname{tr} AFKB U d\mu \right| < 1$ where F varies throughout the unit ball of $H_{n \times n}^2(W)$ and K^* varies throughout the unit ball of $\bar{H}_{0; n \times n}^2(W)$.

For all $F \in H_{n \times n}^2(W)$ there exists a sequence $\{P_s\}_1^\infty$ of elements of $\Delta_{n \times n}$ such that $\lim_{s \rightarrow \infty} \|P_s - F\|_W = 0$. For $s = 1, 2, 3, \dots$ $\|P_s - F\|_W = \|AP_s - AF\|_2$, and so $\lim_{s \rightarrow \infty} \|AP_s - AF\|_2 = 0$. Clearly $AP_s \in H_{n \times n}^2$ for each s ; hence $AF \in H_{n \times n}^2$. Moreover $\|F\|_W = \|AF\|_2$. Thus F in the unit ball of $H_{n \times n}^2(W)$ implies AF is in the unit ball of $H_{n \times n}^2$. In fact, since A is outer, the set of all products AF where $F \in H_{n \times n}^2(W)$, $\|F\|_W \leq 1$, is dense in the unit ball of $H_{n \times n}^2$. A similar argument shows that the set of all products KB where $K^* \in \bar{H}_{0; n \times n}^2(W)$, $\|K^*\|_W \leq 1$, is dense in the unit ball of $H_{0; n \times n}^2$. Thus it follows from 3.9 that $\delta = \sup_R \left| \int \operatorname{tr} RU d\mu \right|$ where R varies throughout the unit ball of $H_{0; n \times n}^1$. Theorem 3.13 implies that $\int \operatorname{tr} RU d\mu$ defines a bounded linear functional on $L_{n \times n}^1$ which, when restricted to $H_{0; n \times n}^1$, has norm δ .

Suppose $S \in L_{n \times n}^\infty$ is an annihilator of $H_{0; n \times n}^1$; that is, for all $R \in H_{0; n \times n}^1$, $\int \operatorname{tr} RS d\mu = 0$. Necessarily, $S \in H_{n \times n}^\infty$. Moreover, all elements of $H_{n \times n}^\infty$ are annihilators of $H_{0; n \times n}^1$.

We now apply a corollary of the Hahn-Banach theorem, see [1], and obtain $\delta = \inf_{S \in H_{n \times n}^\infty} \|U - S\|_\infty$.

Choose $G \in H_{n \times n}^\infty$ such that $\|U - G\|_\infty \leq 1 - \alpha$ where $\alpha > 0$. Since multiplication by a unitary matrix leaves the norm constant, $\|U - G\|_\infty = \|I - U^*G\|_\infty \leq 1 - \alpha$. It follows from the definition of the $L_{n \times n}^\infty$ norm that $|I - U^*(e^{i\theta})G(e^{i\theta})|_B \leq 1 - \alpha$ a.e. This implies that

$$([I - U^*(e^{i\theta})G(e^{i\theta})]X, [I - U^*(e^{i\theta})G(e^{i\theta})]X)^{1/2} \leq 1 - \alpha$$

a.e. for any constant matrix X with $|X| = 1$. From this it follows that $\alpha + G^*G \leq U^*G + G^*U$. We could just as readily obtain that $\alpha + GG^* \leq GU^* + UG^*$.

All the above steps are reversible so the theorem follows.

In order for (4.4) to hold, it is necessary that $\operatorname{ess} \inf \det G > 0$. In the scalar case, 4.2 and 4.3 are equivalent.

We continue to require that $\log \det W$ be summable.

THEOREM 4.5. *The manifolds $H_{n \times n}^2(W)$ and $\bar{H}_{0; n \times n}^2(W)$ are at positive angle if and only if there exists $M > 0$ such that $\|F\|_W \leq M \|F + G^*\|_W$ for all $F \in H_{n \times n}^2(W)$ and $G^* \in \bar{H}_{0; n \times n}^2(W)$.*

A proof of the above will not be given here since it is not significantly different from that of the scalar case.

Consider a trigonometric polynomial $\sum A_k e^{ik\theta}$. If $H_{n \times n}^2(W)$ and $\bar{H}_{0; n \times n}^2(W)$ are at

positive angle, then 4.5 implies the existence of $M > 0$ such that $\|\sum_{k=-s}^s A_k e^{ik\theta}\|_W \leq M \|\sum A_k e^{ik\theta}\|_W$ for $s=0, 1, 2, \dots$

LEMMA 4.6. *If $H_{n \times n}^2(W)$ and $\bar{H}_{0; n \times n}^2(W)$ are at positive angle, then $F \in L_{n \times n}^2(W)$ implies $F \in L_{n \times n}^1$.*

Proof. Suppose C is a constant matrix such that $W_0 C = 0$. Since $(W_0 C, C) = (WC, C) = \|W^{1/2} C\|_2^2 = 0$, $W^{1/2} C = 0$ which implies $C = 0$. Therefore W_0 is nonsingular.

The operator which sends each trigonometric polynomial into its constant term is bounded. Hence the linear functional $I(\cdot)$ defined for all trigonometric polynomials A by $I(A) = (A_0, I)_W = \text{tr } A_0 W_0$ is bounded on a dense subset of $L_{n \times n}^2(W)$. Hence there exists $K \in L_{n \times n}^2(W)$ such that $I(F) = (F, K)_W$ for all $F \in L_{n \times n}^2(W)$. It follows that $WK = W_0$, so $KW_0^{-1} = W^{-1} \in L_{n \times n}^2(W)$ and $\|W^{-1}\|_W^2 = \int \text{tr } W^{-1} d\mu < \infty$. Applying 3.7, we obtain

$$\|F\|_1 = \int \text{tr } (FF^*)^{1/2} W^{1/2} W^{-1/2} d\mu \leq \left(\int \text{tr } FF^* W d\mu \right)^{1/2} \left(\int \text{tr } W^{-1} d\mu \right)^{1/2} < \infty.$$

THEOREM 4.7. *If $H_{n \times n}^2(W)$ and $\bar{H}_{0; n \times n}^2(W)$ are at positive angle, then the Fourier series of $F \in L_{n \times n}^2(W)$ converges to F in norm.*

Proof. For all trigonometric polynomials $\sum A_k e^{ik\theta}$, let $Q_s(\sum A_k e^{ik\theta}) = \sum_{k=-s}^s A_k e^{ik\theta}$; $s=0, 1, 2, \dots$. Then Q_s has a bounded extension to all of $L_{n \times n}^2(W)$. Also for any trigonometric polynomial A , $\|(Q_s - I)A\|_W \rightarrow 0$ as $s \rightarrow \infty$.

Let $\alpha > 0$ and $F \in L_{n \times n}^2(W)$. There exists a trigonometric polynomial F_α such that $\|F - F_\alpha\|_W < \alpha$. Now

$$\|(Q_s - I)F\|_W \leq \|(Q_s - I)(F - F_\alpha)\|_W + \|(Q_s - I)F_\alpha\|_W \leq M\|F - F_\alpha\|_W + \|(Q_s - I)F_\alpha\|_W$$

where $M > 0$. The theorem follows.

5. Invertibility of systems of Toeplitz operators on H^2 .

THEOREM 5.1. *Let $\phi = G^*H$ where $H, H^{-1}, G, G^{-1} \in H_{n \times n}^2$ and $HH^* = (GG^*)^{-1}$. If $W = GG^*$, then T_ϕ is invertible if and only if there exists $B > 0$ such that $\|F\|_W \leq B\|F + K^*\|_W$ for all $F \in H_{n \times n}^2(W)$ and $K^* \in \bar{H}_{0; n \times n}^2(W)$.*

Proof. Suppose $R \in H_{n \times n}^2$ is orthogonal to the range of T_ϕ . Then for all $F \in H_{n \times n}^2$, $0 = (T_\phi F, R) = \int \text{tr } \phi F R^* d\mu = (HF, GR)$. Since H is outer, all products of the form HF , $F \in H_{n \times n}^2$, are dense in $H_{n \times n}^2$. Thus $GR = 0$ and $R = 0$. Hence the range of T_ϕ is dense in $H_{n \times n}^2$.

Let $N \in H_{n \times n}^2$, then $N = H^{-1}F$ where $F = HN \in H_{n \times n}^1$. Similarly, for $M \in H_{0; n \times n}^2$, $M^* = G^*E^*$ where $E = M(G^{-1}) \in H_{0; n \times n}^1$. Since H^{-1} is outer (see 2.3), there exists a sequence $\{P_s\}_{s=1}^\infty \subset \Delta_{n \times n}$ such that $H^{-1}P_s \in H_{n \times n}^2$ for $s=1, 2, 3, \dots$ and $\lim_{s \rightarrow \infty} \|H^{-1}P_s - N\|_2 = 0$. For $s=1, 2, 3, \dots$, $\|H^{-1}P_s - N\|_2^2 = \|P_s - F\|_W^2$ so $\lim_{s \rightarrow \infty} \|P_s - F\|_W = 0$ and $F \in H_{n \times n}^2(W)$. Similarly, $E^* \in \bar{H}_{0; n \times n}^2(W)$.

Now consider $T_\phi(H^{-1}F) = \phi H^{-1}F - (I-P)\phi H^{-1}F$. Since $(I-P)\phi H^{-1}F = G^*E_r^*$ for some $E_r \in H_{0;n \times n}^1$, the expression can be rewritten as $T_\phi(H^{-1}F) = G^*F + G^*E_r^*$. Note that $\|H^{-1}F\|_2 = \|F\|_w$. It follows that $\|N\|_2^2 = \|F\|_w^2 \leq B^2\|F + E_r^*\|_w^2 = B^2\|G^*F + G^*E_r^*\|_2^2 = B^2\|T_\phi(H^{-1}F)\|_2^2 = B^2\|T_\phi(N)\|_2^2$; that is, $\|N\|_2 \leq B\|T_\phi(N)\|_2$.

Conversely, suppose T_ϕ is invertible. Consider $F \in H_{n \times n}^2(W)$. There exists a sequence $\{P_s\}_{s=1}^\infty \subseteq \Delta_{n \times n}$ such that $\lim_{s \rightarrow \infty} \|P_s - F\|_w = 0$. Hence $\lim_{s \rightarrow \infty} \|H^{-1}P_s - H^{-1}F\|_2^2 = 0$ and $H^{-1}F \in H_{n \times n}^2$. Recall that $\|H^{-1}F\|_2 = \|F\|_w$. Note that $G^*E_r^*$ is the projection of G^*F onto $H_{n \times n}^2$. Now for all $E^* \in \bar{H}_{0;n \times n}^2(W)$, $\|F\|_w^2 = \|H^{-1}F\|_2^2 \leq B^2\|T_\phi(H^{-1}F)\|_2^2 = B^2\|G^*F + G^*E_r^*\|_2^2 \leq B^2\|G^*F + G^*E^*\|_2^2 = B^2\|H^{-1}F + H^{-1}E^*\|_2^2 = B^2\|F + E^*\|_w^2$; that is, $\|F\|_w \leq B\|F + E^*\|_w$ for some $B > 0$.

LEMMA 5.2. *Suppose $\phi \in L_{n \times n}^\infty$ is unitary and T_ϕ is invertible. Then there exists $G \in H_{n \times n}^\infty$ and $\alpha > 0$ such that $\alpha + G^*G \leq \phi^*G + G^*\phi$.*

Proof. It follows from 2.12 that $\phi = K^*H$ where $H, H^{-1}, K, K^{-1} \in H_{n \times n}^2$ and $HH^* = (KK^*)^{-1}$. Let $W = KK^*$. Then 5.1 implies that for some $B > 0$, $\|F\|_w \leq B\|F + K^*\|_w$ for all $F \in H_{n \times n}^2(W)$ and $K^* \in \bar{H}_{0;n \times n}^2(W)$. From 4.5, $H_{n \times n}^2(W)$ and $H_{0;n \times n}^2(W)$ are at positive angle. Since $W = K\phi H^{-1}$, ϕ has the same properties relative to W as U in 4.3. The lemma follows.

LEMMA 5.3. *Suppose $\phi \in L_{n \times n}^\infty$ is unitary and $\phi = K^*H$ where $H, H^{-1}, K, K^{-1} \in H_{n \times n}^2$. If there exists $G \in H_{n \times n}^\infty$ and $\alpha > 0$ such that $\alpha + G^*G \leq \phi^*G + G^*\phi$, then T_ϕ is invertible.*

Proof. Since ϕ is unitary, $HH^* = (KK^*)^{-1}$. Let $W = KK^*$, then $W = K\phi H^{-1}$. From 4.3, $H_{n \times n}^2(W)$ and $\bar{H}_{0;n \times n}^2(W)$ are at positive angle. From 4.5, there exists $B > 0$ such that $\|F\|_w \leq B\|F + E^*\|_w$ for all $F \in H_{n \times n}^2(W)$ and $E^* \in \bar{H}_{0;n \times n}^2(W)$. Now 5.1 implies T_ϕ is invertible.

THEOREM 5.4. *Let $\phi \in L_{n \times n}^\infty$ and suppose T_ϕ is invertible. Let $\phi = UK$ be the unitary-outer factorization of ϕ (see 2.11). Then*

- (i) $\text{ess inf } |\det \phi| > 0$;
- (ii) *there exists $G \in H_{n \times n}^\infty$ and $\alpha > 0$ such that $\alpha + G^*G \leq U^*G + G^*U$.*

Proof. For (i), see 2.7. Since 2.11 implies T_U is invertible, the theorem follows from 5.2.

THEOREM 5.5 *Let $\phi \in L_{n \times n}^\infty$ and suppose $\phi^{-1} \in L_{n \times n}^\infty$. Let $\phi = UK$ be the unitary outer factorization of ϕ . If $U = R^*H$ where $R, R^{-1}, H, H^{-1} \in H_{n \times n}^2$, and if there exists $G \in H_{n \times n}^\infty$ and $\alpha > 0$ such that $\alpha + G^*G \leq U^*G + G^*U$, then T_ϕ is invertible.*

Proof. Since $\phi^{-1} \in L_{n \times n}^\infty$, ϕ has a unitary-outer factorization $\phi = UK$ with T_K invertible. From 5.3, T_U is invertible. Since $T_\phi = T_U T_K$, T_ϕ is invertible.

In the scalar case, Devinatz [2] has shown the following.

THEOREM 5.6. Suppose $\phi \in L^\infty$. Then T_ϕ is invertible if and only if

- (i) $\text{ess inf } |\phi| > 0$;
- (ii) there exists $g \in H^\infty$ such that $g^{-1} \in H^\infty$ and $\alpha > 0$ such that $|\arg g\phi| \leq \pi/2 - \alpha$.

BIBLIOGRAPHY

1. F. F. Bonsall, *Dual extremum problems in the theory of functions*, J. London Math. Soc. **31** (1956), 105–110.
2. A. Devinatz, *Toeplitz operator on H^2 spaces*, Trans. Amer. Math. Soc. **112** (1964), 304–317.
3. P. Hartman and A. Wintner, *The spectra of Toeplitz matrices*, Amer. J. Math. **76** (1954), 867–882.
4. H. Helson, *Lectures on invariant subspaces*, Academic Press, New York, 1964.
5. H. Helson and D. Lowdenslager, *Prediction theory and Fourier series in several variables*, Acta Math. **99** (1958), 165–202.
6. H. Helson and G. Szego, *A problem in prediction theory*, Ann. Mat. Pura Appl. **51** (1960), 107–138.
7. P. Masani, *Shift invariant spaces and prediction theory*, Acta Math. **107** (1962), 275–290.
8. H. Pousson, *Systems of Toeplitz operators on H^2* , Proc. Amer. Math. Soc. **19** (1968), 603–608.
9. J. H. M. Wedderburn, *Lectures on matrices*. Amer. Math. Soc. Colloq. Publ., vol. 17, Amer. Math. Soc., Providence, R. I., 1934.
10. H. Widom, *Inversion of Toeplitz matrices*, II, Illinois J. Math. **4** (1960), 88–99.
11. N. Wiener and P. Masani, *Multivariate stochastic processes*. I, Acta Math. **98** (1957), 111–150.

COLLEGE OF WILLIAM AND MARY,
WILLIAMSBURG, VIRGINIA
UNIVERSITY OF KENTUCKY,
LEXINGTON, KENTUCKY