

# ALGEBRAIC MODELS FOR MEASURE PRESERVING TRANSFORMATIONS

BY

N. DINCULEANU AND C. FOIAŞ

**1. Introduction.** The purpose of this paper is to study measure preserving transformations  $T$  on probability measure spaces  $(X, \Sigma, \mu)$  by means of algebraic models  $(\Gamma, U, \varphi)$  (see Definitions 1 and 2).

The results obtained here contain those obtained in [3] concerning algebraic models  $(\Gamma, \varphi)$  of measure spaces  $(X, \Sigma, \mu)$ .

Each transformation possesses algebraic models and conversely every algebraic system is a model for a certain transformation (Theorem 2). Algebraic models determine transformations uniquely up to a conjugacy (Theorem 1).

Transformations with discrete models (see Definition 3) are uniquely determined by  $(\Gamma, U)$  (Theorem 3). Such transformations are characterized by the existence of an orthonormal basis  $\Gamma' \subset L^2(\mu)$  of functions  $|f| \equiv 1$ , which is also a multiplicative group, such that  $U_T \Gamma' \subset C \cdot \Gamma'$  (direct product), where  $C$  is the circle group (Theorem 5). In certain cases, conjugacy does no more involve  $U$  either (Theorem 4). Continuous automorphisms and rotations on an abelian compact group—equipped with Haar measure—are examples of transformations with discrete model (Corollary of Theorem 5), and in fact, every *invertible* transformation with discrete model is a superposition of an automorphism and a rotation (Theorem 6).

The class of transformations with discrete models contains the transformations with quasi-discrete spectrum (see Abramov [1]) and the transformations with discrete spectrum (see Halmos [5]). Necessary and sufficient conditions are given for algebraic systems in order to be models for transformations with quasi-discrete spectrum (Theorem 10) or with discrete spectrum (Theorem 11). We mention also Theorem 12 which gives necessary and sufficient conditions in order that  $\Gamma_1 = \Gamma_\infty$ .

In Theorems 7 and 9, ergodicity of transformations is characterized by means of algebraic models.

**2. Preliminaries.** Let  $(X, \Sigma, \mu)$  be a probability measure space and  $T: X \rightarrow X$  a measure preserving transformation.

(1) We denote by  $\Gamma(\mu)$  the multiplicative group of the (equivalence classes of) functions  $f \in L^\infty(\mu)$  with  $|f| \equiv 1$ , by  $\varphi_\mu$  the function of positive type on  $\Gamma(\mu)$  defined by

$$\varphi_\mu(f) = \int f d\mu, \quad \text{for } f \in \Gamma(\mu)$$

---

Received by the editors February 26, 1967 and, in revised form, April 12, 1967.

and by  $U_T$  the linear isometry defined on  $L^2(\mu)$  by  $U_T f = f \circ T$ . Then [3, Proposition 1],

$$\varphi_\mu(f) = 1 \quad \text{if and only if } f = 1$$

and  $U_T$  (or, more precisely, the restriction of  $U_T$  to  $\Gamma(\mu)$ ) is an *injective homomorphism* of  $\Gamma(\mu)$  into itself, such that

$$U_T c = c, \quad \text{for } c \in C \text{ (the circle group)}$$

and

$$\varphi_\mu(U_T f) = \varphi_\mu(f), \quad \text{for } f \in \Gamma(\mu).$$

If  $T$  is invertible, then  $U_T$  is an *automorphism* of  $\Gamma(\mu)$ .

(2) For every function  $f \in \Gamma(\mu)$  put

$$w_T(f) = U_T f \cdot \bar{f}.$$

Then  $w_T$  is a *homomorphism* of  $\Gamma(\mu)$  into itself and we have

$$U_T f = w_T(f) \cdot f.$$

$w_T(f)$  is called the *generalized proper value* corresponding to the *generalized proper function*  $f$  of  $U_T$ .

A subgroup  $\Gamma \subset \Gamma(\mu)$  is invariant under  $U_T$  (that is  $U_T \Gamma \subset \Gamma$ ) if and only if  $\Gamma$  is invariant under  $w_T$  (that is  $w_T \Gamma \subset \Gamma$ ).

(3) For every integer  $n \geq 0$  put

$$\Gamma_n = \Gamma_n(T) = w_T^{-n}(C) = \{f \in \Gamma(\mu); w_T^n(f) \in C\}.$$

In particular,  $\Gamma_0 = C$  and  $\Gamma_1$  is the set of the proper functions of  $U_T$  belonging to  $\Gamma(\mu)$ . Each  $\Gamma_n$  is a group invariant under  $U_T$  and  $\Gamma_n \subset \Gamma_{n+1}$  for every  $n$ . The set

$$\Gamma_\infty = \Gamma_\infty(T) = \bigcup_{n=0}^{\infty} \Gamma_n$$

is also a subgroup of  $\Gamma(\mu)$  invariant under  $U_T$ . Moreover, if  $\Gamma \subset \Gamma(\mu)$  is a group such that

$$C \subset \Gamma \quad \text{and} \quad w_T^{-1} \Gamma = \Gamma$$

then  $\Gamma_\infty \subset \Gamma$ . (In fact, for every  $n$  we have  $w_T^{-n}(C) \subset w_T^{-n} \Gamma \subset \Gamma$ ).

In particular, if  $\Gamma_{n+1} = \Gamma_n$  for some  $n$ , then  $\Gamma_\infty = \Gamma_n$ .

(4) For every integer  $k \geq 0$ ,  $U_T^k$  is an *injective homomorphism* of  $\Gamma(\mu)$  into itself and

$$\varphi_\mu(U_T^k f) = \varphi_\mu(f), \quad \text{for } f \in \Gamma(\mu).$$

If  $\Gamma \subset \Gamma(\mu)$  is invariant under  $U_T$ , then  $\Gamma$  is invariant under  $U_T^k$ .

For every  $n$  we have

$$\Gamma_n(T) \subset \Gamma_n(T^k)$$

therefore

$$\Gamma_\infty(T) \subset \Gamma_\infty(T^k).$$

(5) Let  $\Gamma$  be an abelian group containing a subgroup  $C'$  of the circle group  $C$ , and suppose that  $\Gamma = C' \cdot \Gamma'$  (direct product), where  $\Gamma'$  is a subgroup of  $\Gamma$ . Let further  $U: \Gamma \rightarrow \Gamma$  be an injective homomorphism such that

$$Uc = c, \quad \text{for } c \in C'.$$

For every  $\gamma \in \Gamma'$  we have  $U\gamma \in \Gamma$ , therefore, there exists a number  $\rho(\gamma) \in C'$  and an element  $V\gamma \in \Gamma'$  such that

$$U\gamma = \rho(\gamma)V\gamma.$$

Then  $\rho: \Gamma' \rightarrow C'$  is a homomorphism and  $V: \Gamma' \rightarrow \Gamma'$  is an injective homomorphism. Moreover, for every  $n$  there exists a homomorphism  $\rho_n: \Gamma' \rightarrow C'$  such that

$$U^n \gamma = \rho_n(\gamma)V^n \gamma, \quad \text{for } \gamma \in \Gamma'.$$

In particular, if  $\Gamma \subset \Gamma(\mu)$  and  $U = U_T$ , then  $V\gamma = \gamma$  and  $\rho(\gamma) = w_T(\gamma)$ , for  $\gamma \in \Gamma' \cap \Gamma_1(T)$ .

Conversely, if  $\rho: \Gamma' \rightarrow C'$  is a homomorphism and  $V: \Gamma' \rightarrow \Gamma'$  is an injective homomorphism, then the equality

$$U(c\gamma) = c\rho(\gamma)V(\gamma), \quad \text{for } c \in C' \quad \text{and} \quad \gamma \in \Gamma'$$

defines an injective homomorphism  $U: \Gamma \rightarrow \Gamma$  which satisfies

$$Uc = c, \quad \text{for } c \in C'$$

and

$$U\gamma = \rho(\gamma)V\gamma, \quad \text{for } \gamma \in \Gamma'.$$

(6) Let  $(X', \Sigma', \mu')$  be a probability measure space and  $T': X' \rightarrow X'$  a measure preserving transformation.

The transformations  $T$  and  $T'$  are *conjugate* (see [5, pp. 44–45]) if there exists a *linear isometry*

$$\phi: L^2(\mu) \rightarrow L^2(\mu')$$

such that

$$\phi L^2(\mu) = L^2(\mu'),$$

$$\phi(fg) = \phi f \cdot \phi g, \quad \text{for } f, g \in L^\infty(\mu)$$

and

$$\phi U_T = U_{T'} \phi.$$

It follows then that  $\phi L^\infty(\mu) = L^\infty(\mu')$  and

$$\|\phi f\|_\infty = \|f\|_\infty, \quad \text{for } f \in L^\infty(\mu).$$

**REMARK.** To say that  $T$  and  $T'$  are conjugate means that the measures  $\mu$  and  $\mu'$  are conjugate (see [3, Definition 1]) by means of a linear isometry  $\phi: L^2(\mu) \rightarrow L^2(\mu')$  which satisfies in addition the equality

$$\phi U_T = U_{T'} \phi.$$

The following proposition gives some conjugacy invariants connected to  $\Gamma(\mu)$ ,  $U_T$  and  $\phi_\mu$ .

**PROPOSITION 1.** *If  $T$  and  $T'$  are conjugate, then there exists an injective homomorphism  $\phi: \Gamma(\mu) \rightarrow \Gamma(\mu')$  having the following properties:*

- (i)  $\phi\Gamma(\mu) = \Gamma(\mu')$ ;
- (ii)  $\phi c = c$ , for  $c \in C$ ;
- (iii) If  $\Gamma \subset \Gamma(\mu)$  generates  $L^2(\mu)$ , then  $\phi\Gamma$  generates  $L^2(\mu')$ ;
- (iv) If  $\Gamma \subset \Gamma(\mu)$  is an orthonormal system in  $L^2(\mu)$  then  $\phi\Gamma$  is orthonormal in  $L^2(\mu')$ ;
- (v)  $\phi\Gamma_n(T) = \Gamma_n(T')$  and  $\phi\Gamma_\infty(T) = \Gamma_\infty(T')$ ;
- (vi)  $\varphi_\mu(f) = \varphi_{\mu'}(\phi f)$ , for  $f \in \Gamma(\mu)$ ;
- (vii)  $\phi U_T = U_{T'}\phi$  and  $\phi w_T = w_{T'}\phi$ .

In fact, if  $\phi$  is a linear isometry of  $L^2(\mu)$  onto  $L^2(\mu')$  realizing the conjugacy between  $T$  and  $T'$ , then the restriction of  $\phi$  to  $\Gamma(\mu)$ , still denoted by  $\phi$ , is the required isomorphism (see also [3, Proposition 2]).

**REMARK.** We shall see (corollary of Theorem 1) that, conversely, if  $\phi$  is an isomorphism of  $\Gamma(\mu)$  onto  $\Gamma(\mu')$  satisfying conditions (vi) and (vii), then  $T$  and  $T'$  are conjugate.

**3. Algebraic models.** The considerations of the preceding section lead to the following

**DEFINITION 1.** *A system  $(\Gamma, U, \varphi)$  consisting of an abelian group  $\Gamma$  with unit 1, an injective homomorphism  $U: \Gamma \rightarrow \Gamma$  and a complex function of positive type  $\varphi$  on  $\Gamma$  such that  $\varphi(\gamma) = 1$  if and only if  $\gamma = 1$  and  $\varphi(U\gamma) = \varphi(\gamma)$ , for  $\gamma \in \Gamma$ , is called an algebraic ergodic system (a.e. system).*

*Two a.e. systems  $(\Gamma, U, \varphi)$  and  $(\Gamma', U', \varphi')$  are said to be isomorphic if there exists an isomorphism  $\phi$  of  $\Gamma$  onto  $\Gamma'$  such that*

$$\varphi(\gamma) = \varphi'(\phi\gamma), \quad \text{for } \gamma \in \Gamma$$

*and*

$$\phi U = U' \phi.$$

If we define the homomorphisms  $w: \Gamma \rightarrow \Gamma$  by

$$w(\gamma) = U\gamma \cdot \gamma^{-1}, \quad \text{for } \gamma \in \Gamma$$

and the homomorphism  $w': \Gamma' \rightarrow \Gamma'$  in a similar way, then condition  $\phi U = U' \phi$  above is equivalent to condition  $\phi w = w' \phi$ .

**EXAMPLE.** If  $T$  is a measure preserving transformation on a probability measure space  $(X, \Sigma, \mu)$ , then  $(C, U_T, \varphi_\mu)$  and  $(\Gamma(\mu), U_T, \varphi_\mu)$  are a.e. systems. More generally, for every group  $\Gamma \subset \Gamma(\mu)$  invariant under  $U_T$ ,  $(\Gamma, U_T, \varphi_\mu)$  is an a.e. system.

We shall see (Theorem 2) that every a.e. system can be obtained in this way.

REMARKS. 1°. To say that  $(\Gamma, U, \varphi)$  is an a.e. system, means that  $(\Gamma, \varphi)$  is a measure system (see [3, Definition 2]) and that  $U: \Gamma \rightarrow \Gamma$  is an injective homomorphism satisfying  $\varphi(U\gamma) = \varphi(\gamma)$  for  $\gamma \in \Gamma$ . Then  $(\Gamma, \varphi \circ U)$  is also a measure system. Moreover, if  $U\Gamma = \Gamma$ , then  $(\Gamma, \varphi)$  and  $(\Gamma, \varphi \circ U)$  are isomorphic measure systems.

Conversely if  $(\Gamma, \varphi)$  and  $(\Gamma, \varphi')$  are isomorphic measure systems by means of an isomorphism  $U: \Gamma \rightarrow \Gamma$ , then  $(\Gamma, U, \varphi)$  is an a.e. system and  $U\Gamma = \Gamma$ .

2°. To say that two a.e. systems  $(\Gamma, U, \varphi)$  and  $(\Gamma', U', \varphi')$  are isomorphic, means that  $(\Gamma, \varphi)$  and  $(\Gamma', \varphi')$  are isomorphic measure systems, by means of an isomorphism  $\phi: \Gamma \rightarrow \Gamma'$  which satisfies  $\phi U = U' \phi$ .

Conversely, if  $(\Gamma, \varphi)$  and  $(\Gamma', \varphi')$  are isomorphic measure systems, then taking  $U: \Gamma \rightarrow \Gamma$  and  $U': \Gamma' \rightarrow \Gamma'$  the identity mappings, the a.e. systems  $(\Gamma, U, \varphi)$  and  $(\Gamma', U', \varphi')$  are isomorphic.

3°. If  $(\Gamma, U, \varphi)$  is an a.e. system, then the set  $C' = \{\gamma \in \Gamma; |\varphi(\gamma)| = 1\}$  is a group, and  $\varphi$  is an injective homomorphism of  $C'$  into the circle group  $C$ . If we identify an element  $\gamma \in C'$  with the number  $\varphi(\gamma) = c$ , we have (see [3, corollary of Proposition 3])

$$\varphi(c\gamma) = c\varphi(\gamma), \quad \text{for } c \in C' \quad \text{and} \quad \gamma \in \Gamma.$$

Moreover,

$$Uc = c, \quad \text{for } c \in C'.$$

In fact, if  $c \in C'$ , then  $\varphi(Uc) = \varphi(c) = c$ , therefore  $Uc \in C'$  and  $Uc = c$ .

If  $C'$  is divisible, then there exists a group  $\Gamma' \subset \Gamma$  such that

$$\Gamma = C' \cdot \Gamma' \quad (\text{direct product}).$$

The a.e. system  $(\Gamma, U, \varphi)$  can be embedded in an a.e. system  $(\Gamma_1, U_1, \varphi_1)$  such that

$$\{\gamma \in \Gamma_1; |\varphi_1(\gamma)| = 1\} = C$$

and then

$$\Gamma_1 = C \cdot \Gamma'_1 \quad (\text{direct product}).$$

In case  $U\gamma = \gamma$  (or, equivalently,  $w(\gamma) = 1$ ) implies  $\gamma \in C$ , the group  $\Gamma'$  can be precised:

PROPOSITION 2. Let  $(\Gamma, U, \varphi)$  be an a.e. system, let  $C' = \{\gamma \in \Gamma; \varphi(\gamma) \in C\}$  and  $w(\gamma) = U\gamma \cdot \gamma^{-1}$ , for  $\gamma \in \Gamma$ .

If  $C'$  is divisible (in particular if  $C' = C$ ) and if  $w(\gamma) = 1$  implies  $\gamma \in C'$ , then every injective homomorphism  $a \rightarrow \gamma_a$  of a group  $G \subset w\Gamma$  into  $\Gamma$  such that  $w(\gamma_a) = a$  for  $a \in G$  (in particular the homomorphism  $1 \rightarrow \gamma_1 = 1$  of  $G = \{1\}$ ) can be extended to an injective homomorphism  $a \rightarrow \gamma_a$  of  $w\Gamma$  into  $\Gamma$ , such that  $w(\gamma_a) = a$ , for  $a \in w\Gamma$ .

If we put  $\Gamma' = \{\gamma_a; a \in w\Gamma\}$ , then  $\Gamma = C' \cdot \Gamma'$  (direct product).

The proof is similar to that given in [5, p. 46], for ergodic transformations with discrete spectrum.

For every  $a \in w\Gamma$  choose  $\mu_a \in \Gamma$  with  $U\mu_a = a\mu_a$ , that is  $w(\mu_a) = a$ . If  $a \in G$  we take  $\mu_a = \gamma_a$ . We have

$$U\mu_{ab} = ab\mu_{ab} \quad \text{and} \quad U\mu_a\mu_b = ab\mu_a\mu_b$$

whence

$$w(\mu_{ab}) = w(\mu_a\mu_b) = ab.$$

By hypothesis, there exists a number  $\gamma(a, b) \in C'$  such that

$$\mu_a\mu_b = \gamma(a, b)\mu_{ab}.$$

If  $a, b \in G$ , then  $\gamma(a, b) = 1$ . Consider the group  $\{c\gamma_a; c \in C', a \in G\}$  and the homomorphism  $p$  of this group into  $C'$  defined by  $p(c\gamma_a) = c$ . We have, in particular,  $p(c) = c$  for  $c \in C'$  and  $p(\gamma_a) = 1$  for  $a \in G$ . Since  $C'$  is divisible,  $p$  can be extended to a homomorphism, still denoted by  $p$ , of  $w\Gamma$  into  $C'$ .

If we now define

$$\gamma_a = \overline{p(\mu_a)}\mu_a, \quad \text{for } a \in w\Gamma$$

then the requirements of the proposition are fulfilled.

REMARK. Condition:  $w(\gamma) = 1$  implies  $\gamma \in C'$ , is satisfied, for example, if  $U = U_T$ , where  $T$  is an ergodic transformation.

DEFINITION 2. Let  $(X, \Sigma, \mu)$  be a probability measure space and  $T: X \rightarrow X$  a measure preserving transformation. We say that an a.e. system  $(\Gamma, U, \varphi)$  is an algebraic model of the transformation  $T$  if there exists an injective homomorphism  $J: \Gamma \rightarrow \Gamma(\mu)$  such that:

- (a)  $J\Gamma$  generates  $L^2(\mu)$ ;
- (b)  $\varphi(\gamma) = \varphi_\mu(J\gamma)$ , for  $\gamma \in \Gamma$ ;
- (c)  $JU = U_T J$ .

It follows that if  $\Gamma \subset \Gamma(\mu)$  is a group generating  $L^2(\mu)$ , and invariant under  $U_T$ , then  $(\Gamma, U_T, \varphi_\mu)$  is an algebraic model for  $T$ .

If  $(\Gamma, U, \varphi)$  is an algebraic model of  $T$  by means of an isomorphism  $J$ , then, identifying  $\Gamma$  and  $J\Gamma$  we can consider that  $\Gamma \subset \Gamma(\mu)$ ,  $U = U_T$  and  $\varphi = \varphi_\mu$ .

If  $(\Gamma, U, \varphi)$  is an algebraic model of  $T$ , then  $T$  is invertible (that is  $U_T L^2(\mu) = L^2(\mu)$ ) if and only if  $U$  is an automorphism of  $\Gamma$  (that is  $U\Gamma = \Gamma$ ). In particular, a transformation  $T$  having  $(\Gamma_1(T), U_T, \varphi_\mu)$  as algebraic model, is always invertible (since  $U_T \Gamma_1 = \Gamma_1$ ).

REMARK. To say that  $(\Gamma, U, \varphi)$  is an algebraic model for  $T$  means that  $(\Gamma, \varphi)$  is an algebraic model for the measure  $\mu$  (see [3, Definition 3]), by means of an isomorphism  $J: \Gamma \rightarrow \Gamma(\mu)$  which satisfies, in addition,  $JU = U_T J$ .

Conversely, if  $(\Gamma, \varphi)$  is an algebraic model of the measure  $\mu$  and if  $U: \Gamma \rightarrow \Gamma$  is

the identity mapping, then  $(\Gamma, U, \varphi)$  is an algebraic model for the identity transformation  $T: X \rightarrow X$ .

Algebraic models determine the transformations uniquely up to a conjugacy:

**THEOREM 1.** *Two measure preserving transformations are conjugate if and only if they possess isomorphic algebraic models.*

Let  $T$  and  $T'$  be two measure preserving transformations on the probability measure spaces  $(X, \Sigma, \mu)$  respectively  $(X', \Sigma', \mu')$ .

If  $T$  and  $T'$  are conjugate, then from Proposition 1 we deduce that their algebraic models  $(\Gamma(\mu), U_T, \varphi_\mu)$  and  $(\Gamma(\mu'), U_{T'}, \varphi_{\mu'})$  are isomorphic.

Conversely, suppose that  $T$  and  $T'$  possess isomorphic models  $(\Gamma, U, \varphi)$  respectively  $(\Gamma', U', \varphi')$ . We may consider  $\Gamma \subset \Gamma(\mu)$ ,  $U = U_T$ ,  $\varphi = \varphi_\mu$  and  $\Gamma' \subset \Gamma(\mu')$ ,  $U' = U_{T'}$  and  $\varphi' = \varphi_{\mu'}$ .

If  $\phi$  is an isomorphism of  $\Gamma$  onto  $\Gamma'$  such that

$$\varphi_{\mu'} = \varphi_{\mu} \circ \phi \quad \text{and} \quad \phi U_T = U_{T'} \phi$$

then (see [3, Theorem 2]),  $\phi$  can be extended to a linear isometry  $\phi: L^2(\mu) \rightarrow L^2(\mu')$  such that

$$\phi L^2(\mu) = L^2(\mu') \quad \text{and} \quad \phi L^\infty(\mu) = L^\infty(\mu'),$$

and

$$\phi(fg) = \phi f \cdot \phi g, \quad \text{for } f, g \in L^\infty(\mu).$$

The equality

$$\phi U_T f = U_{T'} \phi f, \quad \text{for } f \in \Gamma$$

remains true first for linear combinations of functions of  $\Gamma$  and then for every  $f \in L^2(\mu)$ , so that  $T$  and  $T'$  are conjugate.

**COROLLARY.** *The transformations  $T$  and  $T'$  are conjugate if and only if the a.e. systems  $(\Gamma(\mu), U_T, \varphi_\mu)$  and  $(\Gamma(\mu'), U_{T'}, \varphi_{\mu'})$  are isomorphic.*

The following theorem states that every a.e. system is an algebraic model for some transformation.

**THEOREM 2.** *Every a.e. system  $(\Gamma, U, \varphi)$  is an algebraic model for a continuous measure preserving homomorphism  $\tau$  on an abelian compact group  $G$  equipped with a suitable regular Borel measure  $\mu$ .*

*Moreover, if  $U\Gamma = \Gamma$ , then  $\tau$  is an automorphism of  $G$ .*

Consider on  $\Gamma$  the discrete topology and take  $G = \Gamma^\wedge$ . Let  $\mu$  be the unique regular Borel measure on  $G$  such that (Bochner's theorem),

$$\varphi(\gamma) = \int \langle x, \gamma \rangle d\mu(x), \quad \text{for } \gamma \in \Gamma.$$

Then the mapping  $J: \Gamma \rightarrow \Gamma(\mu)$  defined by

$$J\gamma = \langle \cdot, \gamma \rangle, \quad \text{for } \gamma \in \Gamma$$

is an injective homomorphism,  $J\Gamma$  generates  $L^2(\mu)$  and

$$\varphi(\gamma) = \varphi_\mu(J\gamma), \quad \text{for } \gamma \in \Gamma.$$

We define now the mapping  $\tau: G \rightarrow G$  by

$$\langle \tau x, \gamma \rangle = \langle x, U\gamma \rangle, \quad \text{for } x \in G \quad \text{and} \quad \gamma \in \Gamma.$$

Then  $\tau$  is a continuous homomorphism of  $G$  into itself, and

$$JU = U_\tau J.$$

If  $\Gamma U = \Gamma$ , then  $\tau$  is injective and  $\tau G = G$ , therefore  $\tau$  is an automorphism of  $G$ .

It remains to prove that  $\tau$  is measure preserving.

Consider the regular Borel measure  $\nu$  defined on  $G$  by

$$\nu(A) = \mu(\tau^{-1}A), \quad \text{for every Borel set } A \subset G.$$

Then for every  $\gamma \in \Gamma$  we have

$$\varphi(\gamma) = \varphi(U\gamma) = \int \langle x, U\gamma \rangle d\mu(x) = \int \langle \tau x, \gamma \rangle d\mu(x) = \int \langle x, \gamma \rangle d\nu(x).$$

By the uniqueness of  $\mu$  we deduce that  $\mu = \nu$ , therefore  $\mu(\tau^{-1}A) = \mu(A)$ , for every Borel set  $A \subset G$  consequently  $\tau$  is measure preserving.

REMARK. The proof of Theorem 2 was used in [4] to prove the following

**COROLLARY.** *Every measure preserving transformation  $T$  on a probability measure space  $(X, \Sigma, \mu)$  is conjugate to a continuous homomorphism  $\tau$  on an abelian compact group  $G$  equipped with a suitable regular Borel measure. If  $T$  is invertible then  $\tau$  is an automorphism of  $G$ .*

#### 4. Discrete algebraic models.

**DEFINITION 3.** *An a.e. system  $(\Gamma, U, \varphi)$  is said to be discrete if  $C \subset \Gamma$  and*

$$\begin{aligned} \varphi(\gamma) &= \gamma, \quad \text{for } \gamma \in C, \\ &= 0, \quad \text{for } \gamma \notin C. \end{aligned}$$

**REMARKS.** 1°. An a.e. system  $(\Gamma, U, \varphi)$  is discrete if and only if  $(\Gamma, U^n, \varphi)$  is discrete.

2°. We have

$$Uc = c, \quad \text{and} \quad w(c) = 1, \quad \text{for } c \in C$$

where  $w(\gamma) = U\gamma \cdot \gamma^{-1}$  for  $\gamma \in \Gamma$  (see Remark 3 after Definition 1).



3°. Let  $(\Gamma, U, \varphi)$  be an a.e. system with  $C \subset \Gamma$ . Then  $\Gamma = C \cdot \Gamma'$  (direct product) where  $\Gamma'$  is a subgroup of  $\Gamma$ . To say that  $(\Gamma, U, \varphi)$  is discrete, means that

$$\begin{aligned}\varphi(\gamma) &= 1 \quad \text{for } \gamma = 1, \\ &= 0 \quad \text{for } \gamma \in \Gamma', \gamma \neq 1.\end{aligned}$$

4°. Let  $(\Gamma, U, \varphi)$  be an a.e. system such that

$$|\varphi(\gamma)| < 1 \quad \text{implies } \varphi(\gamma) = 0.$$

Then  $(\Gamma, U, \varphi)$  is "essentially" a discrete system. In fact we can consider  $(\Gamma, U, \varphi)$  as a model of a measure preserving transformation  $T$  on a probability measure space, and consider  $\Gamma \subset \Gamma(\mu)$ ,  $U = U_T$  and  $\varphi = \varphi_\mu$ . Consider then the group  $\Gamma_1 = \{c\gamma; c \in C, \gamma \in \Gamma\}$ ; then  $(\Gamma_1, U_T, \varphi_\mu)$  is a discrete model of  $T$  and contains the initial model  $(\Gamma, U, \varphi)$ .

For a discrete system  $(\Gamma, U, \varphi)$ , the function  $\varphi$  is completely determined by  $\Gamma$ , so that the system itself is completely determined by  $(\Gamma, U)$ .

**PROPOSITION 3.** *Let  $\Gamma$  be an abelian group containing  $C$  and let  $U: \Gamma \rightarrow \Gamma$  be an injective homomorphism such that*

$$Uc = c, \quad \text{for } c \in C.$$

*If we define*

$$\begin{aligned}\varphi(\gamma) &= \gamma \quad \text{if } \gamma \in C, \\ &= 0 \quad \text{if } \gamma \notin C,\end{aligned}$$

*then  $(\Gamma, U, \varphi)$  is a discrete system.*

In fact,  $\varphi$  is of positive type:

$$\begin{aligned}\sum_{i,j=1}^n \alpha_i \bar{\alpha}_j \varphi(\gamma_i \gamma_j^{-1}) &= \sum_{\gamma_i \sim \gamma_j} \alpha_i \bar{\alpha}_j \varphi(\gamma_i \gamma_j^{-1}) = \sum_k \sum_{\gamma_i, \gamma_j \in C\gamma_k} \alpha_i \bar{\alpha}_j \varphi(\gamma_i \gamma_j^{-1}) \\ &= \sum_k \sum_{\gamma_i \in C\gamma_k} |\alpha_i \varphi(\gamma_i \gamma_k^{-1})|^2 \geq 0\end{aligned}$$

where  $\gamma_i \sim \gamma_j$  means  $\gamma_i \gamma_j^{-1} \in C$  and  $C\gamma_k$  the equivalence classes.

If  $\gamma \in C$  then  $U\gamma = \gamma$ , therefore

$$\varphi(U\gamma) = \varphi(\gamma);$$

if  $\gamma \notin C$ , then  $U\gamma \notin C$  (since  $U$  is injective), therefore  $\varphi(\gamma) = 0$  and  $\varphi(U\gamma) = 0$ , consequently

$$\varphi(U\gamma) = \varphi(\gamma).$$

Moreover,  $\varphi(\gamma) = 1$ , if and only if  $\gamma = 1$ , so that  $(\Gamma, U, \varphi)$  is a discrete system.

For discrete systems, isomorphism does no more involve functions of positive type.

**THEOREM 3.** *Two discrete systems  $(\Gamma, U, \varphi)$  and  $(\Gamma', U', \varphi')$  are isomorphic if and only if there exists an isomorphism  $\phi$  of  $\Gamma$  onto  $\Gamma'$  such that*

$$\phi c = c, \quad \text{for } c \in C$$

and

$$\phi U = U' \phi.$$

In fact, if the systems are isomorphic by an isomorphism  $\phi$ , then for every  $c \in C$  we have

$$\varphi'(\phi c) = \varphi(c) = c \neq 0$$

therefore  $\phi c \in C$ , and then

$$\varphi'(\phi c) = \phi c$$

consequently  $\phi c = c$ .

Conversely, let  $\phi: \Gamma \rightarrow \Gamma'$  be an isomorphism such that  $\phi c = c$  for  $c \in C$  and  $\phi U = U' \phi$ . We have to prove that  $\varphi = \varphi' \circ \phi$ . For  $c \in C$  we have  $\phi c = c$ , therefore

$$\varphi'(\phi c) = \phi c = c = \varphi(c).$$

If  $\gamma \notin C$ , then  $\phi \gamma \notin C$  (since  $\phi$  is injective), therefore  $\varphi'(\phi \gamma) = 0$  and  $\varphi(\gamma) = 0$ , consequently

$$\varphi'(\phi \gamma) = \varphi(\gamma).$$

**REMARK.** If  $(\Gamma, U, \varphi)$  is a discrete system, we shall say also that  $(\Gamma, U)$  is a discrete system. If  $(\Gamma, U_\gamma)$  is a discrete system and  $\Gamma \subset \Gamma(\mu)$ , for some transformation  $T$  on a measure space  $(X, \Sigma, \mu)$ , we understand that  $\varphi = \varphi_\mu$ .

From Proposition 3 it follows that  $(\Gamma, U)$  is a discrete system provided that  $\Gamma$  is an abelian group containing  $C$  and  $U: \Gamma \rightarrow \Gamma$  is an injective homomorphism such that  $Uc = c$  for  $c \in C$ .

For certain discrete models  $(\Gamma, U)$  isomorphism does no more involve homomorphisms  $U$  either:

**THEOREM 4.** *Let  $(\Gamma_1, U_1)$  and  $(\Gamma_2, U_2)$  be two discrete systems and put*

$$w_i(\gamma) = U_i \gamma \cdot \gamma^{-1}, \quad \text{for } \gamma \in \Gamma_i, \quad i = 1, 2.$$

*Suppose that*

$$\gamma \in \Gamma_i \quad \text{and} \quad w_i(\gamma) = 1 \quad \text{imply} \quad \gamma \in C, \quad i = 1, 2.$$

*Then  $(\Gamma_1, U_1)$  and  $(\Gamma_2, U_2)$  are isomorphic, if and only if the groups  $w_1 \Gamma_1$  and  $w_2 \Gamma_2$  are isomorphic by an isomorphism  $\phi$  such that  $\phi w_1 = w_2 \phi$  and  $\phi c = c$  for  $c \in C \cap w_1 \Gamma_1$ .*

If  $(\Gamma_1, U_1)$  and  $(\Gamma_2, U_2)$  are isomorphic by means of an isomorphism  $\phi: \Gamma_1 \rightarrow \Gamma_2$  such that

$$\phi U_1 = U_2 \phi \quad \text{and} \quad \phi c = c \quad \text{for } c \in C,$$

then we have also

$$\phi w_1 = w_2 \phi.$$

From  $\phi\Gamma_1 = \Gamma_2$  we deduce then  $\phi w_1\Gamma_1 = w_2\Gamma_2$ . The restriction of  $\phi$  to  $w_1\Gamma_1$  is the required isomorphism.

Conversely, suppose that  $w_1\Gamma_1$  and  $w_2\Gamma_2$  are isomorphic by means of an isomorphism  $\phi: w_1\Gamma_1 \rightarrow w_2\Gamma_2$  such that  $\phi w_1 = w_2\phi$  and  $\phi c = c$  for  $c \in C \cap w_1\Gamma_1$ .

By Proposition 2 there exists an injective homomorphism  $a \rightarrow \gamma_a$  of  $w_1\Gamma_1$  into  $\Gamma_1$  such that  $w_1(\gamma_a) = a$  for  $a \in w_1\Gamma_1$ ; then  $\Gamma_1 = C \cdot \Gamma'_1$  (direct product) where  $\Gamma'_1 = \{\gamma_a; a \in w_1\Gamma_1\}$ .

Consider the groups  $G_1 = w_1^2\Gamma_1$  and  $G_2 = w_2^2\Gamma_2$ . Since  $\phi w_1\Gamma_1 = w_2\Gamma_2$  and  $\phi w_1 = w_2\phi$  we have  $G_2 = \phi G_1$ .

If  $a \in G_1$  then  $w_1(\gamma_a) = a$  and  $a = w_1(b)$  for some  $b \in w_1\Gamma_1$ , therefore  $\gamma_a = cb$  for some  $c \in C$ ; if we have also  $\gamma_a = c_1 b_1$  with  $c_1 \in C$  and  $b_1 \in w_1\Gamma_1$ , then  $c\bar{c}_1 = b_1 b^{-1} \in w_1\Gamma_1$ , therefore, by hypothesis,

$$c\bar{c}_1 = \phi(c\bar{c}_1) = \phi(b_1)\overline{\phi(b)},$$

whence  $c\phi b = c_1\phi b_1$ . We define then unambiguously

$$\gamma_{\phi a} = c\phi b, \text{ if } \gamma_a = cb \text{ with } c \in C \text{ and } b \in w_1\Gamma_1.$$

It is easy to see that  $\phi a \rightarrow \gamma_{\phi a}$  is an injective homomorphism of  $G_2$  into  $\Gamma_2$  such that  $w_2(\phi a) = \phi a$ . By Proposition 2, this homomorphism can be extended to an injective homomorphism  $a \rightarrow \gamma_a$  of  $w_2\Gamma_2$  into  $\Gamma_2$  such that  $w_2(\gamma_a) = a$  for  $a \in w_2\Gamma_2$ .

We extend now  $\phi$  from  $w_1\Gamma_1$  to  $\Gamma_1$  by

$$\psi c\gamma_a = c\gamma_{\phi a} \text{ for } c \in C \text{ and } a \in w_1\Gamma_1.$$

$\psi$  is an extension of  $\phi$ , since if  $b \in w_1\Gamma_1$ , then  $b = c\gamma_a$  for some  $c \in C$  and  $a \in w_1\Gamma_1$ , whence  $a = w_1(\gamma_a) = w_1(b) \in w_1^2\Gamma_1$  and  $\gamma_a = \bar{c}b$ , therefore  $\gamma_{\phi a} = \bar{c}\phi b$ ; it follows then that  $\phi b = c\gamma_{\phi a} = \psi(c\gamma_a) = \psi b$ .

Moreover,  $\psi$  is an isomorphism of  $\Gamma_1$  onto  $\Gamma_2$  and  $\psi c = c$  for  $c \in C$ . Finally, if  $c \in C$  and  $\gamma_a \in \Gamma'_1$ , we have

$$\psi U_1 c\gamma_a = \psi c a \gamma_a = \psi a \cdot \psi c \gamma_a = \phi a \cdot c \gamma_{\phi a} = U_2 c \gamma_{\phi a} = U_2 \psi c \gamma_a$$

therefore  $\psi U_1 = U\psi_2$ . By Theorem 3,  $(\Gamma_1, U_1)$  and  $(\Gamma_2, U_2)$  are isomorphic.

For transformations with discrete models we have the following characterization:

**THEOREM 5.** *A measure preserving transformation  $T$  on a probability measure space  $(X, \Sigma, \mu)$  has a discrete model if and only if there exists a set  $\Gamma' \subset \Gamma(\mu)$  such that*

- (a)  $\Gamma'$  is a group;
- (b)  $\Gamma'$  is an orthonormal basis of  $L^2(\mu)$ ;
- (c)  $U_T \Gamma' \subset C\Gamma'$ .

We remark first that if  $\Gamma'$  is a group and an orthonormal basis in  $L^2(\mu)$ , then  $\Gamma'$  contains no constant function except 1, so that  $C \cdot \Gamma'$  is a direct product.

If conditions a, b and c are satisfied, then  $(C \cdot \Gamma', U_T, \varphi_\mu)$  is a discrete algebraic model for  $T$ . In fact,  $C \subset C \cdot \Gamma'$  and  $C \cdot \Gamma'$  generates  $L^2(\mu)$ ; if  $c \in C$ , then

$$\varphi_\mu(c) = \int c \, d\mu = c$$

while if  $\gamma \notin C$ , then  $\gamma = c \cdot \gamma'$  for some  $c \in C$  and  $\gamma' \in \Gamma'$  with  $\gamma' \neq 1$ , therefore

$$\varphi_\mu(\gamma) = c \int \gamma' d\mu = c(\gamma'|1) = 0.$$

Conversely, let  $(\Gamma, U, \varphi)$  be a discrete algebraic model for  $T$ ; we may suppose  $\Gamma \subset \Gamma(\mu)$ ,  $U = U_T$  and  $\varphi = \varphi_\mu$ . Write  $\Gamma$  as a direct product  $\Gamma = C \cdot \Gamma'$ , where  $\Gamma'$  is a subgroup of  $\Gamma$ , containing no constant function except 1. Finally,  $\Gamma'$  is an orthonormal system, since for  $\gamma' \in \Gamma'$  we have

$$\begin{aligned} \int \gamma' d\mu &= \varphi(\gamma') = 1 \quad \text{if } \gamma' = 1, \\ &= 0 \quad \text{if } \gamma' \neq 1. \end{aligned}$$

**COROLLARY.** *If  $G$  is an abelian compact group, equipped with Haar measure  $\mu$ , then continuous automorphisms  $\tau'$  and rotations  $R$  on  $G$ , as well as their superpositions  $\tau = R\tau'$ , have discrete model.*

We remark first that continuous automorphisms  $\tau'$  and rotations  $R$ , therefore, their superpositions  $\tau = R\tau'$ , are measure preserving.

The group of characters  $\Gamma' = G^\wedge$  is an orthogonal system in  $L^2(\mu)$  and  $U_{\tau'}\Gamma' \subset \Gamma'$ ; if  $R$  is defined on  $G$  by  $Rx = cx$ , for some  $c \in G$ , then

$$U_{\tau'}\gamma(x) = \gamma(\tau x) = \gamma(c)\gamma(\tau'x) = \gamma(c)U_{\tau'}\gamma(x)$$

for every  $\gamma \in \Gamma'$ , therefore  $U_{\tau'}\Gamma' \subset C \cdot \Gamma'$ . By Theorem 5,  $\tau$  has discrete model. Conversely:

**THEOREM 6.** *Every invertible measure preserving transformation  $T$ , with discrete model  $(\Gamma, U_T)$ , on a probability measure space  $(X, \Sigma, \mu)$ , is conjugate to the superposition of a continuous automorphism and a rotation on an abelian compact group, equipped with Haar measure.*

Consider  $\Gamma = C \cdot \Gamma'$  (direct product) and

$$U_T\gamma = \rho(\gamma)V\gamma, \quad \text{for } \gamma \in \Gamma'$$

where  $\rho$  is a character of  $\Gamma'$  and  $V$  is an injective homomorphism of  $\Gamma'$ . Since  $T$  is invertible, we have  $U_T L^2(\mu) = L^2(\mu)$ , therefore  $V\Gamma' = \Gamma'$ . Consider  $\Gamma'$  endowed with the discrete topology and consider the Haar measure  $\nu$  on the abelian compact group  $G = \Gamma'^\wedge$ . Then  $\rho \in G$ . We define the continuous homomorphism  $\tau'$  on  $G$  by

$$\langle \tau'x, \gamma \rangle = \langle x, V\gamma \rangle, \quad \text{for } x \in G \text{ and } \gamma \in \Gamma'.$$

Since  $V\Gamma' = \Gamma'$ ,  $\tau'$  is an automorphism. Consider finally the mapping  $\tau: G \rightarrow G$  defined by

$$\tau(x) = \rho\tau'(x), \quad \text{for } x \in G.$$

Then  $(C \cdot G^\wedge, U_\tau)$  is a discrete model for  $\tau$ , and the mapping  $\phi: C \cdot \Gamma' \rightarrow C \cdot G^\wedge$  defined by

$$\phi c\gamma = c\langle \cdot, \gamma \rangle, \quad \text{for } c \in C \text{ and } \gamma \in \Gamma'$$

is an isomorphism such that  $\phi c = c$  for  $c \in C$ . Moreover, for  $\gamma \in \Gamma'$  we have

$$\begin{aligned} \phi U_T \gamma &= \phi \rho(\gamma) V \gamma = \rho(\gamma) \langle \cdot, V \gamma \rangle \\ &= \rho(\gamma) \langle \tau' \cdot, \gamma \rangle = \langle \rho \tau' \cdot, \gamma \rangle = \langle \tau \cdot, \gamma \rangle \\ &= U_\tau \langle \cdot, \gamma \rangle = U_\tau \phi \gamma \end{aligned}$$

and this equality remains valid for  $\gamma \in \Gamma$ , therefore  $\phi U_T = U_\tau \phi$ . By Theorem 3,  $T$  and  $\tau$  are conjugate.

**COROLLARY 1.** *A measure preserving transformation  $T$  on a probability measure space  $(X, \Sigma, \mu)$  is conjugate to a continuous automorphism on a compact abelian group, equipped with Haar measure, if and only if there exists a set  $\Gamma' \subset \Gamma(\mu)$  such that*

- (a)  $\Gamma'$  is a group;
- (b)  $\Gamma'$  is an orthonormal basis of  $L^2(\mu)$ ;
- (c)  $U_T \Gamma' = \Gamma'$ .

**COROLLARY 2.** *A measure preserving transformation  $T$  on a probability measure space  $(X, \Sigma, \mu)$  is conjugate to a rotation on an abelian compact group, equipped with Haar measure, if and only if  $T$  has a discrete model  $(\Gamma, U_T)$  with  $\Gamma \subset \Gamma_1$ .*

We mention also the following property of discrete models.

**PROPOSITION 4.** *Let  $T$  be a measure preserving transformation on a probability measure space  $(X, \Sigma, \mu)$  and let  $(\Gamma, U_T, \varphi_\mu)$ ,  $(\Gamma', U_T, \varphi_\mu)$  be two discrete systems.*

*If  $(\Gamma, U_T, \varphi_\mu)$  is a discrete model for  $T$  and if  $\Gamma' \subset \Gamma$ , then  $\Gamma = \Gamma'$ .*

In fact, let  $f \in \Gamma'$ . If for every  $g \in \Gamma$  we had  $fg \notin C$ , then

$$\int fg \, d\mu = 0$$

therefore  $f \equiv 0$ , which would contradict  $|f| \equiv 1$ .

It follows that there exists  $g \in \Gamma$  with  $fg \in C$ .

Then  $f \in \bar{g}C \subset \Gamma$ , therefore  $\Gamma' = \Gamma$ .

**5. Ergodic transformations.** In this section we give some characterizations of ergodic transformations by means of their algebraic models.

Let  $(X, \Sigma, \mu)$  be a probability measure space and  $T: X \rightarrow X$  a measure preserving transformation. The transformation  $T$  is ergodic if  $f \in L^2(\mu)$  and  $U_T f = f$  imply  $f = \text{constant}$ .

**PROPOSITION 5.** *If  $T$  is ergodic, then  $(\Gamma_1(T), U_T)$  is a discrete system.*

In fact if  $\gamma \in \Gamma_1(T) - C$ , then  $U_T \gamma = c\gamma$  for some  $c \neq 1$  (because  $T$  is ergodic), therefore

$$\int \gamma d\mu = \int U_T \gamma d\mu = c \int \gamma d\mu$$

consequently

$$\int \gamma d\mu = 0.$$

REMARKS. 1°. If  $T^n$  is ergodic for some  $n$ , then  $T$  is ergodic, therefore  $(\Gamma_1(T), U_T)$  is a discrete system. Theorem 7 below states a somewhat converse property.

2°. We shall see (Corollary 1 of Proposition 6) that if  $T^n$  is ergodic for every  $n$ , then  $(\Gamma_\infty(T), U_T)$  is a discrete system.

LEMMA. If  $T$  has a discrete model  $(\Gamma, U_T)$  and if  $\Gamma_1(T) \subset \Gamma$ , then for every natural  $n$  we have

$$\Gamma_1(T^n) \cap \Gamma = \Gamma_1(T).$$

Consider  $\Gamma = C \cdot \Gamma'$ , where  $\Gamma'$  is a group and an orthonormal basis of  $L^2(\mu)$ . Consider the homomorphisms  $\rho_n: \Gamma' \rightarrow C$  and  $V: \Gamma' \rightarrow \Gamma'$  such that  $U_T^n \gamma = \rho_n(\gamma) V^n(\gamma)$ , for  $\gamma \in \Gamma'$ .

Let  $\gamma \in \Gamma_1(T^n) \cap \Gamma'$ . Then  $U_T^n \gamma = c\gamma$ , for some  $c \in C$ , therefore  $\rho_n(\gamma) = c$  and  $V^n \gamma = \gamma$ . Let  $k \leq n$  be the least natural number such that  $V^k \gamma = \gamma$  and consider the  $k$ -dimensional space  $K$  generated by  $\gamma, V\gamma, \dots, V^{k-1}\gamma$ . Then  $K$  is invariant under  $U_T$ , therefore there exists a basis  $f_1, \dots, f_k$  of  $K$  consisting of proper functions of  $U_T$ :

$$U_T f_i = c_i f_i, \quad \text{with } c_i \in C.$$

Then  $f_i \in \Gamma_1(T) \subset \Gamma$ . Moreover, we may take  $f_i \in \Gamma'$  (multiplying each  $f_i$  by a suitable number of  $C$ ). The basis  $(f_1, \dots, f_k)$  must then coincide with the basis  $(\gamma, V\gamma, \dots, V^{k-1}\gamma)$ ; for example  $f_1 = \gamma$ , therefore  $U_T \gamma = c_1 \gamma$ .

It follows that  $\gamma \in \Gamma_1(T)$ , therefore  $\Gamma_1(T^n) \cap \Gamma' \subset \Gamma_1(T)$ , consequently  $\Gamma_1(T^n) \cap \Gamma \subset \Gamma_1(T)$ .

The converse inclusion follows from  $\Gamma_1(T) \subset \Gamma_1(T^n)$ .

THEOREM 7. Suppose that  $T$  has a discrete model  $(\Gamma, U_T)$  and let  $n$  be a natural number. If:

- (a) either  $\Gamma \subset \Gamma_1(T)$ , or  $\Gamma_1(T) \subset \Gamma$ ;
- (b)  $\gamma \in \Gamma$  and  $U_T^n \gamma = \gamma$  imply  $\gamma \in C$ ;

then  $T^n$  is ergodic.

Let  $f \in L^2(\mu)$  be a function such that  $U_T^n f = f$  and prove that  $f$  is constant.

Consider  $\Gamma = C \cdot \Gamma'$ , where  $\Gamma'$  is a group and an orthonormal basis of  $L^2(\mu)$ . Then

$$f = \sum_{\gamma \in \Gamma'} \alpha(\gamma) \gamma$$

where

$$\alpha(\gamma) = \int f \bar{\gamma} d\mu, \quad \text{for every } \gamma \in \Gamma'.$$

For every natural number  $k \in N$  we have

$$U_T^{kn}f = f$$

and

$$U_T^{kn}f = \sum_{\gamma \in \Gamma'} \alpha(\gamma) \rho_{kn}(\gamma) V^{kn}\gamma$$

therefore

$$\alpha(V^{kn}\gamma) = \alpha(\gamma) \rho_{kn}(\gamma), \quad \text{for } k \in N \quad \text{and} \quad \gamma \in \Gamma',$$

whence

$$|\alpha(V^{kn}\gamma)| = |\alpha(\gamma)|, \quad \text{for } k \in N \quad \text{and} \quad \gamma \in \Gamma'.$$

We shall prove that for every element  $\gamma \neq 1$  of  $\Gamma'$  we have  $\alpha(\gamma) = 0$ . Let therefore  $\gamma \in \Gamma'$  be such that  $\gamma \neq 1$ .

If  $V^{kn}\gamma = \gamma$  for some  $k$ , then  $\gamma \in \Gamma_1(T^n)$ . In fact, if  $\Gamma \subset \Gamma_1(T)$ , then  $\gamma \in \Gamma_1(T^n)$  without any other assumption, while if  $\Gamma_1(T) \subset \Gamma$ , then by the preceding lemma

$$\gamma \in \Gamma_1(T^{kn}) \cap \Gamma = \Gamma_1(T) = \Gamma_1(T^n) \cap \Gamma.$$

Writing now the equality  $\alpha(V^{kn}\gamma) = \alpha(\gamma) \rho_{kn}(\gamma)$  for  $k=1$  we obtain

$$\alpha(\gamma) = \alpha(\gamma) \rho_n(\gamma)$$

therefore either  $\alpha(\gamma) = 0$  or  $\rho_n(\gamma) = 1$ . But  $\rho_n(\gamma) = 1$  means  $U_T^n \gamma = \gamma$ , which by hypothesis implies  $\gamma = 1$  and we get a contradiction. It follows that  $\alpha(\gamma) = 0$ .

If  $V^{kn}\gamma \neq \gamma$  for every  $k$ , then the functions  $\gamma, V^n\gamma, V^{2n}\gamma, \dots$  are different from each other, therefore

$$\sum_{k=0}^{\infty} |\alpha(V^{kn}\gamma)|^2 \leq \sum_{\gamma' \in \Gamma'} |\alpha(\gamma')|^2 < \infty$$

consequently  $|\alpha(V^{kn}\gamma)| \rightarrow 0$  as  $k \rightarrow \infty$ , whence  $\alpha(\gamma) = 0$ .

It follows that  $f = \alpha(1)1$ , that is  $f$  is constant, consequently  $T^n$  is ergodic.

REMARKS. 1°. Is it possible to drop condition (a) in the preceding theorem? The answer is positive if condition (b) is satisfied for every  $n$  (see Theorem 9 below).

2°. Is it true that if  $T$  is ergodic, then  $\Gamma_1(T) \subset \Gamma$  for every discrete model  $(\Gamma, U_T)$  of  $T$ ?

The answer is positive if, in addition,  $T^n$  is ergodic for every  $n$ . Moreover, in this case we have  $\Gamma_\infty(T) \subset \Gamma$  for every discrete model  $(\Gamma, U_T)$  of  $T$  (see Corollary 2 of Proposition 6).

For ergodic transformations, we have the following conjugacy criterion:

**THEOREM 8.** *Two ergodic transformations  $T$  and  $T'$  with discrete model, are conjugate if and only if there exist discrete models  $(\Gamma, U_T)$  and  $(\Gamma', U_{T'})$  of  $T$  and  $T'$  respectively, such that the groups  $w_T\Gamma$  and  $w_{T'}\Gamma'$  are isomorphic by an isomorphism  $\phi$  such that  $\phi w_T = w_{T'} \phi$  and  $\phi c = c$  for  $c \in C \cap w_T\Gamma$ .*

We use Theorem 4.

**6. Transformations with ergodic iterates.** Let  $(X, \Sigma, \mu)$  be a probability measure space and  $T: X \rightarrow X$  a measure preserving transformation.

**PROPOSITION 6.** *Suppose that  $T^n$  is ergodic for every  $n$ . If  $(\Gamma, U_T)$  is a discrete system, then  $(\bigcup_{n=0}^{\infty} w_T^{-n}(\Gamma), U_T)$  is again a discrete system.*

We prove first that  $(w_T^{-1}(\Gamma), U_T)$  is a discrete system.

It is clear that  $w_T^{-1}(\Gamma)$  is a subgroup of  $\Gamma(\mu)$  invariant under  $U_T$  and containing  $C$ . We have to prove that

$$\varphi_\mu(f) = \int f d\mu = 0, \quad \text{for } f \in w_T^{-1}(\Gamma) - C.$$

Let  $f \in w_T^{-1}(\Gamma) - C$ . There are two possibilities:

(a)  $\int U_T^n f \cdot \bar{f} d\mu = 0$ , for every  $n \geq 1$ .

Then  $f, U_T f, U_T^2 f, \dots$  is an orthonormal system in  $L^2(\mu)$ . If  $g$  is the projection of 1 on the space generated by this sequence, we have

$$g = \sum_{n=0}^{\infty} a_n U_T^n f \quad \text{with } \sum |a_n|^2 < \infty.$$

Then

$$\int f d\mu = \int U_T^n f d\mu = (U_T^n f | 1) = (U_T^n f | g) = a_n$$

and  $a_n \rightarrow 0$ , therefore  $\int f d\mu = 0$ .

(b) There exists  $n$  such that

$$\int U_T^n f \cdot \bar{f} d\mu \neq 0.$$

Since

$$U_T^n f \cdot \bar{f} = w_T(f \cdot U_T f \cdot \dots \cdot U_T^{n-1} f) \in \Gamma$$

and since  $(\Gamma, U_T)$  is a discrete system, we have

$$U_T^n f \cdot \bar{f} \in C$$

that is

$$U_T^n f = cf \quad \text{for some } c \in C.$$

Since  $f$  is not constant and  $T^n$  is ergodic, we have  $c \neq 1$ .

Suppose that  $n$  is the least natural number satisfying  $U_T^n f = cf$ . The  $n$ -dimensional space  $K$  generated by  $f, U_T f, \dots, U_T^{n-1} f$  is invariant under  $U_T$ , therefore there exists a basis  $f_1, \dots, f_n$  of  $K$  consisting of proper functions of  $U_T$ :

$$U_T f_i = c_i f_i, \quad \text{with } c_i \in C.$$

Each  $f_i$  is of the form

$$f_i = \sum_{k=0}^{n-1} \alpha_{ik} U_T^k f$$

therefore

$$U_T^n f_i = c_i^n f_i.$$



On the other hand

$$U_T^n f_i = c_i^n f_i$$

therefore

$$c_i^n = c, \quad \text{for every } i.$$

Then  $c_i \neq 1$  for each  $i$  and

$$\int f_i d\mu = \int U_T f_i d\mu = c_i \int f_i d\mu$$

therefore  $\int f_i d\mu = 0$ . From

$$f = \sum_{i=1}^n \alpha_i f_i$$

we deduce that  $\int f d\mu = 0$ .

By induction we deduce then that for every  $n$ ,  $(w_T^{-n}(\Gamma), U_T)$  is a discrete system, therefore  $(\bigcup_{n=0}^{\infty} w_T^{-n}(\Gamma), U_T)$  is also a discrete system.

**COROLLARY 1.** *If  $T^n$  is ergodic for every  $n$ , then  $(\Gamma_{\infty}(T), U_T)$  is a discrete system.*

In fact  $(C, U_T)$  is a discrete system, and  $\Gamma_{\infty}(T) = \bigcup_{n=1}^{\infty} w_T^{-n}(C)$ .

**COROLLARY 2.** *Suppose that  $T$  has a discrete model  $(\Gamma, U_T)$ . If  $T^n$  is ergodic for every  $n$ , then*

$$w_T^{-1}\Gamma = \Gamma \quad \text{and} \quad \Gamma_{\infty}(T) \subset \Gamma.$$

In fact, in this case  $(w_T^{-1}\Gamma, U_T)$  is again a discrete system and  $\Gamma \subset w^{-1}\Gamma$ , therefore, by Proposition 3,  $w^{-1}\Gamma = \Gamma$ . Then  $\Gamma_{\infty}(T) \subset \Gamma$ .

**THEOREM 9.** *Suppose that  $T$  has a discrete model  $(\Gamma, U_T)$ .*

*If for every natural number  $n$ , conditions  $\gamma \in \Gamma$  and  $U^n \gamma = \gamma$  imply  $\gamma \in C$ , then  $T^n$  is ergodic for every  $n$ .*

Consider  $\Gamma$  as a direct product  $\Gamma = C \cdot \Gamma'$ , where  $\Gamma'$  is a group and an orthonormal basis of  $L^2(\mu)$ .

Consider the homomorphisms  $\rho_n: \Gamma' \rightarrow C$  and  $V: \Gamma' \rightarrow \Gamma'$  such that  $U_T^n \gamma = \rho_n(\gamma) V^n \gamma$ , for  $\gamma \in \Gamma'$  and  $n \in N$ .

We shall prove first that  $T$  is ergodic. Let  $f \in L^2(\mu)$  be such that  $U_T f = f$  and prove that  $f$  is constant. We have

$$f = \sum_{\gamma \in \Gamma'} \alpha(\gamma) \gamma$$

where  $\alpha(\gamma) = (f | \gamma)$ . For every  $n$  we have  $U_T^n f = f$  and

$$U_T^n f = \sum_{\gamma \in \Gamma'} \alpha(\gamma) \rho_n(\gamma) V^n \gamma$$

therefore

$$\alpha(V^n \gamma) = \alpha(\gamma) \rho_n(\gamma), \quad \text{for } \gamma \in \Gamma' \quad \text{and} \quad n \in N$$

whence

$$|\alpha(V^n \gamma)| = |\alpha(\gamma)|, \quad \text{for } \gamma \in \Gamma' \quad \text{and} \quad n \in N.$$

We shall prove that  $\alpha(\gamma) = 0$  if  $\gamma \neq 1$ . We remark that the hypothesis implies that if  $\gamma \neq 1$ , then  $\rho_n(\gamma) \neq 1$  for every  $n$ .

If  $\gamma \neq 1$  and  $V^n \gamma = \gamma$  for some  $n$ , then the equality  $\alpha(V^n \gamma) = \alpha(\gamma) \rho_n(\gamma)$  becomes

$$\alpha(\gamma) = \alpha(\gamma) \rho_n(\gamma)$$

therefore  $\alpha(\gamma) = 0$ .

If  $\gamma \neq 1$  and  $V^n \gamma \neq \gamma$  for every  $n$ , then the functions  $\gamma, V\gamma, V^2\gamma, \dots$  are different from each other, therefore

$$\sum_{n=1}^{\infty} |\alpha(V^n \gamma)|^2 \leq \sum_{\gamma' \in \Gamma'} |\alpha(\gamma')|^2 = \|f\|_2^2 < \infty$$

consequently  $|\alpha(V^n \gamma)|^2 \rightarrow 0$  as  $n \rightarrow \infty$ , whence  $\alpha(\gamma) = 0$ .

We deduce that  $f = \alpha(1)1$ , that is  $f$  is constant, therefore  $T$  is ergodic.

We remark now that for every  $n$ ,  $(\Gamma, U_T^n)$  is a discrete model for  $T^n$ , satisfying the conditions of the theorem with respect to  $U_T^n$ , therefore  $T^n$  is ergodic.

**7. Transformations with quasi-discrete spectrum.** Let  $(X, \Sigma, \mu)$  be a probability measure space and  $T: X \rightarrow X$  a measure preserving transformation.

**DEFINITION 4.** We say that  $T$  has quasi-discrete spectrum if  $T$  has a discrete model  $(\Gamma, U_T)$  with  $\Gamma \subset \Gamma_{\infty}(T)$ .

To say that  $T$  has quasi-discrete spectrum means that there exists a group  $\Gamma' \subset \Gamma_{\infty}(T)$  which is an orthonormal basis of  $L^2(\mu)$ , such that  $U_T \Gamma' \subset C \cdot \Gamma'$ .

Here are some properties of transformations with quasi-discrete spectrum:

(1) If  $T$  has quasi-discrete spectrum, then  $(\Gamma_{\infty}(T), U_T, \varphi_{\mu})$  is an (not necessarily discrete) algebraic model of  $T$ .

(2) If  $T$  has quasi-discrete spectrum, then  $T^n$  has quasi-discrete spectrum, for every  $n$  (since  $\Gamma_{\infty}(T) \subset \Gamma_{\infty}(T^n)$ ).

(3) If  $T$  is conjugate with a transformation with quasi-discrete spectrum, then  $T$  has itself quasi-discrete spectrum (see Proposition 1).

(4) Two transformations  $T$  and  $T'$  with quasi-discrete spectrum are conjugate if and only if the a.e. systems  $(\Gamma_{\infty}(T), U_T, \varphi_{\mu})$  and  $(\Gamma_{\infty}(T'), U_{T'}, \varphi_{\mu'})$  are isomorphic.

We use Proposition 1 and Theorem 1.

For transformations  $T$  for which  $(\Gamma_{\infty}(T), U_T)$  is itself a discrete model we have, in addition, the following properties:

(5) Let  $T$  and  $T'$  be two measure preserving transformations having  $(\Gamma_{\infty}(T), U_T)$  respectively  $(\Gamma_{\infty}(T'), U_{T'})$  as discrete models.

Then  $T$  and  $T'$  are conjugate if and only if there exists an isomorphism  $J$  of  $\Gamma_{\infty}(T)$  onto  $\Gamma_{\infty}(T')$  such that

$$Jc = c, \quad \text{for } c \in C$$

and

$$JU_T = U_{T'}J.$$

(6) If  $T$  has quasi-discrete spectrum and if there exists a discrete system  $(\Gamma^*, U_T)$  with  $\Gamma_\infty(T) \subset \Gamma^*$ , then  $(\Gamma_\infty(T), U_T)$  is a discrete model of  $T$  and there is no other discrete model of  $T$  containing or contained in  $(\Gamma_\infty(T), U_T)$ .

We use Proposition 3.

For transformations with all iterates ergodic we have some more properties:

(7) If  $T^n$  is ergodic for every  $n$  and if  $\Gamma_\infty(T)$  generates  $L^2(\mu)$ , then  $(\Gamma_\infty(T), U_T)$  is a discrete model of  $T$  and there is no other discrete model of  $T$ .

In fact, by Corollary 1 of Proposition 6,  $(\Gamma_\infty(T), U_T)$  is a discrete system, therefore  $(\Gamma_\infty(T), U_T)$  is a discrete model of  $T$ . By Corollary 2 of Proposition 6, for any other discrete model  $(\Gamma, U_T)$  of  $T$  we have  $\Gamma_\infty(T) \subset \Gamma$ , therefore  $\Gamma_\infty(T) = \Gamma$ .

(8) Let  $T$  and  $T'$  be two transformations with quasi-discrete spectrum and all iterates  $T^n$  and  $T'^n$  ergodic.

Then  $T$  and  $T'$  are conjugate if and only if  $w_T \Gamma_\infty(T)$  and  $w_{T'} \Gamma_\infty(T')$  are isomorphic by an isomorphism  $\phi$  such that  $\phi w_T = w_{T'} \phi$  and  $\phi c = c$  for  $c \in C \cap w_T \Gamma_\infty(T)$ .

We use Theorem 8 and property (5) above.

The following theorem gives a characterization of discrete systems which are models for transformations with quasi-discrete spectrum.

**THEOREM 10.** *If  $(\Gamma, U)$  is a discrete system such that*

$$\Gamma = \bigcup_{n=0}^{\infty} w^{-n}(C), \quad \text{where } w(\gamma) = U\gamma \cdot \gamma^{-1}, \quad \text{for } \gamma \in \Gamma,$$

*then the corresponding transformation  $T$  has quasi-discrete spectrum.*

*If, in addition, for every natural number  $n \in N$ ,  $\gamma \in \Gamma$  and  $U^n \gamma = \gamma$  imply  $\gamma \in C$ , then  $T^n$  is ergodic for every  $n$ .*

In fact  $w$  is the restriction of  $w_T$  to  $\Gamma$ , therefore  $\Gamma \subset \Gamma_\infty(T)$ , consequently  $T$  has quasi-discrete spectrum.

For the second part of the theorem we use Theorem 9 to deduce that all the iterates  $T^n$  are ergodic. In this case we have  $\Gamma = \Gamma_\infty(T)$ .

**REMARK.** Theorem 10 and property (8) were proved by Abramov [1].

*Example of transformation with discrete model but without quasi-discrete spectrum.* Let  $X_n = \{-1, 1\}$  and  $\mu_n(\{-1\}) = \mu_n(\{1\}) = \frac{1}{2}$  for  $n = 0, \pm 1, \pm 2, \dots$ . Consider the product  $X = \prod_{n=-\infty}^{\infty} X_n$ , equipped with the product measure  $\mu$  and the bilateral shift  $T(x_n) = (y_n)$ , where  $y_n = x_{n+1}$  for every  $n$ . Then  $T^n$  is ergodic for every  $n$  and the only proper value of  $T$  is 1, so that  $w_T^{-1}(C) = C$ . It follows that  $\Gamma_\infty(T) = C$  so that  $T$  has not quasi-discrete spectrum.

On the other hand, consider the function  $f_0: X \rightarrow R$  defined by

$$\begin{aligned} f_0((x_n)) &= -1 & \text{if } x_0 &= -1, \\ &= 1 & \text{if } x_0 &= 1, \end{aligned}$$

and the group  $\Gamma$  generated by  $U_T^n f_0$ ,  $n = 0, \pm 1, \pm 2, \dots$  and by the constants. Then  $(\Gamma, U_T)$  is a discrete model of  $T$ .

**8. Transformations with discrete spectrum.** Let  $(X, \Sigma, \mu)$  be a probability measure space and  $T: X \rightarrow X$  a measure preserving transformation.

**DEFINITION 5.** *We say that  $T$  has discrete spectrum if  $T$  has a discrete model  $(\Gamma, U_T)$  with  $\Gamma \subset \Gamma_1(T)$ .*

To say that  $T$  has a discrete spectrum means that there exists a group  $\Gamma' \subset \Gamma_1(T)$  of proper functions of  $U_T$  which is an orthonormal basis of  $L^2(\mu)$ .

Here are some properties of transformations with discrete spectrum:

(1) *Every transformation with discrete spectrum is invertible (since  $U_T \Gamma_1 = \Gamma_1$ ).*  
 (2) *A transformation has discrete spectrum if and only if it is conjugate to a rotation on a compact abelian group equipped with Haar measure (see Corollary 2 of Theorem 6).*

(3) *Every transformation with discrete spectrum has quasi-discrete spectrum.*

(4) *If  $T$  has discrete spectrum, then  $(\Gamma_1(T), U_T, \varphi_\mu)$  is an (not necessarily discrete) algebraic model of  $T$ .*

(5) *If  $T$  has discrete spectrum, then  $T^n$  has discrete spectrum for every  $n$ .*

(6) *If  $T$  is conjugate with a transformation with discrete spectrum, then  $T$  has itself discrete spectrum.*

(7) *Two transformations  $T$  and  $T'$  with discrete spectrum are conjugate if and only if the a.e. systems  $(\Gamma_1(T), U_T, \varphi_\mu)$  and  $(\Gamma_1(T'), U_{T'}, \varphi_{\mu'})$  are isomorphic.*

Transformations  $T$  for which  $(\Gamma_1(T), U_T)$  is itself a discrete model, have additional properties:

(8) *Let  $T$  and  $T'$  be two measure preserving transformations having  $(\Gamma_1(T), U_T)$  respectively  $(\Gamma_1(T'), U_{T'})$  as discrete models. Then  $T$  and  $T'$  are conjugate if and only if there exists an isomorphism  $J$  of  $\Gamma_1(T)$  onto  $\Gamma_1(T')$  such that:*

$$Jc = c, \quad \text{for } c \in C$$

and

$$JU_T = U_{T'}J.$$

(9) *If  $T$  has discrete spectrum and if there exists a discrete system  $(\Gamma^*, U_T)$  with  $\Gamma_1(T) \subset \Gamma^*$  then  $(\Gamma_1(T), U_T)$  is a discrete model of  $T$ , and there is no other discrete model of  $T$  containing or contained in  $(\Gamma_1(T), U_T)$ .*

For ergodic transformations we have some more properties:

(10) *If  $T$  is ergodic and if  $\Gamma_1(T)$  generates  $L^2(\mu)$ , then  $(\Gamma_1(T), U_T)$  is a discrete model of  $T$ , and there is no other discrete model containing or contained in  $(\Gamma_1(T), U_T)$ .*

(11) *Let  $T$  and  $T'$  be two ergodic transformations with discrete spectrum.*

*Then  $T$  and  $T'$  are conjugate if and only if  $U_T$  and  $U_{T'}$  have the same spectrum [4, p. 46].*

We use Theorem 8 remarking that  $w_T \Gamma_1(T)$  is the spectrum of  $U_T$  and  $w_{T'} \Gamma_1(T')$  is the spectrum of  $T'$ .

The characterization of discrete systems which are models for transformations with discrete spectrum, is given by the following:

THEOREM 11. *If  $(\Gamma, U)$  is a discrete system such that*

$$w^{-1}(C) = \Gamma, \text{ where } w(\gamma) = U\gamma \cdot \gamma^{-1}, \text{ for } \gamma \in \Gamma$$

*then the corresponding transformation  $T$  has discrete spectrum.*

*If, in addition, there exists  $n \in N$  such that  $\gamma \in \Gamma$  and  $U^n \gamma = \gamma$  imply  $\gamma \in C$ , then  $T^n$  is ergodic.*

In fact,  $w$  is the restriction of  $w_T$  to  $\Gamma$ , therefore  $\Gamma \subset \Gamma_1(T)$ , consequently  $T$  has discrete spectrum. The second part follows from Theorem 9. In this case we have  $\Gamma = \Gamma_1(T)$ .

When does  $\Gamma_1$  coincide with  $\Gamma_\infty$ ?

THEOREM 12. *Let  $T$  be an ergodic transformation with discrete spectrum, on a probability measure space  $(X, \Sigma, \mu)$ . We have  $\Gamma_1(T) = \Gamma_\infty(T)$  if and only if the point spectrum of  $U_T$  contains no root of 1 (except 1 itself).*

Suppose first that  $\Gamma_1 = \Gamma_\infty$ . Let  $\xi$  be a proper value of a function  $f \in \Gamma_1$ :

$$U_T f = \xi f.$$

We shall prove that if  $\xi^n = 1$  for some  $n$ , then  $\xi = 1$ . In fact, suppose that  $N$  is the least natural number with  $\xi^N = 1$ . We have then

$$U_T f^N = f^N$$

therefore, (since  $T$  is ergodic)  $f^N$  is constant, and we may suppose that  $f^N \equiv 1$ , multiplying  $f$  by a suitable number, if necessary. Then  $f$  takes on the values  $1, \xi, \dots, \xi^{N-1}$  on the corresponding sets  $A_0, A_1, \dots, A_{N-1}$

$$f = \sum_{k=0}^{N-1} \xi^k \varphi_{A_k}.$$

Since  $f(Tx) = \xi f(x)$ , we have

$$\sum_{k=0}^{N-1} \xi^k \varphi_{T^{-1}A_k} = \sum_{k=0}^{N-1} \xi^{k+1} \varphi_{A_k}$$

therefore  $TA_k = A_{k+1}$  for  $k=0, 1, \dots, N-1$ , where  $A_N = A_0$ . It follows that  $\mu(A_k) = \mu(A_0) > 0$  for every  $k$ .

If  $N$  is odd we take  $\nu$  such that  $\nu^N = 1$ ; if  $N$  is even, we take  $\nu$  such that  $\nu^N = -1$ .

Define now the function

$$g = \sum_{k=0}^{N-1} \nu^k \xi^{k(k-1)/2} \varphi_{A_k}.$$

Then we have

$$Ug = \nu g.$$

In fact, if  $x \in A_k$  and  $k=0, 1, \dots, N-2$ , we have  $Tx \in A_{k+1}$ , therefore

$$Ug(x) = g(Tx) = \nu^{k+1} \xi^{k(k+1)/2} = \nu \xi^k \nu^k \xi^{k(k-1)/2} = \nu f(x)g(x)$$

and for  $x \in A_{N-1}$  we have  $Tx \in A_N = A_0$ , therefore

$$Ug(x) = g(Tx) = 1 = \nu^N \xi^{N(N-1)/2} = \nu \xi^{N-1} \nu^{N-1} \xi^{(N-1)(N-2)/2} = \nu f(x)g(x)$$

since  $\xi^{N(N-1)/2}$  is equal to 1 if  $N$  is odd and to  $-1$  if  $N$  is even.

It follows that  $g \in \Gamma_2 = \Gamma_1$ , therefore  $\nu f = \alpha \in C$ , consequently  $f \in C$ . We deduce then that  $Uf = f$ , therefore  $\xi = 1$ .

Conversely, suppose that  $\Gamma_1 \neq \Gamma_2$  and prove that  $U_T$  has at least a proper value  $\alpha \neq 1$  such that  $\alpha^N = 1$  for some  $N$ .

Let  $g_0 \in \Gamma_2 - \Gamma_1$ . Since  $\Gamma_1$  is an orthonormal basis of  $L^2(\mu)$  and  $g_0 \neq 0$ , there exists  $h \in \Gamma_1$  such that

$$(g_0, h) = \int g_0 h \, d\mu \neq 0.$$

If we put  $g = g_0 h$ , we have  $g \in \Gamma_2 - \Gamma_1$  and  $\int g \, d\mu \neq 0$ . There exists a function  $f \in \Gamma_1$  such that

$$U_T g = fg.$$

There exists also a number  $\lambda \in C$  such that

$$U_T f = \lambda f.$$

By induction, we deduce that for every  $n$  we have

$$U_T^n g = \lambda^{n(n-1)/2} f^n g.$$

We have  $f^n \in C$  for some  $n$ . In fact, if we had  $f^n \notin C$  for every  $n$ , then (since  $f^n \in \Gamma_1$ ),

$$\int U_T^n g \bar{g} \, d\mu = \lambda^{n(n-1)/2} \int f^n \, d\mu = 0$$

therefore, the sequence  $g, Ug, U^2g, \dots$  would be orthonormal, consequently

$$\int g \, d\mu = \int U_T^n g \, d\mu = (U_T^n g, 1) \rightarrow 0$$

and we would get a contradiction.

Let  $N$  be the least integer  $\geq 0$  such that

$$f^N = \mu \in C.$$

Then

$$U^N g = \lambda^{N(N-1)/2} \mu g = \xi g.$$

We have

$$\xi \int g \, d\mu = \int U_T^N g \, d\mu = \int g \, d\mu \neq 0$$

therefore  $\xi = 1$ , consequently

$$U^N g = g.$$

Since  $g$  is not constant and since  $\Gamma_1$  generates  $L^2(\mu)$ , there exists a proper function  $k \neq 1$  of  $\Gamma_1$  such that  $(g, k) \neq 0$ . If  $\alpha$  is the corresponding proper value:

$$U_T k = \alpha k$$

we have  $\alpha \neq 1$  and

$$(g, k) = (U^N g, k) = (g, U^{-N} k) = (g, \alpha^{-N} k) = \alpha^N (g, k)$$

therefore  $\alpha^N = 1$ .

## BIBLIOGRAPHY

1. L. M. Abramov, *Metric automorphisms with quasi-discrete spectrum*, Izv. Akad. Nauk SSSR Ser. Mat. **26** (1962), 513–530. (Russian)
2. N. Dinculeanu and C. Foiaş, *A universal model for ergodic transformations on separable measure spaces*, Michigan Math. J. **13** (1966), 109–117.
3. ———, *Algebraic models for measures*, Illinois J. Math. **12** (1968), 340–351.
4. C. Foiaş, *Automorphisms of compact abelian groups, as models for measure-preserving transformations*, Michigan Math. J. **13** (1966), 349–352.
5. P. R. Halmos, *Lectures on ergodic theory*, Publ. Math. Soc. Japan, No. 3, Tokyo, 1956.
6. A. Ionescu Tulcea and C. Ionescu Tulcea, *On the lifting property. I*, J. Math. Anal. Appl. **3** (1961), 537–546.
7. K. Jacobs, *Lecture notes on ergodic theory*, Aarhus Universitet, Aarhus, 1962/1963.

UNIVERSITY OF BUCAREST,  
BUCAREST, ROMANIA