

# A COMPATIBILITY CONDITION BETWEEN INVARIANT RIEMANNIAN METRICS AND LEVI-WHITEHEAD DECOMPOSITIONS ON A COSET SPACE

BY  
JOSEPH A. WOLF<sup>(1)</sup>

**0. Introduction.** Let  $M=G/K$  be an effective coset space of a connected Lie group by a compact subgroup. Then there may be many  $G$ -invariant riemannian metrics on  $M$ . But one expects the algebraic structure of the pair  $(G, K)$  to have a strong influence on the curvatures of  $M$  relative to any  $G$ -invariant riemannian metric. For example

(1) if  $G$  is semisimple with finite center and  $K$  is a maximal compact subgroup, then it is classical from symmetric space theory that all  $G$ -invariant riemannian metrics on  $M$  have every sectional curvature  $\leq 0$ ;

(2) if  $G$  is commutative then every  $G$ -invariant riemannian metric on  $M$  is flat; and

(3) if  $G$  is noncommutative and nilpotent then [7] every  $G$ -invariant riemannian metric on  $M$  has sectional curvatures of both signs.

Those results are proved by choosing an  $\text{ad}_G(K)$ -stable complement  $\mathfrak{M}$  to the Lie algebra  $\mathfrak{K}$  of  $K$  inside the Lie algebra  $\mathfrak{G}$  of  $G$ , and by performing calculations in  $\mathfrak{M}$  and in  $\mathfrak{G}$  in a manner justified by embedding  $G$  in the orthonormal frame bundle of  $M$ . But at certain crucial parts of those calculations one must have  $\mathfrak{G}$  either semisimple or nilpotent. The idea in this paper is to create a setup in which the calculations can still be carried out, by requiring that the complement  $\mathfrak{M}$  split as

$$\mathfrak{M} = (\mathfrak{M} \cap \mathfrak{K}) + (\mathfrak{M} \cap \mathfrak{L}) \text{ orthogonal direct sum,}$$

where

$$\mathfrak{G} = \mathfrak{K} + \mathfrak{L} \text{ is a Levi-Whitehead decomposition}$$

and

$$\mathfrak{M} \cap \mathfrak{K} \text{ contains the nilpotent radical of } \mathfrak{G}.$$

§2 is a study of the circumstances under which  $\mathfrak{M}$  can be chosen, and the Levi-Whitehead decomposition  $\mathfrak{G} = \mathfrak{K} + \mathfrak{L}$  can be chosen, so that  $M$  has such an orthogonal splitting. We describe those circumstances by the condition (2.2) that the invariant riemannian metric on  $M$  be "consistent" with  $\mathfrak{G} = \mathfrak{K} + \mathfrak{L}$ .

Given the consistency condition (2.2), our main result (Theorem 3.9) says that every unit vector  $X \in \mathfrak{M} \cap \mathfrak{K}$ , orthogonal to the nilpotent radical  $\mathfrak{N}$  of  $\mathfrak{G}$ , is a

---

Received by the editors May 21, 1968.

<sup>(1)</sup> Research supported by an Alfred P. Sloan Research Fellowship and by National Science Foundation Grant GP-8008.

direction of negative mean curvature on  $M$ . Our applications (§4) essentially consist of observing that, if  $M$  has mean curvature  $\geq 0$  everywhere, then the consistency condition implies  $\mathfrak{N} = (\mathfrak{M} \cap \mathfrak{N})$ , i.e.  $\mathfrak{N} = \mathfrak{N} + (\mathfrak{N} \cap \mathfrak{R})$  semidirect sum. The most striking of the applications is Theorem 4.4, which says:

*Let  $M$  be a connected Riemannian manifold that has a solvable transitive group of isometries. Then the following conditions are equivalent.*

- (i)  $M$  has mean curvature  $\geq 0$  everywhere.
- (ii)  $M$  has every sectional curvature  $\geq 0$ .
- (iii)  $M$  has every sectional curvature zero.
- (iv)  $M$  is isometric to the product of an euclidean space and a flat riemannian torus.

Theorem 4.7 adds negative curvature conditions in case  $M$  has a transitive nilpotent group of isometries, extending the results of [7] to mean curvature.

**1. Definitions and notation.**  $\mathfrak{G}$  is a real Lie algebra. We have the *nilpotent radical*  $\mathfrak{N}$  and the *solvable radical*  $\mathfrak{R}$ , characteristic nilpotent and solvable ideals in  $\mathfrak{G}$  defined by

$\mathfrak{N}$  is the union of the nilpotent ideals of  $\mathfrak{G}$ ,

$\mathfrak{R}$  is the union of the solvable ideals of  $\mathfrak{G}$ .

The basic facts on  $\mathfrak{N}$  and  $\mathfrak{R}$  are the following.

(1.1) *If  $C$  is a fully reducible group of automorphisms of  $\mathfrak{G}$ , then there are  $C$ -invariant semisimple subalgebras  $\mathfrak{L} \subset \mathfrak{G}$  that map isomorphically onto  $\mathfrak{G}/\mathfrak{R}$  under the projection  $\varphi: \mathfrak{G} \rightarrow \mathfrak{G}/\mathfrak{R}$ , and any two such subalgebras are conjugate by an automorphism  $\text{ad}_{\mathfrak{G}}(\exp n)$  of  $\mathfrak{G}$  where  $n \in \mathfrak{N}$  is left fixed by every  $c \in C$ .*

The existence is due to G. D. Mostow [3, Corollary 5.2], and the conjugacy statement is the result [5, Theorem 4] of E. J. Taft. In general a semisimple subalgebra  $\mathfrak{L} \subset \mathfrak{G}$  such that  $\varphi: \mathfrak{L} \cong \mathfrak{G}/\mathfrak{R}$  is called a *Levi factor* of  $\mathfrak{G}$ , and the Levi factors of  $\mathfrak{G}$  are just the maximal semisimple subalgebras.

(1.2) *If  $\mathfrak{L}$  is a Levi factor of  $\mathfrak{G}$ , then  $\mathfrak{G} = \mathfrak{R} + \mathfrak{L}$  semidirect sum,  $\mathfrak{R} + \mathfrak{L}$  (semidirect) is an ideal in  $\mathfrak{G}$ , and the derived algebra  $[\mathfrak{G}, \mathfrak{G}] \subset \mathfrak{R} + \mathfrak{L}$ .*

The first assertion is immediate and the second follows from  $\mathfrak{R} \subset \mathfrak{R}$ . For the third, one notes that  $[\mathfrak{R}, \mathfrak{R}] \subset \mathfrak{R}$  by Ado's Theorem and that  $\text{ad}_{\mathfrak{G}}(\mathfrak{L})$  normalizes  $\text{ad}_{\mathfrak{R}}(\mathfrak{R})$  in the derivation algebra of  $\mathfrak{R}$ .

Let  $M = G/K$  be a coset space of a Lie group by a closed subgroup.  $\mathfrak{R} \subset \mathfrak{G}$  are the Lie algebras of  $K \subset G$ . An  $\text{ad}_G(K)$ -invariant subspace  $\mathfrak{M} \subset \mathfrak{G}$  such that  $\mathfrak{G} = \mathfrak{M} + \mathfrak{R}$  (vector space direct sum), is called an *invariant complement* for  $K$ . If an invariant complement for  $K$  exists, then  $M = G/K$  is called a *reductive coset space*.

$K$  is called a *reductive subgroup* of  $G$  in case the group  $\text{ad}_G(K)$  of linear transformations of  $\mathfrak{G}$  is fully reducible. If  $K$  is a reductive subgroup of  $G$ , then  $\text{ad}_G(K)\mathfrak{R}$

$=\mathfrak{K}$  implies that  $M=G/K$  is a reductive coset space. The converse fails in the example

$$G = SL(2, \mathbf{R}) \quad \text{and} \quad K = \left\{ \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} : n \in \mathbf{Z} \right\}.$$

However, compact subgroups, and semisimple subgroups with only finitely many components, are reductive subgroups.

**2. The compatibility condition.** We can now define the compatibility conditions with which we will operate.  $M=G/K$  is a coset space of a Lie group by a closed subgroup.  $\mathfrak{L}$  is a Levi factor of  $\mathfrak{G}$  and  $\mathfrak{M}$  is an invariant complement for  $K$ . If

$$(2.1) \quad \mathfrak{M} = (\mathfrak{M} \cap \mathfrak{K}) + (\mathfrak{M} \cap \mathfrak{L}) \quad \text{and} \quad \mathfrak{K} = (\mathfrak{M} \cap \mathfrak{K}) + (\mathfrak{K} \cap \mathfrak{K})$$

then we say that  $\mathfrak{L}$  splits  $\mathfrak{M}$  and that  $\mathfrak{M}$  is split by the Levi-Whitehead decomposition  $\mathfrak{G}=\mathfrak{K}+\mathfrak{L}$ . Suppose further that we have a  $G$ -invariant pseudo-riemannian metric  $ds^2$  on  $M$ . Represent  $ds^2$  by an  $\text{ad}_G(K)$ -invariant inner product  $\langle \cdot, \cdot \rangle$  on  $\mathfrak{M}$ . If

$$(2.2) \quad \begin{aligned} \mathfrak{M} &= (\mathfrak{M} \cap \mathfrak{K}) + (\mathfrak{M} \cap \mathfrak{L}), \\ \mathfrak{K} &= (\mathfrak{M} \cap \mathfrak{K}) + (\mathfrak{K} \cap \mathfrak{K}) \quad \text{and} \quad \langle \mathfrak{M} \cap \mathfrak{K}, \mathfrak{M} \cap \mathfrak{L} \rangle = 0, \end{aligned}$$

then we say that  $\mathfrak{L}$  splits  $\mathfrak{M}$  orthogonally and that  $ds^2$  is consistent with the Levi-Whitehead decomposition  $\mathfrak{G}=\mathfrak{K}+\mathfrak{L}$ .

**2.3. PROPOSITION.** *Let  $K$  be a closed reductive subgroup of a Lie group  $G$ . Then for every  $\text{ad}_G(K)$ -invariant Levi factor  $\mathfrak{L}$  of  $\mathfrak{G}$ , there exists an invariant complement  $\mathfrak{M}$  for  $K$ , such that  $\mathfrak{L}$  splits  $\mathfrak{M}$ . If  $ds^2$  is a  $G$ -invariant pseudo-riemannian metric on  $G/K$ ,  $\varphi: \mathfrak{G} \rightarrow \mathfrak{G}/\mathfrak{K}$  is the projection, and the representations of  $K$  on  $\mathfrak{K}/(\mathfrak{K} \cap \mathfrak{K})$  and  $\mathfrak{L}/(\mathfrak{L} \cap \varphi^{-1} \varphi \mathfrak{K})$  are disjoint, then it is automatic that  $\mathfrak{L}$  splits  $\mathfrak{M}$  orthogonally and  $ds^2$  is consistent with  $\mathfrak{G}=\mathfrak{K}+\mathfrak{L}$ .*

**Proof.** Mostow's result (1.1) provides an  $\text{ad}_G(K)$ -invariant Levi factor  $\mathfrak{L}$  of  $\mathfrak{G}$ . As  $K$  is reductive in  $G$  we have  $\text{ad}_G(K)$ -invariant direct sum decompositions

$$\mathfrak{K} = \mathfrak{M}_1 + (\mathfrak{K} \cap \mathfrak{K}) \quad \text{and} \quad \mathfrak{G}/\mathfrak{K} = \mathfrak{M}'_2 + \varphi(\mathfrak{K}).$$

Now define

$$\mathfrak{M}_2 = \mathfrak{L} \cap \varphi^{-1}(\mathfrak{M}'_2) \quad \text{and} \quad \mathfrak{M} = \mathfrak{M}_1 + \mathfrak{M}_2.$$

Then  $\text{ad}_G(K)\mathfrak{M}_i = \mathfrak{M}_i$  so  $\mathfrak{M}$  is an  $\text{ad}_G(K)$ -invariant subspace of  $\mathfrak{G}$  that satisfies (2.1). If  $x \in \mathfrak{M} \cap \mathfrak{K}$  then  $\varphi(x) \in \mathfrak{M}'_2 \cap \varphi(\mathfrak{K}) = 0$  so  $x \in \mathfrak{K}$ ; then  $x \in \mathfrak{M}_1 \cap (\mathfrak{K} \cap \mathfrak{K}) = 0$  so  $x = 0$ ; thus  $\mathfrak{M} \cap \mathfrak{K} = 0$ . On the other hand

$$\dim \mathfrak{K} = \dim \mathfrak{M}_1 + \dim(\mathfrak{K} \cap \mathfrak{K})$$

and

$$\begin{aligned} \dim \mathfrak{G}/\mathfrak{K} &= \dim \mathfrak{M}'_2 + \dim \varphi(\mathfrak{K}) = \dim \mathfrak{M}_2 + \dim \varphi(\mathfrak{K}) \\ &= \dim \mathfrak{M}_2 - \dim(\mathfrak{K} \cap \mathfrak{K}) + \dim \mathfrak{K} \end{aligned}$$

so

$$\dim \mathfrak{G} = \dim \mathfrak{K} + \dim \mathfrak{G}/\mathfrak{K} = \dim \mathfrak{M} + \dim \mathfrak{K}.$$

Thus  $\mathfrak{M}$  is a vector space complement to  $\mathfrak{R}$  in  $\mathfrak{G}$ . Now  $\mathfrak{M}$  is an invariant complement for  $K$  such that  $\mathfrak{L}$  splits  $\mathfrak{M}$ .

Let  $ds^2$  and the inner product  $\langle \cdot, \cdot \rangle$  on  $\mathfrak{M}$  be given. The representation of  $K$  on  $\mathfrak{R}/(\mathfrak{R} \cap \mathfrak{R})$  is the representation  $\text{ad}_G|_K$  on  $\mathfrak{M}_1$ ; the representation of  $K$  on  $\mathfrak{L}/(\mathfrak{L} \cap \varphi^{-1}\varphi\mathfrak{R})$  is  $\text{ad}_G|_K$  on  $\mathfrak{M}_2$ . If those two are disjoint then necessarily  $\langle \mathfrak{M}_1, \mathfrak{M}_2 \rangle = 0$ . Q.E.D.

We reformulate the metric portion of Proposition 2.3.

**2.4. PROPOSITION.** *Let  $M=G/K$  be a homogeneous pseudo-riemannian manifold with metric  $ds^2$ , where  $K$  is a reductive subgroup of  $G$ . Let  $\mathcal{O}$  be the base point,  $M_{\mathcal{O}}$  the tangent space at  $\mathcal{O}$ ,  $\chi$  the linear isotropy representation of  $K$  on  $M_{\mathcal{O}}$ , and  $R_{\mathcal{O}}$  the subspace of  $M_{\mathcal{O}}$  spanned by vector fields from elements of the solvable radical of  $\mathfrak{G}$ . Suppose that the representations of  $K$  induced by  $\chi$ , on  $R_{\mathcal{O}}$  and on  $M_{\mathcal{O}}/R_{\mathcal{O}}$ , are disjoint. Then  $M_{\mathcal{O}} = R_{\mathcal{O}} + R_{\mathcal{O}}^{\perp}$  and  $ds^2$  is consistent with every Levi-Whitehead decomposition  $\mathfrak{G} = \mathfrak{R} + \mathfrak{L}$  for which  $\text{ad}_G(K) \cdot \mathfrak{L} = \mathfrak{L}$ .*

For, in the notation of the proof of Proposition 2.3,  $R_{\mathcal{O}}$  is spanned by the vector fields from  $\mathfrak{M}_1$  while  $R_{\mathcal{O}}^{\perp}$  is spanned by those from  $\mathfrak{M}_2$ .

In the riemannian case we will be able to arrange that  $\mathfrak{M}_1 = \mathfrak{M} \cap \mathfrak{R}$  contain the nilpotent radical  $\mathfrak{N}$ . For that, we need a technical lemma.

**2.5. LEMMA.** *In a connected nilpotent Lie group every compact subgroup is central.*

**Proof.** Let  $N$  be the connected nilpotent Lie group,  $\pi: \tilde{N} \rightarrow N$  the universal Lie group covering, and  $\Gamma$  the kernel of  $\pi$ . Then  $\Gamma$  is a discrete central subgroup of  $\tilde{N}$ . Let  $C$  be a maximal compact subgroup of  $N$  and  $\tilde{C} = \pi^{-1}(C)$ . Then  $C$  is a torus group,  $\tilde{C}$  is a simply connected commutative subgroup of  $\tilde{N}$ , and  $\Gamma \subset \tilde{C}$  such that  $C = \tilde{C}/\Gamma$  compact. As  $\Gamma$  is central in  $\tilde{N}$ , and as  $\tilde{N}$  is nilpotent, now  $\tilde{C}$  is central in  $\tilde{N}$ , so  $C$  is central in  $N$ . If  $E$  is any compact subgroup of  $N$  we have  $n \in N$  such that  $nEn^{-1} \subset C$ , so  $E \subset n^{-1}Cn = C$ , proving  $E$  central in  $N$ . Q.E.D.

**2.6. PROPOSITION.** *Let  $M=G/K$  be an effective coset space of a connected Lie group by a compact subgroup. Then, for any  $\text{ad}_G(K)$ -invariant Levi factor  $\mathfrak{L}$  of  $\mathfrak{G}$ , there is an invariant complement  $\mathfrak{M}$  for  $K$  such that*

$$(2.7) \quad \mathfrak{M} = (\mathfrak{M} \cap \mathfrak{R}) + (\mathfrak{M} \cap \mathfrak{L}), \quad \mathfrak{R} = (\mathfrak{M} \cap \mathfrak{R}) + (\mathfrak{R} \cap \mathfrak{R}), \quad \text{and} \quad \mathfrak{N} \subset \mathfrak{M} \cap \mathfrak{R}.$$

*In particular, if  $G$  is a group of isometries for a riemannian metric on  $M$ , and if the metric is consistent with the Levi-Whitehead decomposition  $\mathfrak{G} = \mathfrak{R} + \mathfrak{L}$ , then in addition we can choose  $\mathfrak{M}$  so that  $\mathfrak{M} \cap \mathfrak{R}$  has a subspace  $\mathfrak{A}$  such that*

$$(2.8) \quad \mathfrak{M} = \mathfrak{R} + \mathfrak{A} + (\mathfrak{M} \cap \mathfrak{L}), \quad \mathfrak{M} \cap \mathfrak{R} = \mathfrak{R} + \mathfrak{A}, \quad \text{orthogonal direct sums.}$$

**Proof.** Following Proposition 2.3 we take  $\mathfrak{M} = \mathfrak{M}_1 + \mathfrak{M}_2$ , invariant complement for  $K$  split by  $\mathfrak{L}$ , where  $\mathfrak{M}_1$  is any  $\text{ad}_G(K)$ -invariant complement to  $\mathfrak{R} \cap \mathfrak{R}$  in  $\mathfrak{R}$ .

Let  $N$  be the analytic subgroup of  $G$  for  $\mathfrak{N}$ . Then  $N$  is closed in  $G$ , so  $N \cap K$  is compact. Let  $T$  be a maximal compact subgroup of  $N$ . It contains  $N \cap K$  and is central in  $N$  by Lemma 2.5. Thus  $T$  is unique, hence normal in  $G$ . As  $T$  is a torus and  $G$  is connected now  $T$  is central in  $G$ . Thus  $N \cap K$  is central in  $G$ . But  $G$  acts effectively on  $M$ , so  $K$  contains no nontrivial normal subgroup of  $G$ . That proves  $N \cap K = \{1\}$ . In particular  $\mathfrak{N} \cap \mathfrak{K} = 0$ . Thus  $\mathfrak{R} = \mathfrak{N} + \mathfrak{A} + (\mathfrak{R} \cap \mathfrak{K})$ ,  $\text{ad}_G(K)$ -invariant direct sum, where  $\mathfrak{A}$  is any invariant complement to  $\mathfrak{N} + (\mathfrak{R} \cap \mathfrak{K})$ . For (2.7) we just choose  $\mathfrak{M}_1 = \mathfrak{N} + \mathfrak{A}$ .

Suppose further that  $ds^2$  is consistent with  $\mathfrak{G} = \mathfrak{R} + \mathfrak{L}$ . Then we have another choice, say  $\mathfrak{M}^* = \mathfrak{M}_1^* + \mathfrak{M}_2^*$ , of invariant complement for  $K$ , with  $\langle \mathfrak{M}_1^*, \mathfrak{M}_2^* \rangle = 0$  and  $\mathfrak{R} = \mathfrak{M}_1^* + (\mathfrak{R} \cap \mathfrak{K})$ . If  $\psi: \mathfrak{M} \cong \mathfrak{G}/\mathfrak{K}$  and  $\psi^*: \mathfrak{M}^* \cong \mathfrak{G}/\mathfrak{K}$  are induced by the projection  $\mathfrak{G} \rightarrow \mathfrak{G}/\mathfrak{K}$ , now  $\psi^{-1}\psi^*: \mathfrak{M}^* \rightarrow \mathfrak{M}$  is a linear isometry carrying  $\mathfrak{M}_i^*$  to  $\mathfrak{M}_i$ . Thus  $\langle \mathfrak{M}_1, \mathfrak{M}_2 \rangle = 0$ , and we obtain (2.8) by choosing  $\mathfrak{A}$  to be the orthocomplement of  $\mathfrak{N}$  in  $\mathfrak{M}_1$ . Q.E.D.

We will view the space  $\mathfrak{A}$  of (2.8) as the "gap" between nilpotent and solvable radicals of  $\mathfrak{G}$ , taken modulo  $\mathfrak{K}$ .

**3. Mean curvature along the gap between the nilpotent and solvable radicals.** We compute the mean curvature of a homogeneous riemannian manifold along a direction in the solvable radical complementary to the nilpotent radical. This is done by specializing the following general calculation to the case where the riemannian metric is consistent with a Levi-Whitehead decomposition.

**3.1. LEMMA.** *Let  $(M, ds^2)$  be a connected  $n$ -dimensional riemannian homogeneous space. Let  $\mathcal{O} \in M$ . Let  $G$  be a connected transitive group of isometries of  $M$  and let  $M'_\mathcal{O}$  denote the subspace of  $M_\mathcal{O}$  consisting of tangent vectors  $Y_\mathcal{O}$  where  $Y$  is in the derived algebra  $[\mathfrak{G}, \mathfrak{G}]$ . If  $X_\mathcal{O} \in M_\mathcal{O}$  is a unit vector orthogonal to  $M'_\mathcal{O}$ , then the mean curvature*

$$(3.2) \quad \begin{aligned} (n-1)k(X_\mathcal{O}) &= \sum_i \langle \{[X, E_i]_\mathfrak{K} + \frac{1}{4}[X, E_i]_\mathfrak{M}\}, X \rangle_\mathfrak{M}, E_i \rangle \\ &\quad - \frac{1}{2} \sum_i \|[X, E_i]_\mathfrak{M}\|^2 \\ &\quad - \frac{1}{4} \sum_{i,j} \langle [X, E_i]_\mathfrak{M}, E_j \rangle \cdot \langle [X, E_j]_\mathfrak{M}, E_i \rangle, \end{aligned}$$

where  $\mathfrak{M} \subset \mathfrak{G}$  is an invariant complement to  $K$ ,  $X \in \mathfrak{M}$  represents  $X_\mathcal{O}$ ,  $\langle \cdot, \cdot \rangle$  is the inner product on  $\mathfrak{M}$  from  $ds^2$ , and  $\{E_i\}$  is any orthonormal basis of  $\mathfrak{M}$  containing  $X$ .

**Proof.** We follow the method of Nomizu [4], using the notation

$\alpha: \mathfrak{M} \times \mathfrak{M} \rightarrow \mathfrak{M}$  for the connection function,

$\mathcal{U}: \mathfrak{M} \times \mathfrak{M} \rightarrow \mathfrak{M}$  for the symmetric part of  $\alpha$ ,

$\mathcal{R}: \mathfrak{M} \times \mathfrak{M} \times \mathfrak{M} \rightarrow \mathfrak{M}$  for the curvature tensor.

Now  $(n-1)k(X) = \sum_{i \neq i_0} K_i = -\sum_{i \neq i_0} \langle \mathcal{R}(X, E_i)X, E_i \rangle = -\sum_i \langle \mathcal{R}(X, E_i)X, E_i \rangle$  where

$\{E_i\}$  is an orthonormal basis of  $\mathfrak{M}$ , where  $X = E_{i_0}$ , and where  $K_i$  is the sectional curvature of the tangent 2-plane spanned by  $X$  and  $E_i$ . Thus

$$(3.3) \quad (n-1)k(X) = -\sum_i \langle \mathcal{R}(X, E_i)X, E_i \rangle.$$

Using [4, formulae 9.6 and 13.1] and correcting a misprint in the latter,

$$\mathcal{R}(X, E_i)X = \alpha(X, \alpha(E_i, X)) - \alpha(E_i, \alpha(X, X)) - \alpha([X, E_i]_{\mathfrak{M}}, X) - [[X, E_i]_{\mathfrak{K}}, X].$$

$$\alpha(S, T) = \frac{1}{2}[S, T]_{\mathfrak{M}} + \mathcal{U}(S, T).$$

$$\mathcal{U}(S, T) = -\frac{1}{2} \sum_j \{ \langle [S, E_j]_{\mathfrak{M}}, T \rangle + \langle [T, E_j]_{\mathfrak{M}}, S \rangle \} E_j.$$

Our hypothesis on  $X$  and  $X_\emptyset$  is that  $\langle X, [A, B]_{\mathfrak{M}} \rangle = 0$  for all  $A, B \in \mathfrak{M}$ . In particular

$$(3.4) \quad \mathcal{U}(X, S) = \mathcal{U}(S, X) = -\frac{1}{2} \sum_j \langle [X, E_j]_{\mathfrak{M}}, S \rangle E_j.$$

Thus  $\alpha(X, X) = 0$ . Substituting that into (3.3) we have

$$(3.5) \quad \begin{aligned} (n-1)k(X) &= -\sum_i \langle \alpha(X, \alpha(E_i, X)), E_i \rangle \\ &\quad + \sum_i \langle \alpha([X, E_i]_{\mathfrak{M}}, X), E_i \rangle \\ &\quad + \sum_i \langle [[X, E_i]_{\mathfrak{K}}, X], E_i \rangle. \end{aligned}$$

In order to evaluate the right-hand side of (3.5) we define coefficients  $b_{jk}$  by  $[X, E_j]_{\mathfrak{M}} = \sum_k b_{jk} E_k$ . Then, using (3.4),

$$\begin{aligned} 2 \sum_i \langle [X, \mathcal{U}(E_i, X)]_{\mathfrak{M}}, E_i \rangle &= -\sum_{i,j} \langle [X, \langle [X, E_j]_{\mathfrak{M}}, E_i \rangle E_j]_{\mathfrak{M}}, E_i \rangle \\ &= -\sum_{i,j} \langle [X, E_j]_{\mathfrak{M}}, E_i \rangle^2 = -\sum_{i,j} b_{ji}^2 \\ &= -\sum_j \left( \sum_i b_{ji}^2 \right) = -\sum_j \|[X, E_j]_{\mathfrak{M}}\|^2 \\ &= -\sum_i \|[X, E_i]_{\mathfrak{M}}\|^2 = \sum_i \langle [X, E_i]_{\mathfrak{M}}, [E_i, X]_{\mathfrak{M}} \rangle \\ &= +\sum_{i,j} \langle \{ \langle [X, E_j]_{\mathfrak{M}}, [E_i, X]_{\mathfrak{M}} \rangle \} E_j, E_i \rangle \\ &= -2 \sum_i \langle \mathcal{U}(X, [E_i, X]_{\mathfrak{M}}), E_i \rangle. \end{aligned}$$

In other words

$$(3.6) \quad -\frac{1}{2} \sum_i \langle [X, \mathcal{U}(E_i, X)]_{\mathfrak{M}}, E_i \rangle - \frac{1}{2} \sum_i \langle \mathcal{U}(X, [E_i, X]_{\mathfrak{M}}), E_i \rangle = 0.$$

Using (3.4) and (3.6) we compute

$$\begin{aligned}
 & -\sum_i \langle \alpha(X, \alpha(E_i, X)), E_i \rangle \\
 &= -\frac{1}{4} \sum_i \langle [X, [E_i, X]_{\mathfrak{M}}]_{\mathfrak{M}}, E_i \rangle - \sum_i \langle \mathcal{U}(X, \mathcal{U}(E_i, X)), E_i \rangle \\
 &\quad -\frac{1}{2} \sum_i \langle [X, \mathcal{U}(E_i, X)]_{\mathfrak{M}}, E_i \rangle - \frac{1}{2} \sum_i \langle \mathcal{U}(X, [E_i, X]_{\mathfrak{M}}), E_i \rangle \\
 &= -\frac{1}{4} \sum_i \langle [X, [E_i, X]_{\mathfrak{M}}]_{\mathfrak{M}}, E_i \rangle - \sum_i \langle \mathcal{U}(X, \mathcal{U}(E_i, X)), E_i \rangle \\
 &= -\frac{1}{4} \sum_i \langle [X, [E_i, X]_{\mathfrak{M}}]_{\mathfrak{M}}, E_i \rangle - \frac{1}{4} \sum_{i,j} \langle [X, E_i]_{\mathfrak{M}}, E_j \rangle \langle [X, E_j]_{\mathfrak{M}}, E_i \rangle.
 \end{aligned}$$

That gives us the first summand of the right-hand side of (3.5):

$$\begin{aligned}
 (3.7) \quad & -\sum_i \langle \alpha(X, \alpha(E_i, X)), E_i \rangle = -\frac{1}{4} \sum_i \langle [[X, E_i]_{\mathfrak{M}}, X]_{\mathfrak{M}}, E_i \rangle \\
 & \quad -\frac{1}{4} \sum_{i,j} \langle [X, E_i]_{\mathfrak{M}}, E_j \rangle \langle [X, E_j]_{\mathfrak{M}}, E_i \rangle.
 \end{aligned}$$

The second summand of the right-hand side of (3.5) is, again using (3.4),

$$\begin{aligned}
 & \sum_i \langle \alpha([X, E_i]_{\mathfrak{M}}, X), E_i \rangle \\
 &= \frac{1}{2} \sum_i \langle [[X, E_i]_{\mathfrak{M}}, X]_{\mathfrak{M}}, E_i \rangle - \frac{1}{2} \sum_{i,j} \langle \langle [X, E_j]_{\mathfrak{M}}, [X, E_i]_{\mathfrak{M}} \rangle E_j, E_i \rangle \\
 &= \frac{1}{2} \sum_i \langle [[X, E_i]_{\mathfrak{M}}, X]_{\mathfrak{M}}, E_i \rangle - \frac{1}{2} \sum_i \|[X, E_i]_{\mathfrak{M}}\|^2.
 \end{aligned}$$

Using (3.7) now, the sum of the first two summands of the right-hand side of (3.5) is

$$\begin{aligned}
 (3.8) \quad & -\sum_i \langle \alpha(X, \alpha(E_i, X)), E_i \rangle + \sum_i \langle \alpha([X, E_i]_{\mathfrak{M}}, X), E_i \rangle \\
 &= +\frac{1}{4} \sum_i \langle [[X, E_i]_{\mathfrak{M}}, X]_{\mathfrak{M}}, E_i \rangle - \frac{1}{2} \sum_i \|[X, E_i]_{\mathfrak{M}}\|^2 \\
 &\quad -\frac{1}{4} \sum_{i,j} \langle [X, E_i]_{\mathfrak{M}}, E_j \rangle \cdot \langle [X, E_j]_{\mathfrak{M}}, E_i \rangle.
 \end{aligned}$$

Adding  $\sum_i \langle [[X, E_i]_{\mathfrak{M}}, X]_{\mathfrak{M}}, E_i \rangle$  to both sides of (3.8), our assertion (3.2) follows from (3.5). Q.E.D.

We apply Lemma 3.1 to the gap between the nilpotent and solvable radicals of  $G$ .

**3.9. THEOREM.** *Let  $(M, ds^2)$  be a riemannian homogeneous space,  $G$  a transitive Lie group of isometries,  $K$  the isotropy subgroup at a point  $\mathcal{O} \in M$ , and  $\mathfrak{L}$  an  $\text{ad}_G(K)$ -invariant Levi factor of  $\mathfrak{G}$ , such that  $ds^2$  is consistent with the Levi-Whitehead decomposition  $\mathfrak{G} = \mathfrak{K} + \mathfrak{L}$ . Choose an invariant complement  $\mathfrak{M} = \mathfrak{K} + \mathfrak{A} + (\mathfrak{M} \cap \mathfrak{L})$*

for  $K$  that satisfies (2.8). Let  $X \in \mathfrak{M}$  be a unit vector,  $X_\theta \in M_\theta$  the corresponding unit tangent vector.

1. If  $X \perp [\mathfrak{G}, \mathfrak{G}]_{\mathfrak{M}}$  then the mean curvature  $k(X_\theta) \leq 0$ , and  $k(X_\theta) = 0$  if and only if (a)  $X \in \mathfrak{R}$  and (b)  $[X, \mathfrak{M}] = 0$ .

2. If  $X \in \mathfrak{A}$  then  $k(X_\theta) < 0$ .

**Proof.** By choice of  $\mathfrak{M}$  and by  $[\mathfrak{G}, \mathfrak{G}] \subset \mathfrak{R} + \mathfrak{L}$  we have an orthogonal direct sum decomposition

$$(3.10) \quad \begin{aligned} \mathfrak{M} &= \mathfrak{M}' + \mathfrak{B} + (\mathfrak{M} \cap \mathfrak{L}), \quad \mathfrak{M}' + \mathfrak{B} = \mathfrak{M} \cap \mathfrak{R}, \quad \mathfrak{M}' \subset \mathfrak{R}, \\ [\mathfrak{G}, \mathfrak{G}]_{\mathfrak{M}} &= \mathfrak{M}' + (\mathfrak{M} \cap \mathfrak{L}), \quad \mathfrak{A} \subset \mathfrak{B}. \end{aligned}$$

Let  $X \in \mathfrak{B}$ . Then we have an orthonormal basis  $\{E_i\}$  of  $\mathfrak{M}$  containing  $X$ , such that each  $E_i$  is in  $\mathfrak{M}'$ ,  $\mathfrak{M} \cap \mathfrak{L}$  or  $\mathfrak{B}$ . We apply Lemma 3.1 with that basis.

Define coefficients by  $[X, E_j]_{\mathfrak{M}} = \sum_k a_{jk} E_k$  and let  $A = (a_{jk})$ . Then

$$\langle [X, E_i]_{\mathfrak{M}}, E_j \rangle = a_{ij} \quad \text{and} \quad \langle [X, E_j]_{\mathfrak{M}}, E_i \rangle = a_{ji}$$

so

$$\sum_{i,j} \langle [X, E_i]_{\mathfrak{M}}, E_j \rangle \langle [X, E_j]_{\mathfrak{M}}, E_i \rangle = \text{trace}(A \cdot {}^t A).$$

Take polar decomposition  $A = ST$  with  $S$  symmetric and  $T$  orthogonal. Then  $A \cdot {}^t A = S \cdot {}^t S$ . Let  $S = (s_{ij})$ , so

$$\text{trace}(A \cdot {}^t A) = \text{trace}(S \cdot {}^t S) = \sum_{i,j} s_{ij}^2 \geq 0,$$

and note that  $\sum s_{ij}^2 = 0$  if and only if  $S = 0$ , which is equivalent to  $A = 0$ . Thus

$$(3.11) \quad -\frac{1}{4} \sum_{i,j} \langle [X, E_i]_{\mathfrak{M}}, E_j \rangle \langle [X, E_j]_{\mathfrak{M}}, E_i \rangle \leq 0$$

with equality if and only if  $[X, \mathfrak{M}]_{\mathfrak{M}} = 0$ .

That takes care of the last summand of (3.2). For the first two summands we define

$$(3.12) \quad k_i = \langle \{[X, E_i]_{\mathfrak{R}} + \frac{1}{4}[X, E_i]_{\mathfrak{M}}\}, X \rangle_{\mathfrak{M}} - \frac{1}{2} \| [X, E_i]_{\mathfrak{M}} \|^2.$$

If  $E_i \in \mathfrak{B}$  then  $\langle [\mathfrak{G}, \mathfrak{G}]_{\mathfrak{M}}, \mathfrak{B} \rangle = 0$  implies  $k_i = -\frac{1}{2} \| [X, E_i]_{\mathfrak{M}} \|^2$ . If  $E_i \in \mathfrak{M} \cap \mathfrak{L}$  then  $[\mathfrak{G}, X]_{\mathfrak{M}} \subset [\mathfrak{G}, \mathfrak{R}]_{\mathfrak{M}} \subset (\mathfrak{M} \cap \mathfrak{R}) \perp (\mathfrak{M} \cap \mathfrak{L})$  implies  $\langle [\mathfrak{G}, X]_{\mathfrak{M}}, E_i \rangle = 0$  so

$$k_i = -\frac{1}{2} \| [X, E_i]_{\mathfrak{M}} \|^2.$$

Thus

$$(3.13) \quad \text{if } E_i \in \mathfrak{B} + (\mathfrak{M} \cap \mathfrak{L}) \text{ then } k_i = -\frac{1}{2} \| [X, E_i]_{\mathfrak{M}} \|^2 \leq 0.$$

If  $E_i \in \mathfrak{M}'$  then  $E_i \in \mathfrak{R}$  so  $[X, E_i] \in \mathfrak{R} \subset \mathfrak{M}$ . Thus

$$(3.14) \quad \text{if } E_i \in \mathfrak{M}' \text{ then } k_i = -\frac{1}{4} \langle \text{ad}(X)^2 E_i, E_i \rangle - \frac{1}{2} \| [X, E_i] \|^2.$$



As  $(\text{ad } X)\mathfrak{N}' \subset \mathfrak{N}'$  we can stipulate that, for numbers  $\{\lambda_b\}$  such that  $\{\lambda_b, \bar{\lambda}_b\}$  are the eigenvalues of  $(\text{ad } X)|_{\mathfrak{N}'}$ , each  $E_i \in \mathfrak{N}'$  is contained in the sum of the subspaces of  $\mathfrak{N}'^c$  on which (for some  $b = b_i$ )  $\text{ad } X - \lambda_b$  and  $\text{ad } X - \bar{\lambda}_b$  are nilpotent. That stipulation made,  $\|[X, E_i]\|^2 \geq |\lambda_b|^2$  and  $|\langle \text{ad } (X)^2 E_i, E_i \rangle| \leq |\lambda_b|^2$ . So (3.14) implies

$$(3.15) \quad \text{if } E_i \in \mathfrak{N}' \text{ then } k_i \leq -\frac{1}{4}|\lambda_b|^2 \leq 0.$$

Combining (3.13) and (3.15) we have  $\sum_i k_i \leq 0$ . Adding that inequality to (3.11), and applying Lemma 3.1, we conclude

$$(3.16) \quad k(X_\theta) \leq 0 \text{ with equality if and only if } [X, \mathfrak{M}]_{\mathfrak{M}} = 0.$$

If  $[X, \mathfrak{M}]_{\mathfrak{M}} = 0$  then  $[X, \mathfrak{M}] \subset \mathfrak{R}$ . As  $\mathfrak{N} \subset \mathfrak{M}$  and  $[X, \mathfrak{N}] \subset \mathfrak{N}$  it follows that  $[X, \mathfrak{N}] = 0$ . Then  $\mathfrak{S} = X\mathfrak{R} + \mathfrak{N}$  is a nilpotent subalgebra of  $\mathfrak{G}$ . But  $X \in \mathfrak{N}$  and  $[\mathfrak{N}, \mathfrak{N}] \subset \mathfrak{N}$ , so  $\mathfrak{S}$  is a nilpotent ideal in  $\mathfrak{N}$ . As  $\mathfrak{N}$  is the maximal nilpotent ideal of  $\mathfrak{N}$  it follows that  $X \in \mathfrak{N}$ . This fact and (3.16) imply the first statement of Theorem 3.9. If  $X \in \mathfrak{A}$  then  $X \notin \mathfrak{N}$ , so  $k(X_\theta) < 0$ . That completes the proof of Theorem 3.9. Q.E.D.

**4. Application to manifolds of nonnegative mean curvature.** We first apply Theorem 3.9 to homogeneous riemannian manifolds.

**4.1. THEOREM.** *Let  $(M, ds^2)$  be a connected homogeneous riemannian manifold,  $G$  a transitive Lie group of isometries,  $K$  an isotropy subgroup, and  $\mathfrak{L}$  an  $\text{ad}_G(K)$ -invariant Levi factor of  $\mathfrak{G}$  such that  $ds^2$  is consistent with the Levi-Whitehead decomposition  $\mathfrak{G} = \mathfrak{R} + \mathfrak{L}$ .*

1. *If  $(M, ds^2)$  has mean curvature  $\geq 0$  everywhere, then the solvable radical  $\mathfrak{R}$  and the nilpotent radical  $\mathfrak{N}$  of  $\mathfrak{G}$  satisfy  $\mathfrak{R} = \mathfrak{N} + (\mathfrak{R} \cap \mathfrak{N})$  semidirect sum.*

2. *If  $(M, ds^2)$  has mean curvature  $> 0$  everywhere, then the derived group  $[G, G]$  of  $G$  is transitive on  $M$ .*

**Proof.** If  $(M, ds^2)$  has mean curvature  $\geq 0$  everywhere, then, in the notation (2.8), Theorem 3.9 says  $\mathfrak{A} = 0$ , so  $\mathfrak{M} \cap \mathfrak{R} = \mathfrak{N}$ ; thus  $\mathfrak{R} = (\mathfrak{M} \cap \mathfrak{R}) + (\mathfrak{R} \cap \mathfrak{R}) = \mathfrak{N} + (\mathfrak{R} \cap \mathfrak{R})$ . If further  $(M, ds^2)$  has mean curvature  $> 0$  everywhere, then Theorem 3.9 says  $\mathfrak{M} = [\mathfrak{G}, \mathfrak{G}]_{\mathfrak{M}}$ , so the derived group  $G' = [G, G]$  has an open orbit  $G'(\theta) \subset M$ . As  $G'(\theta)$  is complete and  $M$  is connected,  $G'(\theta) = M$ , so  $G'$  is transitive on  $M$ . Q.E.D.

**4.2. COROLLARY.** *Let  $(M, ds^2)$  be a connected riemannian homogeneous manifold,  $G$  a transitive Lie group of isometries of  $M$ ,  $\theta \in M$ , and  $K$  the isotropy subgroup of  $G$  at  $\theta$ . Let  $R_\theta$  denote the subspace of the tangent space  $M_\theta$  consisting of vectors  $Y_\theta$  where  $Y$  is contained in the solvable radical  $\mathfrak{R}$  of  $\mathfrak{G}$ . Suppose that the linear isotropy representation of  $K$  splits into disjoint representations on  $R_\theta$  and  $R_\theta^\perp$ .*

1. *If  $(M, ds^2)$  has mean curvature  $\geq 0$  everywhere, then  $\mathfrak{R}$  is related to the nilpotent radical  $\mathfrak{N}$  of  $\mathfrak{G}$  by  $\mathfrak{R} = \mathfrak{N} + (\mathfrak{R} \cap \mathfrak{R})$ .*

2. If  $(M, ds^2)$  has mean curvature  $> 0$  everywhere, then the derived group of  $G$  is transitive on  $M$ .

**Proof.** Let  $\mathfrak{L}$  be any  $\text{ad}_G(K)$ -invariant Levi factor of  $\mathfrak{G}$ . Proposition 2.4 says that  $ds^2$  is consistent with the Levi-Whitehead decomposition  $\mathfrak{G} = \mathfrak{K} + \mathfrak{L}$ . Our assertions now follow from Theorem 4.1. Q.E.D.

In order to apply Theorem 4.1 to the case of a transitive solvable group of isometries, we must first prove the following lemma about a simply transitive nilpotent group of isometries. Note that the lemma extends the positive curvature portion of [7].

4.3. LEMMA. *Let  $(N, ds^2)$  be a connected nilpotent Lie group with a left invariant riemannian metric. Then the following conditions are equivalent.*

- (i)  $(N, ds^2)$  has mean curvature  $\geq 0$  everywhere.
- (ii)  $(N, ds^2)$  has every sectional curvature zero.
- (iii)  $N$  is commutative.

**Proof.** As (iii)  $\Rightarrow$  (ii)  $\Rightarrow$  (i) trivially we need only check that (i)  $\Rightarrow$  (iii). So assume that  $(N, ds^2)$  has mean curvature  $\geq 0$  everywhere. In the context of Theorem 3.9,

$$G = N, \quad K = \{1\}, \quad \mathfrak{L} = 0, \quad \mathfrak{M} = \mathfrak{N},$$

and consistency of  $ds^2$  with  $\mathfrak{G} = \mathfrak{K} + \mathfrak{L}$  is automatic. Now Theorem 3.9 says that there is no noncentral element  $X \in \mathfrak{N}$  such that  $X \perp [\mathfrak{N}, \mathfrak{N}]$ . But nilpotence of  $\mathfrak{N}$  implies that in the lower central series

$$\mathfrak{N} = \mathfrak{N}_0 \supset \mathfrak{N}_1 \supset \dots \supset \mathfrak{N}_s \supsetneq \mathfrak{N}_{s+1} = 0, \quad \mathfrak{N}_{n+1} = [\mathfrak{N}, \mathfrak{N}_k],$$

any vector space complement to  $\mathfrak{N}_1 = [\mathfrak{N}, \mathfrak{N}]$  generates  $\mathfrak{N}$ . Let  $[\mathfrak{N}, \mathfrak{N}]^\perp$  be the complement. As it consists of central elements of  $\mathfrak{N}$  (our application of Theorem 3.9), it must be all of  $\mathfrak{N}$ . Thus  $N$  is commutative. Q.E.D.

Now we have a general result on the curvature of riemannian solvmanifolds.

4.4. THEOREM. *Let  $(M, ds^2)$  be a connected riemannian manifold that has a solvable transitive group of isometries. Then the following conditions are equivalent.*

- (i)  $(M, ds^2)$  has mean curvature  $\geq 0$  everywhere.
- (ii)  $(M, ds^2)$  has every sectional curvature  $\geq 0$ .
- (iii)  $(M, ds^2)$  has every sectional curvature zero.
- (iv)  $(M, ds^2)$  is isometric to the product of an euclidean space and a flat riemannian torus.

**Proof.** As (iv)  $\Rightarrow$  (iii)  $\Rightarrow$  (ii)  $\Rightarrow$  (i) trivially we need only check that (i)  $\Rightarrow$  (iv). So assume that  $(M, ds^2)$  has mean curvature  $\geq 0$ .

$G$  denotes the closure of a solvable transitive group of isometries of  $(M, ds^2)$  in the full group of isometries. So  $G$  is a solvable transitive Lie group of isometries. Let  $K$  be an isotropy subgroup.  $\mathfrak{G}$  is its own solvable radical  $\mathfrak{R}$ , so the  $\text{ad}_G(K)$ -invariant Levi factor  $\mathfrak{L}=0$ , and  $ds^2$  is consistent with  $\mathfrak{G}=\mathfrak{R}+\mathfrak{L}=\mathfrak{R}$ . Our invariant complement  $\mathfrak{M}$  for  $K$  satisfying (2.8) now has the form  $\mathfrak{M}=\mathfrak{R}+\mathfrak{A}$ , and Theorem 3.9 says  $\mathfrak{A}=0$ .

Let  $N$  be the analytic subgroup of  $G$  for  $\mathfrak{R}$ . Now we have an open orbit  $N(\mathcal{O}) \subset M$ . As  $N(\mathcal{O})$  is complete and  $M$  is connected, the two are equal. Thus  $N$  is transitive on  $M$ . As  $G$  acts effectively on  $M$ , also  $N$  acts effectively, so Lemma 2.5 says  $K \cap N = \{1\}$ . That proves  $N$  simply transitive on  $M$ . Lemma 4.3 says that  $(M, ds^2)$  is flat and  $N$  is commutative. It follows [6, Théorème 4] that  $(M, ds^2)$  is the product of an euclidean space and a flat riemannian torus. Q.E.D.

**4.5. COROLLARY.** *Let  $(M, ds^2)$  be a connected riemannian Einstein manifold that has a solvable transitive group of isometries. Then either  $(M, ds^2)$  has vanishing Ricci tensor and is isometric to the product of an euclidean space with a flat riemannian torus, or  $(M, ds^2)$  has negative definite Ricci tensor.*

**Proof.** The Einstein homogeneous hypothesis says that  $(M, ds^2)$  has constant mean curvature, say  $k$ . If  $k \geq 0$  then Theorem 4.4 says that  $(R_{ij}) \equiv 0$  and that  $(M, ds^2)$  is the product of an euclidean space with a flat riemannian torus. If  $k < 0$  then  $(R_{ij})$  is negative definite. Q.E.D.

For examples of the latter case of Corollary 4.5, let  $(M, ds^2)$  be a noncompact irreducible riemannian symmetric space,  $G$  the largest connected group of isometries,  $K$  an isotropy subgroup, and  $G=NAK$  an Iwasawa decomposition. Then  $S=NA$  is a simply transitive solvable Lie group of isometries of  $(M, ds^2)$ , and  $(M, ds^2)$  is a connected riemannian Einstein manifold with negative definite Ricci tensor. G. Jensen [2] has shown that this example is essentially exhaustive in dimensions  $\leq 4$ .

The following lemma is similar to results of G. Jensen [2].

**4.6. LEMMA.** *Let  $G$  be a Lie group, let  $ds^2$  be a left invariant riemannian metric on  $G$ , and let  $X$  be a nonzero central element of the Lie algebra  $\mathfrak{G}$ . Then the mean curvature  $k(X) \geq 0$ , and  $k(X)=0$  if and only if  $X$  is orthogonal to the derived algebra of  $\mathfrak{G}$ .*

**Proof.** We use the notation of the proof of Lemma 3.1. Note  $\mathfrak{M}=\mathfrak{G}$ . We take  $X$  to be a unit vector and  $\{E_i\}$  to be an orthonormal basis of  $\mathfrak{G}$  that contains  $X$ . Then (3.3) holds. As  $X$  is central in  $\mathfrak{G}$ , the analog of (3.4) is

$$\mathcal{U}(S, X) = \mathcal{U}(X, S) = -\frac{1}{2} \sum_j \langle [S, E_j], X \rangle E_j.$$

We still have  $\mathcal{U}(X, X)=0$ , so  $\alpha(X, X)=0$  and (3.5) holds. But  $[X, E_i]=0$  simplifies (3.5) to

$$\begin{aligned}
 (n-1)k(X) &= -\sum_i \langle \alpha(X, \alpha(E_i, X)), E_i \rangle \\
 &= \frac{1}{2} \sum_{i,j} \langle \alpha(X, \langle [E_i, E_j], X \rangle E_j), E_i \rangle \\
 &= \frac{1}{2} \sum_{i,j} \langle [E_i, E_j], X \rangle \cdot \langle \alpha(X, E_j), E_i \rangle \\
 &= -\frac{1}{4} \sum_{i,j,k} \langle [E_i, E_j], X \rangle \langle [E_j, E_k], X \rangle E_k, E_i \rangle \\
 &= -\frac{1}{4} \sum_{i,j} \langle [E_i, E_j], X \rangle \langle [E_j, E_i], X \rangle \\
 &= \frac{1}{4} \sum_{i,j} \langle [E_i, E_j], X \rangle^2.
 \end{aligned}$$

Thus  $k(X) \geq 0$ , and  $k(X)=0$  if and only if each  $\langle [E_i, E_j], X \rangle = 0$ , which is equivalent to  $\langle [\mathfrak{G}, \mathfrak{G}], X \rangle = 0$ . Q.E.D.

We now combine Lemmas 4.3 and 4.6, extending our calculations [7] from sectional curvature to mean curvature, and sharpening Theorem 4.4 in the case of a nilpotent group. After hearing the result, G. Jensen gave another proof of Theorem 4.7 [2, Theorem 4].

**4.7. THEOREM.** *Let  $(M, ds^2)$  be a connected riemannian manifold that has a nilpotent transitive group of isometries. Then the following conditions are equivalent.*

- (i)  $(M, ds^2)$  has mean curvature  $\geq 0$  everywhere.
- (ii)  $(M, ds^2)$  has mean curvature  $= 0$  everywhere.
- (iii)  $(M, ds^2)$  has mean curvature  $\leq 0$  everywhere.
- (iv)  $(M, ds^2)$  has every sectional curvature  $\geq 0$ .
- (v)  $(M, ds^2)$  has every sectional curvature  $= 0$ .
- (vi)  $(M, ds^2)$  has every sectional curvature  $\leq 0$ .
- (vii)  $(M, ds^2)$  is isometric to the product of an euclidean space and a flat riemannian torus.

**Proof.** Let  $N$  denote the identity component of the closure of a nilpotent transitive group of isometries. Then  $N$  is a connected nilpotent transitive Lie group of isometries of  $(M, ds^2)$ . Its isotropy subgroups are central by Lemma 2.5, hence trivial; thus  $N$  is simply transitive on  $(M, ds^2)$ . Now we may view  $ds^2$  as a left invariant riemannian metric on  $N$ .

Lemma 4.3 says that (i) implies (v); so (ii) implies (v).

We use Lemma 4.6 to prove that (iii) implies (v). Let  $\mathfrak{Z}$  be the last nonzero term of the lower central series of  $\mathfrak{N}$ . Then  $\mathfrak{Z}$  is central in  $\mathfrak{N}$ . Let  $0 \neq X \in \mathfrak{Z}$ . Assume (iii), so  $k(X) \leq 0$ . Lemma 4.6 says  $k(X) \geq 0$ . Thus  $k(X)=0$  and Lemma 4.6 says  $\langle [\mathfrak{N}, \mathfrak{N}], X \rangle = 0$ . If  $\mathfrak{N}$  is noncommutative then  $X \in \mathfrak{Z} \subset [\mathfrak{N}, \mathfrak{N}]$  and so  $\langle [\mathfrak{N}, \mathfrak{N}], X \rangle$

$\neq 0$ . That proves  $N$  commutative, so every sectional curvature of  $(N, ds^2)$  is zero. Thus (iii) implies (v).

Now (i), (ii) and (iii) each implies (v). It follows that (iv) and (vi) each implies (v). But (v) implies (vii) by [6, Théorème 4], and (vii) clearly implies each of (i), (ii), (iii), (iv), (v) and (vi). Q.E.D.

**4.8. COROLLARY.** *Let  $(M, ds^2)$  be a connected riemannian Einstein manifold that has a nilpotent transitive group of isometries. Then  $(M, ds^2)$  is isometric to the product of an euclidean space and a flat riemannian torus.*

**Proof.** We have the hypothesis of Theorem 4.7 as well as condition (i), (ii) or (iii); thus we have condition (vii) of Theorem 4.7. Q.E.D.

Our last application is a refinement of [8, Corollary 5.8].

**4.9. THEOREM.** *Let  $(M, ds^2)$  be a compact connected locally homogeneous riemannian manifold with mean curvature  $\geq 0$  everywhere. Suppose that the fundamental group  $\pi_1(M)$  has a solvable subgroup of finite index.*

*Let  $\pi: \tilde{M} \rightarrow M$  be the universal riemannian covering,  $G$  the largest connected group of isometries of  $(\tilde{M}, \pi^* ds^2)$ ,  $K$  an isotropy subgroup of  $G$ ,  $\mathfrak{L}$  an  $\text{ad}_G(K)$ -invariant Levi factor of  $\mathfrak{G}$  and  $L$  its analytic subgroup of  $G$ , and  $\mathfrak{R}$  and  $\mathfrak{N}$  the solvable and nilpotent radicals of  $\mathfrak{G}$  and  $R$  and  $N$  their analytic subgroups of  $G$ . Let  $\Gamma$  denote the group  $\cong \pi_1(M)$  of deck transformations of  $\tilde{M} \rightarrow M$ . Suppose that  $\pi^* ds^2$  is consistent with  $\mathfrak{G} = \mathfrak{R} + \mathfrak{L}$ .*

1.  $L$  is compact.
2.  $R = N \cdot (K \cap R)_0$  semidirect product, where  $(K \cap R)_0$  is a torus group whose Lie algebra  $\mathfrak{K} \cap \mathfrak{R}$  acts effectively on  $\mathfrak{R}$  in the adjoint representation.
3.  $\Gamma$  has a torsion free normal nilpotent subgroup  $\Delta$  of finite index,  $\Delta \subset N \cdot Z_L(N)_0$  where  $Z_L(N)$  is the centralizer of  $N$  in  $L$ , and  $\Delta$  projects isomorphically to a discrete subgroup with compact quotient in  $N$ .

**Proof.** Compactness of  $L$  is part of [8, Corollary 5.8], and the decomposition  $R = N \cdot (K \cap R)_0$  follows from Theorem 4.1. In the proof of [8, Corollary 5.8] it is shown that  $\Gamma$  has a nilpotent subgroup  $\Delta$  of finite index, and that the identity component of the closure of  $R\Delta$  in  $G$  has form

$$F = R \cdot U \text{ for some torus } U \subset L.$$

Enlarge  $U$  to a maximal torus  $T$  of  $F$ . Then  $T = T_N \cdot T_{R/N} \cdot U$  local direct product, where  $T_N$  is a maximal torus of  $N$  and  $T_{R/N}$  is an  $\text{ad}_G(R)$ -conjugate of  $(K \cap R)_0$ . These constructions are not changed if  $\Delta$  is cut down to a subgroup of finite index. So we first cut  $\Delta$  down to  $\Delta \cap F$ , then [8, (4.5)] to a torsion free group, and finally to a normal subgroup of  $\Gamma$ .

Let  $V$  be the kernel of the action of  $T$  on  $N$ , i.e. the centralizer  $Z_T(N)$ . Then  $T_N \subset V_0 \subset T_N \cdot U$ . Let  $F^* = F/V$ ; then  $F^* = N^* \cdot T^*$  where  $N^* = N/T_N$  simply connected nilpotent group and

$$T^* = (T_{R/N} \cdot U) / \{(T_{R/N} \cdot U) \cap V\}.$$

$\Delta^*$  is the projection of  $\Delta$  to  $F^*$ . As  $\Delta$  is a discrete subgroup with compact quotient in  $F$ , the same is true of  $\Delta^*$  in  $F^*$ . By construction of  $F^*$  and the fact that  $T_N$  is central in  $F$ , conjugation represents  $T^*$  faithfully as a group of automorphisms of  $N^*$ , so L. Auslander's result [1] says that  $\Delta^* \cap N^*$  has finite index in  $\Delta^*$ . Again cutting  $\Delta$  down, we may assume  $\Delta^* \subset N^*$ , i.e. that  $\Delta \subset N \cdot Z_U(N)_0 \subset N \cdot Z_L(N)_0$ . Q.E.D.

#### REFERENCES

1. L. Auslander, *Bieberbach's theorems on space groups and discrete uniform subgroups of Lie groups*, Ann. of Math. **71** (1960), 579–590.
2. G. Jensen, *Homogeneous Einstein spaces*, Thesis, University of California, Berkeley, 1968.
3. G. D. Mostow, *Fully reducible subgroups of algebraic groups*, Amer. J. Math. **78** (1956), 200–221.
4. K. Nomizu, *Invariant affine connections on homogeneous spaces*, Amer. J. Math. **76** (1954), 33–65.
5. E. J. Taft, *Orthogonal conjugacies in associative and Lie algebras*, Trans. Amer. Math. Soc. **113** (1964), 18–29.
6. J. A. Wolf, *Sur la classification des variétés riemanniennes homogènes à courbure constante*, C. R. Acad. Sci. Paris **250** (1960), 3443–3445.
7. ———, *Curvature in nilpotent Lie groups*, Proc. Amer. Math. Soc. **15** (1964), 271–274.
8. ———, *Growth of finitely generated solvable groups and curvature of riemannian manifolds*, J. Differential Geometry, (to appear).

UNIVERSITY OF CALIFORNIA,  
BERKELEY, CALIFORNIA