

# UNIVALENT FUNCTIONS WITH UNIVALENT DERIVATIVES. II

BY

S. M. SHAH<sup>(1)</sup> AND S. Y. TRIMBLE

**1. Introduction.** Let  $D$  denote the open unit disc with center at the origin. It is known that if  $f$  and all its derivatives are univalent in  $D$ , then  $f$  must be an entire function of exponential type [11]. However, to conclude that  $f$  is entire, we shall show it is not necessary to suppose that each derivative is univalent in  $D$ .

Let  $\rho_n$  be the largest number with the property that  $f^{(n)}$  is univalent in an open disc about the origin of radius  $\rho_n$ . (Note that  $\rho_n$  is finite unless  $f^{(n)}(z) = az + b$ . We shall exclude this possibility by always assuming that  $f$  is not a polynomial.) In this paper, we investigate the relation between the growth of  $\{\rho_n\}_{n=0}^{\infty}$  and the radius of convergence of  $f$  about the origin. In particular, we show that if  $\rho_n$  converges to zero slowly enough, then  $f$  must still be an entire function. (See the corollary to Theorem 1.) Further, if  $f$  is entire, we exhibit relations between the growth of  $\{\rho_n\}_{n=0}^{\infty}$  and the order and type of  $f$ .

It is well known [12, p. 212], [3] that if  $f$  is defined in a disc about the origin of radius  $\rho$  by

$$f(z) = \sum_{n=0}^{\infty} a_n z^n,$$

and if  $a_1 \neq 0$ , then  $f$  is univalent in this disc if

$$(1.1) \quad \sum_{n=2}^{\infty} n|a_n|\rho^{n-1} \leq |a_1|.$$

It is also well known [6, p. 213], [4, p. 3] that if  $f$  is univalent in  $D$ , then  $a_1 \neq 0$  and

$$(1.2) \quad |a_2| \leq 2|a_1|.$$

We shall use both these facts in the proofs below. For convenience of notation, we shall sometimes write  $f^{(0)} = f$  and  $a_n = a(n)$ .

**2. Derivatives with varying radii of univalence.** We single out two kinds of functions. Let  $f$  be an analytic function defined in  $|z| < R$ . (We allow  $R = \infty$ .) We shall say that  $f$  has property (A) at  $N$  if there is a nonnegative integer,  $N$ , such that for  $n \geq N$ ,  $\rho_n > 0$ . Note that this implies that if  $n \geq N$ , then  $a_{n+1} \neq 0$ . We shall say that  $f$  has property (B) at  $N$  if there is a positive integer,  $N$ , such that

---

Presented to the Society, January 24, 1969; received by the editors February 18, 1969.

<sup>(1)</sup> The research work of the first author was supported by the National Science Foundation under Grant GP-7544.

$\{|a_{n-1}/a_n\}_{n=N}^{\infty}$  is a positive and nondecreasing sequence. Note that property (B) implies property (A).

**THEOREM 1.** *Let  $f$  be defined by  $f(z) = \sum_{k=0}^{\infty} a_k z^k$ . Let  $R$  denote the radius of convergence of  $f$  and  $\rho_n$  the radius of univalence of  $f^{(n)}$ . Then*

$$(2.1) \quad \liminf_{n \rightarrow \infty} n\rho_n \leq 4R$$

and

$$(2.2) \quad R \log 2 \leq \limsup_{n \rightarrow \infty} n\rho_n.$$

If  $f$  has property (A) at  $N$ , then

$$(2.3) \quad \liminf_{n \rightarrow \infty} [n(\rho_N \rho_{N+1} \cdots \rho_n)^{1/n}] \leq 4eR.$$

If  $f$  has property (B), then

$$(2.4) \quad R \log 2 \leq \liminf_{n \rightarrow \infty} n\rho_n \leq \limsup_{n \rightarrow \infty} n\rho_n \leq 4R.$$

**Proof.** If an infinite number of the  $\rho_n$  are zero, then (2.1) is obviously true. So, suppose  $f$  has property (A) at  $N$ . Let  $F_n(z) = f^{(n)}(\rho_n z)$ . Using (1.2) on  $F_n$  and assuming  $n \geq N$ , we have

$$(2.5) \quad |a_{n+2}| \leq 4|a_{n+1}|/(n+2)\rho_n.$$

Since  $\liminf_{n \rightarrow \infty} |a_n/a_{n+1}| \leq R$ , (2.1) is established. If  $f$  has property (B), then  $\lim_{n \rightarrow \infty} |a_n/a_{n+1}| = R$  and the right-hand part of (2.4) is established.

Using (2.5), an induction argument shows that if  $k \geq N+2$ , then

$$|a_k| \leq \frac{4^{k-N-1} |a_{N+1}| (N+1)!}{(\rho_N \rho_{N+1} \cdots \rho_{k-2}) k!}.$$

So,

$$\frac{1}{R} = \limsup_{k \rightarrow \infty} |a_k|^{1/k} \leq \frac{4e}{\liminf_{k \rightarrow \infty} [k(\rho_N \cdots \rho_k)^{1/k}]}.$$

This proves (2.3).

We no longer assume that  $f$  has property (A). Let  $0 < r < R$ . Since  $\sum |a_n| r^n < \infty$ , there is an increasing sequence,  $\{n_p\}_{p=1}^{\infty}$ , of positive integers such that for  $p=1, 2, \dots$ , and  $k=2, 3, \dots$ , we have  $|a(n_p+1)| \geq |a(n_p+k)| r^{k-1}$ . For  $n=1, 2, \dots$ , let  $x_n = n(1-2^{-1/(n+2)})$ . Then

$$(2.6) \quad \sum_{k=2}^{\infty} \frac{(k+n_p)! |a(n_p+k)| r^k [x(n_p)]^k}{(k-1)! n_p^k} \leq r |a(n_p+1)| \sum_{k=2}^{\infty} \frac{(k+n_p)!}{(k-1)!} \left( \frac{x(n_p)}{n_p} \right)^k \\ = \frac{(n_p+1)! |a(n_p+1)| r x(n_p)}{n_p}.$$

Define  $F_p$  in  $D$  by

$$F_p(z) = f^{(n_p)}(r z x(n_p)/n_p).$$

From (1.1) and (2.6), it follows that  $F_p$  is univalent in  $D$ . Hence,  $rx(n_p)/n_p \leq \rho(n_p)$ . Since  $\lim_{n \rightarrow \infty} x_n = \log 2$ , (2.2) is proved.

Now assume that  $f$  has property (B) at  $N$ . If  $n \geq N$ , let  $r_n = |a_{n-1}/a_n|$ . Then for  $k=2, 3, \dots$ , we have  $|a_{n+k}| \geq |a_{n+k}| r_n^{k-1}$ . Using the preceding argument, it follows that if  $n \geq N$ , then

$$(2.7) \quad r_n x_n \leq n \rho_n.$$

Since  $\lim_{n \rightarrow \infty} r_n x_n = R \log 2$ , (2.4) follows and the entire theorem is proved.

**COROLLARY.** *If  $\lim_{n \rightarrow \infty} n \rho_n = \infty$ , then  $f$  is a transcendental entire function. If  $f$  is a transcendental entire function, then  $\limsup_{n \rightarrow \infty} n \rho_n = \infty$ . If  $f$  is a transcendental entire function with property (B), then  $\lim_{n \rightarrow \infty} n \rho_n = \infty$ .*

The converse of the first part of this corollary is false. In fact, if  $\{b_n\}_{n=0}^{\infty}$  is any sequence of positive numbers such that  $\liminf_{n \rightarrow \infty} b_n > 0$ , then there is an entire function,  $f$ , such that each  $\rho_n > 0$  but  $\liminf_{n \rightarrow \infty} b_n \rho_n = 0$ . For instance, consider the following: If  $n=0, 1, \dots$ , let  $a_{2n} = 1/((2n)!)^{1/2}$  and  $a_{2n+1} = 1/b_{2n}((2n+1)!)^{1/2}$ . (If  $b_{2n}=0$ , let  $a_{2n+1}=0$ . There can only be a finite number of these.) Define  $f_1$  by  $f_1(z) = \sum_{n=0}^{\infty} a_n z^n$ . Then  $f_1$  is certainly entire, and for large  $n$ , (2.5) becomes  $b_{2n} \rho_{2n} \leq 4/(2n+2)^{1/2}$ .

The converse of the second part of the corollary is also false. Let  $n_1=2$ . Suppose that  $p \geq 1$  and that  $n_p$  has been chosen. Let  $n_{p+1}$  be an integer such that  $n_{p+1} > n_p$  and if  $j \geq n_{p+1} + 1 - n_p$ , then

$$j^2(j+n_p)^{n_p+1} \leq n_p^{(j-1)/2}.$$

If  $n=n_p$  for some  $p$ , let  $a_{n+1}=1$ . Otherwise, let  $a_n=0$ . Define  $f_2$  by  $f_2(z) = \sum_{j=1}^{\infty} a_j z^j$ . We use (1.1) to show that  $f_2^{(n_p)}$  is univalent in a disc about the origin of radius  $1/(n_p)^{1/2}$ :

$$\begin{aligned} \sum_{j=2}^{\infty} j \left| \frac{(j+n_p)! a(j+n_p)}{j!} \right| \frac{1}{n_p^{(j-1)/2}} &\leq \sum_{j=n_{p+1}+1-n_p}^{\infty} \frac{(j+n_p)!}{(j-1)! n_p^{(j-1)/2}} \\ &< \sum_{j=n_{p+1}+1-n_p}^{\infty} \frac{(j+n_p)^{n_p+1}}{n_p^{(j-1)/2}} \leq \sum_{j=2}^{\infty} \frac{1}{j^2} < (n_p+1)!. \end{aligned}$$

Hence, for  $f_2$ ,  $\limsup_{n \rightarrow \infty} n \rho_n \geq \limsup_{p \rightarrow \infty} n_p/(n_p)^{1/2} = \infty$ , but the radius of convergence of  $f_2$  about the origin is 1.

**3. Entire functions and univalent derivatives.** Next, we obtain relations between the radii of univalence of the derivatives of an entire function,  $f$ , and the order and type of  $f$ . In this and the following section, we shall let  $\Lambda$  be the order and  $\lambda$  be the lower order of  $f$ . If  $0 < \Lambda < \infty$ , we let  $T$  be the type and  $t$  be the lower type of  $f$ .

Several results connecting the order and type of an entire function with the sequence,  $\{\rho_n\}_{n=1}^{\infty}$ , already exist. Boas has shown [1] that if  $f$  is a transcendental entire function of exponential type less than  $\log 2$ , then there is a subsequence,

$\{\rho(n_p)\}_{p=1}^\infty$ , such that  $\rho(n_p) \geq 1$  for all  $p$ . (Levinson [5] supplied a second proof of this.) Boas also pointed out [1] that if the order of  $f$  is less than one, or if  $f$  is of order one but minimal type, then  $\limsup_{n \rightarrow \infty} \rho_n = \infty$ . Pólya [7, p. 18] has stated that

$$\liminf_{n \rightarrow \infty} \frac{\log \rho_n}{\log n} \leq \frac{1 - \Lambda}{\Lambda} \leq \limsup_{n \rightarrow \infty} \frac{\log \rho_n}{\log n}.$$

We shall improve these results and establish several more as well.

**LEMMA.** Let  $f$  be defined on  $|z| < R$  by  $f(z) = \sum_{k=0}^\infty a_k z^k$ . (We allow  $R = \infty$ .) Let  $v(r)$  denote its central index. For  $n = 1, 2, \dots$ , let  $x_n = n(1 - 2^{-1/(n+2)})$ . Then if  $R > r > 0$  and  $v(r) \geq 2$ ,

$$(3.1) \quad rx(v(r) - 1) < (v(r) - 1)\rho(v(r) - 1).$$

**Proof.** Let  $0 < r < R$ . Then  $|a(v(r))| > |a(v(r) + k)|r^k$  for  $k = 1, 2, \dots$ . Suppose  $n = v(r) - 1$ . Using the same argument that proved (2.2), it follows that  $rx_n < n\rho_n$ .

**THEOREM 2.** Let  $f$  be a transcendental entire function defined by  $f(z) = \sum_{n=0}^\infty a_n z^n$ . Let  $\delta = \liminf_{r \rightarrow \infty} v(r)/r$ . Then

$$(3.2) \quad \liminf_{n \rightarrow \infty} \frac{\log(\max\{1, n\rho_n\})}{\log n} \leq \frac{1}{\Lambda},$$

$$(3.3) \quad \frac{1 - \lambda}{\lambda} \leq \limsup_{n \rightarrow \infty} \frac{\log \rho_n}{\log n},$$

and

$$(3.4) \quad \frac{\log 2}{\delta} \leq \limsup_{n \rightarrow \infty} \rho_n.$$

Suppose that  $0 < \Lambda < \infty$  and that  $f$  has property (A) at  $N$ . In this case

$$(3.5) \quad e^{\Lambda-1} \liminf_{n \rightarrow \infty} n^{\Lambda-1} \rho_n^\Lambda \leq \liminf_{n \rightarrow \infty} (\rho_N \rho_{N+1} \cdots \rho_n)^{\Lambda/n} (n^{\Lambda-1}) \leq \frac{4^\Lambda e^{\Lambda-1}}{\Lambda T}.$$

In any case,

$$(3.6) \quad \liminf_{n \rightarrow \infty} n^{\Lambda-1} \rho_n^\Lambda \leq \frac{4^\Lambda}{\Lambda T}.$$

**Proof.** To prove (3.2), we may assume that  $n\rho_n \geq 1$  for all  $n$  and that  $\Lambda > 0$ . It is known [8] that

$$\frac{1}{\Lambda} \geq \liminf_{n \rightarrow \infty} \frac{\log |a_n/a_{n+1}|}{\log n}.$$

From (2.5), we have that

$$\log n\rho_n < \log 4 + \log |a_{n+1}/a_{n+2}|.$$

Hence,

$$\frac{1}{\Lambda} \geq \liminf_{n \rightarrow \infty} \frac{\log n}{\log(n+1)} \left( \frac{\log n\rho_n}{\log n} - \frac{\log 4}{\log n} \right) = \liminf_{n \rightarrow \infty} \frac{\log n\rho_n}{\log n}.$$

It has been shown [13] that

$$\frac{1}{\lambda} = \limsup_{r \rightarrow \infty} \frac{\log r}{\log v(r)}.$$

Since  $f$  is entire,  $\lim_{r \rightarrow \infty} v(r) = \infty$ . So, from (3.1), for all large  $r$ , we have

$$\frac{\log r}{\log v(r)} \leq \frac{\log \rho(v(r)-1)}{\log (v(r)-1)} + 1 + \frac{\log x(v(r)-1)}{\log v(r)}.$$

Hence,

$$\begin{aligned} \frac{1}{\lambda} &\leq \limsup_{r \rightarrow \infty} \frac{\log \rho(v(r)-1)}{\log (v(r)-1)} + 1 \\ &\leq 1 + \limsup_{n \rightarrow \infty} \frac{\log \rho_n}{\log n}. \end{aligned}$$

This establishes (3.3).

Using (3.1) again, for large  $r$  we have

$$\frac{r}{v(r)} < \frac{\rho(v(r)-1)}{x(v(r)-1)}.$$

Therefore,

$$\frac{1}{\delta} \leq \limsup_{r \rightarrow \infty} \frac{\rho(v(r)-1)}{x(v(r)-1)} \leq \frac{\limsup_{n \rightarrow \infty} \rho_n}{\log 2}.$$

This proves (3.4).

Now suppose that  $f$  has property (A) at  $N$ . It is known [2, p. 11] that

$$(3.7) \quad e\Lambda T = \limsup_{n \rightarrow \infty} n|a_n|^{\Lambda/n}.$$

But

$$\limsup_{n \rightarrow \infty} n|a_n|^{\Lambda/n} = \limsup_{n \rightarrow \infty} n \prod_{k=N+2}^n \left| \frac{a_k}{a_{k-1}} \right|^{\Lambda/n}.$$

From (2.5),

$$\begin{aligned} e\Lambda T &\leq \limsup_{n \rightarrow \infty} n \prod_{k=N+2}^n \left( \frac{4}{k\rho_{k-2}} \right)^{\Lambda/n} \\ &= (4e)^\Lambda \limsup_{n \rightarrow \infty} \frac{n^{1-\Lambda}}{(\rho_N \rho_{N+1} \cdots \rho_n)^{\Lambda/n}}. \end{aligned}$$

Since the left-hand inequality of (3.5) is always true, (3.5) is proved. If an infinite number of the  $\rho_n$  are zero, then (3.6) is trivial. If only a finite number of the  $\rho_n$  are zero, then (3.6) becomes (3.5). This establishes the theorem.

Suppose that  $0 < \Lambda < \infty$ , and let  $\delta' = \liminf_{r \rightarrow \infty} v(r)/r^\Lambda$ . In [9], it was shown that  $\delta' \leq \Lambda t \leq \Lambda T$ . Hence, if  $\Lambda = 1$ , (3.4) can be written as

$$(3.8) \quad \frac{\log 2}{t} \leq \frac{\log 2}{\delta} \leq \limsup_{n \rightarrow \infty} \rho_n.$$

Further, if  $\lambda < 1$ , then  $\delta = 0$ , and either (3.3) or (3.4) imply that  $\limsup_{n \rightarrow \infty} \rho_n = \infty$ . This, together with (3.8), yield improvements on the results of Boas. We note that

there are transcendental entire functions of order one such that  $\delta=0$  and  $T>0$ . For example, let

$$\psi(z) = \sum_{n=1}^{\infty} \frac{z^{p_n}}{p_n!}$$

where  $p_1=3$  and if  $n>1$ ,  $p_n=[p_{n-1} \log p_{n-1}]$ . Then  $\Lambda=T=1$ , but  $\delta=0$ . We summarize these results in a corollary.

**COROLLARY 1.** *If  $\lambda < 1$ , then  $\limsup_{n \rightarrow \infty} \rho_n = \infty$ . If  $\Lambda = 1$ , then*

$$\frac{\log 2}{t} \leq \frac{\log 2}{\delta} \leq \limsup_{n \rightarrow \infty} \rho_n.$$

*In particular, if  $\delta=0$ , then  $\limsup_{n \rightarrow \infty} \rho_n = \infty$ .*

From (3.2) and (3.4), we get a second corollary.

**COROLLARY 2.** *If  $\Lambda > 1$ , then  $\liminf_{n \rightarrow \infty} \rho_n = 0$ . If  $\Lambda = 1$ , then*

$$\frac{4}{T} \geq \liminf_{n \rightarrow \infty} \rho_n.$$

*In particular, if  $\Lambda=1$  and  $T=\infty$ , then  $\liminf_{n \rightarrow \infty} \rho_n = 0$ .*

If  $\Lambda=1$  and  $0 < T < \infty$ , then it may be the case that

$$0 < \liminf_{n \rightarrow \infty} \rho_n \leq \limsup_{n \rightarrow \infty} \rho_n < \infty.$$

The function,  $\phi(z)=e^z$ , is an example of this.

**THEOREM 3.** *Let  $f$  be an entire function with property (B) at  $N$ . Let*

$$\gamma = \limsup_{r \rightarrow \infty} v(r)/r.$$

*Then*

$$(3.9) \quad \liminf_{n \rightarrow \infty} \frac{\log \rho_n}{\log n} = \frac{1-\Lambda}{\Lambda} \leq \frac{1-\lambda}{\lambda} = \limsup_{n \rightarrow \infty} \frac{\log \rho_n}{\log n},$$

$$(3.10) \quad \frac{\log 2}{\gamma} \leq \liminf_{n \rightarrow \infty} \rho_n \leq \frac{4}{\gamma},$$

*and*

$$(3.11) \quad \frac{\log 2}{\delta} \leq \limsup_{n \rightarrow \infty} \rho_n \leq \frac{4}{\delta}.$$

*Suppose  $0 < \Lambda < \infty$ . Then*

$$(\log 2)^\Lambda \leq \Lambda t e^{1-\Lambda} \limsup_{n \rightarrow \infty} (\rho_N \rho_{N+1} \cdots \rho_n)^{\Lambda/n} (n^{\Lambda-1}) \leq 4^\Lambda$$

*and*

$$(\log 2)^\Lambda \leq \Lambda T e^{1-\Lambda} \liminf_{n \rightarrow \infty} (\rho_N \rho_{N+1} \cdots \rho_n)^{\Lambda/n} (n^{\Lambda-1}) \leq 4^\Lambda.$$

**Proof.** It is known [8] that

$$(3.12) \quad \liminf_{n \rightarrow \infty} \frac{\log |a_n/a_{n+1}|}{\log n} = \frac{1}{\Lambda} \leq \frac{1}{\lambda} = \limsup_{n \rightarrow \infty} \frac{\log |a_n/a_{n+1}|}{\log n}.$$

From (2.5) and (2.7), it follows that for  $n \geq N$

$$(3.13) \quad x_n \left| \frac{a_{n-1}}{a_n} \right| < n\rho_n < 4 \left| \frac{a_{n+1}}{a_{n+2}} \right|.$$

Hence, from (3.12) and (3.13), we get (3.9).

Since  $f$  has property (B), we have that

$$\gamma = \limsup_{n \rightarrow \infty} n \left| \frac{a_n}{a_{n-1}} \right|$$

and

$$\delta = \liminf_{n \rightarrow \infty} n \left| \frac{a_n}{a_{n-1}} \right|.$$

From this, (2.5), and (2.7), parts (3.10) and (3.11) can be proved.

Assume now that  $0 < \Lambda < \infty$ . From [10], it follows that

$$e\Lambda t = \liminf_{n \rightarrow \infty} n|a_n|^{\Lambda/n}.$$

We use this, (3.7), and (3.13) to establish the last two parts of the theorem. The proofs are similar to the proof of (3.5), and so are omitted.

**COROLLARY.** Let  $f$  be an entire function with property (B).

- (i) If  $\Lambda < 1$ , then  $\lim_{n \rightarrow \infty} \rho_n = \infty$ .
- (ii) If  $\lambda > 1$ , then  $\lim_{n \rightarrow \infty} \rho_n = 0$ .
- (iii) If  $\Lambda = 1$ , then  $\lim_{n \rightarrow \infty} \rho_n = \infty$  if and only if  $T = 0$ . If  $\lim_{n \rightarrow \infty} \rho_n = 0$ , then  $t = \infty$ .

**Proof.** Parts (i) and (ii) follow from (3.9). From [9], we have that if  $\Lambda = 1$ , then  $\delta \leq t \leq T \leq \gamma \leq eT$ . Part (iii) follows from this, (3.10), and (3.11).

**4. Conclusion.** So far, all of the work has been done with functions defined in discs centered at the origin. However, this work immediately carries over to functions defined in a disc centered at any point in the plane. To be specific, let  $f$  be analytic on  $\Delta = \{z : |z - z_0| < r\}$  and let  $g(z) = f(rz + z_0)$  for  $z \in D$ . Then  $f^{(n)}$  is univalent on  $\{z : |z - z_0| < \rho_n \leq r\}$  if and only if  $g^{(n)}$  is univalent on  $\{z : |z| < \rho_n/r\}$ .

The authors would like to thank Professor J. D. Buckholtz for reading the manuscript and for helpful suggestions.

#### REFERENCES

1. R. P. Boas, *Univalent derivatives of entire functions*, Duke Math. J. **6** (1940), 719–721
2. ———, *Entire functions*, Academic Press, New York, 1954.
3. A. W. Goodman, *Univalent functions and nonanalytic curves*, Proc. Amer. Math. Soc. **8** (1957), 598–601.
4. W. K. Hayman, *Multivalent functions*, Cambridge Univ. Press, Cambridge, 1958.

5. N. Levinson, *A theorem of Boas*, Duke Math. J. **8** (1941), 181–182.
6. Zeev Nehari, *Conformal mapping*, McGraw-Hill, New York, 1952.
7. G. Pólya, *On the zeros of the derivatives of a function and its analytic character*, Bull. Amer. Math. Soc. **49** (1943), 178–191.
8. S. M. Shah, *On the lower order of integral functions*, Bull. Amer. Math. Soc. **52** (1946), 1046–1052.
9. ———, *The maximum term of an entire series (III)*, Quart. J. Math. Oxford Ser. **19** (1948), 220–223.
10. ———, *On the coefficients of an entire series of finite order*, J. London Math. Soc. **26** (1951), 45–46.
11. S. M. Shah and S. Y. Trimble, *Univalent functions with univalent derivatives*, Bull. Amer. Math. Soc. **75** (1969), 153–157.
12. E. C. Titchmarsh, *Theory of functions*, Oxford Univ. Press, Oxford, 1944.
13. J. M. Whittaker, *The lower order of integral functions*, J. London Math. Soc. **8** (1933), 20–27.

UNIVERSITY OF KENTUCKY,  
LEXINGTON, KENTUCKY