

SLICES OF MAPS AND LEBESGUE AREA⁽¹⁾

BY
WILLIAM P. ZIEMER

Abstract. For a large class of k dimensional surfaces, S , it is shown that the Lebesgue area of S can be essentially expressed in terms of an integral of the $k-1$ area of a family, F , of $k-1$ dimensional surfaces that cover S . The family F is regarded as being composed of the slices of F . The definition of the $k-1$ area of a surface restricted to one of its slices is formulated in terms of the theory developed by H. Federer, [F3].

1. Introduction. There are many results in geometric measure theory that yield an inequality between the k dimensional measure of a set $R \subset E^n$ and the integral of the $k-1$ dimensional measure of the intersection of R with the level surfaces of certain real-valued functions defined on E^n . There are similar results in the theory of Lebesgue area.

In this paper we will show that the Lebesgue area of a large class of k dimensional surfaces, S , can be essentially expressed in terms of an integral of the $k-1$ area of a family, F , of $k-1$ dimensional surfaces that cover S . An analogous result is obtained for the k dimensional Hausdorff measure of a Hausdorff k -rectifiable set. We regard the family F as being composed of the slices of the surface and one of the essential parts of our problem is to determine the appropriate definition of the $k-1$ area of a surface restricted to one of its slices. The definition that we employ is expressed in terms of the theory developed by H. Federer, [F3]. In case $k=2$, there are two other definitions considered and we show that they lead to the same results. One definition was created by L. Cesari [C] and it employs the concepts of the Carathéodory theory of prime ends. The other definition depends upon the notion of the length of a light mapping that was introduced in [F2].

For 2 dimensional surfaces, R. Rishel [RL] and R. Fullerton [FN] obtained results that are in the same spirit as ours by employing Cesari's definition of generalized length. However, Rishel's definition of the slice of a surface is different from ours and Fullerton's premature death precluded him from developing his results beyond polyhedral parametric surfaces and continuously differentiable non-parametric surfaces.

Received by the editors June 23, 1970.

AMS 1970 subject classifications. Primary 2880; Secondary 2850.

Key words and phrases. Lebesgue area, slice of a surface, current valued measure, rectifiable set.

⁽¹⁾ This work was supported in part by NSF grant GP 19694.

Copyright © 1972, American Mathematical Society

2. Preliminaries and definitions. Euclidean n -space will be denoted by E^n . Lebesgue measure on E^n will be denoted by L_n and H^k will stand for k dimensional Hausdorff measure. If $A \subset E^n$, then the measure $H^k \llcorner A$ is defined by $H^k \llcorner A(E) = H^k(A \cap E)$. A set $R \subset E^n$ is called *Hausdorff k -rectifiable* if there is a Lipschitzian function f on E^k to E^n such that

$$H^k(R - \text{range } f) = 0.$$

We refer the reader to [F4] for a thorough investigation of the properties possessed by Hausdorff k -rectifiable sets. Many of these properties will be employed below and the following will be especially useful. If $R \subset E^n$ is a Hausdorff k -rectifiable set, $g: R \rightarrow E^m$ is Lipschitzian, and $j = \min[k, m]$, then g possesses an $(H^k \llcorner R, k)$ approximate j dimensional Jacobian $H^k \llcorner R$ almost everywhere, [F4, 3.2.19]. Denote this Jacobian by $\text{ap } J_j g$ and in case $k > m = 1$, we will use the notation $\text{ap } |\nabla g|$. The following theorem is of particular importance, [F4, 3.2.20, 3.2.22].

2.1. THEOREM. *If $R \subset E^n$ is Hausdorff k -rectifiable, $g: R \rightarrow E^m$ is Lipschitzian, and $j = \min[k, m]$, then*

$$\int_R \text{ap } J_j g \, dH^k = \int_{E^m} H^\alpha[g^{-1}(y) \cap R] \, dH^j(y)$$

where $\alpha = \max[0, k - m]$.

Finally, we recall the definition of Hausdorff k dimensional density

$$\Theta^k(H^k \llcorner R, y) = \lim_{r \rightarrow 0} \alpha(k)^{-1} r^{-k} H^k[R \cap B(y, r)]$$

where $B(y, r)$ denotes the open n -ball of radius r with center at y and $\alpha(k)$ is the volume of the unit k -ball in E^k . If R is Hausdorff k -rectifiable, then

$$(1) \quad \Theta^k(H^k \llcorner R, y) = 1$$

for H^k almost all $y \in R$.

We will now show that the Hausdorff k measure of a rectifiable set can be expressed in terms of the integral of the $k-1$ measure of its slices. The proof of this result will establish the method that is basic in the demonstration of the main theorem, 3.3, that appears below.

2.2. THEOREM. *Suppose $R \subset E^n$ is a Hausdorff k -rectifiable set. Then*

$$H^k(R) = \sup \left\{ \int_{E^1} H^{k-1}[u^{-1}(r) \cap R] \, dL_1(r) \right\}$$

where the supremum is taken over all Lipschitz functions $u: E^n \rightarrow E^1$ that have Lipschitz constant 1.

Proof. If u is such a function, then $\text{ap } |\nabla u|$ exists and is dominated by 1 at $H^k \llcorner R$ almost all points. Therefore, by appealing to Theorem 2.1 it is clear that the above supremum is no more than $H^k(R)$.

In order to establish the opposite inequality, it will suffice to show that for every $\varepsilon > 0$, there is a function $u: E^n \rightarrow E^1$ with Lipschitz constant 1 which satisfies

$$(2) \quad \int_R \text{ap } |\nabla u| dH^k > (1 - \varepsilon)H^k(R).$$

To this end, first recall that R has an H^k approximate tangent k plane, $P(y)$, at H^k almost all $y \in R$, [F4, 3.3.18]. Moreover, there exists a $H^k \llcorner R$ measurable function α defined on R with values in the space of Grassmann simple k vectors of unit norm. In addition, for H^k almost all $y \in R$, $P(y)$ is the k dimensional vector-subspace of E^n associated with $\alpha(y)$, [F4, 3.2.25]. Consequently, from Lusin's Theorem, the density theorem [F4, 2.9.11], and (1), it follows that at H^k almost all $y \in R$ there is a set $A \subset R$ such that

$$(3) \quad \begin{aligned} & \text{(i) } \Theta^k(H^k \llcorner A, y) = 1 \text{ and} \\ & \text{(ii) } \alpha \text{ is continuous at } y \text{ relative to } A. \end{aligned}$$

Consider a point $y \in R$ where $P(y)$ exists and let $N(y)$ be the $n - k$ plane passing through y that is perpendicular to $P(y)$. For every real number $\beta > 0$, let

$$C(\beta) = \{x : \text{dist } [x, N(y)] > \beta \text{ dist } [x, P(y)]\}.$$

It follows from the definition of the H^k approximate tangent k plane, $P(y)$, that

$$(4) \quad \lim_{r \rightarrow 0} r^{-k} H^k[R \cap B(y, r) - C(\beta)] = 0$$

for every $\beta > 0$.

Choose $y \in R$ such that $P(y)$ exists, and where (1) and (3) hold. This will be true at H^k almost all $y \in R$. Let $\varepsilon > 0$. Choose $\zeta(\varepsilon) = \zeta$ (which will be determined later) and select β large enough to ensure that

$$(5) \quad |\pi_y[(z - y)/|z - y||] > 1 - \zeta$$

whenever $z \in C(\beta)$. Here $\pi_y: E^n \rightarrow P(y)$ denotes the orthogonal projection. From (4), (3), and the equality

$$\Theta^k(H^k \llcorner R, y) = \Theta^k(H^k \llcorner A, y) = 1$$

follows the existence of $r^*(y, \varepsilon) = r^*$ such that, for $0 < r < r^*$,

$$(6) \quad \begin{aligned} & H^k[A \cap B(y, r) \cap C(\beta)] > (1 - \varepsilon)^{1/2} H^k[R \cap B(y, r)] \quad \text{and} \\ & |\alpha(x) - \alpha(y)| < \zeta \quad \text{whenever } x \in A \cap B(y, r). \end{aligned}$$

Select $0 < r < r^*$ and define $u(x) = \text{dist } [x, E^n - B(y, r)]$. Since $|\nabla u(x)| = 1$ in $B(y, r)$, (5) implies that

$$(7) \quad |\pi_y[\nabla u(x)]| > 1 - \zeta \quad \text{for } x \in B(y, r) \cap C(\beta).$$

If ζ is chosen sufficiently small, then from (6) and (7) clearly follows $|\pi_x[\nabla u(x)]| > (1-\varepsilon)^{1/2}$ for $x \in A \cap B(y, r) \cap C(\beta)$ where $P(x)$ exists. Therefore,

$$\int_{R \cap B(y, r)} \text{ap } |\nabla u| \, dH^k \geq (1-\varepsilon)^{1/2} H^k[A \cap B(y, r) \cap C(\beta)]$$

and thus (6) leads to

$$(8) \quad \int_{R \cap B(y, r)} \text{ap } |\nabla u| \, dH^k \geq (1-\varepsilon) H^k[R \cap B(y, r)]$$

whenever $0 < r < r^*$.

For H^k almost every $y \in R$, consider the family of all n -balls $B(y, r)$ where $0 < r < r^*(y, \varepsilon)$. Then by a covering theorem due to Besicovitch and A. P. Morse that has been generalized in [F4, 2.8.15], there exist a countable number of n -balls, B_1, B_2, \dots , whose closures are disjoint such that $H^k \llcorner R(E^n - \bigcup_{i=1}^{\infty} B_i) = 0$. With each n -ball B_i is associated the function $u_i(x) = \text{dist}[x, E^n - B_i]$ so that (8) is satisfied. Letting $u(x) = \sum_{i=1}^{\infty} u_i(x)$, $x \in E^n$, it is clear that u has Lipschitz constant 1 and that

$$\begin{aligned} (1-\varepsilon)H^k(R) &= (1-\varepsilon)H^k \llcorner R(E^n) = (1-\varepsilon) \sum_{i=1}^{\infty} H^k \llcorner R(B_i) \\ &\leq \sum_{i=1}^{\infty} \int_{R \cap B_i} \text{ap } |\nabla u_i| \, dH^k = \int_R \text{ap } |\nabla u| \, dH^k. \end{aligned}$$

According to (2), this concludes the proof.

3. Slices and Lebesgue area. In this section we will establish a result concerning Lebesgue area that is analogous to Theorem 2.2. *Throughout this section we will consider a continuous map $f: X \rightarrow E^n$ of finite Lebesgue area, where X is a k dimensional smooth manifold, $k \leq n$. We will also assume that $k=2$ or $H^{k+1}[f(X)] = 0$.*

Our treatment relies heavily on the work of Federer [F3] and the notation and results of that paper will be employed here without change. Thus, the *monotone-light factorization* $f = l_f \circ m_f$ will be considered where l_f is defined on the *middle space*, M_f . Moreover, there is a unique current-valued measure μ over M_f whose total variation, $\|\mu\|$, is equal to the Lebesgue area of f . If T is a current, then $M(T)$ is the *mass* of T and $F(T)$ denotes the *flat norm* of T . The *boundary* of T is denoted by ∂T . Define a measure ν over E^n by $\nu(A) = \|\mu\|(l_f^{-1}(A))$.

One of the main results of [F3] states that there is an integer valued function, K , such that

$$(9) \quad \mathcal{L}(f) = \int_{E^n} K(y) \, dH^k(y)$$

where $\mathcal{L}(f)$ is the Lebesgue area of f . For $y \in E^n$ and $r > 0$, let $Z(r)$ be the family of components of $l_f^{-1}[B(y, r)]$. For ν almost all $y \in E^n$, there are a finite number of

“essential” points, z , of $I_f^{-1}(y)$ with the property that if r is sufficiently small, then with each $V \in Z(r)$ that contains an essential z is associated an integer $m(V)$ such that

$$(10) \quad K(y) = \sum_{V \in Z(r)} |m(V)|.$$

Moreover, there is an oriented k dimensional plane, $P(y)$, passing through y such that, for each such $V \in Z(r)$,

$$(11) \quad \lim_{r \rightarrow 0} r^{-k} F[\mu(V) - m(V) \cdot P(y) \cap B(y, r)] = 0.$$

3.1. DEFINITION. Let $u: E^n \rightarrow E^1$ be Lipschitz and let $Z(r)$ be the set of components of $M_f \cap \{z : u \circ I_f(z) < r\}$. Define

$$\lambda(f; u, r) = \sum_{V \in Z(r)} M[\partial\mu(V)].$$

As a function of r , $\lambda(f; u, r)$ is L_1 measurable; indeed,

$$\liminf_{t \rightarrow r^-} \lambda(f; u, t) \geq \lambda(f; u, r).$$

To see this, choose $r > 0$ and let V_1, V_2, \dots be the components that constitute $Z(r)$. For $t > r$ let $W_i(t)$ be the union of those components of $Z(t)$ that are contained in V_i , $i = 1, 2, \dots$. The sets $W_i(t)$ are nested and increase with t . Therefore, since $\bigcup_{t < r} W_i(t) = V_i$, it follows that, as currents, $\mu[W_i(t)] \rightarrow \mu(V_i)$ weakly as $t \rightarrow r^-$. From the facts that ∂ is continuous and mass is lower semicontinuous with respect to weak convergence follows

$$\liminf_{t \rightarrow r^-} M[\partial\mu(W_i(t))] \geq M[\partial\mu(V_i)], \quad i = 1, 2, \dots$$

Hence,

$$\begin{aligned} \liminf_{t \rightarrow r^-} \lambda(f; u, t) &\geq \liminf_{t \rightarrow r^-} \sum_{i=1}^{\infty} M[\partial\mu(W_i(t))] \\ &\geq \sum_{i=1}^{\infty} M[\partial\mu(V_i)] = \lambda(f; u, r). \end{aligned}$$

3.2. LEMMA. If $f: X \rightarrow E^n$ and $u: E^n \rightarrow E^1$ has Lipschitz constant N , then

$$\int_{-\infty}^{\infty} \lambda(f; u, r) dL_1(r) \leq N \mathcal{L}(f).$$

Proof. Let

$$\gamma(r) = \|\mu\|(\{z : u \circ I_f(z) < r\}).$$

Observe, for L_1 almost all r , that $\gamma'(r) < \infty$ and for each $V \in Z(r)$ that $\mu(V)$ is an integral current, [F3, 3.4]. Now by applying [FF, 3.9] to $T = \mu(V)$, the proof proceeds as in [F3, 3.2] and we obtain

$$M[\partial\mu(V)] \leq N \liminf_{h \rightarrow 0^+} h^{-1} \|\mu(V)\|(\{y : r - h \leq u(y) < r\}).$$

From this follows

$$\begin{aligned}\lambda(f; u, r) &\leq N \liminf_{h \rightarrow 0^+} h^{-1} \sum_{V \in Z(r)} \|\mu(V)\|(\{y : r-h \leq u(y) < r\}) \\ &\leq N \liminf_{h \rightarrow 0^+} h^{-1} \|\mu(V)\|(\{z : r-h \leq u \circ l_r(z) < r\}) \leq N\gamma'(r).\end{aligned}$$

Our purpose now is to show that the supremum of the left side of 3.2 over all functions u with $N=1$ equals the Lebesgue area of f .

To this end let R be the set of those $y \in E^n$ for which (10) and (11) hold and for which $0 < K(y) < \infty$. Then R is a Hausdorff k -rectifiable set and, without loss of generality, we may assume that

$$(12) \quad \Theta^k(H^k \llcorner R, y) = 1, \quad y \in R.$$

Choose $y \in R$ and to simplify notation, take $y=0$. Assume r to be taken small enough so that (10) holds. Select an essential $z \in l_r^{-1}(0)$ and consider those $V \in Z(r)$ that contain z . Let $T_r = \mu(V)$ and $P = P(0)$. Let $\pi: E^n \rightarrow P$ be the orthogonal projection and define $h_r: P \rightarrow P$ by $h_r(x) = r^{-1} \cdot x$. The flat norm of a current is not increased under a projection and, for k dimensional currents in E^k , the flat norm and the mass norm agree. Therefore, it follows from (11) that

$$\lim_{r \rightarrow 0} r^{-k} M[\pi_{\#}(T_r) - m(V) \cdot B(0, r) \cap P] = 0.$$

This implies

$$\lim_{r \rightarrow 0} M[(h_r \circ \pi)_{\#}(T_r) - m(V) \cdot B(0, 1) \cap P] = 0.$$

Again, from the continuity of ∂ and the lower semicontinuity of mass, for $\eta > 0$ and all sufficiently small r ,

$$M[(h_r \circ \pi)_{\#}(\partial T_r)] \geq |m(V)| M[\partial(B(0, 1) \cap P)] - \eta.$$

Thus, for all small r ,

$$(13) \quad \begin{aligned}M(\partial T_r) &\geq M[\partial \pi_{\#}(T_r)] = M[(h_r \circ \pi)_{\#}(\partial T_r)] r^{k-1} \\ &\geq [|m(V)| - \eta \alpha(k-1)^{-1}] M[\partial(B(0, r) \cap P)].\end{aligned}$$

Choose $\delta > 0$. It follows from (10) and (13) that, for $y \in R$, there is $r^* = r^*(y, \delta)$ such that if $B = B(y, r)$, $0 < r < r^*$, and $u(x) = -\text{dist}(x, E^n - B)$, then

$$\lambda(f; u, -t) = \sum_{V \in Z(-t)} M[\partial \mu(V)] \geq [K(y) - \delta] M[\partial(B(y, t) \cap P)]$$

where $Z(-t)$ is the set of components of $\{z : u \circ l_r(z) < -t\}$ and $0 < t < r$. Therefore,

$$(14) \quad \int_{-\infty}^{\infty} \lambda(f; u, t) dL_1(t) \geq [K(y) - \delta] \alpha(k) r^k.$$

Define a measure ζ over E^n by

$$\zeta(E) = \int_{E \cap R} K(y) dH^k(y)$$

for every Borel set E . Then

$$(15) \quad \zeta(E^n) = \mathcal{L}(f);$$

indeed, $\zeta = \nu$, but we will not use this fact. Appealing to [F4, 2.9.8] we have for H^k almost all $y \in R$, that

$$(16) \quad \lim_{r \rightarrow 0} \zeta[B(y, r)]/H^k \llcorner R[B(y, r)] = K(y).$$

For $\varepsilon > 0$ and $y \in R$, it follows from (12), (14), and (16) that there exists $r^*(y, \varepsilon) = r^*$ such that

$$(17) \quad \int_{-\infty}^{\infty} \lambda(f; u, t) dL_1(t) \geq (1 - \varepsilon)\zeta[B(y, r)]$$

whenever $0 < r < r^*$. Consider the family of n -balls, $B(y, r)$, $y \in R$ and $0 < r < r^*(y, \varepsilon)$. Appealing to the covering theorem [F4, 2.8.15], there exist balls B_1, B_2, \dots whose closures are disjoint and that have the property

$$(18) \quad \zeta\left(E^n - \bigcup_{i=1}^{\infty} B_i\right) = 0.$$

As in the proof of 2.2 we define

$$u(x) = \sum_{i=1}^{\infty} u_i(x), \quad x \in E^n,$$

where $u_i(x) = -\text{dist}(x, E^n - B_i)$, $i = 1, 2, \dots$. Then u has Lipschitz constant 1 and $\lambda(f; u, r) = \sum_{i=1}^{\infty} \lambda(f; u_i, r)$, $r \in E^1$. Therefore, (17) and (18) yield

$$\begin{aligned} \int_{-\infty}^{\infty} \lambda(f; u, r) dL_1(r) &= \sum_{i=1}^{\infty} \int_{-\infty}^{\infty} \lambda(f; u_i, r) dL_1(r) \\ &= \sum_{i=1}^{\infty} (1 - \varepsilon)\zeta(B_i) = (1 - \varepsilon)\zeta(E^n) = (1 - \varepsilon)\mathcal{L}(f). \end{aligned}$$

Thus, the following theorem has been established.

3.3. THEOREM. *Suppose $f: X \rightarrow E^n$ has finite Lebesgue area and suppose $k=2$ or $H^{k+1}[f(X)] = 0$. Then $\sup \{ \int_{-\infty}^{\infty} \lambda(f; u, r) dL_1(r) \} = \mathcal{L}(f)$ where the supremum is taken over all functions $u: E^n \rightarrow E^1$ that have Lipschitz constant 1.*

4. Mappings from a 2-cell. The results of [F3] are also valid in case X is a manifold with boundary, [G], [M], and therefore Theorem 3.3 also holds in this case. In this section we consider the special situation when X is the unit square, Q , in E^2 for then there are two other reasonable definitions for the 1-area or "length" of f restricted to a slice determined by u . In order for 3.3 to remain valid, Definition 3.1 of $\lambda(f; u, r)$ is modified so that only those components $V \in Z(r)$ are considered for which closure $V \cap m_r(\text{bdry } Q) = 0$. Let $C(r)$ be the set of such components.

4.1. DEFINITION. If U is a subset of a separable metric space, let

$$\delta(U) = \text{bdry } U \cap \{z : \text{dimension}(\text{bdry } U, z) > 0\}.$$

4.2. DEFINITION. For $V \in C(r)$, let

$$|V| = \int_{\mathbb{E}^n} N[l_r, \delta(V), y] dH^1(y)$$

where $N[l_r, \delta(V), y]$ denotes the number of points (possibly ∞) in $l_r^{-1}(y) \cap \delta(V)$. Define

$$\rho(f; u, r) = \sum_{V \in C(r)} |V|.$$

Similarly, for $V \in C(r)$, let $\|V\|$ denote the length of f restricted to the boundary of $m_r^{-1}(V)$ as defined by Cesari in [C, 20.2]. Define

$$\sigma(f; u, r) = \sum_{V \in C(r)} \|V\|.$$

Finally, define

$$\mathcal{C}(f) = \sup \left\{ \int_{-\infty}^{\infty} \sigma(f; u, r) dL_1(r) \right\}$$

where the supremum is taken over all functions $u: E^n \rightarrow E^1$ that have Lipschitz constant 1. By replacing $\sigma(f; u, r)$ by $\rho(f; u, r)$, we define $\mathcal{D}(f)$ in a similar manner.

The following result is an immediate consequence of [C, 20.5] and [RL, 1.6].

4.2. THEOREM. If $f: Q \rightarrow E^n$ is continuous, then $\mathcal{L}(f) \geq \mathcal{C}(f) \geq \mathcal{D}(f)$.

In order to prove that $\mathcal{D}(f) \geq \mathcal{L}(f)$, we first observe that if f and g are Fréchet equivalent maps of Q into E^n , then $\mathcal{D}(f) = \mathcal{D}(g)$. The same is true for the functionals \mathcal{L} and \mathcal{C} , [C, 31.7]. These facts will be very useful since we later employ Morrey's representation theorem. Undoubtedly, it is possible to prove $\mathcal{D}(f) \geq \mathcal{L}(f)$ without resorting to Morrey's theorem, but we prefer this method since we anticipate an application of (24) below in future work.

In this regard recall that a continuous map $f: Q \rightarrow E^n$ is *almost conformal* provided that

- (i) the coordinate functions of f are absolutely continuous in the sense of Tonelli (ACT) on Q , and their partial derivatives are square integrable on Q , and
- (ii) the partial derivatives $D_1 f, D_2 f$ satisfy $D_1 f(x) \cdot D_2 f(x) = 0$ for L_2 almost all $x \in Q$.

The formal differential, df , which is defined as the linear transformation associated with the matrix of partial derivatives of f , exists for L_2 almost all points in Q . Thus, for all such x , $df(x)$ is a linear transformation from E^2 into E^n and it induces a linear transformation on the space of Grassmann 2-vectors. Thus, if α is a 2-vector in E^2 , then $df(x)(\alpha) = df(x, \alpha)$ is a 2-vector in E^n . Later, we will also use the fact that if $\{f_i\}$ is a sequence of mollifiers of f , then

- (i) $f_i \rightarrow f$ uniformly on compact subsets of Q and
- (19) (ii) $\int_Q |df_i - df|^2 dL_2 \rightarrow 0$ where $|df(x)|$ denotes the norm of $df(x)$.

If $u: E^n \rightarrow E^1$ is Lipschitz and $f: Q \rightarrow E^n$ is almost conformal, then it is not difficult to show that $F = u \circ f$ is ACT on Q and $|\nabla F|$ is square integrable on Q . Moreover, the following facts were established in [Z]:

(i) If g is square integrable, then

$$\int_Q |\nabla F| g \, dL_2 = \int_{E^1} \int_{F^{-1}(r)} g(y) \, dH^1(y) \, dL_1(r).$$

- (ii) If W is a component of $\{x : F(x) < r\}$, then for L_1 almost all r , W has finite perimeter. That is, if W is considered as a current, then in fact, it is an integral current.
- (iii) Let $\beta(W)$ be the reduced boundary of W , that is, $\beta(W)$ consists of those points x where $n(x)$, the exterior normal to W , exists at x . Observe that $\beta(W) \subset \text{bdry } W \subset F^{-1}(r)$. Following the proof of [Z, 3.3], we have $H^1[\beta(W) - \delta(W)] = 0$.

4.3. LEMMA. If $f: Q \rightarrow E^n$ is almost conformal and $u: E^n \rightarrow E^1$ is Lipschitz, then for L_1 almost all r , $f^{-1}(y)$ is totally disconnected for H^1 almost all $y \in u^{-1}(r)$.

Proof. In view of the fact that the coordinate functions of f are ACT on Q , it follows that there is a countable set of vertical line segments in Q , Λ_1 , such that Λ_1 is dense in Q and if $\lambda \in \Lambda_1$ then $H^1[f(\lambda)] < \infty$. Likewise, there is a set Λ_2 corresponding to the horizontal direction. Setting $\Lambda = \Lambda_1 \cup \Lambda_2$, it follows that if $N = f(\Lambda)$ then $H^2[N] = 0$. However, [F1, 3.2] or 2.1 implies that $H^1[N \cap u^{-1}(r)] = 0$ for L_1 almost all r . Hence, if $y \notin N \cap u^{-1}(r)$, then $f^{-1}(y)$ has only point components.

4.4. LEMMA. With the same hypotheses as 4.3, $\lambda(f; u, r) \leq \rho(f; u, r)$ for L_1 almost all $r \in E^1$.

Proof. Let α be a continuous function on Q whose values are unit Grassmann 2-vectors in E^2 . Select $r \in E^1$ so that the results of (20) and 4.3 hold, and let W be a component of $\{x : F(x) < r\}$ such that $(\text{closure } W) \cap \text{bdry } Q = 0$. Since the mollifiers, f_i , of f are C^∞ , (19) implies for every C^∞ differential 2-form φ that

$$\begin{aligned} \lim_{i \rightarrow \infty} f_{i\#}(W)(\varphi) &= \lim_{i \rightarrow \infty} \int_W \varphi[f_i(x)] \cdot df_i(x, \alpha(x)) \, dL_2(x) \\ (21) \qquad &= \int_W \varphi[f(x)] \cdot df(x, \alpha(x)) \, dL_2(x). \end{aligned}$$

On the other hand, [F3, 3.4] implies that, for L_1 almost all r , $\lim_{i \rightarrow \infty} f_{i\#}(W)(\varphi) = \mu(V)(\varphi)$ where $V = m_f(W)$. Thus,

$$(22) \qquad \mu(V)(\varphi) = \int_W \varphi[f(x)] \cdot df(x, \alpha(x)) \, dL_2(x).$$

Since $\int_Q |df_i - df|^2 dL_2 \rightarrow 0$ it follows from [(20)(i)] that there exists a subsequence (which will still be denoted as $\{df_i\}$) such that

$$(23) \quad \int_{F^{-1}(r)} |df_i - df| dH^1 \rightarrow 0.$$

From Hölder's inequality and (i) of (20), it is clear that (23) holds for L_1 almost all r . Since $\beta(W) \subset F^{-1}(r)$ for all r under consideration, (19), (iii) of (20), (21), (22), (23), and the Gauss-Green Theorem [F4, 4.5.6] imply that, for L_1 almost all r ,

$$\begin{aligned} \partial\mu(V)(\varphi) &= \lim_{i \rightarrow \infty} \partial f_{i\#}(W)(\varphi) = \lim_{i \rightarrow \infty} f_{i\#}(\partial W)(\varphi) \\ &= \lim_{i \rightarrow \infty} \int_{\beta(W)} \varphi[f_i(x)] \cdot df_i(x, v(x)) dH^1 \\ (24) \quad &= \int_{\beta(W)} \varphi[f(x)] \cdot df(x, v(x)) dH^1(x) \\ &= \int_{\beta(W) \cap \delta(W)} \varphi[f(x)] \cdot df(x, v(x)) dH^1(x) \end{aligned}$$

where $v(x)$ is the unit vector perpendicular to $n(x)$ chosen so that $v(x) \wedge n(x) = \alpha(x)$. Since the partial derivatives of the coordinate functions of f exist L_2 almost everywhere in Q , we apply [F4, 3.1.8] to find disjoint sets $Q = \bigcup_{i=0}^{\infty} A_i$ such that $L_2(A_0) = 0$ and f restricted to A_i , $i > 0$, is Lipschitzian. From [FU, Theorem 3(d)] and (i) of (20), we may assume that $H^1[\delta(W) \cap A_0] = 0$. Thus, [F4, 3.2.20] and (24) yield

$$M[\partial\mu(V)] \leq \int_{E^n} N[f, \delta(W), y] dH^1(y).$$

However, 4.3 allows us to assume that

$$\int_{E^n} N[f, \delta(W), y] dH^1(y) = \int_{E^n} N[l_f, \delta(V), y] dH^1(y)$$

and the lemma follows directly from this.

4.5. THEOREM. *Suppose $f: Q \rightarrow E^n$ has finite Lebesgue area and, in addition, suppose f has the property that for each $y \in E^n$, no component of $f^{-1}(y)$ disconnects Q . Then $\mathcal{C}(f) = \mathcal{D}(f) = \mathcal{L}(f)$.*

Proof. With the conditions imposed on f it follows that the middle space, M_f , is either a 2-cell or a 2-sphere, [R, II.2.91].

In the event that M_f is a 2-cell, Morrey's representation theorem [MO1], [MO2] asserts the existence of an almost conformal map $g: Q \rightarrow E^n$ that is Fréchet equivalent to f . Now $\mathcal{D}(f) = \mathcal{D}(g)$ and $\mathcal{L}(f) = \mathcal{L}(g)$ while 4.2, 4.4, and 3.3 imply that $\mathcal{D}(g) = \mathcal{L}(g)$.

If M_f is a 2-sphere, then f has a Fréchet equivalent $g: Q \rightarrow E^n$ with the property that g is constant on the boundary of Q while f is not constant on any non-degenerate continuum in the interior of Q , [R, II.3.28]. However, this case is

treated essentially the same way as the preceding one and, thus, the proof is complete.

In order to establish Theorem 4.5 for any map $f: Q \rightarrow E^n$ of finite Lebesgue area, it will suffice to prove that \mathcal{D} is cyclically additive because it is known that Lebesgue area possesses this property, [R, V.2.55].

Let C_1, C_2, \dots be the cyclic elements of M_f and let $r_i: M_f \rightarrow C_i$ be the monotone retraction, [R, II.2.40]. The mappings $f_i = l_f \circ r_i \circ m_f$ are the cyclic components of f and they satisfy the hypotheses of Theorem 4.5. Let C_1, C_2, \dots, C_k be any finite number of cyclic elements of M_f . It will be sufficient to show that

$$(25) \quad \mathcal{D}(f) \geq \sum_{i=1}^k \mathcal{D}(f_i)$$

in view of the fact that $\mathcal{D}(f) \leq \sum_{i=1}^{\infty} \mathcal{D}(f_i)$, which follows from the cyclic additivity of \mathcal{L} , 4.2, and 4.5.

Choose $\varepsilon > 0$. From the construction that appears in the proof of Theorem 3.3, it follows that there is a function $u: E^n \rightarrow E^1$ with Lipschitz constant 1 such that, for $i=1, 2, \dots, k$,

$$(26) \quad \int_{-\infty}^{\infty} \lambda(f_i; u, r) dL_1(r) > \mathcal{L}(f_i) - \varepsilon/k.$$

In order to see this we will only consider the case of $k=2$, since the general case is handled in the same way and is no more difficult except for complications in notation. Let ζ_i be the measure associated with f_i as in (15), $i=1, 2$, and let R_i be the Hausdorff 2-rectifiable set that is determined by f_i as in the proof of Theorem 3.3. Choose compact sets $K_1 \subset R_1 - R_2$, $K_2 \subset R_2 - R_1$, and $K_3 \subset R_1 \cap R_2$ such that

$$\begin{aligned} \zeta_1(K_1) &> \zeta_1(R_1 - R_2) - \varepsilon/3, & \zeta_2(K_2) &> \zeta_2(R_2 - R_1) - \varepsilon/3, \\ \zeta_i(K_3) &> \zeta_i(R_1 \cap R_2) - \varepsilon/3, & i &= 1, 2. \end{aligned}$$

Let U_i be open sets that are mutually disjoint that contain K_i , $i=1, 2, 3$. For each $y \in K_i$ and all sufficiently small balls centered at y , (17) holds with f and ζ replaced by f_i and ζ_i , $i=1, 2$. Similarly, at H^2 almost all $y \in K_3$, (17) is satisfied simultaneously for $i=1, 2$ for all sufficiently small balls centered at y . We may assume that all balls considered are contained in some U_i , $i=1, 2, 3$. Thus, we have a Vitali cover of $K = \bigcup_{i=1}^3 K_i$ and, therefore, there exist balls B_1, B_2, \dots such that

$$H^2 \llcorner K \left[E^n - \bigcup_{i=1}^{\infty} B_i \right] = 0.$$

As in the proof of Theorem 3.3, we define a Lipschitz function $u: E^n \rightarrow E^1$ in terms of the B_i . Since ζ_i is absolutely continuous with respect to H^2 , it follows for $i=1, 2$ that

$$\begin{aligned} \int_{-\infty}^{\infty} \lambda(f_i; u, r) dL_1(r) &\geq \sum_{j=1}^{\infty} (1-\varepsilon) \zeta_i(B_j) = (1-\varepsilon) \zeta_i(K_i \cup K_3) \\ &\geq (1-\varepsilon) [\zeta_i(R_i) - 2\varepsilon/3] = (1-\varepsilon) [\mathcal{L}(f_i) - 2\varepsilon/3]. \end{aligned}$$

By redefining ε appropriately, (26) is now established. Therefore, (26), 4.4, and 4.5 imply that

$$(27) \quad \int_{-\infty}^{\infty} \rho(f_i; u, r) dL_1(r) > \mathcal{D}(f_i) - \varepsilon/k, \quad i = 1, 2, \dots, k.$$

Let W be a component of $\{x : u \circ f(x) < r\}$ and let $V = m_f(W)$. Now $r_i(V) = V \cap C_i$ and the boundary of $V \cap C_i$ relative to C_i is contained in $(\text{bdry } V) \cap C_i$. Consequently, if $V_i = V \cap C_i$, $\delta(V_i) \subset \delta(V)$, $i = 1, 2, \dots, k$. Any two cyclic elements intersect at most in one point, [R, II.2.24], and therefore

$$N[l_f, \delta(V), y] \geq \sum_{i=1}^k N[l_f, \delta(V_i), y]$$

for all but finitely many $y \in E^n$. Hence,

$$\rho(f; u, r) \geq \sum_{i=1}^k \rho(f_i; u, r)$$

and (27) now leads to

$$\begin{aligned} \mathcal{D}(f) &\geq \int_{-\infty}^{\infty} \rho(f; u, r) dL_1(r) \\ &\geq \sum_{i=1}^k \int_{-\infty}^{\infty} \rho(f_i; u, r) dL_1(r) > \sum_{i=1}^k \mathcal{D}(f_i) - \varepsilon. \end{aligned}$$

Since ε was arbitrarily chosen, (25) is now established and we have established that if $f: Q \rightarrow E^n$ has finite Lebesgue area, then

$$(28) \quad \mathcal{L}(f) = \mathcal{C}(f) = \mathcal{D}(f).$$

Moreover, by considering $u: E^n \rightarrow E^1$ as the distance function from a fixed hyperplane in E^n , it follows from Theorems 4.6 and 7.16 of [F2] that there is a constant k such that $\mathcal{L}(f) \leq k\mathcal{D}(f)$. Thus, if $\mathcal{D}(f) < \infty$ then so is $\mathcal{L}(f) < \infty^{(2)}$. Therefore 4.2 and (28) yield the following result.

4.6. THEOREM. *If $f: Q \rightarrow E^n$ is continuous, then $\mathcal{L}(f) = \mathcal{C}(f) = \mathcal{D}(f)$.*

The results and techniques employed by Slepian [S1], [S2] will allow the extension of this theorem to maps defined on any 2-dimensional manifold.

BIBLIOGRAPHY

- [C] L. Cesari, *Surface area*, Ann. of Math. Studies, no. 35, Princeton Univ. Press, Princeton, N. J., 1956. MR 17, 596.
 [F1] H. Federer, *Some integral geometric theorems*, Trans. Amer. Math. Soc. 77 (1954), 238–261. MR 16, 163.
 [F2] ———, *On Lebesgue area*, Ann. of Math. (2) 61 (1955), 289–353. MR 16, 683.
 [F3] ———, *Currents and area*, Trans. Amer. Math. Soc. 98 (1961), 204–233. MR 23 #A1006.

(²) The author is indebted to the referee for this observation.

[F4] H. Federer, *Geometric measure theory*, Die Grundlehren der math. Wissenschaften, Band 153, Springer-Verlag, New York, 1969. MR 41 #1976.

[FF] H. Federer and W. H. Fleming, *Normal and integral currents*, Ann. of Math. (2) 72 (1960), 458–520. MR 23 #A588.

[FU] B. Fuglede, *Extremal length and functional completion*, Acta Math. 98 (1957), 171–219. MR 20 #4187.

[FN] R. E. Fullerton, *The Cesari-Cavalieri area of a surface*, Duke Math. J. 30 (1963), 151–160. MR 26 #2581.

[G] R. F. Gariépy, *Current valued measures and Geöcze area*, Ph.D. Thesis, Wayne State University, Detroit, Mich., 1969; see also Trans. Amer. Math. Soc. (to appear).

[M] J. H. Michael, *The convergence of measures on parametric surfaces*, Trans. Amer. Math. Soc. 107 (1963), 140–152. MR 26 #3874.

[MO1] C. B. Morrey, *A class of representations of manifolds*. I, II, Amer. J. Math. 55 (1933), 683–707; *ibid.* 56 (1934), 275–293.

[MO2] ———, *An analytic characterization of surfaces of finite Lebesgue area*. I, II, Amer. J. Math. 57 (1935), 692–702; *ibid.* 58 (1936), 313–322.

[R] T. Radó, *Length and area*, Amer. Math. Soc. Colloq. Publ., vol. 30, Amer. Math. Soc., Providence, R. I., 1948. MR 9, 505.

[RL] R. Rishel, *Area as the integral of lengths of contours*, Trans. Amer. Math. Soc. 97 (1960), 95–119. MR 22 #9574.

[S1] P. Slepian, *Theory of Lebesgue area of continuous maps of 2-manifolds into n -space*, Ann. of Math. (2) 68 (1958), 669–689. MR 20 #5271.

[S2] ———, *On the Lebesgue area of a doubled map*, Pacific J. Math. 8 (1958), 613–620. MR 24 #A2003.

[Z] W. Ziemer, *Some lower bounds for Lebesgue area*, Pacific J. Math. 19 (1966), 381–390. MR 34 #2830.

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, INDIANA 47401

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MICHIGAN 48104