

MAPPINGS FROM 3-MANIFOLDS ONTO 3-MANIFOLDS⁽¹⁾

BY
ALDEN WRIGHT

Abstract. Let f be a compact, boundary preserving mapping from the 3-manifold M^3 onto the 3-manifold N^3 . Let Z_p denote the integers mod a prime p , or, if $p=0$, the integers. (1) If each point inverse of f is connected and strongly 1-acyclic over Z_p , and if M^3 is orientable for $p > 2$, then all but a locally finite collection of point inverses of f are cellular. (2) If the image of the singular set of f is contained in a compact set each component of which is strongly acyclic over Z_p , and if M^3 is orientable for $p \neq 2$, then N^3 can be obtained from M^3 by cutting out of $\text{Int } M^3$ a compact 3-manifold with 2-sphere boundary, and replacing it by a Z_p -homology 3-cell. (3) If the singular set of f is contained in a 0-dimensional set, then all but a locally finite collection of point inverses of f are cellular.

I. Introduction. We suppose throughout the introduction that $f: M^3 \twoheadrightarrow N^3$ is a compact, boundary preserving mapping from the 3-manifold M^3 onto the 3-manifold N^3 (where M^3 and N^3 may or may not have boundary). Let Z_p denote the integers modulo a prime p , or, if $p=0$, the integers.

If $f^{-1}(x)$ is connected and strongly 1-acyclic over Z_p for all $x \in N^3$, and if M^3 is orientable for $p > 2$, then in Corollary 1 it is shown that all but a locally finite collection of point inverses are cellular. This implies that N^3 can be obtained from M^3 by cutting out of $\text{Int } M^3$ a locally finite collection of compact 2-manifolds, each bounded by a 2-sphere, and replacing them by a 3-cell (see Corollary 3). Thus, if M^3 is compact, N^3 is a factor in a connected sum decomposition of M^3 .

Now suppose that the image of the singular set of f is contained in a compact set X each component of which is strongly acyclic over Z_p . If M^3 is orientable for $p \neq 2$, then N^3 can be obtained from M^3 by cutting out of M^3 a finite number of compact 3-manifolds, each bounded by a 2-sphere, and replacing each by a Z_p -homology 3-cell. In particular, if X has a neighborhood which is an irreducible 3-manifold with boundary (or if N^3 is irreducible), then N^3 is a factor in a connected sum decomposition of M^3 . This extends Theorem 1 of Lambert in [9]. In the special case where the image of the singular set is contained in a Cantor set,

Received by the editors November 18, 1970.

AMS 1969 subject classifications. Primary 5542, 5705; Secondary 5460, 5478, 5560.

Key words and phrases. Monotone mapping, 3-manifold, acyclic mapping, decomposition space, strongly acyclic.

⁽¹⁾ Some of these results are contained in the author's Ph.D. thesis written under the direction of Professor D. R. McMillan at the University of Wisconsin. The author would like to thank Professor McMillan for many helpful suggestions. Partially supported by N.S.F. Grant GP-12025.

we can say in addition that all but a finite number of point inverses are cellular. This was previously proved by the author using other techniques.

Lemma 5 restates one of Armentrout's results on approximating cellular maps with homeomorphisms. Using this lemma, we combine the results of Theorems 1 and 3 in Theorem 5. Thus if M^3 is compact and orientable for $p \neq 2$, and if the image of the point inverses of f which are not connected and strongly 1-acyclic over Z_p is contained in a compact set X each component of which is strongly acyclic over Z_p , then N^3 can be obtained from M^3 by cutting out of $\text{Int } M^3$ a finite number of 3-manifolds each bounded by a 2-sphere, and replacing each by a Z_p -homology 3-cell. Theorem 6 combines Theorems 1 and 4 in a similar fashion.

In Theorem 7, we extend a result of McMillan [13] to show that if the image of the singular set of f is contained in a (nonclosed) 0-dimensional set, then all but a locally finite collection of point inverses are cellular.

Let G be a nontrivial abelian group. A compact set $X \subset M$ is *strongly k -acyclic over G* if for each open set $U \subset M$ containing X , there is an open set V such that $X \subset V \subset U$ and such that the inclusion induced homomorphism $i_*: H_k(V; G) \rightarrow H_k(U; G)$ is zero. (If X is connected and strongly k -acyclic over G for $1 \leq k \leq n$, then $X \subset M$ has property $uv^n(G)$ in the sense of [8].) The compact set $X \subset M$ is *strongly acyclic over G* if it is connected and strongly k -acyclic over G for all $k \geq 1$. We refer the reader to [13] (especially Lemma 1) for further facts about strong acyclicity. In particular, for any positive integer k , a compact set X in the interior of a 3-manifold M^3 is strongly k -acyclic over G if and only if each component of X is strongly k -acyclic over G . Also X is strongly acyclic over Z if and only if X is connected and $H^*(X; Z) = 0$ (see [7]).

The compact set $X \subset M$ has property UV^∞ if for each open set $U \subset M$ containing X , there is an open set V such that $X \subset V \subset U$ and such that V is contractible in U . A set X in a 3-manifold M^3 is *cellular* in M^3 if $X = \bigcap_{i=1}^\infty F_i$ where each F_i is a 3-cell, and $F_{i+1} \subset \text{Int } F_i$ for all i .

If σ is a loop in a space M , we will denote its homology class in $H_1(M; G)$ by $[\sigma]$. The symbol Z_p for $p > 0$ will denote the finite cyclic group of order p . The symbol Z_0 will denote the integers.

A manifold will be assumed to be connected and to have no boundary unless otherwise specified. We assume that all manifolds have a piecewise-linear structure. A 3-manifold is *irreducible* if every polyhedral 2-sphere in it bounds a polyhedral 3-cell. If M^3 and N^3 are 3-manifolds, possibly with boundary, the *connected sum* $M^3 \# N^3$ of M^3 and N^3 is obtained by removing the interior of a 3-cell from the interior of each, and then sewing the two manifolds together along the resulting boundary components, using an orientation reversing homeomorphism if M^3 and N^3 are oriented.

A *map* or *mapping* is a continuous function. A *monotone map* is a map all of whose point inverses are connected. A map $f: M \rightarrow N$ is *compact* (proper) if, for any compact set K in N , $f^{-1}(K)$ is compact. If $f: M \rightarrow N$ is a compact monotone

map, then the point inverses of M form a monotone upper semicontinuous decomposition of M whose associated decomposition space is homeomorphic to N . Conversely, if G is a monotone upper semicontinuous decomposition of M , the projection map $p: M \twoheadrightarrow M/G$ is a compact monotone map.

Let $\{X_\alpha\}_{\alpha \in A}$ be a collection of compact subsets of a space M . Then $\{X_\alpha\}_{\alpha \in A}$ is a *locally finite collection* if for $y \in M$, y has a neighborhood U which intersects only a finite number of elements of the collection.

II. Maps all of whose point inverses are strongly acyclic.

LEMMA 1. *If X is a compact connected subset of a space M and if X is strongly k -acyclic over Z in M for $1 \leq k \leq n$, then X is strongly k -acyclic over Z_p in M for $1 \leq k \leq n$ and for any prime $p > 1$.*

Proof. Let W and V be chosen so that $X \subset W \subset V \subset U$ and so that the inclusion induced homomorphisms $i_*: H_k(V; Z) \rightarrow H_k(U; Z)$ and $j_*: H_k(W; Z) \rightarrow H_k(V; Z)$ are zero for $1 \leq k \leq n$. Consider the following commutative diagram:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H_k(W; Z) \otimes Z_p & \longrightarrow & H_k(W; Z_p) & \longrightarrow & \text{Tor}_1(H_{k-1}(W; Z), Z_p) \longrightarrow 0 \\
 & & \downarrow i_* \otimes \text{id} & & \downarrow i'_* & & \downarrow \\
 0 & \longrightarrow & H_k(V; Z) \otimes Z_p & \longrightarrow & H_k(V; Z_p) & \longrightarrow & \text{Tor}_1(H_{k-1}(V; Z), Z_p) \longrightarrow 0 \\
 & & \downarrow j_* \otimes \text{id} & & \downarrow j'_* & & \downarrow \\
 0 & \longrightarrow & H_k(U; Z) \otimes Z_p & \longrightarrow & H_k(U; Z_p) & \longrightarrow & \text{Tor}_1(H_{k-1}(U; Z), Z_p) \longrightarrow 0
 \end{array}$$

The horizontal rows, which are exact, are from the universal coefficient theorem. By our choice of W and V , the outer vertical maps are zero. Using a diagram chasing argument, we see that $j'_* i'_*$ is the zero homomorphism.

LEMMA 2. *Let M^3 and N^3 be 3-manifolds, and let $f: M^3 \twoheadrightarrow N^3$ be a compact, monotone, onto map. Let p be 0 or a prime, and suppose M^3 is orientable if $p \neq 2$. If $f^{-1}(y)$ is strongly 1-acyclic over Z_p for every $y \in N^3$, then each $f^{-1}(y)$ is strongly acyclic over Z_p in M^3 .*

Proof. By Alexander duality and Theorem 3 of [8] we see that $H^k(f^{-1}(y); Z_p) = 0$ for $k \geq 2$. Then the continuity of H^* and the universal coefficient theorem for cohomology show that $f^{-1}(y)$ is strongly acyclic over Z_p for all $y \in N^3$. (For more details, see Theorems 4.4 and 3.2 of [7].)

LEMMA 3. *Let M^3 and N^3 be 3-manifolds, and let $f: M^3 \twoheadrightarrow N^3$ be a compact, monotone, onto map such that $f^{-1}(y)$ is strongly 1-acyclic over G for each $y \in N^3$. If $H_1(N^3; G) = 0$, then $H_1(M^3; G) = 0$.*

The proof of Lemma 3 is similar to the proof of Theorem 2.1 of [15].

If M^n and N^n are n -manifolds with boundary, a map $f: M^n \rightarrow N^n$ is said to be *boundary preserving* if $f|_{\text{Bd } M^n}$ is a homeomorphism of $\text{Bd } M^n$ onto $\text{Bd } N^n$, and if $f^{-1}(\text{Bd } N^n) = \text{Bd } M^n$. A 2-manifold with boundary S is *properly embedded* in a 3-manifold with boundary M^3 if $S \cap \text{Bd } M^3 = \text{Bd } S$.

A Z_p -homology (homotopy) 3-cell is a compact Z_p -acyclic (contractible) 3-manifold with boundary. A cube-with-handles is obtained by adding orientable 1-handles to a 3-cell. We define a Z_p -homology (homotopy) cube-with-handles similarly. We will say that a set X is the intersection of a decreasing sequence of (Z_p -homology, homotopy) cubes-with-handles if $X = \bigcap_{i=1}^{\infty} K_j^3$ where each K_j^3 is a (Z_p -homology, homotopy) cube-with-handles and $K_{j+1}^3 \subset \text{Int } K_j^3$.

THEOREM 1. *Let p denote 0 or a prime, and let M^3 and N^3 be compact 3-manifolds, possibly with boundary, where M^3 is orientable if $p > 2$. Let $f: M^3 \rightarrow N^3$ be a monotone, onto, boundary preserving map. Let U be an open subset of N^3 . If $f^{-1}(x)$ is strongly 1-acyclic over Z_p for all $x \in U$, then $\{x \in U : f^{-1}(x) \text{ is not cellular}\}$ is a finite set.*

REMARK. This theorem was first proved for $p=0, 2$ in [16]. It has since been generalized by D. R. McMillan in [13].

Proof. The case where $p=0$ reduces to the case where $p=2$ by Lemma 1. By the proofs of Theorems 1 and 2 of [11] and by Kneser's Theorem [6] it is sufficient to prove that $\{x \in U : f^{-1}(x) \text{ is not } UV^{\infty}\}$ is finite.

We can apply Lemma 2 to see that $f^{-1}(x)$ is strongly acyclic over Z_p for each $x \in U$. By Theorem 2 of [12], $f^{-1}(x)$ is the intersection of a decreasing sequence of Z_p -homology cubes-with-handles.

Let q be the rank (i.e. the minimum number of generators) of $\pi_1(M^3)$. By a corollary to the Grushko-Neumann Theorem (p. 192 of [10]), there are at most q disjoint Z_p -homology 3-cells in M^3 which are not homotopy 3-cells. Thus there are at most q points in U whose inverse images are not the intersection of a decreasing sequence of homotopy cubes-with-handles.

Let $x \in U$, where $f^{-1}(x)$ is the intersection of a decreasing sequence of homotopy cubes-with-handles. We will complete the proof by showing that $f^{-1}(x)$ is UV^{∞} . Let U' be an open set in M^3 containing $f^{-1}(x)$. There is a homotopy cube-with-handles H^3 such that

$$f^{-1}(x) \subset \text{Int } H^3 \subset H^3 \subset U' \cap f^{-1}(U).$$

Let W be an open 3-cell in U such that $x \in W$ and $f^{-1}(W) \subset \text{Int } H^3$. Define inductively $G_0, G_1, G_2, \dots$ by letting $G_0 = \pi_1(f^{-1}(W))$, and by letting

$$G_i = G_{i-1}(X_1 X_2 X_1^{-1} X_2^{-1} X_3^p).$$

(See p. 74 of [10] for notation.) In other words, G_i is the subgroup of G_{i-1} generated by all elements of the form $uvu^{-1}v^{-1}\tau^p$ where $u, v, \tau \in G_{i-1}$. Let $F_0, F_1, F_2, \dots$ be the corresponding subgroups of $\pi_1(H^3)$.

The subgroup G_1 certainly contains the commutator subgroup of G_0 . The image of G_1 in $H_1(f^{-1}(W); Z)$ is $p \cdot H_1(f^{-1}(W); Z)$. Thus

$$\pi_1(f^{-1}(W))/G_1 \cong H_1(f^{-1}(W); Z)/p \cdot H_1(f^{-1}(W); Z) \cong H_1(f^{-1}(W); Z_p).$$

Let $\delta \in \pi_1(f^{-1}(W))$. Since $H_1(f^{-1}(W); Z_p) = 0$ (by Lemma 3), $\delta \in G_1$. Thus δ is a product of elements of the form $uvu^{-1}v^{-1}\tau^p$ where $u, v, \tau \in G_0$. By applying the same argument to u, v , and τ , we see that $u, v, \tau \in G_1$. Thus $\delta \in G_2$. By repeating this argument, $\delta \in \bigcap_{i=0}^{\infty} G_i$. By Corollary 2.12 on p. 109 of [10], $\bigcap_{i=1}^{\infty} F_i = 1$. Thus $\delta = 1$ in $\pi_1(H^3)$, and $f^{-1}(x)$ is UV^{∞} .

COROLLARY 1. *Let M^3 and N^3 be 3-manifolds, possibly with boundary, and let $f: M^3 \twoheadrightarrow N^3$ be a compact, monotone, boundary preserving, onto map. Let p denote 0 or a prime, and suppose that M^3 is orientable if $p > 2$. If $f^{-1}(x)$ is strongly 1-acyclic over Z_p in M^3 for all $x \in U$, then $\{x \in U : f^{-1}(x) \text{ is not cellular}\}$ is a locally finite set in N^3 .*

III. Maps where the image of the singular set lies in a strongly acyclic set. We state below a slightly strengthened version of Theorem 2 of [13]: here we assume that M^3 is orientable only if $p > 2$, and thus the 1-handles which are attached to $\text{Bd } Q_i$ to obtain H_i may be attached in a nonorientable fashion. (See the statement of Theorem 2 for the definition of Q_i and H_i .) The only additional difficulty in the proof is when we have $S_i \subset \text{Bd } Z_i^*$ and $S_k \subset \text{Bd } Z_k^*$ topologically parallel. (See p. 133 of [12].) As before, each loop in S_i Z_p -bounds in Z_i^* , and the same argument shows that S_i is a 2-sphere if S_i is not homeomorphic to a projective plane. But if S_i is a projective plane, it must contain an orientation-reversing simple closed curve since S_i is two-sided. This contradicts the fact that every simple closed curve in S_i Z_p -bounds in Z_i^* , since $p = 0, 2$.

THEOREM 2. *Let p denote 0 or a prime. Let X be a compact, proper subset of $\text{Int } M^3$, where M^3 is a 3-manifold, possibly with boundary. Suppose M^3 is orientable if $p > 2$, and suppose that X has the following property relative to M^3 and p . For each open set $U \subset M^3$ with $X \subset U$, there is an open set V , $X \subset V \subset U$, such that, under inclusion, $H_1(V - X; Z_p) \rightarrow H_1(U; Z_p)$ is zero. Then $X = \bigcap_{i=1}^{\infty} H_i$, where H_i is a compact polyhedron in M^3 , each component of H_i is a 3-manifold with nonempty boundary, $H_{i+1} \subset \text{Int } H_i$ and each H_i has the following structure: it is obtained from a compact polyhedron Q_i , each component of which is a 3-manifold whose boundary consists entirely of 2-spheres, by adding to $\text{Bd } Q_i$ a finite number of (solid, possibly nonorientable) 1-handles.*

Let $f: M \rightarrow N$ be a map. Then let $S_f = \{x \in M : f^{-1}f(x) \text{ is nondegenerate}\}$.

THEOREM 3. *Let p denote 0 or a prime. Let M^3 and N^3 be piecewise-linear 3-manifolds, possibly with boundary, where M^3 is orientable if $p \neq 2$. Let X be a compact subset of $\text{Int } N^3$ such that each component of X is strongly acyclic over Z_p . Let $f: M^3 \twoheadrightarrow N^3$ be a compact, boundary preserving map with $f(S_f) \subset X$. Then N^3 can*

be obtained from M^3 by cutting out of $\text{Int } M^3$ a finite number of polyhedral 3-manifolds which are each bounded by a 2-sphere, and replacing each by a polyhedral Z_p -homology 3-cell.

Proof. By Theorem 2 of [12], X is the intersection of a decreasing sequence of Z_p -homology cubes-with-handles. Thus we can assume that N^3 is a Z_p -homology cube-with-handles, and that each two-sided surface in $\text{Int } N^3$ separates N^3 .

The first half of the proof will be to show that $f^{-1}(X)$ has the following property in $\text{Int } M^3$: for each open set $U \subset \text{Int } M^3$ with $f^{-1}(X) \subset U$, there is an open set V , with $f^{-1}(X) \subset V \subset U$, such that, under inclusion, $H_1(V - f^{-1}(X); Z_p) \rightarrow H_1(U; Z_p)$ is zero.

Let U be an open set in $\text{Int } M^3$ with $f^{-1}(X) \subset U$. Since $\text{Cl}(S_f) \subset U$, $f(U)$ is open. Let Z^3 be a compact polyhedron in $f(U)$ such that each component of Z^3 is a 3-manifold with boundary, and such that $X \subset \text{Int } Z^3$. Since X is strongly 1-acyclic over Z_p , there is an open set W containing X such that, under inclusion

$$H_1(W - X; Z_p) \rightarrow H_1(Z^3; Z_p)$$

is zero.

Let $V = f^{-1}(W)$, and let $[\sigma] \in H_1(V - f^{-1}(X); Z_p)$ where we can assume that σ is a finite, pairwise disjoint collection of (oriented, if $p \neq 2$) simple closed curves such that $f(\sigma)$ is polyhedral in Z^3 . Let F^3 be a regular neighborhood of $f(\sigma)$ in $(\text{Int } Z^3) - X$. We can triangulate Z^3 so that F^3 and $f(\sigma)$ are subcomplexes of the triangulation. Then the homeomorphism $f^{-1}|(\text{Bd } Z^3 \cup F^3)$ induces a triangulation of $f^{-1}(\text{Bd } Z^3 \cup F^3)$. Since each of the finite number of components of $f^{-1}(Z^3)$ is a 3-manifold with boundary, by Theorem 5 of [2] there is a triangulation of $f^{-1}(Z^3)$ which is compatible with the above triangulation of $f^{-1}(\text{Bd } Z^3 \cup F^3)$. Using the relative simplicial approximation theorem, there is a piecewise-linear, nondegenerate map g from $f^{-1}(Z^3)$ onto Z^3 such that

$$\begin{aligned} g|f^{-1}(\text{Bd } Z^3 \cup F^3) &= f|f^{-1}(\text{Bd } Z^3 \cup F^3), \\ g^{-1}(\text{Bd } Z^3 \cup F^3) &= f^{-1}(\text{Bd } Z^3 \cup F^3). \end{aligned}$$

By subdividing we can assume that g is simplicial.

At this point we divide the remainder of the first half of the proof into three cases: Case 1 ($p=0$), Case 2 ($p=2$), and Case 3 ($p>2$).

Case 1 ($p=0$). Since $f(\sigma) \subset W - X$, $[f(\sigma)] = 0$ in $H_1(Z^3; Z)$. Thus $f(\sigma)$ must bound a 2-complex L^2 in Z^3 where each component of L^2 is an orientable, two-sided 2-manifold with boundary. We can adjust L^2 slightly so that it is in general position mod $f(\sigma)$ with respect to our last triangulation of Z^3 . Then $g^{-1}(L^2)$ will be a 2-complex in $f^{-1}(Z^3) \subset U$, where each component of $g^{-1}(L^2)$ is a two-sided 2-manifold with boundary. Thus, since M^3 is orientable, each component of $g^{-1}(L^2)$ is orientable. Since σ bounds $g^{-1}(L^2)$, $[\sigma] = 0$ in $H_1(U; Z)$, and the inclusion-induced homomorphism $H_1(V - f^{-1}(X); Z) \rightarrow H_1(U; Z)$ is trivial.

Case 2 ($p=2$). The proof is essentially the same as Case 1, except that L^2 and $g^{-1}(L^2)$ may not be orientable.

Case 3 ($p>2$). Note that

$$H_1(Z^3; Z)/G \cong H_1(Z^3; Z) \otimes Z_p \cong H_1(Z^3; Z_p)$$

where G is the subgroup of $H_1(Z^3; Z)$ generated by elements of the form $p[\gamma]$ where $[\gamma] \in H_1(Z^3; Z)$. Since $[f(\sigma)] = 0$ in $H_1(Z^3; Z_p)$, there is a 1-cycle $[\tau] \in H_1(Z^3; Z)$ so that $[f(\sigma)] = p[\tau]$ in $H_1(Z^3; Z)$. We can assume that τ is a finite, pairwise disjoint collection of polyhedral, oriented, simple closed curves which are in general position with respect to our last triangulation of Z^3 . Then $g^{-1}(\tau)$ is a finite, pairwise disjoint collection of simple closed curves in $f^{-1}(Z^3)$. We can find a regular neighborhood T^3 of τ so close to τ that $g^{-1}(T^3)$ is a regular neighborhood of $g^{-1}(\tau)$. We can find a 1-cycle $[\delta] \in H_1(\text{Bd } T^3; Z)$ so that $[f(\sigma)] = [\delta]$ in $H_1(Z^3 - \text{Int } T^3; Z)$. We can assume that δ is a finite collection of mutually exclusive, oriented, simple closed curves on $\text{Bd } T^3$. Then there is a 2-complex $L^2 \subset Z^3 - \text{Int } T^3$ where each component of L^2 is a two-sided, orientable, 2-manifold, and where $\text{Bd } L^2 = f(\sigma) \cup \delta$ (homologically $f(\sigma) - \delta$). We can assume that L^2 is in general position mod $f(\sigma)$ with respect to our last triangulation of Z^3 . Then $g^{-1}(L^2)$ will be a 2-complex where each component of $g^{-1}(L^2)$ is a two-sided 2-manifold with boundary. Thus $g^{-1}(L^2)$ is orientable.

Since L^2 is two-sided in Z^3 , δ is two-sided in $\text{Bd } T^3$. Thus $g^{-1}(\delta)$ is two-sided in $g^{-1}(\text{Bd } T^3)$, and using this two-sidedness, we can induce an orientation of $g^{-1}(\delta)$ which is consistent with that on $g^{-1}(L^2)$. Thus $[g^{-1}(\delta)] = [\sigma]$ in $H_1(f^{-1}(Z^3); Z)$.

Let α be a meridional curve on $\text{Bd } T^3$ which is in general position with respect to δ . Then α will intersect δ algebraically $\pm p$ times. Since the two-sidedness of δ is preserved by g^{-1} , each component of $g^{-1}(\alpha)$ which is a meridional curve must intersect $g^{-1}(\delta)$ algebraically $\pm p$ times. Thus, $[g^{-1}(\delta)] = p[g^{-1}(\tau)]$ in $H_1(T^3; Z)$.

Therefore, $[\sigma] = p[g^{-1}(\tau)]$ in $H_1(Z^3; Z)$, and the inclusion-induced homomorphism $H_1(V - X; Z_p) \rightarrow H_1(U; Z_p)$ is trivial. This completes Case 3.

By Theorem 2, we can find a compact polyhedron H_0^3 , where each component of H_0^3 is a 3-manifold with nonempty boundary, and where H_0^3 has the following structure: it is obtained from a compact polyhedron Q_0^3 , each component of which is a 3-manifold whose boundary consists entirely of 2-spheres, by adding to $\text{Bd } Q_0^3$ a finite number of (solid, possibly nonorientable) 1-handles.

We can also assume that each 1-handle is attached to only one boundary component of $\text{Bd } Q_0^3$ since we can add 1-handles to $\text{Bd } Q_0^3$ which join different components of $\text{Bd } Q_0^3$ without destroying the property that $\text{Bd } Q_0^3$ consists entirely of 2-spheres.

We claim that each component of $\text{Bd } Q_0^3$ separates M^3 . For suppose that S_0 is a component of $\text{Bd } Q_0^3$ that does not separate M^3 . Then there is a polyhedral simple closed curve J which intersects S_0 at exactly one point which is a piercing point. It is easy to see that we can choose J so that it does not intersect any of the 1-handles

which are added to Q_0^3 to obtain H_0^3 . Let S_1 be the component of $\text{Bd } H_0^3$ which is obtained from S_0 by adding handles. Then J intersects S_1 only in the same piercing point. Since $f^{-1}|f(\text{Bd } H_0^3)$ is a homeomorphism, $f(J)$ is a loop in N^3 which intersects $f(S_1)$ in exactly one piercing point. Thus $f(S_1)$ does not separate N^3 . But $f(S_1)$ is a 2-sided surface in N^3 , so $f(S_1)$ must separate N^3 . This is a contradiction, so S_0 does separate M^3 .

Let Q^3 be the closure of the "inside" complementary domains of the "outermost" boundary components of Q_0^3 . (Here, "inside" and "outermost" are relative to $\text{Bd } M^3$, which is connected.) Thus we have "filled in the holes" in Q_0^3 to obtain Q^3 , and each component of Q^3 has connected boundary. We define H^3 to be Q^3 union the 1-handles of $H_0^3 - Q_0^3$ which are not already contained in Q^3 .

There are properly embedded polyhedral disks $B_1^2, \dots, B_r^2$ in H^3 such that the 1-handles which are added to Q^3 to obtain H^3 are regular neighborhoods of $B_1^2, \dots, B_r^2$ in H^3 . Let these 1-handles be $N(B_1^2), \dots, N(B_r^2)$. Since $S_r \subset f^{-1}(X) \subset \text{Int } H^3$, each component of $f(H^3)$ is a 3-manifold with boundary in $\text{Int } N^3$. Each B_i^2 is mapped properly into $f(H^3)$ by f , and furthermore, $f|B_i^2$ has no singularities near $\text{Bd } B_i^2$. So by Dehn's Lemma, there exist nonsingular properly embedded polyhedral disks $D_1^2, \dots, D_r^2$ in $f(H^3)$ with $\text{Bd } D_i^2 = f(\text{Bd } B_i^2)$. By a cutting and pasting argument, we can choose $D_1^2, \dots, D_r^2$ to be disjoint. We can also find disjoint regular neighborhoods $N(D_1^2), \dots, N(D_r^2)$ of $D_1^2, \dots, D_r^2$ in $f(H^3)$ so that

$$f(N(B_i^2) \cap \text{Bd } H^3) = N(D_i^2) \cap \text{Bd } f(H^3).$$

For each i , there is a homeomorphism $h_i: N(B_i^2) \rightarrow N(D_i^2)$ such that

$$h_i|(\text{Bd } H^3 \cap N(B_i^2)) = f|(\text{Bd } H^3 \cap N(B_i^2)).$$

We define a homeomorphism

$$h: M^3 - \text{Int } Q^3 \rightarrow (N^3 - \text{Int } f(H^3)) \cup \left(\bigcup_{i=1}^r N(D_i^2) \right)$$

by $h|(M^3 - \text{Int } H^3) = f|(M^3 - \text{Int } H^3)$, and by $h|N(B_i^2) = h_i$ for each $i = 1, \dots, r$.

Then $h(\text{Bd } Q^3)$ is a finite disjoint collection of 2-spheres in N^3 each of which bounds a Z_p -homology 3-cell. Furthermore, these homology 3-cells are disjoint since each component of $h(\text{Bd } Q^3)$ is outermost in the sense that it can be joined to $\text{Bd } N^3$ with an arc which misses $h(\text{Bd } Q^3)$ except at one end point.

Let $K_1^3, \dots, K_m^3$ be these homology 3-cells, and let $Q_1^3, \dots, Q_m^3$ be the corresponding components of Q^3 so that $h^{-1}(\text{Bd } K_i^3) = \text{Bd } Q_i^3$. Each Q_i^3 is a 3-manifold with 2-sphere boundary. Then h is a homeomorphism from $M^3 - (\bigcup_{i=1}^m Q_i^3)$ onto $N^3 - (\bigcup_{i=1}^m K_i^3)$. Thus we obtain N^3 from M^3 by cutting out the Q_i^3 's and replacing each with the corresponding K_i^3 .

REMARK. If we define $*Q_i^3$ to be the closed 3-manifold obtained from Q_i^3 by sewing a 3-cell onto $\text{Bd } Q_i^3$, and if we define $*K_i^3$ to be the closed 3-manifold obtained from K_i^3 in the same way, then

$$M^3 \# *K_1^3 \# \dots \# *K_m^3 \cong N^3 \# *Q_1^3 \# \dots \# *Q_m^3.$$

We should also note that we have shown that for any open set U in M^3 which contains X , then $f^{-1}(X)$ has a polyhedral neighborhood $H^3 \subset U$ where each component of H^3 is formed by adding 1-handles to a 3-manifold with 2-sphere boundary. Furthermore, we have shown that these 1-handles are attached in an orientable fashion to the 2-sphere boundary.

COROLLARY 2. *Let M^3 and N^3 be compact 3-manifolds, possibly with boundary. Let X be a compact proper set in $\text{Int } N^3$ with the following property: For each open set $U \subset \text{Int } N^3$ with $X \subset U$, there is an open set V , $X \subset V \subset U$, such that under inclusion $H_1(V - X; \mathbb{Z}_p) \rightarrow H_1(U; \mathbb{Z}_p)$ is zero. Suppose also that X has a polyhedral neighborhood each component of which is an orientable, irreducible 3-manifold with boundary. If there is a boundary preserving map f from M^3 onto N^3 such that $f(S_f) \subset X$, then M^3 can be obtained from N^3 by removing the interiors of a finite number of 3-manifolds each of which is bounded by a 2-sphere, and by replacing each by a 3-cell.*

Proof. By using Theorem 2 and the fact that X has a polyhedral neighborhood each component of which is an irreducible 3-manifold with boundary, we see that X has a polyhedral neighborhood each component of which is a cube-with-handles. Thus we can assume that N^3 is a cube-with-handles. The remainder of the proof of Theorem 3 now goes through with the weaker hypothesis on X .

THEOREM 4. *Let M^3 and N^3 be 3-manifolds, possibly with boundary, and let $f: M^3 \rightarrow N^3$ be an onto, compact, boundary preserving mapping from M^3 onto N^3 such that $f(S_f) \subset X$ where X is a closed 0-dimensional set in N^3 . Then f is monotone, and $\{x \in N^3 : f^{-1}(x) \text{ is not cellular in } M^3\}$ is a locally finite subset of N^3 .*

Proof. Let $x \in X$, and let U be an arbitrarily small open 3-cell containing x . Then there is a polyhedral 3-manifold with boundary K^3 so that $x \in \text{Int } K^3 \subset K^3 \subset U$ and so that $\text{Bd } K^3 \cap X = \emptyset$. In fact, using Theorem 2 of [12] and the fact that U is irreducible, we can see that K^3 can be chosen to be a cube-with-handles. Then $f^{-1}(K^3)$ is a connected neighborhood of $f^{-1}(x)$ which can be chosen "arbitrarily close" to $f^{-1}(x)$. Thus f is monotone.

We can cover X with the interiors of a locally finite collection of mutually exclusive collection of cubes-with-handles. Thus, in order to prove the theorem, it suffices to consider the case where N^3 is a cube-with-handles, and where M^3 is a compact 3-manifold with connected boundary. In this case, we will prove that all but a finite number of point inverses of f are cellular.

The set X is strongly 1-acyclic over $\mathbb{Z}_2$ in N^3 , and thus by the remark following the proof of Theorem 3, we have $f^{-1}(X) = \bigcap_{i=1}^{\infty} H_i^3$, where H_i^3 is a 3-manifold with connected boundary, and where $H_i^3 \subset \text{Int } H_{i-1}^3$. We can assume that H_i^3 is obtained from a compact polyhedron Q_i^3 where each component of Q_i^3 is a 3-manifold with 2-sphere boundary, by adding to $\text{Bd } Q_i^3$ a finite number of (orientable, solid) 1-handles. We also have that each 1-cycle in $\text{Bd } H_i^3$ bounds in $\text{Int } H_{i-1}^3$. We have assumed that M^3 is compact and that $H_1(M^3; \mathbb{Z}_2)$ is finitely generated;

so it is easy to show that there is an integer N so that there are not more than N disjoint 3-manifolds with 2-sphere boundary and nontrivial Z_2 -homology in $\text{Int } M^3$. Therefore, all but at most N components of $f^{-1}(X)$ are the intersection of a decreasing sequence of Z_2 -homology cubes-with-handles.

If Z_i^3 is a Z_2 -homology cube-with-handles, the inclusion-induced homomorphism $H_1(\text{Bd } Z_i^3; Z_2) \rightarrow H_1(Z_i^3; Z_2)$ is onto. Thus, if $Z_i^3 \subset \text{Int } Z_{i-1}^3$ where Z_{i-1}^3 is another Z_2 -homology cube-with-handles, and if each 1-cycle in $\text{Bd } Z_i^3$ Z_2 -bounds in $\text{Int } Z_{i-1}^3$, then the inclusion-induced homomorphism $H_1(Z_i^3; Z_2) \rightarrow H_1(Z_{i-1}^3; Z_2)$ is trivial. Therefore, each component of $f^{-1}(X)$ which is the intersection of Z_2 -homology cubes-with-handles must be strongly 1-acyclic over Z_2 . This shows that at most a finite number of point inverses of f are not strongly 1-acyclic over Z_2 .

We can now apply Theorem 1 which implies that only a finite number of the strongly 1-acyclic over Z_2 point inverses of f are not cellular.

IV. Maps almost all of whose point inverses are strongly 1-acyclic over Z_p .

LEMMA 4. Let $f: M \twoheadrightarrow N$ be a compact map from a metric space M onto a metric space N . Let X be a closed set in N . Let G be a decomposition of M defined by

$$G = \{f^{-1}(y) : y \in X\} \cup \{x \in M : f(x) \notin X\}.$$

Let $Q = M/G$ and let $\pi: M \rightarrow Q = M/G$ be the projection map for the decomposition G . Let $p: Q \rightarrow N$ be defined so as to make the following diagram commute:

$$\begin{array}{ccc} M & \xrightarrow{\pi} & Q \\ & \searrow f & \swarrow p \\ & N & \end{array}$$

Then

- (1) G is upper semicontinuous and hence π is continuous and compact.
- (2) The decomposition $\{p^{-1}(y) : y \in N\}$ is upper semicontinuous and hence p is continuous and compact.

Proof. Lemma 4 follows from the fact that $\{f^{-1}(y) : y \in N\}$ is an upper semicontinuous decomposition of M .

LEMMA 5. Let $p: Q \twoheadrightarrow N^3$ be a compact, monotone map from a metric space Q onto a 3-manifold N^3 , possibly with boundary. Let X be a closed set in N^3 containing $\text{Bd } N^3$. Suppose that $p|_{p^{-1}(X)}$ is a homeomorphism, and that $W = Q - p^{-1}(X)$ is an open 3-manifold. If $p^{-1}(x)$ is cellular for all $x \in N^3 - X$, then there is a homeomorphism $h: N^3 \twoheadrightarrow Q$ such that $h|_X = p^{-1}|_X$.

The proof of Lemma 9 is the same as the proof of Theorem 1 of [1].

Suppose $f: M^3 \twoheadrightarrow N^3$ is a mapping. We let $A_f^p = \{x \in M^3 : f^{-1}f(x) \text{ is either not connected or is not strongly 1-acyclic over } Z_p\}$.

THEOREM 5. *Let p denote 0 or a prime, and let M^3 and N^3 be compact 3-manifolds, possibly with boundary, where M^3 is orientable if $p \neq 2$. Let Y be a compact set in $\text{Int } N^3$ each component of which is strongly acyclic over Z_p . Let $f: M^3 \twoheadrightarrow N^3$ be an onto, boundary preserving map such that $f(A_p^?) \subset Y$. Then N^3 can be obtained from M^3 by cutting out of M^3 a finite number of polyhedral 3-manifolds, each bounded by a 2-sphere, and replacing each by a Z_p -homology 3-cell.*

Proof. By Theorem 1 there are only a finite number of points $x_1, x_2, \dots, x_n$ in $N^3 - Y$ whose inverses under f are not cellular in M^3 . Let

$$X = Y \cup \{x_1, x_2, \dots, x_n\} \cup \text{Bd } N^3.$$

We use this X to define $Q, \pi: M^3 \twoheadrightarrow Q$, and $p: Q \twoheadrightarrow N^3$ as in Lemma 4. Since $\pi|(M^3 - f^{-1}(X))$ is a homeomorphism from $M^3 - f^{-1}(X)$ onto $W = Q - p^{-1}(X)$, W is an open 3-manifold. And since $p|p^{-1}(X)$ is one-to-one and continuous, $p|p^{-1}(X)$ is a homeomorphism. Therefore, by Lemma 5, there is a homeomorphism $h: N^3 \twoheadrightarrow Q$. In particular, Q is a 3-manifold Q^3 . Let

$$X' = Y \cup \{x_1, \dots, x_n\}.$$

Then $\pi(S_\pi) \subset p^{-1}(X') = h(X')$, and X' is strongly acyclic over Z_p , so the map π satisfies the hypotheses of Theorem 3.

THEOREM 6. *Let p denote 0 or a prime, and let M^3 and N^3 be 3-manifolds, possibly with boundary, where M^3 is orientable if $p > 2$. Let Y be a closed 0-dimensional set in $\text{Int } N^3$, and let $f: M^3 \twoheadrightarrow N^3$ be an onto, compact, boundary preserving map such that $f(A_p^?) \subset Y$. Then $\{x \in N^3 : f^{-1}(x) \text{ is not cellular in } M^3\}$ is a locally finite subset of N^3 .*

Proof. By Corollary 1, the set $\{x \in N^3 - Y : f^{-1}(x) \text{ is not cellular in } M^3\}$ is a locally finite subset of N^3 .

Let

$$X = Y \cup \text{Bd } N^3 \cup \{x \in N^3 - Y : f^{-1}(x) \text{ is not cellular}\}.$$

Let $Q, \pi: M^3 \twoheadrightarrow Q, p: Q \twoheadrightarrow N^3$, and $h: N^3 \twoheadrightarrow Q$ be defined as in Lemmas 4 and 5. Let

$$X' = Y \cup \{x \in N^3 - Y : f^{-1}(x) \text{ is not cellular}\}.$$

Then $\pi(S_\pi) \subset p^{-1}(X') = h(X')$, and thus $\pi(S_\pi)$ is contained in a closed 0-dimensional set in Q . Theorem 4 can be applied to the map $\pi: M^3 \twoheadrightarrow Q^3$ to say that

$$\{y \in Q^3 : \pi^{-1}(y) \text{ is not cellular in } M^3\}$$

is a locally finite subset of Q^3 . The image under p (or h^{-1}) of this set is

$$\{x \in N^3 : f^{-1}(x) \text{ is not cellular in } M^3\}$$

which must then be a locally finite subset of N^3 .

V. Further applications. The following lemma is a slight generalization of Lemma 5 of [13]. While the proof of Lemma 5 of [13] suffices to prove our Lemma 6, a proof is included here for completeness and since part of the proof will be needed to prove Theorem 7.

LEMMA 6. *Let M^3 and N^3 be 3-manifolds. Let $f: M^3 \rightarrow N^3$ be a compact, monotone mapping so that $f(S_f)$ is 0-dimensional. Let $x \in N^3$. If there is an open set U containing $f^{-1}(x)$ so that the inclusion-induced homomorphism from $H_1(U; Z)$ into $H_1(M^3; Z)$ is trivial, then $f^{-1}(x)$ is strongly 1-acyclic over Z .*

Proof. Let B^3 be an open 3-cell in N^3 with compact closure so that $x \in B^3$ and $W = f^{-1}(B^3)$ is contained in U . Let $K_1, K_2, K_3, \dots$ be a locally finite collection of compact sets in W so that $\bigcup_{i=1}^{\infty} K_i = W$ and each K_i is contained in an open 3-cell $B_i^3 \subset W$. Let

$$\varepsilon_i = \inf \{ \rho(x, y) : x \in K_i \text{ and } y \in W - B_i^3 \}$$

where ρ is a metric on M^3 . Let

$$C_i = \{ x \in N^3 : \text{diam}(f^{-1}(x)) \geq \varepsilon_i \text{ and } f^{-1}(x) \cap K_i \neq \emptyset \}.$$

It is easy to see that each C_i is a closed set. Let $C = \bigcup_{i=1}^{\infty} C_i$.

We will show that $\{f(K_i)\}$ is a locally finite collection in B^3 . Let $x_0 \in B^3$ and let V be a neighborhood of x_0 in B^3 with compact closure. Since f is a compact map, $f^{-1}(V)$ has compact closure. Since $\{K_i\}$ is a locally finite collection in W , $f^{-1}(V)$ intersects only a finite number of the K_i 's, and thus V intersects only a finite number of the $f(K_i)$'s. Using the fact the $\{f(K_i)\}$ is a locally finite collection, we see that C is a closed 0-dimensional subset of B^3 .

Consider the following commutative diagram where the horizontal maps are induced by inclusion, and the vertical maps are induced by f .

$$\begin{array}{ccc} H_1(W - f^{-1}(C); Z) & \xrightarrow{\alpha} & H_1(W; Z) \\ \downarrow & & \downarrow \\ H_1(B^3 - C; Z) & \longrightarrow & H_1(B^3; Z) \end{array}$$

First, we claim that α is an epimorphism. Let $[\delta] \in H_1(W; Z)$ where δ is a simple closed curve. Let O be an open set in B^3 so that $f(\delta) \subset O$ and $(\text{Bd } O) \cap C = \emptyset$. By applying Lemma 2 of [13], we see that δ is homologous in $f^{-1}(O)$ to a 1-cycle in $f^{-1}(O) - f^{-1}(O \cap C) \subset W - f^{-1}(C)$.

Finally, we claim that α is the zero homomorphism. Let $[\tau] \in H_1(W - f^{-1}(C); Z)$ where τ is a simple closed curve. We can also suppose that $f(\tau)$ is a simple closed curve, and that $f(\tau)$ bounds an orientable surface S in $B^3 - C$. By our choice of the ε_i 's, for each $y \in B^3 - C$, there is an open set V_y so that $f^{-1}(V_y)$ is contractible in W . Let $\mathcal{V} = \{V_y : y \in B^3 - C\}$. We can find a triangulation T of S which is so fine that

for each 2-simplex $\sigma \in T$, there is a $V_\sigma \in \mathcal{V}$ so that $\sigma \subset V_\sigma$. Using the fact that f is monotone, we can find a map h from the 1-skeleton of T into $W - f^{-1}(C)$ so that, if σ is a 2-simplex of T , $h(\partial\sigma) \subset f^{-1}(V_\sigma)$. (See the proof of Theorem 2.1 of [15] for details.) We can also suppose that $hf|_\tau$ is the identity. Since each V_σ is contractible in W , h can be extended to a map H which takes the surface S into W and which takes ∂S onto τ . Thus, $\alpha[\tau] = 0$ in $H_1(W; Z)$.

THEOREM 7. *Let M^3 and N^3 be 3-manifolds, possibly with boundary. Let f be a compact, monotone, boundary preserving mapping from M^3 onto N^3 such that $f(S_f)$ is 0-dimensional. Then $\{x \in N^3 : f^{-1}(x) \text{ is not cellular}\}$ is a locally finite subset of N^3 .*

Proof. By a procedure similar to the first part of the proof of Lemma 6, we can find a closed set $C \subset f(S_f) \subset N^3$ so that, if $x \notin C$, then there is an open set U_x where $f^{-1}(x) \subset U_x$ and U_x is contractible in M^3 . By Lemma 6, if $x \in N^3 - C$, then $f^{-1}(x)$ is strongly 1-acyclic over Z . Thus $f(A_f^0) \subset C$, and C is a closed 0-dimensional set. Theorem 7 now follows from Theorem 6.

Let $f: M^3 \rightarrow N^3$ be an onto, compact, boundary preserving map as before. Many of our earlier results have shown that $\{x \in N^3 : f^{-1}(x) \text{ is not cellular in } M^3\}$ is a locally finite subset of N^3 . The following three corollaries concern mappings of this type.

COROLLARY 3. *Let M^3 and N^3 be 3-manifolds, possibly with boundary. Let $f: M^3 \rightarrow N^3$ be a compact, monotone, boundary preserving map such that $\{x \in N^3 : f^{-1}(x) \text{ is not cellular}\}$ is a locally finite subset of N^3 . Then*

(i) *For each $x \in N^3$ and each open set U containing $f^{-1}(x)$, there is an open set V with $f^{-1}(x) \subset V \subset U$, such that $V - f^{-1}(x)$ is homeomorphic to $S^2 \times (0, 1)$.*

(ii) *N^3 can be obtained from M^3 by cutting out of M^3 a locally finite collection of mutually exclusive, polyhedral 3-manifolds, each with 2-sphere boundary, and replacing each by a 3-cell.*

Proof. (i) If $f^{-1}(x)$ is cellular, this follows from Theorem 1 of [3].

Let $x_1, x_2, x_3, \dots$ be the points in N^3 such that $f^{-1}(x_i)$ is not cellular for $i = 1, 2, 3, \dots$. Let $X = \{x_1, x_2, x_3, \dots\} \cup \text{Bd } N^3$. Let the 3-manifold Q^3 , the maps $\pi: M^3 \rightarrow Q^3$, $p: Q^3 \rightarrow N^3$, and the homeomorphism $h: N^3 \rightarrow Q^3$ be defined as in Lemmas 4 and 5. It will be sufficient to show that $f^{-1}(x_1)$ has the required neighborhood. We are given an open set $U \supset f^{-1}(x_1)$. Let U' be an open set in M^3 so that $f^{-1}(x_1) \subset U' \subset U$ and $U' \cap f^{-1}(x_i) = \emptyset$ for $i \geq 2$. Then $h^{-1}\pi(U')$ is an open set containing x_1 in N^3 . Let W be an open 3-cell so that $x_1 \subset W \subset h^{-1}\pi(U')$. Let $V = \pi^{-1}h(W)$. Then $V - f^{-1}(x_1)$ is homeomorphic by $\pi^{-1}h$ to $W - \{x_1\}$ which is homeomorphic to $S^2 \times (0, 1)$.

(ii) As in part (i) let $x_1, x_2, x_3, \dots$ be the points of N^3 whose inverses are not cellular. We can find pairwise disjoint closed neighborhoods $K_1, K_2, K_3, \dots$ of $f^{-1}(x_1), f^{-1}(x_2), f^{-1}(x_3), \dots$ respectively so that $K_i - f^{-1}(x_i)$ is homeomorphic to $S^2 \times (0, 1]$. Then each K_i is a 3-manifold with 2-sphere boundary, and $\pi|_{K_i}$ is a

boundary preserving map of K_i onto a 3-cell. Furthermore, $\pi|M^3 - \bigcup_{i=1}^{\infty} K_i$ is a homeomorphism. Thus Q^3 can be obtained by cutting $K_1, K_2, K_3, \dots$ out of M^3 , and replacing each by a 3-cell.

COROLLARY 4. *Let M^3 and N^3 be compact 3-manifolds, possibly with boundary. Let $f: M^3 \rightarrow N^3$ be a boundary preserving, onto map such that $\{x \in N^3 : f^{-1}(x) \text{ is not cellular in } M^3\}$ is a finite set. If M^3 is homeomorphic to N^3 , then $f^{-1}(x)$ is cellular for every $x \in N^3$.*

Proof. By Corollary 3, part (ii), there are closed 3-manifolds $*K_0^3, \dots, *K_n^3$ such that

$$M^3 = N^3 \# *K_0^3 \# \dots \# *K_n^3.$$

By a corollary to the Grushko-Neumann Theorem (see p. 192 of [10]), the rank of $\pi_1(M^3)$ is equal to the sum of the ranks of $\pi_1(N^3), \pi_1(K_0^3), \dots, \pi_1(K_n^3)$. Therefore

$$\pi_1(*K_0^3) = \dots = \pi_1(*K_n^3) = 1,$$

and each $*K_i^3$ ($i=0, \dots, n$) is a homotopy 3-sphere.

If M^3 is closed and orientable, we use the unique decomposition theorem of Milnor [14] to show that $*K_0^3, \dots, *K_n^3$ are all 3-spheres. This shows that $f^{-1}(x)$ is cellular for every $x \in N^3$.

If M^3 is orientable with boundary, we can sew a cube-with-handles onto each boundary component of M^3 to obtain a closed manifold M_0^3 . The homeomorphism from M^3 to N^3 induces a similar sewing of cubes-with-handles onto $\text{Bd } N^3$ to give a closed 3-manifold N_0^3 which is homeomorphic to M_0^3 . We have

$$M_0^3 = N_0^3 \# *K_0^3 \# \dots \# *K_n^3$$

and the argument for the closed orientable case applies.

If M^3 is nonorientable, we apply the previous argument to the orientable double covering of M^3 .

COROLLARY 5. *Let M^3 and N^3 be compact (i.e., closed) 3-manifolds. Let $f: M^3 \rightarrow N^3$ be an onto map such that $\{x \in N^3 : f^{-1}(x) \text{ is not cellular in } M^3\}$ is finite, and let $g: N^3 \rightarrow M^3$ be an onto map such that $\{x \in M^3 : g^{-1}(x) \text{ is not cellular in } N^3\}$ is finite. Then M^3 is homeomorphic to N^3 .*

Proof. By Corollary 2, we have $M^3 = N^3 \# *K_0^3 \# \dots \# *K_n^3$ and $N^3 = M^3 \# *Q_0^3 \# \dots \# *Q_m^3$. By a corollary to the Grushko-Neumann Theorem (p. 192 of [10]) we see that $*K_0^3, \dots, *K_n^3, *Q_0^3, \dots, *Q_m^3$ are all homotopy 3-spheres. This implies that all of the point inverses of f and g have property UV^∞ . Then Corollary 5 follows from Corollary 2.3 of [11].

VI. On Haken's finiteness theorem. In [5], Wolfgang Haken stated a finiteness theorem for incompressible surfaces in a compact 3-manifold M^3 . We are interested here only in the special case of the theorem where the surfaces are closed: this

case is stated as Theorem C. Some difficulties arise with Haken's proof in the case where M^3 is not irreducible. Haken's proof is correct and can be simplified considerably in the case where M^3 is irreducible. We give here an argument due to John Hempel to show that the finiteness theorem holds in the case where M^3 may not be irreducible. Haken intended to prove Kneser's Theorem [7] as a special case of the finiteness theorem; our argument uses Kneser's Theorem. The previous results of this paper depend on the finiteness theorem directly through Theorem 2 of [12].

In this section we will be working in the piecewise-linear category. A *surface* is a 2-manifold. If F^2 is a surface in a 3-manifold M^3 , and if F^2 is not a 2-sphere, then F^2 is *incompressible* in M^3 if every simple closed curve in F^2 that bounds an (open) disk in $M^3 - F^2$ also bounds a disk in F^2 . A 2-sphere is *incompressible* in M^3 if it does not bound a 3-cell in M^3 . A 3-manifold M^3 is irreducible if every 2-sphere in M^3 bounds a 3-cell in M^3 .

Two surfaces F_0^2 and F_1^2 in a 3-manifold M^3 are *parallel* in M^3 if there is an embedding $\alpha: F_0^2 \times [0, 1] \rightarrow M^3$ such that $\alpha_0: F_0^2 \rightarrow M^3$ is the inclusion map, and $\alpha_1: F_0^2 \rightarrow M^3$ takes F_0^2 homeomorphically onto F_1^2 . If $F_1^2, \dots, F_n^2$ are disjoint surfaces in a 3-manifold M^3 , and if L^3 is the closure of a complementary domain of $M^3 - \bigcup_{i=1}^n F_i^2$, then L^3 is a *parallelity component* if, for some $i=1, \dots, n$, there is a homeomorphism $h: F_i^2 \times [0, 1] \rightarrow L^3$ such that $h_0: F_i^2 \rightarrow L^3$ is the inclusion map, and $h_1: F_i^2 \rightarrow L^3$ takes F_i^2 homeomorphically onto F_j^2 for some $j=1, \dots, n, j \neq i$.

If C^3 is a 3-manifold, possibly with boundary, we define $\hat{C}^3$ to be the 3-manifold, possibly with boundary, obtained from C^3 by capping off each 2-sphere boundary component of C^3 with a 3-cell.

If B^3 is a 3-cell, and if $B_1^3, \dots, B_k^3$ are disjoint polyhedral 3-cells in $\text{Int } B^3$, then we call the manifold-with-boundary $B^3 - (\bigcup_{i=1}^k \text{Int } B_i^3)$ a *punctured 3-cell*.

LEMMA A. If F^2 is an incompressible surface in the product $M^2 \times [0, 1]$, where M^2 is a compact 2-manifold, then F^2 is parallel to $M^2 \times \{0\}$ and $M^2 \times \{1\}$.

This lemma is stated and proved by Haken on pp. 91–96 of [5].

LEMMA B. If C^3 is a 3-manifold, possibly with boundary, and $\hat{C}^3$ is irreducible, then the finiteness theorem holds for C^3 . In other words, there is an integer $n=n(C^3)$ such that if $F_1^2, \dots, F_{n+1}^2$ are $n+1$ disjoint incompressible polyhedral surfaces in C^3 , then two of these surfaces are parallel.

Proof. We have assumed the finiteness theorem for irreducible 3-manifolds, so there is an integer $n(\hat{C}^3)$ such that if there are more than $n(\hat{C}^3)$ disjoint incompressible surfaces in $\hat{C}^3$, then two of them are parallel. There are disjoint 3-cells $B_1^3, \dots, B_k^3$ such that $C^3 = \hat{C}^3 - [\bigcup_{i=1}^k \text{Int } B_i^3]$. Let $n=n(C^3)=n(\hat{C}^3)+2k$. Let $F_1^2, \dots, F_{n+1}^2$ be $n+1$ disjoint incompressible surfaces in C^3 . Then $n-k+1$ of these surfaces are incompressible in $\hat{C}^3$. There are $k+1$ distinct pairs from $F_1^2, \dots, F_{n+1}^2$ which are parallel in $\hat{C}^3$. (We say that the pair (F_i^2, F_j^2) is distinct from the pair

(F_r^2, F_s^2) if either $i \neq r, s$ or $j \neq r, s$.) Then, using Lemma A, we see that there are $k+1$ parallelity components in $\hat{C}^3$ whose interiors are disjoint. Thus, one of these parallelity components does not contain any B_i^3 , and is a parallelity component in C^3 .

THEOREM C. *Let M^3 be a compact 3-manifold, possibly with boundary. Then there is an integer $n_0 = n(M^3)$ such that if $F_1^2, \dots, F_{n_0+1}^2$ are n_0+1 disjoint polyhedral-incompressible surfaces in M^3 , then two of these surfaces are parallel.*

Proof. Let $\Sigma = \{S_1^2, \dots, S_l^2\}$ be a disjoint collection of 2-spheres in M^3 . Let $N_1^3, \dots, N_l^3$ be disjoint regular neighborhoods of $S_1^2, \dots, S_l^2$ respectively. Let $C_1^3, \dots, C_k^3$ be the components of $\text{Cl}(M^3 - \bigcup_{i=1}^l N_i^3)$. (The C_i^3 's are determined up to homeomorphism by the S_i^2 's and do not depend on the choice of the N_i^3 's. Note that k may not equal l since some of the S_i^2 's may not separate M^3 .) We will call Σ a complete system of 2-spheres in M^3 if $\hat{C}_1^3, \dots, \hat{C}_k^3$ are each irreducible. We will let $n(M^3, \Sigma) = \sum_{i=1}^k n(C_i^3)$ where $n(C_i^3)$ ($i=1, \dots, k$) is defined in Lemma B.

Kneser's Theorem [7] shows that there is a complete system Σ_0 of 2-spheres in M^3 . We will assume Σ_0 is a fixed complete system and we will let $n_0 = n(M^3, \Sigma_0)$.

Let $F_1^2, \dots, F_{n_0+1}^2$ be disjoint incompressible surfaces in M^3 . Let $F^2 = \bigcup_{i=1}^{n_0+1} F_i^2$. Suppose $\Sigma = \{S_1^2, \dots, S_l^2\}$ is a complete system of 2-spheres in M^3 , each of which is in general position with respect to F^2 , and suppose that $n(M^3, \Sigma) = n_0$. Let $m(M^3, \Sigma, F^2)$ be the number of components of $(\bigcup_{i=1}^l S_i^2) \cap F^2$. (Each of these components is a simple closed curve.) We can suppose $m(M^3, \Sigma, F^2)$ is minimal over all such complete systems of 2-spheres in M^3 . Theorem C will be proved if $m(M^3, \Sigma, F^2)$ is zero. For then there will be more than $n(C_j^3)$ of the surfaces $F_1^2, \dots, F_{n_0+1}^2$ in one of the components C_j^3 , and two of these surfaces must be parallel in C_j^3 by Lemma B. (Let $N_1^3, \dots, N_l^3$ and $C_1^3, \dots, C_k^3$ be defined as before.)

So we suppose that $m(M^3, \Sigma, F^2) > 0$. Any simple closed curve of $(\bigcup_{i=1}^l S_i^2) \cap F^2$ must bound a disk in F^2 , since F^2 is incompressible. Therefore, we can choose an "innermost" (on F^2) simple closed curve J of $(\bigcup_{i=1}^l S_i^2) \cap F^2$; suppose $J \subset S_r^2 \cap F_s^2$ for some $r=1, \dots, l$ and $s=1, \dots, n_0+1$. Let D^2 be the disk that J bounds in F_s^2 . Then D^2 is contained in some C_q^3 (where $q=1, \dots, k$) except for a regular neighborhood of $\text{Bd } D^2$.

Let E_1^2 and E_2^2 be the two disks bounded by J in S_r^2 . We can push each of the 2-spheres $E_1^2 \cup D^2$ and $E_2^2 \cup D^2$ to one side so that they each miss D^2 , and so that they are each contained in C_q^3 . Then one of these 2-spheres must be in the boundary of a punctured cube P^3 in C_q^3 since $\hat{C}_q^3$ is irreducible. Let $S_r'^2$ be the 2-sphere that is not in the boundary of P^3 , and let $\Sigma' = \{S_1^2, \dots, S_{r-1}^2, S_r'^2, S_{r+1}^2, \dots, S_l^2\}$. We will show that Σ' is a complete system of 2-spheres in M^3 , that $n(M^3, \Sigma') = n_0$, and that $m(M^3, \Sigma', F^2) < m(M^3, \Sigma, F^2)$.

Let C_t^3 ($t=1, \dots, k$) be the component of $\text{Cl}(M^3 - \bigcup_{i=1}^l N_i^3)$ on the "other side" of S_r^2 . (If S_r^2 does not separate M^3 , then C_q^3 may equal C_t^3 .) If we choose a small regular neighborhood $N_r'^3$ of $S_r'^2$ (so that $N_r'^3 \cap D^2 = \emptyset$) and let $N_i'^3 = N_i^3$ for

$i \neq r$, we can define $C_q'^3$ and $C_t'^3$ to be components of $\text{Cl}(M^3 - \bigcup_{i=1}^l N_i'^3)$. A sub-disk D_0^2 of D^2 is a spanning disk of C_q^3 and if we remove the interior of a regular neighborhood of D_0^2 , this separates C_q^3 into two components, one homeomorphic to $C_q'^3$, and the other homeomorphic to the punctured cube P^3 . Thus $\hat{C}_q^3$ is homeomorphic to $\hat{C}_q'^3$. Furthermore, $C_t'^3$ is homeomorphic to the manifold obtained by sewing P^3 to C_t^3 along a disk on the boundary of each. Thus $\hat{C}_t^3$ is homeomorphic to $\hat{C}_t'^3$. We also have $n(C_q^3) + n(C_t^3) = n(C_q'^3) + n(C_t'^3)$ since the 2-sphere boundary components of $C_q^3 \cap P^3$ which were removed from C_q^3 to obtain $C_q'^3$ were added to C_t^3 to obtain $C_t'^3$. Thus $n(M^3, \Sigma') = n_0$.

Since $S_r'^2 \cap D^2 = \emptyset$, $m(M^3, \Sigma', F^2) < m(M^3, \Sigma, F^2)$, and this contradicts our assumption that $m(M^3, \Sigma, F^2)$ was minimal.

BIBLIOGRAPHY

1. S. Armentrout, *Concerning cellular decompositions of 3-manifolds with boundary*, Trans. Amer. Math. Soc. **137** (1969), 231–236. MR **38** #5224.
2. R. H. Bing, *Locally tame sets are tame*, Ann. of Math. (2) **59** (1954), 145–158. MR **15**, 816.
3. M. Brown, *A proof of the generalized Schoenflies theorem*, Bull. Amer. Math. Soc. **66** (1960), 74–76. MR **22** #8470b.
4. P. T. Church, *Differentiable and monotone maps on manifolds. II*, Trans. Amer. Math. Soc. **158** (1971), 493–503.
5. W. Haken, *Some results on surfaces in 3-manifolds*, Studies in Modern Topology, Math. Assoc. Amer., distributed by Prentice-Hall, Englewood Cliffs, N. J., 1968, pp. 39–98. MR **36** #7118.
6. H. Kneser, *Geschlossen Flächen in dreidimensionalen Mannigfaltigkeiten*, Jber. Deutsch. Math.-Verein. **38** (1929), 248–260.
7. R. C. Lacher, *Cellularity criteria for maps*, Michigan Math. J. **17** (1970), 385–396.
8. R. C. Lacher and A. H. Wright, *Mapping cylinders and 4-manifolds*, Topology of Manifolds, Markham, Chicago, Ill., 1970.
9. H. W. Lambert, *Replacing certain maps of 3-manifolds by homeomorphisms*, Proc. Amer. Math. Soc. **23** (1969), 676–678. MR **41** #1058.
10. W. Magnus, A. Karrass and D. Solitar, *Combinatorial group theory: Presentations of groups in terms of generators and relations*, Pure and Appl. Math., vol. 13, Interscience, New York, 1966. MR **34** #7617.
11. D. R. McMillan, Jr., *Strong homotopy equivalence of 3-manifolds*, Bull. Amer. Math. Soc. **73** (1967), 718–722. MR **37** #4817.
12. ———, *Compact acyclic subsets of three-manifolds*, Michigan Math. J. **16** (1969), 129–136. MR **39** #4822.
13. ———, *Acyclicity in three-manifolds*, Bull. Amer. Math. Soc. **76** (1970), 942–964.
14. J. W. Milnor, *A unique decomposition theorem for 3-manifolds*, Amer. J. Math. **84** (1962), 1–7. MR **25** #5518.
15. T. M. Price, *A necessary condition that a cellular upper semicontinuous decomposition of E^n yield E^n* , Trans. Amer. Math. Soc. **122** (1966), 427–435. MR **33** #1843.
16. A. H. Wright, *Monotone mappings of compact 3-manifolds*, Ph.D. Thesis, University of Wisconsin, Madison, Wis., 1969.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UTAH 84112

Current address: Department of Mathematics, Western Michigan University, Kalamazoo, Michigan 49001