

HOMOMORPHISMS OF COMMUTATIVE CANCELLATIVE SEMIGROUPS INTO NONNEGATIVE REAL NUMBERS

BY

MOHAN S. PUTCHA AND TAKAYUKI TAMURA

ABSTRACT. Let S be a commutative cancellative semigroup and T_0 be a cofinal subsemigroup of S . Let h_0 be a homomorphism of T_0 into the semigroup of nonnegative real numbers under addition. We prove that Kobayashi's condition [2] is necessary and sufficient for h_0 to be extended to S . Further, we find a necessary and sufficient condition in order that the extension be unique. Related to this, the "boundedness condition" is introduced. For further study, several examples are given.

1. Introduction. A commutative cancellative archimedean idempotent-free semigroup is called an \mathfrak{N} -semigroup. Kobayashi [2] proved the following:

THEOREM 1.1. *Let T_0 be a subsemigroup of an \mathfrak{N} -semigroup S and let h_0 be a homomorphism of T_0 into the semigroup \mathbb{R}_+^0 of nonnegative real numbers under addition. Then h_0 can be extended to a homomorphism of S into \mathbb{R}_+^0 if and only if the pair $\langle T_0, h_0 \rangle$ satisfies the following condition: if $x, y \in T_0$ and $x \mid y$ (x divides y) in S , then $h_0(x) \leq h_0(y)$.*

One of the authors [4] has studied the homomorphisms of T_0 into \mathbb{R}_+ from the viewpoint of positive quasi-orders. In this paper, we treat the homomorphisms of T_0 into the nonnegative real numbers in the case when S is a commutative cancellative semigroup and T is its subsemigroup. Theorem 2.1 will be a straightforward generalization of the classical result that characters can be extended from a subgroup of an abelian group G to G itself. In §2, we will show that Theorem 1.1 holds if T_0 is cofinal in S . In §3, we will introduce a "boundedness condition" and discuss the relation between this condition and the extension of a homomorphism beyond a filter. In §4, we will give a few examples, which show that Theorem 2.1 does not necessarily hold if T_0 is not cofinal.

A subsemigroup U of a commutative semigroup S is called unitary in S if

Received by the editors November 13, 1974.

AMS (MOS) subject classifications (1970). Primary 20M99.

Key words and phrases. Commutative cancellative (idempotent-free) semigroup, cofinal subsemigroup, unitary subsemigroup, homomorphism, K -condition, \mathcal{B} -condition (boundedness condition), unitary closure, filter, filter closure.

$x \in S$, $a \in U$ and $ax \in U$ imply $x \in U$. U is called cofinal in S if, for every $x \in S$, there is a $y \in S$ such that $xy \in U$. As is well known, see [1] or [5], a unitary cofinal subsemigroup U induces a group congruence ρ_U on S defined by $x \rho_U y$ if and only if $ax = by$ for some $a, b \in U$. We denote S/ρ_U by S/U . Furthermore the kernel of $S \rightarrow S/U$ coincides with U .

Let T be a nonempty subsemigroup of S . The smallest unitary subsemigroup \bar{T} of S containing the subsemigroup T is called the unitary closure of T in S . \bar{T} is given by

$$\bar{T} = \{x \in S: xt \in T \text{ for some } t \in T\}.$$

A nonempty subsemigroup F of S is called a filter of S [3] if $x, y \in S$ and $xy \in F$ implies $x, y \in F$. The smallest filter \tilde{T} of S containing the subsemigroup T is called the filter closure of T in S . Then

$$\tilde{T} = \{x \in S: xy \in T \text{ for some } y \in S\}.$$

(1.2) The following hold.

(1.2.1) $T \rightarrow \bar{T}$ and $T \rightarrow \tilde{T}$ are closure mappings, that is, $T \subseteq \bar{T}$, $T \subseteq \tilde{T}$.

$T_1 \subseteq T_2$ implies $\bar{T}_1 \subseteq \bar{T}_2$ and $\tilde{T}_1 \subseteq \tilde{T}_2$.

$\bar{\bar{T}} = \bar{T}$, $\tilde{\tilde{T}} = \tilde{T}$.

(1.2.2) \bar{T} is unitary in S , \tilde{T} is a filter in S , and T is cofinal in \tilde{T} .

(1.2.3) $\tilde{T} = \bar{\tilde{T}} = \tilde{\bar{T}}$.

(1.2.4) $\bar{T} \subseteq \tilde{T}$ and \bar{T} is unitary cofinal in \tilde{T} .

Throughout this paper, \mathbf{R} denotes the set of real numbers, R the set of rational numbers, \mathbf{R}_+ (\mathbf{R}_-) the set of positive (negative) real numbers; \mathbf{R}_+^0 (\mathbf{R}_-^0) the set of nonnegative (nonpositive) real numbers; \mathbf{Z}_+ (\mathbf{Z}_-) the set of positive (negative) integers and \mathbf{Z}_+^0 (\mathbf{Z}_-^0) the set of nonnegative (nonpositive) integers.

If S is a semigroup and if X is a subsemigroup of the additive group \mathbf{R} , then the notation $\text{Hom}(S, X)$ denotes the semigroup of homomorphisms of S into X under the usual operation. Let X_1, X_2, Y_1 and Y_2 be commutative semigroups such that $X_1 \subseteq X_2$ and $Y_1 \subseteq Y_2$. Let $h_1 \in \text{Hom}(X_1, Y_1)$ and $h_2 \in \text{Hom}(X_2, Y_2)$. If $h_2|_{X_1} = h_1$, we say that h_1 of $\text{Hom}(X_1, Y_1)$ is extended to h_2 of $\text{Hom}(X_2, Y_2)$; in particular, if $Y_1 = Y_2$, we say that h_1 of $\text{Hom}(X_1, Y_1)$ is extended to X_2 . If the extension h_2 of h_1 of $\text{Hom}(X_1, Y_1)$ to X_2 is unique, we say that h_1 of $\text{Hom}(X_1, Y_1)$ is uniquely extended to X_2 . Let $h \in \text{Hom}(S, \mathbf{R})$. h is called trivial if $h(x) = 0$ for all $x \in S$.

In this paper the binary operation in a commutative semigroup will be denoted by addition, i.e. $+$.

2. Extensions from cofinal subsemigroups. In this section, we will prove the following generalization of Theorem 1.1.

THEOREM 2.1. *Let T_0 be a cofinal subsemigroup of a commutative cancellative semigroup S and let h_0 be a homomorphism of T_0 into the additive semigroup \mathbb{R}_+^0 of nonnegative real numbers. Then h_0 can be extended to S if and only if*

$$(K) \quad t_1 \in S + t_2 \text{ implies } h_0(t_1) \geq h_0(t_2) \text{ for all } t_1, t_2 \in T_0.$$

In this paper, the condition (K) will be called the K -condition. It is obvious that if h_0 can be extended to S then the K -condition must hold. We will prove sufficiency. Let X denote the set of pairs $\langle T, h \rangle$ where T is a subsemigroup of S containing T_0 and $h \in \text{Hom}(T, \mathbb{R}_+^0)$ such that $h|_{T_0} = h_0$ and $\langle T, h \rangle$ satisfies the K -condition.

Let $[a]$ be the cyclic subsemigroup generated by a and let $[T, a]$ be the subsemigroup generated by T and a , i.e.,

$$[T, a] = T \cup (T + [a]) \cup [a].$$

LEMMA 2.2. *Let $\langle T, h \rangle \in X$ and suppose that $a \in S$ and $(T + [a]) \cap T \neq \emptyset$. Then there exists $h': [T, a] \rightarrow \mathbb{R}_+^0$ such that $\langle [T, a], h' \rangle \in X$. Further, h' is unique.*

PROOF. There exist $t_1, t_2 \in T, N \in \mathbb{Z}_+$ such that $t_1 = N \cdot a + t_2$. Then $h(t_1) \geq h(t_2)$ by the K -condition. Define $h': [T, a] \rightarrow \mathbb{R}_+^0$ by

$$h'(t + n \cdot a) = h(t) + \frac{n}{N}[h(t_1) - h(t_2)], \quad t \in T, n \in \mathbb{Z}_+^0,$$

$$h'(na) = \frac{n}{N}[h(t_1) - h(t_2)], \quad n \in \mathbb{Z}_+.$$

First we show that h' is well defined: $t + n \cdot a = t' + n' \cdot a, t, t' \in T, n, n' \in \mathbb{Z}_+$, implies $N \cdot t + Nn \cdot a + (n + n') \cdot t_2 = N \cdot t' + Nn' \cdot a + (n + n') \cdot t_2$, that is, $N \cdot t + n \cdot t_1 + n' \cdot t_2 = N \cdot t' + n' \cdot t_1 + n \cdot t_2$. This shows $h'(t + n \cdot a) = h'(t' + n' \cdot a)$, hence h' is well defined. From its definition, h' is clearly a homomorphism into \mathbb{R}_+^0 and $h'|_T = h$. Assume $t + n \cdot a = s + t' + n' \cdot a$ for some $a \in S$. Then $N \cdot t + n \cdot t_1 + n' \cdot t_2 = N \cdot s + N \cdot t' + n' \cdot t_1 + n \cdot t_2$ which implies

$$N \cdot h(t) + n \cdot h(t_1) + n' \cdot h(t_2) \geq N \cdot h(t') + n' \cdot h(t_1) + n \cdot h(t_2)$$

by the K -condition. This gives $h'(t + n \cdot a) \geq h'(t' + n' \cdot a)$. Hence $\langle [T, a], h' \rangle \in X$. If h'' is any extension of h to a homomorphism of $[T, a]$ into \mathbb{R}_+^0 , we must have

$$h(t_1) = h''(t_1) = N \cdot h''(a) + h''(t_2) = N \cdot h''(a) + h(t_2)$$

so that $h''(a) = N^{-1}[h(t_1) - h(t_2)] = h'(a)$. It follows that $h''(t + n \cdot a) = h'(t + n \cdot a)$ for all $t \in T$, all $n \in Z_+$, that is, $h'' = h'$. \square

To consider the case when $(T + [a]) \cap T = \emptyset$, we need a lemma. From now on, t_1, t_2, t_3 and t_4 will denote arbitrary elements of T .

Let

$$A(a) = \{n^{-1}[h(t_2) - h(t_1)]: t_1 + n \cdot a \in t_2 + S\},$$

$$B(a) = \{n^{-1}[h(t_2) - h(t_1)]: t_2 \in t_1 + n \cdot a + S\},$$

where $a \in S$, $A(a)$ and $B(a)$ mean the sets depending on a . Note that $0 \in A(a)$ and hence $A(a) \neq \emptyset$.

LEMMA 2.3. *Let $\langle T, h \rangle \in X$ and suppose that $(S + [a]) \cap T \neq \emptyset$. Then $\text{Sup } A(a) \leq \text{Inf } B(a) < \infty$.*

PROOF. Since $(S + [a]) \cap T \neq \emptyset$, there are $t_1, t_2 \in T, x \in S$ and $n \in Z_+$ such that $t_2 = t_1 + n \cdot a + x$. Hence $B(a) \neq \emptyset$ and $\text{Inf } B(a) < \infty$. Suppose $t_1 + n_1 \cdot a = t_2 + s_1$ and $t_4 = t_3 + n_2 \cdot a + s_2$ where $n_1, n_2 \in Z_+, s_1, s_2 \in S$. Then $n_1 \cdot t_4 + n_2 \cdot t_1 = n_1 \cdot t_3 + n_2 \cdot t_2 + n_2 \cdot s_1 + n_1 \cdot s_2$. By the K-condition, $n_1 \cdot h(t_4) + n_2 \cdot h(t_1) \geq n_1 \cdot h(t_3) + n_2 \cdot h(t_2)$. Hence $n_2^{-1}[h(t_4) - h(t_3)] \geq n_1^{-1}[h(t_2) - h(t_1)]$. Thus we get $\text{Sup } A(a) \leq \text{Inf } B(a)$. \square

LEMMA 2.4. *Let $\langle T, h \rangle \in X$ and suppose that $(S + [a]) \cap T \neq \emptyset$ but $(T + [a]) \cap T = \emptyset$. Then h can be extended to a homomorphism $h': [T, a] \rightarrow R_+^0$ and $\langle [T, a], h' \rangle \in X$. The h' is determined by choosing $h'(a)$ such that $\text{Sup } A(a) \leq h'(a) \leq \text{Inf } B(a)$. Moreover, every extension h'' of h to $[T, a]$ such that $\langle [T, a], h'' \rangle \in X$ is obtained in this way.*

PROOF. Choose $b \in R_+^0$ such that

$$(2.4.1) \quad \text{Sup } A(a) \leq b \leq \text{Inf } B(a).$$

Define

$$(2.4.2) \quad \begin{aligned} h'(t + n \cdot a) &= h(t) + n \cdot b & \text{for } t \in T, n \in Z_+^0, \\ h'(na) &= nb & \text{for } n \in Z_+. \end{aligned}$$

Since S is cancellative and $(T + [a]) \cap T = \emptyset$, every element of $T + [a]$ is uniquely expressed as $t + n \cdot a$ and hence h' is well defined. Then h' is clearly a homomorphism $[T, a] \rightarrow R_+^0$ and $h'|_T = h$. Suppose that $t_1 + n_1 \cdot a = t_2 + n_2 \cdot a + s, n_1, n_2 \in Z_+^0, s \in S$. Then there are three possibilities: $n_1 = n_2, n_1 > n_2$ and $n_1 < n_2$. If $n_1 = n_2$, then, since S is cancellative, $t_1 = t_2 + s$, hence $h(t_1) \geq h(t_2)$ by the K-condition. This implies $h'(t_1 + n_1 \cdot a) \geq h'(t_2 + n_2 \cdot a)$.

If $n_1 > n_2$, then $t_1 + (n_1 - n_2) \cdot a = t_2 + s$ and, by the choice of b ,

$$h(t_2) - h(t_1) \leq (n_1 - n_2) \cdot b.$$

This implies

$$h'(t_2 + n_2 \cdot a) = h(t_2) + n_2 \cdot b \leq h(t_1) + n_1 \cdot b = h'(t_1 + n_1 \cdot a).$$

If $n_1 < n_2$, then $t_1 = t_2 + (n_2 - n_1) \cdot a + s$. By the choice of b ,

$$(n_2 - n_1) \cdot b \leq h(t_1) - h(t_2).$$

This gives

$$h'(t_2 + n_2 \cdot a) = h(t_2) + n_2 \cdot b \leq h(t_1) + n_1 \cdot b = h'(t_1 + n_1 \cdot a).$$

Therefore $\langle [T, a], h' \rangle \in X$.

Assume that h'' is an extension of h to $[T, a]$ and that $t_1 + n_1 \cdot a = t_2 + s_1$ and $t_4 = t_3 + n_2 \cdot a + s_2$, $n_1, n_2 \in \mathbb{Z}_+$, $s_1, s_2 \in S$. Using the assumption that h'' obeys the K -condition, $t_1 + n_1 a = t_2 + s$ gives $h''(t_1) + n_1 h''(a) \geq h''(t_2)$, so that $h''(a) \geq (h(t_2) - h(t_1))/n_1$, hence $h''(a) \geq \sup A(a)$. Likewise we have $h''(a) \leq (h(t_4) - h(t_3))/n_2$, hence $h''(a) \leq \inf B(a)$. By the former half of the lemma, $\langle [T, a], h'' \rangle \in X$. \square

COROLLARY 2.5. *In Lemma 2.4, the extension h' is unique if and only if*

$$(2.5.1) \quad \sup A(a) = \inf B(a).$$

PROOF OF SUFFICIENCY OF THEOREM 2.1. Define the partial order in X by $\langle T_1, h_1 \rangle \leq \langle T_2, h_2 \rangle$ if and only if $T_1 \subseteq T_2$ and h_2 is an extension of h_1 to T_2 . Then it is easy to see that X satisfies the condition for Zorn's lemma and so X has maximal members. To show that any such maximal member has domain S , it suffices to show that if $\langle T, h \rangle \in X$ and $a \notin T$, then h can be extended to $h': [T, a] \rightarrow \mathbb{R}_+^0$ such that $\langle [T, a], h' \rangle \in X$. Since T is cofinal, $(S + [a]) \cap T \neq \emptyset$, furthermore there are two possibilities: $(T + [a]) \cap T \neq \emptyset$ and $(T + [a]) \cap T = \emptyset$. Lemma 2.2 has dealt with the first case; Lemma 2.4 has done the second case. Thus the theorem has been proved. \square

COROLLARY 2.6. *Let S be a commutative cancellative semigroup and T_0 a unitary cofinal subsemigroup of S . Then every homomorphism h of T_0 into \mathbb{R}_+^0 can be extended to S .*

PROOF. Every h satisfies the K -condition.

COROLLARY 2.7. *Let T_0 be an ideal of S . Then every homomorphism h of T_0 into \mathbb{R}_+^0 can be uniquely extended to S .*

PROOF. Lemma 2.2 is applied to this case since $(T_0 + [a]) \cap T_0 \neq \emptyset$ for each $a \in S$. The direct alternate proof of this corollary is left for the reader's exercise. \square

Since every subsemigroup of a commutative archimedean semigroup is cofinal, Theorem 1.1 is a special case of Theorem 2.1.

THEOREM 2.8. *Let T be a cofinal subsemigroup of a commutative cancellative subsemigroup S , and let $h: T \rightarrow \mathbb{R}_+^0$ be a homomorphism. Then h admits a unique extension to S if and only if, for each $a \in S$, $\text{Sup } A(a) = \text{Inf } B(a)$.*

PROOF. Assume h admits a unique extension to S . Then $\langle T, h \rangle$ satisfies the K -condition. Suppose that h_1 and h_2 are distinct extensions such that $\langle [T, a], h_1 \rangle$ and $\langle [T, a], h_2 \rangle$ obey the K -condition for some $a \notin T$. Then $(S + [a]) \cap T \neq \emptyset$ since T is cofinal in S ; $(T + [a]) \cap T = \emptyset$ by Lemma 2.2. Now Lemma 2.4 shows that $\langle [T, a], h_1 \rangle$ and $\langle [T, a], h_2 \rangle$ are in \mathcal{X} . By Theorem 2.1, h_1 and h_2 can be extended to homomorphisms $h'_1: S \rightarrow \mathbb{R}_+^0$ and $h'_2: S \rightarrow \mathbb{R}_+^0$ respectively; but $h'_1 \neq h'_2$. This contradicts the assumption. Therefore h admits a unique extension to $[T, a]$ for each $a \notin T$. If $(T + [a]) \cap T \neq \emptyset$, we can easily show that if $n \in \mathbb{Z}_+$, $t_1, t_2 \in T$ and $t_1 + n \cdot a = t_2$, then

$$\text{Sup } A(a) = \text{Inf } B(a) = (h(t_2) - h(t_1))/n.$$

If $(T + [a]) \cap T = \emptyset$, then Corollary 2.5 shows $\text{Sup } A(a) = \text{Inf } B(a)$.

Conversely, suppose $\text{Sup } A(a) = \text{Inf } B(a)$ for every $a \in S$. If $t_2 = t_1 + s$, $t_1, t_2 \in T$, $s \in S$, then $2t_2 \in 2t_1 + s + S$, which implies

$$\text{Inf } B(s) \leq 2[h(t_2) - h(t_1)].$$

As $t_1 + 2 \cdot s \in t_2 + S$, $\text{Sup } A(s) \geq \frac{1}{2}[h(t_2) - h(t_1)]$. Hence

$$\frac{1}{2}[h(t_2) - h(t_1)] \leq 2[h(t_2) - h(t_1)].$$

It follows that $h(t_2) \geq h(t_1)$. Hence h satisfies the K -condition, and so h is extended to S . By Lemma 2.2 and Corollary 2.5, the extension is unique since $\text{Sup } A(a) = \text{Inf } B(a)$ for each $a \in S$. \square

3. Boundedness condition. In Lemma 2.3, we see that the set A is bounded. In light of this, we will introduce the boundedness condition (\mathcal{B} -condition). In this section, we assume that S is a commutative cancellative semigroup and let $P = S \setminus F$ where P is a prime ideal, $P \neq \emptyset$, and F is a filter [3], $F \neq \emptyset$. Let $a \in P$. The subsemigroup of S generated by a and F is denoted by $P_F(a)$ or $P(a)$ if F is fixed. We define the relation ρ on P as follows: $c \rho d$ if and only if $m \cdot c + s = n \cdot d + t$ for some $s, t \in F$ and some $m, n \in \mathbb{Z}_+$. Then ρ is an equiva-

lence relation on P and each ρ -class is a subsemigroup of P ; i.e., ρ has the following properties:

$$(3.1.1) \quad x \rho y \text{ implies } x \rho x + y \text{ for all } x, y \in P.$$

$$(3.1.2) \quad x \rho m \cdot x + t \text{ for all } t \in F \text{ and all } m \in Z_+.$$

Let $Q(a)$ denote the ρ -class containing $a \in P$. Let $U_F(a)$ or $U(a) = [Q(a), F]$, i.e., the subsemigroup of S generated by $Q(a)$ and F . By (3.1.1) and (3.1.2), we see $Q(a) + F \subseteq Q(a)$. If $s, t \in F$ and $b \in Q(a)$, then $k \cdot b + s = l \cdot b + t$, ($k, l \in Z_+^0$), implies $k = l$ and $s = t$. In fact, if $k > l$, $(k - l) \cdot b + s = t$ by cancellation, but this is impossible since F is a filter and $b \in P$. Hence $k \leq l$. Likewise $k \geq l$. Therefore, $k = l$, and hence $s = t$ by cancellation. Thus we have

(3.2) Each element of $Q(a) + F$ has a unique expression as the sum of an element of $Q(a)$ and an element of F .

As defined in §1, \bar{X} denotes the unitary closure of X and \widetilde{X} denotes the filter closure of X .

LEMMA 3.3.

$$(3.3.1) \quad U(a) = \{x \in S: m \cdot x \in \overline{P(a)} \text{ for some } m \in Z_+\}$$

and

$$(3.3.2) \quad \overline{P(a)} \subseteq U(a) = \overline{U(a)} \subseteq \widetilde{P(a)} = \widetilde{U(a)}.$$

PROOF. (3.3.1) If $x \in F$, then $x \in P(a) \subseteq \overline{P(a)}$. If $x \in Q(a)$, then $m \cdot x + s = n \cdot a + t$ for some $s, t \in F$, some $m, n \in Z_+$; hence $m \cdot x \in \overline{P(a)}$. Therefore $U(a)$ is contained in the set at the right-hand side. To prove the other direction, let $m \cdot x \in \overline{P(a)}$. By definition, $n \cdot a + s + m \cdot x = l \cdot a + t$ for some $s, t \in F$, and some $n, l \in Z_+^0$. Suppose $n > l$. Then $a + z = t$ for some $z \in S$. This contradicts $a \in P$. Hence $n \leq l$. If $n = l$, then $x \in F$. If $n < l$, then $m \cdot x + s = (l - n) \cdot a + t$ which implies $x \in Q(a)$, hence $x \in U(a)$. Thus we have (3.3.1).

(3.3.2) It immediately follows from (3.3.1) and the definition that $\overline{P(a)} \subseteq U(a) \subseteq \widetilde{P(a)}$. Taking their filter closures, we get $\widetilde{P(a)} = \widetilde{U(a)}$ by (1.2.3). It remains to show $\overline{U(a)} \subseteq U(a)$. Let $x \in \overline{U(a)}$. Then $b = c + x$ for some $b, c \in U(a)$. By (3.3.1), we can choose $m \in Z_+$ such that $m \cdot b, m \cdot c \in \overline{P(a)}$. Since $m \cdot b = m \cdot c + m \cdot x$ and $\overline{P(a)}$ is unitary by (1.2.2), we see that $m \cdot x \in \overline{P(a)}$. So $x \in U(a)$. Therefore $\overline{U(a)} \subseteq U(a)$. This completes the proof. \square

Let T be a subsemigroup of a commutative cancellative semigroup S and let $h \in \text{Hom}(T, R_+^0)$. We say that $\langle T, h \rangle$ satisfies the β -condition (boundedness condition) in S if, for each $a \in S$, there is an $M \in R_+^0$ such that

$$(\beta) \quad x, y \in T, m \in Z_+^0 \text{ and } y + m \cdot a \in x + S \text{ implies } h(x) - h(y) \leq m \cdot M.$$

Here M is required to be independent of x, y and m . The notation $0 \cdot a + y$ expresses y itself, and hence the \mathcal{B} -condition implies the \mathcal{K} -condition. The \mathcal{B} -condition is equivalent to the combination of the \mathcal{K} -condition and the following:

For each $a \in S$, the set
 $(\mathcal{B}') \quad \{m^{-1}[h(x) - h(y)]: x, y \in T, m \in Z_+, y + m \cdot a \in x + S\}$
 is bounded.

LEMMA 3.4. *The following are equivalent:*

(3.4.1) $\langle T, h \rangle$ satisfies the \mathcal{K} -condition in S .

(3.4.2) h is extended to $\bar{h} \in \text{Hom}(\bar{T}, \mathbb{R}_+^0)$.

(3.4.3) h is extended to $\tilde{h} \in \text{Hom}(\tilde{T}, \mathbb{R}_+^0)$.

PROOF. (3.4.1) \Rightarrow (3.4.2). This follows from Theorem 2.1 since T is cofinal in \bar{T} .

(3.4.2) \Rightarrow (3.4.3). Since \bar{T} is unitary cofinal in \tilde{T} by (1.2.4), \bar{h} can be extended to $\tilde{h} \in \text{Hom}(\tilde{T}, \mathbb{R}_+^0)$ by Corollary 2.6, and hence h is extended to \tilde{T} .

(3.4.3) \Rightarrow (3.4.1). This is obvious from the definition of \tilde{T} . \square

LEMMA 3.5. *Let T be a filter of S , $T \neq S$, and let $h \in \text{Hom}(T, \mathbb{R}_+^0)$. Let $a \in S \setminus T$ and $r \in \mathbb{R}_+^0$. Define $h_r: P_T(a) \rightarrow \mathbb{R}_+^0$ by*

$$h_r(x) = \begin{cases} m \cdot r + h(s) & \text{if } x = m \cdot a + s \text{ where } m \in Z_+, s \in T, \\ m \cdot r & \text{if } x = m \cdot a \text{ where } m \in Z_+. \end{cases}$$

Every extension of h to $P_T(a)$ is obtained as h_r for some $r \in \mathbb{R}_+^0$.

PROOF. Since the expression of x is unique, h_r is well defined. The proof of the lemma is easy. \square

THEOREM 3.6. *Let T be a filter of a commutative cancellative semigroup S , $T \neq S$, and let $h \in \text{Hom}(T, \mathbb{R}_+^0)$. Then the following are equivalent:*

(3.6.1) $\langle T, h \rangle$ satisfies the \mathcal{B} -condition.

(3.6.2) h can be extended to $\widetilde{U_T(a)}$ for each $a \in S \setminus T$.

(3.6.3) For each $a \in S \setminus T$, $\langle P_T(a), h_r \rangle$ satisfies the \mathcal{K} -condition in S for some $r \in \mathbb{R}_+^0$.

PROOF. (3.6.1) \Rightarrow (3.6.2). Choose $r \in \mathbb{R}_+^0$ such that

$$r \geq \sup \{m^{-1}[h(x) - h(y)]: y + m \cdot a \in x + S\}$$

and then define $\bar{h}: U_T(a) \rightarrow \mathbb{R}_+^0$ by

$$\bar{h}(b) = \begin{cases} h(b) & \text{if } b \in T, \\ \frac{m \cdot r + h(s) - h(t)}{n} & \text{if } b \in Q(a) \text{ and } n \cdot b + t = m \cdot a + s \\ & \text{for some } s, t \in T. \end{cases}$$

By the choice of r , $\bar{h}(b) \geq 0$ for all $b \in U_T(a)$. To show \bar{h} is well defined, let $n \cdot b + t = m \cdot a + s$ and $n_1 \cdot b + t_1 = m_1 \cdot a + s_1$ where $s, t, s_1, t_1 \in T$. Then we have $(mn_1) \cdot a + n_1 \cdot s + n \cdot t_1 = (m_1n) \cdot a + n \cdot s_1 + n_1 \cdot t$ which implies $mn_1 = m_1n$ and $n_1 \cdot s + n \cdot t_1 = n \cdot s_1 + n_1 \cdot t$ since T is a filter. Then it follows that \bar{h} is well defined. Next we show that \bar{h} is a homomorphism. If $b, c \in Q(a)$, then $n \cdot b + t = m \cdot a + s$, $k \cdot c + u = l \cdot a + v$ for some $t, s, u, v \in T$, $n, m, k, l \in Z_+$; so $(nk) \cdot (b + c) + k \cdot t + n \cdot u = (mk + ln) \cdot a + k \cdot s + n \cdot v$ which implies $\bar{h}(b) + \bar{h}(c) = \bar{h}(b + c)$. If $b \in Q(a)$ and $c \in T$, then $n \cdot b + t = m \cdot a + s$ and $n \cdot (b + c) + t = m \cdot a + s + n \cdot c$, so the same result follows, and we see $\bar{h}(b) + \bar{h}(c) = \bar{h}(b + c)$ for all $b, c \in U_T(a)$. Since $U_T(a)$ is unitary cofinal in $\widetilde{U_T(a)}$ by Lemma 3.3 and (1.2.4), \bar{h} can be extended to $\tilde{h} \in \text{Hom}(\widetilde{U_T(a)}, \mathbf{R}_+^0)$ by Corollary 2.6.

(3.6.2) \Rightarrow (3.6.1). Let $x, y \in T$ and assume $b + x = m \cdot a + y$ for some $b \in S$. Hence $b \in Q(a)$. By assumption, h is extended to $\tilde{h} \in \text{Hom}(\widetilde{U_T(a)}, \mathbf{R}_+^0)$, and $m \cdot \tilde{h}(a) + h(y) - h(x) = \tilde{h}(b) \geq 0$ which implies the conclusion.

(3.6.2) \Rightarrow (3.6.3). By Lemma 3.3, $\widetilde{U_T(a)} = \widetilde{P_T(a)}$. Let \tilde{h} be the extension of h to $\widetilde{U_T(a)}$. Then $\tilde{h}|_{P_T(a)} = h_r$ for some $r \in \mathbf{R}_+^0$ by Lemma 3.5. By Lemma 3.4, $\langle P_T(a), h_r \rangle$ satisfies the K -condition.

(3.6.3) \Rightarrow (3.6.2). Again use Lemma 3.3 and Lemma 3.4. \square

4. Examples. Examples 4.1, 4.2 and 4.3 show that the K -condition does not imply the B -condition; Theorem 2.1 is not true in general if T_0 is not cofinal.

EXAMPLE 4.1. A commutative cancellative idempotent-free semigroup S is defined by

$$S = \{(x, y): y \in Z_+ \text{ if } x = 0, y \in Z \text{ if } x \in Z_+\}$$

in which the operation is

$$(x, y) + (z, u) = (x + z, y + u).$$

Let $T_0 = \{(0, y): y \in Z_+\}$. T_0 is not cofinal in S . Define $h_0 \in \text{Hom}(T_0, \mathbf{R}_+^0)$ by $h_0(0, y) = y$. Suppose h_0 is extended to $h \in \text{Hom}(S, \mathbf{R}_+^0)$. Let $x_0 \in Z_+$ be fixed and let $\lambda = h(x_0, 0)$, $\lambda \in \mathbf{R}_+^0$. Choose $y \in Z_+$ such that $y > \lambda$. Then $h(x_0, -y) + h(0, y) = h(x_0, 0) = \lambda$, hence $h(x_0, -y) = \lambda - y \geq 0$. This is a contradiction. Therefore h_0 cannot be extended to any element of $\text{Hom}(S, \mathbf{R}_+^0)$.

Let $(b, c) \in S \setminus T_0$ be fixed. Take arbitrarily $x, y, m \in Z_+$. Let $p = mb$, and $q = mc + y - x$. Then $p \in Z_+$, $q \in Z$ and $(0, x) + (p, q) = m \cdot (b, c) + (0, y)$. Since $(x - y)/m = (h_0(0, x) - h_0(0, y))/m$ can be arbitrarily large, $\langle T_0, h_0 \rangle$ does not satisfy the \mathcal{B} -condition. Since T_0 is a filter of S , the \mathcal{K} -condition is satisfied by $\langle T_0, h_0 \rangle$.

EXAMPLE 4.2. Let $S = \{(x, y): x \in Z_+^0, y \in Z, y \geq 1 - x^2\}$ and define T_0 and h_0 by

$$T_0 = \{(0, y): y \in Z_+\}, \quad h_0(0, y) = y.$$

T_0 is not cofinal but is a filter in S . Suppose that h_0 can be extended to a homomorphism h of S into R_+ . For each $n \in Z_+^0$, let

$$\varphi(n) = h_0(n, 1 - n^2).$$

Then $\varphi(1) + \varphi(n - 1) - \varphi(n) = h_0(0, 2n - 1) = 2n - 1$ for each $n \in Z_+$. From this recurrence relation, we have

$$n\varphi(1) + \varphi(0) - \varphi(n) = \sum_{i=1}^n (2i - 1) = n(n + 1) - n = n^2.$$

Since $\varphi(0) = h(0, 1) = 1$, it follows that $n\varphi(1) - \varphi(n) = n^2 - 1$. By the assumption $\varphi(n) \geq 0$ for all $n \in Z_+$, we have

$$n\varphi(1) \geq n^2 - 1 \quad \text{for all } n \in Z_+$$

hence $\varphi(1) \geq n - 1/n$ for all $n \in Z_+$. This is impossible. It follows that h_0 cannot be extended to an element of $\text{Hom}(S, R_+^0)$. We show that the \mathcal{B} -condition is not satisfied. Let $m \in Z_+$, $m > 1$, and choose $y, z \in Z_+$ such that $z - y = m^2 - 1$. Then

$$(0, z) + (m, 1 - m^2) = m \cdot (1, 0) + (0, y)$$

but $(z - y)/m = m - 1/m$ can be taken arbitrarily large.

EXAMPLE 4.3. Let π be the transcendental real number and let $a = \pi/4$. Then $0 < \pi/4 < 1$, and a is transcendental over the field R of rational numbers. If $a_k = a^k$ ($k = 1, 2, \dots$) (a^k is the usual k th power of a), then $1, a_1, a_2, \dots$ are linearly independent over R and $0 < a_k < 1$ ($k = 1, 2, \dots$). Let T_0 be the additive semigroup of R_+ generated by a_1, \dots, a_k, \dots . T_0 is actually a free commutative semigroup over a_1, \dots, a_k, \dots . Let $b_k = 1 - a_k > 0$ ($k = 1, 2, \dots$), and let S be the subsemigroup of R_+ generated by T and $b_1, b_2, \dots, b_k, \dots$. T_0 is a filter of S . Define $h_0: T_0 \rightarrow R_+$ by the homomorphism given by $h_0(a_k) = k$. Since T_0 is free, h_0 is well defined. Then h_0 cannot be extended to $h \in \text{Hom}(S, R_+^0)$. For suppose h_0 is extended to h . Then $1 = a_k + b_k \in S$, which implies $h(1) = h(a_k + b_k) \geq h(a_k) = k$ for all k . This is a

contradiction. Finally we show that $\langle T_0, h_0 \rangle$ does not satisfy the \mathcal{B} -condition. Let $m \in Z_+$ and $a_k \in T$. As $a_k | 1, a_k | m$ and so $a_k | (m + a_i)$ in S for all $m \in Z_+$, all $a_i, a_k \in T$. Then

$$(h_0(a_k) - h_0(a_i))/m = (k - i)/m$$

is not bounded.

ACKNOWLEDGEMENT. The authors thank the referee for his kind suggestion on this paper.

ADDENDUM. The assumption of cancellation does not restrict our discussion by the following reason. Let S be a commutative semigroup, S_0 the greatest cancellative homomorphic image of S , and $g_0: S \rightarrow S_0$ the homomorphism. If f is a homomorphism of S_0 into R_+^0 , then $h = fg_0$ is a homomorphism of S into R_+^0 . Every homomorphism h of S into R_+^0 can be obtained in this manner. Accordingly the results in this paper are extended to the case in which cancellation is not assumed.

REFERENCES

1. A. H. Clifford and G. B. Preston, *The algebraic theory of semigroups*, vol. II, Math. Surveys, no. 7, Amer. Math. Soc., Providence, R. I., 1967. MR 36 #1558.
2. Y. Kobayashi, *Homomorphisms on N -semigroups into R_+ and the structure of N -semigroups*, J. Math. Tokushima Univ. 7 (1973), 1–20. MR 48 #3831.
3. M. Petrich, *Introduction to semigroups*, Merrill, Columbus, Ohio, 1973.
4. M. S. Putcha, *Positive quasi-orders on semigroups*, Duke Math. J. 40 (1973), 857–869. MR 49 #2998.
5. T. Tamura and H. B. Hamilton, *The study of commutative semigroups with greatest group homomorphism*, Trans. Amer. Math. Soc. 175 (1972), 401–419. MR 47 #3581.

DEPARTMENT OF MATHEMATICS, NORTH CAROLINA STATE UNIVERSITY,
RALEIGH, NORTH CAROLINA 27607

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, DAVIS, CALIFORNIA 95616