

ON THE ZEROS OF STIELTJES AND VAN VLECK POLYNOMIALS

BY

NEYAMAT ZAHEER AND MAHFOOZ ALAM⁽¹⁾

DEDICATED TO THE UNIVERSITY OF WISCONSIN-MILWAUKEE DISTINGUISHED PROFESSOR
MORRIS MARDEN ON THE OCCASION OF HIS RETIREMENT ON MAY 9, 1975

ABSTRACT. Stieltjes and Van Vleck polynomials arise in the study of the polynomial solutions of the generalized Lamé differential equation. Our object is to generalize a theorem due to Marden on the location of the zeros of Stieltjes and Van Vleck polynomials. In fact, our generalization is two-fold: Firstly, we employ sets which are more general than the ones used by Marden for prescribing the location of the complex constants occurring in the Lamé differential equation; secondly, Marden deals only with the standard form of the said differential equation, whereas our result is equally valid for yet another form of the same differential equation. The part of our main theorem concerning Stieltjes polynomials may also be regarded as a generalization of Lucas' theorem to systems of partial fraction sums.

1. Introduction. A *generalized Lamé differential equation* is a second order linear differential equation of the form

$$(1.1) \quad \frac{d^2 w}{dz^2} + \left[\frac{\sum_{j=1}^p \alpha_j}{(z - a_j)} \right] \cdot \frac{dw}{dz} + \frac{\Phi(z)}{\prod_{j=1}^p (z - a_j)} \cdot w = 0,$$

where $\Phi(z)$ is a polynomial of degree at most $(p - 2)$ and α_j, a_j are complex constants. Heine [3] showed that there exist at most $C(n + p - 2, p - 2)$ polynomials $V(z)$ such that for $\Phi(z) = V(z)$ equation (1.1) has a polynomial solution $S(z)$ of degree n . Each polynomial $V(z)$ and the corresponding polynomial $S(z)$, associated with the differential equation (1.1), are, respectively, called [1, p. 37] a *Van Vleck polynomial* and a *Stieltjes polynomial*.

We shall be primarily interested in determining the location of the zeros of the systems of polynomials that arise in the study of the polynomial solutions of the differential equation

Received by the editors March 17, 1975 and, in revised form, December 8, 1975.

AMS (MOS) subject classifications (1970). Primary 30A08; Secondary 33A70.

Key words and phrases. Generalized Lamé differential equations, Stieltjes polynomials, Van Vleck polynomials.

⁽¹⁾ Work of the second author was supported by the University Grants Commission (Government of India) Research Fellowship.

$$(1.2) \quad \frac{d^2 w}{dz^2} + \left[\sum_{j=1}^p \alpha_j \left\{ \frac{\prod_{i=1}^{n_j-1} (z - b_{ji})}{\prod_{s=1}^{n_j} (z - a_{js})} \right\} \right] \cdot \frac{dw}{dz} + \left[\frac{\Phi(z)}{\prod_{j=1}^p \prod_{s=1}^{n_j} (z - a_{js})} \right] \cdot w = 0,$$

where $\Phi(z)$ is a polynomial of degree at most $(n_1 + n_2 + \dots + n_p - 2)$ and a_{js} , b_{ji} and α_j are suitable complex constants. We may, however, note that the differential equation (1.2) can always be written in the form (1.1) by expressing each fraction (in the coefficient of dw/dz) into its partial fractions, and that (1.2) is indeed of the form (1.1) in case $n_j = 1$ for all values of j . Consequently, there do exist Stieltjes polynomials $S(z)$ and Van Vleck polynomials $V(z)$ associated with the differential equation (1.2).

In order to avoid repetitions of long expressions and unnecessary details in the treatment of our proofs throughout the next section, we shall freely use the following abbreviations and notations: We shall denote by $S(z)$ the n th-degree Stieltjes polynomial corresponding to a Van Vleck polynomial $V(z)$ associated with the differential equation (1.2) or (1.1); the zeros of $V(z)$, $S(z)$, and $S'(z)$ will be denoted by t_j , z_j , and z'_j , respectively. We shall write

$$(1.3) \quad f_j(z) = \prod_{i=1}^{n_j-1} (z - b_{ji}), \quad g_j(z) = \prod_{s=1}^{n_j} (z - a_{js}), \quad h_j(z) = \frac{f_j(z)}{g_j(z)}$$

for every $j = 1, 2, \dots, p$ (with the convention that $f_j(z) \equiv 1$ for $n_j = 1$), and

$$(1.4) \quad F(z) = \sum_{j=1}^p \alpha_j \cdot h_j(z).$$

2. A generalization of Marden's theorem. In this section we generalize a theorem due to Marden ([1, Theorem (9, 1)] or [2]) on the location of the zeros of Stieltjes polynomials and of Van Vleck polynomials associated with the differential equation (1.1). Our generalization is two-fold: Firstly, we use sets which are more general than the ones used by Marden for prescribing the location of the points a_j (cf. (1.1)); secondly, Marden's theorem deals with the polynomials $S(z)$ and $V(z)$ in relation to (1.1) only, whereas our result is equally valid for both forms (1.1) and (1.2).

In order to prove our main theorem, we need to establish the following lemmas on the zeros of the polynomials $S(z)$ and $V(z)$.

LEMMA (2.1). *Every zero z_k of an n th-degree Stieltjes polynomial $S(z)$, associated with the differential equation (1.2), is either one of the points a_{js} ($1 \leq j \leq p$, $1 \leq s \leq n_j$) or satisfies the equation*

$$(2.1) \quad \frac{1}{2} F(z_k) + \sum_{j \neq k; j=1}^n \frac{1}{z_k - z_j} = 0,$$

where $F(z)$ is as defined by (1.4).

PROOF. Let $S(z) = \prod_{j=1}^n (z - z_j)$ be a Stieltjes polynomial corresponding to a Van Vleck polynomial $V(z)$. Then (cf. abbreviations (1.3)–(1.4)) the differential equation (1.2) can be written as

$$(2.2) \quad S''(z) + F(z) \cdot S'(z) + \left[\frac{V(z)}{\prod_{j=1}^p g_j(z)} \right] \cdot S(z) = 0.$$

If $z_k \neq a_{js}$ ($1 \leq j \leq p$, $1 \leq s \leq n_j$) is a zero of $S(z)$, then equation (2.2) becomes

$$(2.3) \quad S''(z_k) + F(z_k) \cdot S'(z_k) = 0.$$

If $z_k \neq a_{js}$, then two cases are to be considered:

Case 1. $S'(z_k) = 0$. Using equation (2.3) and successively differentiating equation (2.2) we observe that $S^{(m)}(z_k) = 0$ for all m , which contradicts the fact that the degree of $S(z)$ is n . Hence, this case is impossible to happen.

Case 2. $S'(z_k) \neq 0$. Since z_k is a nonrepeated zero of $S(z)$ in this case, we can write $S(z) = (z - z_k)T(z)$, where $T(z_k) \neq 0$. Therefore $S'(z_k) = T(z_k)$ and $S''(z_k) = 2T'(z_k)$. Consequently, due to (2.3), we have

$$-F(z_k) = \frac{S''(z_k)}{S'(z_k)} = 2 \frac{T'(z_k)}{T(z_k)} = 2 \cdot \sum_{j \neq k; j=1}^n \frac{1}{z_k - z_j}.$$

That is,

$$\frac{1}{2} \cdot F(z_k) + \sum_{j \neq k; j=1}^n \frac{1}{z_k - z_j} = 0.$$

This establishes our claim regarding the zeros of $S(z)$.

LEMMA (2.2). Every zero t_k of a Van Vleck polynomial $V(z)$, associated with the differential equation (1.2), if not an a_{js} is either one of the zeros of $S'(z)$ or satisfies the equation

$$(2.4) \quad F(t_k) + \sum_{j=1}^{n-1} \frac{1}{t_k - z'_j} = 0,$$

z'_j being the zeros of $S'(z)$.

PROOF. If t_k is a zero of $V(z)$ and if $S(z)$ is a corresponding Stieltjes polynomial, then from (1.2) we get

$$(2.5) \quad S''(t_k) + F(t_k) \cdot S'(t_k) = 0.$$

If $t_k \neq a_{js}$ and $S'(t_k) \neq 0$, then

$$\frac{S''(t_k)}{S'(t_k)} = \sum_{j=1}^{n-1} \frac{1}{(t_k - z'_j)},$$

where z'_j ($1 \leq j \leq n-1$) are the zeros of $S'(z)$. Thus, from (2.5) it follows that

$$-F(t_k) = \sum_{j=1}^{n-1} \frac{1}{t_k - z'_j}.$$

That is, t_k satisfies equation (2.4). The proof is now complete.

We now proceed to prove our main theorem.

THEOREM (2.3). *Let E be an ellipse with centre at the point c , semimajor axis as a , and eccentricity e . Let $0 \leq \gamma < \pi/2$ and*

$$\lambda = \max\{1, (1 - e^2/2)^{1/2} \sec \gamma\}.$$

If $|\arg \alpha_j| \leq \cos^{-1}(1/\lambda)$ for every $j = 1, 2, \dots, p$, and if all the points a_{js}, b_{jt} (occurring in equation (1.2)) lie on or inside E , then the zeros of each Stieltjes polynomial $S(z)$ and the zeros of each Van Vleck polynomial $V(z)$, associated with the differential equation (1.2), lie in the circular region

$$|z - c| \leq a \sec\left[\{(q-1)\pi + \cos^{-1}(1/\lambda)\} / (2q-1)\right],$$

where $q = \max\{n_1, n_2, \dots, n_p\}$.




FIGURE 1

PROOF. Let us draw a circle C with centre at the point c and radius as

$$R' = a \sec \left[\{ (q-1)\pi + \cos^{-1}(1/\lambda) \} / (2q-1) \right].$$

Suppose, on the contrary, that one or more zeros z_k of $S(z)$ lie outside the circle C and that z_1 is the one farthest from the centre c , so that all the zeros of $S(z)$ lie on or inside the circle C' given by $|z - c| = |z_1|$. Let θ, θ' , respectively, denote the acute angles which the tangents from z_1 to the ellipse E make with the tangent at the point z_1 on C' (see Figure 1). Now join z_1 to the centre c of the ellipse by a straight line, cutting the circle C at a point P (say). Now draw the tangents PA' and PB' from the point P to the ellipse E . Let ψ, ψ' (resp. φ, φ') denote, respectively, the angles which the tangents from the point z_1 (resp. P) to the ellipse make with the line joining the points z_1 and c (Figure 1). Since z_1 is outside C , we see that $\psi < \varphi$ and $\psi' < \varphi'$. Let φ_0 (resp. φ'_0) be the maximum value of φ (resp. φ') corresponding to all possible positions that z_1 could take on the circle C' . If ρ (resp. ρ') denotes the length of the perpendicular dropped from the centre c upon the tangent PA' (resp. PB'), we see that $\sin \varphi = \rho/R'$ (resp. $\sin \varphi' = \rho'/R'$) and that φ (resp. φ') takes a maximum value when $\rho = a$ (resp. $\rho' = a$). Therefore,

$$(2.6) \quad \sin \varphi_0 = \sin \varphi'_0 = \cos \{ (q-1)\pi + \cos^{-1}(1/\lambda) \} / (2q-1).$$

Now, the definition of λ implies that $0 \leq \cos^{-1}(1/\lambda) \leq \gamma < \pi/2$. Using these inequalities, we can easily verify that

$$(2.7) \quad 0 \leq \{ (q-1)\pi + \cos^{-1}(1/\lambda) \} / (2q-1) < \pi/2$$

and that

$$(2.8) \quad \frac{(q-1)\pi + \cos^{-1}(1/\lambda)}{(2q-1)} \geq \frac{(n_j-1)\pi + \cos^{-1}(1/\lambda)}{(2n_j-1)}, \quad 1 \leq j \leq p.$$

Since, $\psi < \varphi \leq \varphi_0$ and $\psi' < \varphi' \leq \varphi'_0$, from (2.6) and (2.7) we conclude that

$$\theta = (\pi/2 - \psi) > (\pi/2 - \varphi_0) = \{ (q-1)\pi + \cos^{-1}(1/\lambda) \} / (2q-1)$$

and

$$\theta' = (\pi/2 - \psi') > (\pi/2 - \varphi'_0) = \{ (q-1)\pi + \cos^{-1}(1/\lambda) \} / (2q-1).$$

Consequently, due to inequalities (2.8), we have

$$(2.9) \quad \theta, \theta' > \frac{(n_j-1)\pi + \cos^{-1}(1/\lambda)}{(2n_j-1)} = \mu_j \quad (\text{say}) \quad \forall j = 1, 2, \dots, p.$$

Let $S(z) = \prod_{j=1}^n (z - z_j)$. If any zero z_k of $S(z)$ is one of the points a_{js} , then it lies in C and we are done. However, if a zero z_k is none of the points a_{js} , then (cf. Lemma (2.1)) z_k satisfies equation (2.1). In particular (for $k = 1$), we have

$$\frac{1}{2} \cdot F(z_1) + \sum_{j=2}^n \frac{1}{z_1 - z_j} = 0.$$

Therefore,

$$\frac{1}{2} \cdot F(z_1) \cdot (z_1 - u_1) + \sum_{j=2}^n \left(\frac{z_1 - u_1}{z_1 - z_j} \right) = 0,$$

where u_1 is any point on the tangent drawn to the circle $|z - c| = |z_1|$ at the point z_1 , i.e.,

$$(2.10) \quad \sum_{j=1}^p \left[\frac{\alpha_j}{2} \cdot \prod_{t=1}^{n_j-1} \left(\frac{z_1 - b_{jt}}{z_1 - u_1} \right) \cdot \prod_{s=1}^{n_j} \left(\frac{z_1 - u_1}{z_1 - a_{js}} \right) \right] + \sum_{j=2}^n \left(\frac{z_1 - u_1}{z_1 - z_j} \right) = 0.$$

Since all the zeros z_k lie in $|z - c| < |z_1|$, we get

$$(2.11) \quad 0 < \arg \left(\frac{z_1 - u_1}{z_1 - z_j} \right) < \pi \quad \forall j = 2, 3, \dots, n.$$

Next, we also observe that the inequalities

$$(2.12) \quad -(\pi - \theta') < \arg \left(\frac{z_1 - b_{jt}}{z_1 - u_1} \right) < -\theta$$

and

$$(2.13) \quad \theta < \arg \left(\frac{z_1 - u_1}{z_1 - a_{js}} \right) < \pi - \theta'$$

hold for every $j = 1, 2, \dots, p$; $s = 1, 2, \dots, n_j$; $t = 1, 2, \dots, n_j - 1$. Inequalities (2.9), (2.12), and (2.13) imply that

$$-(n_j - 1)(\pi - \mu_j) < \arg \left[\prod_{t=1}^{n_j-1} \left(\frac{z_1 - b_{jt}}{z_1 - u_1} \right) \right] < -(n_j - 1)\mu_j$$

and

$$n_j \mu_j < \arg \left[\prod_{s=1}^{n_j} \left(\frac{z_1 - u_1}{z_1 - a_{js}} \right) \right] < n_j(\pi - \mu_j)$$

for all values of $j = 1, 2, \dots, p$. Using the above inequalities and the value of μ_j from (2.9), a simple calculation yields the following inequalities:

$$(2.14) \quad \begin{aligned} \cos^{-1}(1/\lambda) &< \arg \left[\prod_{t=1}^{n_j-1} \left(\frac{z_1 - b_{jt}}{z_1 - u_1} \right) \cdot \prod_{s=1}^{n_j} \left(\frac{z_1 - u_1}{z_1 - a_{js}} \right) \right] \\ &< \pi - \cos^{-1}(1/\lambda) \end{aligned}$$

for every $j = 1, 2, \dots, p$. In view of (2.14) and the hypothesis on α_j , we conclude that

$$(2.15) \quad 0 < \arg \left[\frac{\alpha_j}{2} \cdot \prod_{i=1}^{n_j-1} \left(\frac{z_1 - b_{ji}}{z_1 - u_1} \right) \cdot \prod_{s=1}^{n_j} \left(\frac{z_1 - u_1}{z_1 - a_{js}} \right) \right] < \pi$$

for every $j = 1, 2, \dots, p$. Hence, in the light of inequalities (2.11) and (2.15), the imaginary part of each term on the left-hand side of equation (2.10) is positive. This contradicts the fact that z_1 satisfies equation (2.10). Therefore, all the zeros of $S(z)$ lie on or inside the circle C .

Now, we prove the second part of our theorem for a Van Vleck polynomial $V(z)$ corresponding to an n th-degree Stieltjes polynomial $S(z)$. Proceeding as in the case of Stieltjes polynomials and assuming t_1 to be a zero of $V(z)$ lying outside the circle C and farthest from the centre c , we see that all the zeros of $V(z)$ lie on or inside the circle C' given by $|z - c| = |t_1|$. Our previous diagram (Figure 1) remains the same except that t_1 replaces z_1 . If $t_1 = a_{js}$, then t_1 is in E and, hence, in C . In case t_1 is a zero of $S'(z)$, then the Lucas theorem [1, Theorem (6, 2)], [4], [5], [6], together with the first part of our theorem (just proved), implies that t_1 lies in C and, hence, the theorem follows. In case t_1 is not a zero of $S'(z)$, then (cf. Lemma (2.2)) it satisfies equation (2.4) with $k = 1$. Hence

$$F(t_1) + \sum_{j=1}^{n-1} \frac{1}{t_1 - z'_j} = 0.$$

Therefore,

$$F(t_1) \cdot (t_1 - u_1) + \sum_{j=1}^{n-1} \frac{t_1 - u_1}{t_1 - z'_j} = 0,$$

where u_1 is any point on the tangent to the circle $|z - c| = |t_1|$ at the point t_1 , i.e.,

$$(2.16) \quad \sum_{j=1}^p (\alpha_j) \left\{ \prod_{i=1}^{n_j-1} \left(\frac{t_1 - b_{ji}}{t_1 - u_1} \right) \cdot \prod_{s=1}^{n_j} \left(\frac{t_1 - u_1}{t_1 - a_{js}} \right) \right\} + \sum_{j=1}^{n-1} \frac{t_1 - u_1}{t_1 - z'_j} = 0.$$

Since the points z'_j lie in C , we may replace z_j by z'_j and z_1 by t_1 in inequalities (2.11) and (2.15) and obtain

$$0 < \arg \left(\frac{t_1 - u_1}{t_1 - z'_j} \right) < \pi \quad \forall j = 1, 2, \dots, n-1,$$

and

$$0 < \arg \left[(\alpha_j) \cdot \prod_{i=1}^{n_j-1} \left(\frac{t_1 - b_{ji}}{t_1 - u_1} \right) \cdot \prod_{s=1}^{n_j} \left(\frac{t_1 - u_1}{t_1 - a_{js}} \right) \right] < \pi, \quad 1 \leq j \leq p.$$

Consequently, the imaginary part of each term on the left-hand side of equation (2.16) is positive, which contradicts the fact that t_1 satisfies equation (2.16). Hence, all the zeros of $V(z)$ also lie on or inside the circle C .

This completes the proof of Theorem (2.3).

An immediate consequence of Theorem (2.3) is the following result, exclusively in terms of the differential equation (1.1).

COROLLARY (2.4). *Let E be an ellipse with centre at the point c , semimajor axis as a , and eccentricity e . Let $0 \leq \gamma < \pi/2$ and*

$$\lambda = \max\{1, (1 - e^2/2)^{1/2} \sec \gamma\}.$$

If $|\arg \alpha_j| \leq \cos^{-1}(1/\lambda)$ for every $j = 1, 2, \dots, p$ and if all the points a_j ($1 \leq j \leq p$) lie on or inside E , then the zeros of each Stieltjes polynomial $S(z)$ and the zeros of each Van Vleck polynomial $V(z)$, associated with the differential equation (1.1), lie in the circular region $|z - c| \leq \lambda a$.

PROOF. In Theorem (2.3), if we take all n_j 's to be 1 (i.e. $q = 1$), then the differential equation (1.2) reduces to the differential equation (1.1), with the constants a_j corresponding to the constants a_{j1} , and Corollary (2.4) is then obvious.

The following corollary is a special case of Theorem (2.3) for circles.

COROLLARY (2.5). *If $|\arg \alpha_j| \leq \gamma < \pi/2$ for every $j = 1, 2, \dots, p$ and if all the points a_{js}, b_{jt} (occurring in the differential equation (1.2)) lie on or inside the circle $C: |z - c| = a$, then the zeros of each Stieltjes polynomial and the zeros of each Van Vleck polynomial, associated with the differential equation (1.2), lie in the circular region $|z - c| \leq a \sec\{[(q - 1)\pi + \gamma]/(2q - 1)\}$.*

PROOF. In Theorem (2.3), if we take $e = 0$ (i.e. if E is taken as the circle C), then $\lambda = \sec \gamma$, and the corollary follows at once.

If we put $q = 1$ in Corollary (2.5), we deduce the following well-known result due to Marden [1, Theorem (9.1)], [2], the proof being similar to that of Corollary (2.4).

COROLLARY (2.6). *In the differential equation (1.1), if $|\arg \alpha_j| \leq \gamma < \pi/2$ for every $j = 1, 2, \dots, p$ and if all the points a_j lie on or inside a circle $C: |z - c| = a$, then the zeros of each Stieltjes polynomial and the zeros of each Van Vleck polynomial, associated with the differential equation (1.1), lie on or inside a concentric circle $C': |z - c| = a \sec \gamma$.*

REMARK. It may be noted that the theorem as stated by Marden [1, Theorem (9, 1)], [2] is, in fact, the same as Corollary (2.6) for the case when the centre $c = 0$. But the same proof as given by him is valid also for the general case when the centre c is not necessarily at the origin.

As an application of Theorem (2.3), we prove

THEOREM (2.7). *Let $0 \leq \gamma < \pi/2$ and $\mu = \max\{1, (\sec \gamma)/\sqrt{2}\}$. If $|\arg \alpha_j| \leq \cos^{-1}(1/\mu)$ for every $j = 1, 2, \dots, p$ and if all the points a_{js}, b_{jt} (occurring in equation (1.2)) lie in the real interval $[-a, a]$, then the zeros of each Stieltjes polynomial $S(z)$ and the zeros of each Van Vleck polynomial $V(z)$, associated with the differential equation (1.2), lie in*

$$|z| \leq a \sec \left[\{(q-1)\pi + \cos^{-1}(1/\mu)\} / (2q-1) \right],$$

where $q = \max\{n_1, n_2, \dots, n_p\}$.

PROOF. Let S_e denote the closed interior of an ellipse E_e with centre at the origin, semimajor axis as a , and eccentricity e , and let

$$\lambda_e = \max \left\{ 1, (1 - e^2/2)^{1/2} \sec \gamma \right\}.$$

Then $[-a, a] = \bigcap_{e \in [0,1)} S_e$ and $(1 - e^2/2)^{1/2} \cdot \sec \gamma$ decreases continuously and monotonically to $(1/\sqrt{2}) \cdot \sec \gamma$ as e increases in $[0, 1)$. Obviously, then $[-a, a] \subseteq S_e$ for every $e \in [0, 1)$ and λ_e decreases continuously and monotonically to μ as e increases in $[0, 1)$ (easy to verify this statement for the cases when $\gamma \leq \pi/4$ and $\gamma > \pi/4$ respectively). Therefore,

$$(2.17) \quad |\arg \alpha_j| \leq \cos^{-1}(1/\mu) \leq \cos^{-1}(1/\lambda_e) \quad \forall e \in [0, 1)$$

and

$$(2.18) \quad a_{js}, b_{jt} \in S_e \quad \forall e \in [0, 1), \quad 1 \leq j \leq p; 1 \leq s \leq n_j; 1 \leq t \leq n_j - 1.$$

In view of (2.17) and (2.18), we can apply Theorem (2.3) and conclude that the zeros of each $S(z)$ and of each $V(z)$ lie in

$$|z| \leq a \sec \left[\frac{(q-1)\pi + \cos^{-1}(1/\lambda_e)}{(2q-1)} \right] = R_e \quad (\text{say}) \quad \forall e \in [0, 1).$$

Since λ_e decreases continuously and monotonically to μ as e increases in $[0, 1)$, we conclude that R_e decreases continuously and monotonically to

$$a \sec \left[\{(q-1)\pi + \cos^{-1}(1/\mu)\} / (2q-1) \right] = R \quad (\text{say}).$$

Finally, it is easy to see that the zeros of each $S(z)$ and those of each $V(z)$ lie in $|z| \leq R$. For, otherwise, it would contradict that the disc $|z| \leq R_e$ contains all the zeros of $S(z)$ and those of $V(z)$ for all values of e . This completes our proof.

The following corollary is an immediate consequence of the above theorem for the case when $q = 1$ (use the same explanation as in the proof of Corollary (2.4)).

COROLLARY (2.8). Let $0 < \gamma < \pi/2$ and $\mu = \max\{1, (1/\sqrt{2})\sec \gamma\}$. In equation (1.1), if $|\arg \alpha_j| < \cos^{-1}(1/\mu)$ for every $j = 1, 2, \dots, p$ and if all the points a_j lie in the real interval $[-a, a]$, then the zeros of each Stieltjes polynomial $S(z)$ and the zeros of each Van Vleck polynomial $V(z)$, associated with the differential equation (1.1), lie in $|z| < \mu a$.

Finally, it may be remarked that the part of our Theorem (2.3) that concerns Stieltjes polynomials may be regarded as a generalization of Lucas' theorem [1, Theorems (6, 1), (6, 1)', (6, 2)] to systems of partial fraction sums.

REFERENCES

1. M. Marden, *Geometry of polynomials*, 2nd ed., Math. Surveys, No. 3, Amer. Math. Soc., Providence, R. I., 1966. MR 37 #1562.
2. ———, *On Stieltjes polynomials*, Trans. Amer. Math. Soc. 33 (1931), 934–944.
3. E. Heine, *Handbuch der Kugelfunctionen*, Bd. I, 2nd ed., Springer, Berlin, 1878, pp. 472–476.
4. F. Lucas, *Propriétés géométriques des fractions rationnelles*, C. R. Acad. Sci. Paris 77 (1874), 431–433; *ibid.* 78 (1874), 140–144; *ibid.* 78 (1874), 180–183; *ibid.* 78 (1874), 271–274.
5. ———, *Géométrie des polynômes*, J. Ecole Polytech. (1) 46 (1879), 1–33.
6. ———, *Statique des polynômes*, Bull. Soc. Math. France 17 (1888), 2–69.

DEPARTMENT OF MATHEMATICS, ALIGARH MUSLIM UNIVERSITY, ALIGARH 202001, U. P., INDIA