

## ON THE EMBEDDING PROBLEM FOR 1-CONVEX SPACES

BY

VO VAN TAN<sup>1</sup>

**ABSTRACT.** In this paper we provide a necessary and sufficient condition for 1-convex spaces (i.e., strongly pseudoconvex spaces) which can be realized as closed analytic subvarieties in some  $\mathbb{C}^N \times \mathbb{P}_M$ . A construction of some normal 3-dimensional 1-convex space which cannot be embedded in any  $\mathbb{C}^N \times \mathbb{P}_M$  is given. Furthermore, we construct explicitly a non-kählerian 3-dimensional 1-convex manifold which answers a question posed by Grauert.

Unless otherwise specified, all  $\mathbb{C}$ -analytic spaces considered here will be noncompact, countable at infinity, reduced  $\mathbb{C}$ -analytic spaces of bounded Zariski dimension. Furthermore, the category of analytic coherent sheaves on a  $\mathbb{C}$ -analytic space  $X$  will be denoted by  $\text{Coh}(X)$ .

**0. Introduction.** On the one hand, it is well known that all Stein spaces can be embedded in  $\mathbb{C}^N$  and any compact  $\mathbb{C}$ -analytic space carrying a positive line bundle is embeddable into a complex projective space  $\mathbb{P}_M$  for arbitrary large integers  $N$  and  $M$ . On the other hand, topologically any 1-convex space is obtained by “welding” some compact analytic space onto some Stein space. Therefore, one naturally raises the question of embedding 1-convex spaces into  $\mathbb{C}^N \times \mathbb{P}_M$ . From now on, such 1-convex spaces will be called “embeddable 1-convex spaces”.

Our purpose here is twofold. First of all, we will provide a necessary and sufficient condition for embedding 1-convex spaces. Second, we will construct some 1-convex spaces (resp., 1-convex manifolds) which are not embeddable, therefore providing us some peculiar aspect of 1-convex spaces (resp., 1-convex manifolds).

The organization of this paper is as follows: In §I, all the basic definitions will be given. With those notions in hand, the statement of our problem will be formulated. Next, §II is devoted to the study of 1-convex spaces. The

---

Received by the editors May 25, 1978 and, in revised form, June 11, 1978.

*AMS (MOS) subject classifications* (1970). Primary 32E05, 32F10; Secondary 14F05, 32J20.

*Key words and phrases.* 1-convex spaces, blowing down process, positive line bundles, precise vanishing theorem, Moishezon spaces, compact cycle homologous to zero, kähler manifolds.

<sup>1</sup>The major part of this work was done while the author was supported by a C.N.R. fellowship at the University of Florence. He also gratefully acknowledges the hospitality of the University of Calabria.

construction of some nonembeddable 1-convex space (resp., nonembeddable 1-convex manifold) will be taken up in §III. We will end our discussion with some open problems.

### I. The context of the problem.

DEFINITION 1 [2]. Let  $S$  be a compact analytic subvariety in a  $\mathbb{C}$ -analytic space  $X$ . Then  $S$  is said to be *exceptional* if

- (i)  $\dim S_x > 0$  for all  $x \in S$ ,
- (ii) there exist a  $\mathbb{C}$ -analytic space  $Y$  and a proper, surjective and holomorphic map  $\mathbb{P}: X \rightarrow Y$ , inducing a biholomorphism  $X \setminus S \simeq Y \setminus T$ , where  $T$  consists of finitely many points and
- (iii)  $\mathbb{P}_* \mathcal{O}_X \simeq \mathcal{O}_Y$ .

EXAMPLE 1. Let  $X$  be the blowing up of  $\mathbb{C}^2$  at the origin and let  $S$  be the proper transform of the origin. Clearly,  $S \simeq \mathbb{P}^1$  is exceptional in  $X$ .

DEFINITION 2 [6a]. Let  $X$  be a given  $\mathbb{C}$ -analytic space with its exceptional subvariety  $S$ .  $X$  is said to be *1-convex* if  $Y$  is Stein. ( $Y$  is called the *Remmert reduction* of  $X$ . Sometimes we will use the notation  $(X, S)$  to denote 1-convex spaces.)

In Example 1,  $(X, S)$  is a 1-convex manifold, since  $Y \simeq \mathbb{C}^2$ . In fact, this example is the prototype for our next investigation.

REMARK. Notice that the definition of 1-convex spaces given here is not a standard one. However, it is known that, in fact, it is equivalent to the usual definitions (see [2] or [6a]). We adopt such a definition here because, as we will see, it is very convenient in our context.

DEFINITION 3 [2]. Let  $S$  be a compact  $\mathbb{C}$ -analytic space and let  $L$  be a holomorphic line bundle on  $S$ . Let us identify the zero section  $\Sigma$  of  $L$  with  $S$ . Then  $L$  is said to be *weakly negative* if  $\Sigma$  (as a compact analytic subvariety in  $L$ ) is exceptional.  $L$  is said to be *weakly positive* if  $L^*$  is weakly negative.

EXAMPLE. The hyperplane section bundle on  $\mathbb{P}_M$  is weakly positive.

DEFINITION 3' [0]. Let  $X$  be a  $\mathbb{C}$ -analytic space, let  $L$  be a holomorphic line bundle on  $X$  and let  $(U_i, e_{ij})$  be a system of 1-cocycles determining  $L$ . Then  $L$  is said to be *positive* if there exists a system  $\{h_i\}$  of smooth and positive functions on  $(U_i)$  such that on  $U_i \cap U_j$ ,

$$h_j = |e_{ij}|^2 h_i$$

and such that the functions  $g_i := -\log h_i$  are strongly pseudoconvex on  $U_i$ .

REMARK. For  $X$  compact, it has been proved that the notions of weakly positive and positive are equivalent (see [2]). In [6b] the relationship between these two concepts on 1-convex spaces is studied.

It is well known (see [7] and the references there) that any Stein space can be embedded in  $\mathbb{C}^N$ . Also, it is known (see [2]) that any compact  $\mathbb{C}$ -analytic space carrying a weakly positive line bundle can be embedded biholomorphically into some  $\mathbb{P}_M$ . Therefore, one is led to:

*Problem.* When is it possible to embed a given 1-convex space into  $\mathbb{C}^N \times \mathbb{P}_M$ ?

Notice that in our Example 1, in view of the definition of the blow-up, the 2-dimensional 1-convex manifold

$$X = \{(x_0, x_1; z_0, z_1) \in \mathbb{C}^2 \times \mathbb{P}_1 \mid x_0 z_1 - x_1 z_0 = 0\}$$

is actually a closed submanifold in  $\mathbb{C}^2 \times \mathbb{P}_1$ .

**II. The embeddable 1-convex spaces.** The first result in this direction in the nonsingular case was established in [1].

**THEOREM 1 [1].** *Let  $X$  be a 1-convex manifold and let us assume that there exists a positive line bundle  $L$  on  $X$ . Then  $X$  is embeddable.*

By slightly modifying the proof in [1] and by using some standard techniques, we shall generalize Theorem 1 in two directions. First of all we do not require  $X$  to be nonsingular, and secondly the line bundle  $L$  does not need to be positive on the whole space  $X$ .

As a common philosophy in this kind of business, any embedding theorem is preceded, in general, by a vanishing theorem. So to begin, let us mention the following version.

**THEOREM 2 [0].** *Let  $X$  be a 1-convex space, let  $L$  be a positive line bundle on  $X$  and let  $\mathcal{F} \in \text{Coh}(X)$ . Then there exists an integer  $k_0 = k_0(L, \mathcal{F})$  such that*

$$H^i(X, \mathcal{F} \otimes L^k) = 0$$

*for all  $k > k_0$  and all  $i \geq 1$ .*

Actually in [0] (see also [1]), the previous result was proved for  $q$ -convex spaces, for any  $q > 1$ . However, in the special case of 1-convex spaces, Theorem 2 can be sharpened as follows.

**THEOREM I.** *Let  $(X, S)$  be a 1-convex space and let us assume that there exists a holomorphic line bundle  $L$  on  $X$  such that  $L|_S$  (sheaf restriction) is positive. Then, for any  $\mathcal{F} \in \text{Coh}(X)$ , there exists an integer  $k_0 := k_0(L, \mathcal{F})$  such that*

$$H^i(X, L^k \otimes \mathcal{F}) = 0$$

*for all  $k > k_0$  and all  $i \geq 1$ .*

The proof of Theorem I is based on the following important result.

**EXTENSION LEMMA.** *Let  $(X, S)$  and  $L$  be as in the hypothesis of Theorem I. Then, after modifying the metric of  $L$ , one can find a 1-convex neighborhood  $\Xi$  of  $S$ , with  $\Xi \subset X$ , such that  $L|_\Xi$  is positive.*

**PROOF.** We shall denote the Zariski tangent space of  $X$  at a point  $x \in X$  by  $T_{X,x}$ . Let  $(h_i, U_i)$  be the metric associated to the line bundle  $L$ . Since by

hypothesis,  $L|S$  is positive, i.e. for all  $x \in V_i := U_i \cap S$ ,

$$-\partial\bar{\partial}\log h_i(x) > 0 \quad \text{on } T_{V_i, x}. \quad (1)$$

In view of Definition 2 for 1-convex spaces, one can find a smooth function  $\Psi$  on  $X$  such that

$$\partial\bar{\partial}\Psi(x) \geq 0 \quad \text{on } T_{X, x} \quad \text{for } x \in X, \quad (2)$$

$$\partial\bar{\partial}\Psi(x) > 0 \quad \text{on } T_{X, x} \quad \text{for } x \in X \setminus S, \quad (3)$$

$$\partial\bar{\partial}\Psi(x) > 0 \quad \text{on } N_x \quad \text{for } x \in S, \quad (4)$$

where  $N_x$  is the complementary space of  $T_{S, x}$  in  $T_{X, x}$ .

Now on each open covering  $U_i$  with  $U_i \cap S \neq \emptyset$  and for any integer  $k_i$  the smooth function

$$A_i := h_i e^{-k_i \Psi}: U_i \rightarrow \mathbb{R}^+$$

is well defined.

In view of (1), (2) and (4), one can choose a  $k_i \gg 0$ , such that

$$-\partial\bar{\partial}\log A_i(x) > 0 \quad \text{on } T_{U_i, x} \quad \text{for all } x \in V_i. \quad (5)$$

Meanwhile, in view of (3), again with a suitable  $k_i \gg 0$ , one has

$$-\partial\bar{\partial}\log A_i(x) > 0 \quad \text{on } T_{U_i, x} \quad \text{for } x \in U_i \setminus V_i. \quad (6)$$

In view of the compactness of  $S$ , (5) and (6), an integer  $k := \max_i k_i$  can be selected such that, on the relative compact neighborhood  $N := \bigcup_i U_i$  of  $S$  in  $X$ ,

$$-\partial\bar{\partial}\log(h_i e^{-k\Psi}) > 0.$$

In other words, with the new metric  $g_i := h_i e^{-k\Psi}$ ,  $L|N$  is positive. Now, since  $S$  is exceptional, it admits a fundamental system of 1-convex neighborhoods; let  $\Xi$  be one of them such that  $\Xi \subset N$ . Clearly,  $L|\Xi$  is positive. Q.E.D.

**PROOF OF THEOREM I.** In view of the Extension Lemma and Theorem 2 above, it suffices to prove that the restriction map

$$\lambda_i: H^i(X, \mathcal{F}) \rightarrow H^i(\Xi, \mathcal{F}) \quad (*)$$

is injective for all  $i \geq 1$  and  $\mathcal{F} \in \text{Coh}(X)$ .

In fact our present situation can be summarized by the following diagram

$$\begin{array}{ccccc} S & \hookrightarrow & \Xi & \hookrightarrow & X \\ \downarrow & & \downarrow & & \downarrow \P \\ T & \hookrightarrow & V & \hookrightarrow & Y \end{array}$$

where  $Y$ ,  $T$  and  $\P$  are as in Definition 2 and  $V$  is an arbitrary, small Stein neighborhood (disconnected) of  $T$  in  $Y$ . In fact without loss of generality, one can choose  $V$  such that  $\Xi = \P^{-1}(V)$ .

Now notice that  $X = (X \setminus S) \cup \Xi$  and  $(X \setminus S) \cap \Xi = \Xi \setminus S$ . Similarly  $Y = (Y \setminus T) \cup V$  and  $(Y \setminus T) \cap V = V \setminus T$ . Hence we can apply the standard technique of Mayer-Vietoris exact sequences in this context; namely let us look at the following commutative diagrams with exact rows.

$$\begin{array}{ccccccc}
 \cdots & \longrightarrow & H^i(\Xi, \mathcal{F}) \oplus H^i(X \setminus S, \mathcal{F}) & \xrightarrow{\delta_i} & H^i(\Xi \setminus S, \mathcal{F}) & \xrightarrow{\epsilon_i} & \cdots \\
 & & \uparrow \alpha_i & & \uparrow \beta_i & & \uparrow \gamma_i \\
 \cdots & \longrightarrow & H^i(V, \hat{\mathcal{F}}) \oplus H^i(Y \setminus T, \hat{\mathcal{F}}) & \xrightarrow{\delta'_i} & H^i(V \setminus T, \hat{\mathcal{F}}) & \longrightarrow & \cdots \\
 & & & & & & \\
 \xrightarrow{\epsilon_i} & H^{i+1}(X, \mathcal{F}) & \xrightarrow{\theta_{i+1}} & H^{i+1}(\Xi, \mathcal{F}) \oplus H^{i+1}(X \setminus S, \mathcal{F}) & \longrightarrow & & \\
 & & & \uparrow \alpha_{i+1} & & \uparrow \beta_{i+1} & \\
 \longrightarrow & H^{i+1}(Y, \hat{\mathcal{F}}) & \longrightarrow & H^{i+1}(V, \hat{\mathcal{F}}) \oplus H^{i+1}(Y \setminus T, \hat{\mathcal{F}}) & \longrightarrow & & 
 \end{array}$$

Now for  $i \geq 0$ ,  $H^{i+1}(Y, \hat{\mathcal{F}}) = 0$  since  $Y$  is Stein and  $\hat{\mathcal{F}} := \mathbb{P}_*(\mathcal{F}) \in \text{Coh}(Y)$ . Hence  $\delta'_i$  is surjective. Furthermore,  $\gamma_i$  is bijective, in view of (ii) in Definition 1. Therefore  $\delta_i$  is also surjective. Consequently  $\epsilon_i$  is a zero map and this implies that  $\theta_{i+1}$  is injective. But the latter map factors via  $\lambda_{i+1}$  which is therefore injective. Hence (\*) is proved. Q.E.D.

REMARKS. (a) The theorem above is often alluded to as the *imprecise vanishing theorem* since it holds in general for some power  $k \gg 0$ . However, if one is willing to deal only with C-analytic manifold, then the so-called *precise vanishing theorem* can be obtained (see [6a] for complete proof and more related results), namely:

PROPOSITION 1. *Let  $X$  be a 1-convex manifold and let  $L$  be a holomorphic line bundle on  $X$  such that  $L|_S$  is weakly positive ( $S$  is singular in general). Then*

$$H^p(X, \Omega^q(L)) = 0 \quad \text{for all } p + q \geq \dim X + 1.$$

(b) Actually, there is more than one way to prove Theorem I above. Another proof can be found in [6b] where a direct argument is used in order to avoid a detour through Theorem 2.

Using the standard technique to derive Theorem A of Cartan from his Theorem B, we can deduce from our previous Theorem I the following:

THEOREM II. *Let  $(X, S)$  and  $L$  be as in Theorem I. Then, for any  $x \in X$ , there exists an integer  $k_x$ , such that the stalk  $(L^k \otimes \mathcal{F})_x$  is generated by its global sections for any  $k \geq k_x$ .*

PROOF. Let  $x \in X$  and let  $I_x$  be the ideal sheaf of germs of holomorphic functions vanishing at  $x$ . Then for any  $\mathcal{F} \in \text{Coh}(X)$ , one has the following exact sequence:

$$0 \rightarrow L^k \otimes I_x \mathcal{F} \rightarrow L^k \otimes \mathcal{F} \rightarrow L^k \otimes \mathcal{F}/I_x \mathcal{F} \rightarrow 0,$$

which in turn induces the following exact sequence:

$$\cdots \rightarrow H^0(X, L^k \otimes \mathcal{F}) \xrightarrow{\alpha} H^0(X, L^k \otimes \mathcal{F}/I_x \mathcal{F}) \rightarrow H^1(X, L^k \otimes I_x \mathcal{F}).$$

In view of Theorem I, there exists an integer  $k_x$ , such that for all  $k \geq k_x$ ,  $H^1(X, L^k \otimes I_x \mathcal{F}) = 0$ . Therefore  $\alpha$  is surjective. Nakayama's lemma tells us that  $(L^k \otimes \mathcal{F})_x$  is generated by its global sections. Q.E.D.

REMARK. Theorem II was also proved in [1]. Our proof here is simpler. From Theorem II, we can deduce the following useful result.

COROLLARY 1. *Let  $(X, S)$ ,  $L$  and  $\mathcal{F}$  be as in Theorem I. Then for any relative compact domain  $D$  in  $X$ , there exist finitely many global sections  $\{f_i\} \in H^0(X, L^k \otimes \mathcal{F})$ , with  $k \gg 0$ , such that the stalk  $(L^k \otimes \mathcal{F})_x$  is generated by those  $\{f_i\}$  for any  $x \in D$ .*

We are now in a position to state the main result of this section.

THEOREM III. *Let  $(X, S)$  be a given 1-convex space. There exists a line bundle  $L$  on  $X$  such that  $L|S$  is positive if and only if  $X$  is embeddable.*

The following result will be needed later.

LEMMA 1 [1]. *Let  $X$  be a Stein space and let  $L$  be a holomorphic line bundle on  $X$ . Then there exist finitely many global sections  $f_1, \dots, f_m \in H^0(X, L)$  such that the set  $\{x \in X | f_1(x) = \dots = f_m(x) = 0\}$  is empty.*

PROOF OF THEOREM III. (i) *Sufficiency.*

(a) Let  $D$  be a relative compact domain in  $X$  such that  $S \subset D$ . For any point  $x \in X$  and any integer  $r$ , one has the following exact sequence:

$$0 \rightarrow I_x^2 \otimes L' \rightarrow I_x \otimes L' \rightarrow I_x/I_x^2 \otimes L' \rightarrow 0.$$

In view of the compactness of  $D$ , Theorem I tells us that there exists an integer  $r_D$  such that the differential map

$$H^0(X, I_x \otimes L') \rightarrow I_x/I_x^2 \otimes L'_x \quad (*)$$

is surjective for any point  $x \in D$  and any integer  $r \geq r_D$ .

Similarly, by considering the following exact sequence:

$$0 \rightarrow I_{x,y} \otimes L^s \rightarrow L^s \rightarrow L_x^s \oplus L_y^s \rightarrow 0,$$

one can prove that there exist an integer  $s_D$  such that the restriction map

$$H^0(X, L^s) \rightarrow L_x^s \oplus L_y^s \quad (**)$$

is surjective for any points  $x \neq y \in D$  and any integer  $s \geq s_D$ .

Now let  $t_D := r_D \cdot s_D$ . Corollary 1 tells us that there exists an integer  $p$  which is a multiple of  $t_D$  and sections  $f_0, \dots, f_q \in H^0(X, L^p)$  which give rise to the well-defined holomorphic map

$$\Sigma := (f_0, \dots, f_q): D \rightarrow \mathbf{P}_q.$$

In view of (\*) and (\*\*),  $\Sigma$  is clearly regular and injective.

(b) Now let  $Z := \{x \in X | f_0(x) = \dots = f_q(x) = 0\}$ . From the construction of the  $f_i$ , clearly  $Z \cap S = \emptyset$ , i.e.,  $Z$  is a Stein space. Let  $J$  be the ideal sheaf determined by  $Z$ , then Theorem I tells us that there exists an integer  $k_0(L, J)$  such that  $H^1(X, L^k \otimes J) = 0$  for all  $k \geq k_0$ . Therefore the map

$$\Lambda: H^0(X, L^k) \rightarrow H^0(Z, O_Z \otimes L^k)$$

is surjective. In view of Lemma 1, there exist global sections  $g_1, \dots, g_r \in H^0(Z, O_Z \otimes L^k)$  which do not have any common zeroes in  $Z$ . From the surjectivity of  $\Lambda$  one can assume that  $g_i \in H^0(X, L^k)$ . Clearly

$$\Phi := (f_0^k, \dots, f_q^k; g_1^p, \dots, g_r^p): X \rightarrow \mathbf{P}_M$$

is a well-defined holomorphic map which embedded  $D$  biholomorphically into  $\mathbf{P}_M$  as a locally closed subspace where  $M := q + r$ .

(c) Let  $Y$  be the Remmert reduction of  $X$  and let  $\rho$  be the map which embeds  $Y$  into some  $\mathbb{C}^n$  [7]. Let us look at the following diagram.

$$\begin{array}{ccc} X & \xrightarrow{\Phi} & \mathbf{P}_M \\ \pi \swarrow & \searrow \Upsilon & \uparrow \Pi_2 \\ Y & & \mathbb{C}^N \times \mathbf{P}_M \\ \rho \searrow & \swarrow \Pi_1 & \\ \mathbb{C}^N & & \end{array}$$

(Note: The diagram shows a commutative diagram with  $X$  at the top left,  $\mathbf{P}_M$  at the top right,  $\mathbb{C}^N$  at the bottom left, and  $\mathbb{C}^N \times \mathbf{P}_M$  at the bottom right. Arrows are:  $X \xrightarrow{\Phi} \mathbf{P}_M$ ,  $X \xrightarrow{\Psi} \mathbb{C}^N$ ,  $X \xrightarrow{\Upsilon} \mathbb{C}^N \times \mathbf{P}_M$ ,  $\mathbf{P}_M \xrightarrow{\Pi_2} \mathbb{C}^N \times \mathbf{P}_M$ ,  $\mathbb{C}^N \xrightarrow{\Pi_1} \mathbb{C}^N \times \mathbf{P}_M$ ,  $Y \xleftarrow{\pi} X$ ,  $Y \xrightarrow{\rho} \mathbb{C}^N$ , and  $\mathbb{C}^N \times \mathbf{P}_M \xrightarrow{\Upsilon} X$ .)

Clearly the composed map  $\Psi := \rho \circ \pi$  is a proper holomorphic map which is biholomorphic outside of  $S$ . Consequently the map  $\Upsilon := \Psi \times \Phi: X \rightarrow \mathbb{C}^N \times \mathbf{P}_M$  is proper since  $\Psi$  is and since  $\Upsilon = \Pi_1 \circ \Upsilon$ . Moreover,  $\Psi$  (resp.,  $\Phi$ ) is injective and regular on  $X \setminus S$  (resp.,  $D$ ); one can then easily check, using the commutativity of the diagrams, that  $\Upsilon$  is also injective and regular on  $X = (X \setminus S) \cup D$ .

(ii) *Necessity*. Let us look at the following diagram

$$\begin{array}{ccc} X & \xrightarrow{\iota} & \mathbb{C}^N \times \mathbf{P}_M \\ \uparrow & & \downarrow \Pi_2 \\ S & \xrightarrow{\quad} & \mathbf{P}_M \end{array} \quad \begin{array}{c} \text{H} \\ \swarrow \end{array}$$

(Note: The diagram shows a commutative diagram with  $X$  at the top left,  $\mathbb{C}^N \times \mathbf{P}_M$  at the top right,  $S$  at the bottom left, and  $\mathbf{P}_M$  at the bottom right. Arrows are:  $X \xrightarrow{\iota} \mathbb{C}^N \times \mathbf{P}_M$ ,  $S \xrightarrow{\quad} \mathbf{P}_M$ ,  $X \xrightarrow{\quad} S$ ,  $\mathbb{C}^N \times \mathbf{P}_M \xrightarrow{\Pi_2} \mathbf{P}_M$ , and  $\mathbf{P}_M \xrightarrow{\quad} \text{H}$ .)

where the maps (except  $\Pi_2$ ) are embedding maps. Let  $\mathbf{H}$  be the hyperplane section bundle on  $\mathbf{P}_M$  and let us lift  $\mathbf{H}$ , via  $\Pi_2$ , to a holomorphic line bundle

$E := \Pi_2^*(H)$  on  $C^N \times P_M$ . Now let us pull back  $E$  to obtain a line bundle  $L := \iota^*E$  on  $X$ . By construction, clearly,  $L|S \simeq H|S$  is positive. Q.E.D.

REMARKS. (a) With some further work, the Extension Lemma can be strengthened as follows (see [6b] for complete proof).

**THEOREM IV.** *Let  $(X, S)$  be a given 1-convex space and let  $L$  be a line bundle on  $X$  such that  $L|S$  is weakly positive. By modifying the metric,  $L$  is actually positive on all of  $X$ .*

(b) In order to illustrate the argument in Part (i) (a) and (b) above, let us consider the following example.

**EXAMPLE 2.** Let  $F \simeq P_1$  be a line in  $P_2$  and let us take a point  $x \notin F$ . By blowing up  $P^2$  at  $x$ , one obtains a 2-dimensional projective manifold  $W$ . Let  $F'$  (resp.,  $S$ ) be the proper transform of  $F$  (resp.,  $x$ ). Then clearly  $X := W \setminus F'$  is a 2-dimensional 1-convex manifold with its exceptional subvariety  $S$ . It is well known that one can embed  $W$  biholomorphically into  $P_5$ . Therefore  $X$  is biholomorphic to a Zariski open submanifold in  $P_5$ .

**III. The nonembeddable 1-convex spaces.** Our previous Theorem III suggests the following:

**Problem 0.** Let  $(X, S)$  be a given 1-convex space. Is it always possible to find a holomorphic line bundle  $L$  on  $X$  such that  $L|S$  is positive?

We are going to tackle this problem following two simple observations:

(A) Let  $(X, S)$  be an embeddable 1-convex space. Then necessarily  $S$  is projective algebraic.

(B) Let  $X$  be an embeddable 1-convex manifold. Then necessarily  $X$  is kählerian.

Hence these two facts lead us to the following:

**Question A.** Do there exist 1-convex spaces  $(X, S)$  such that the exceptional subvariety  $S$  is not projective algebraic?

**Question B.** Do there exist non-kählerian 1-convex manifolds?

(Questions A and B were first raised to the author by H. Grauert.) In this section, we shall provide satisfactory answers for both Questions A and B as well as for Problem 0.

( $\hat{A}$ ) Construction of 3-dimensional normal 1-convex spaces  $(X, S)$  such that  $S$  is not projective.

First we shall need the following result:

**LEMMA 2.** *Let  $L$  be a holomorphic line bundle over a  $C$ -analytic manifold  $X$  and let us identify  $X$  with the zero section of  $L$ . Then there exists a canonical isomorphism (holomorphic) from the normal bundle of  $X$  in  $L$  onto the line bundle  $L$ .*

The proof of Lemma 2 is purely topologic so we leave it to the reader. Now we are in a position to begin our construction.

*Step 1.* Let us consider the following special projective algebraic 2-fold exhibited by Hironaka (see [4, Chapter V]). Let  $C$  be a nonsingular cubic curve in  $\mathbf{P}_2$ . It is known that  $C$  also acquires a group structure on its set of points. Fix an inflection point  $p_0 \in C$  to be the origin of the group law on  $C$ . Since the torsion-free part of the abelian group  $C$  has infinite rank, therefore one can select ten points, say  $p_1, \dots, p_{10} \in C$  which are linearly independent over  $\mathbf{Z}$  in the group law. Now blow up  $p_1, \dots, p_{10}$  in  $\mathbf{P}_2$  successively, let  $Z$  be the resulting manifold and let  $W$  be the strict transform of  $C$ . Since  $C^2 = 9$ , we have  $W^2 = 9 - 10 = -1$ . Following [2],  $W$  is exceptional in  $Z$ .




FIGURE 1

*Step 2.* Let us put on  $Z$  a weakly negative line bundle, say  $L$ , which always exists since  $Z$  is projective. In view of Lemma 2,  $N_{Z/L}$ , the normal bundle of  $Z$  in  $L$ , is also weakly negative. Following [2, Satz 5]  $L$  is a 1-convex manifold admitting  $Z$  as its exceptional subvariety. Furthermore, since  $W$  is exceptional in  $Z$  and  $\dim W = 1$ , a result in [2, Satz 9] tells us that the normal bundle  $N_{W/Z}$  of  $W$  in  $Z$  is actually weakly negative (see Figure 1).

*Step 3.* It is known that the line bundle  $L$  is weakly negative iff  $L^*$  is ample in the sense of Grothendieck [3]. Now, let us look at the following exact sequence of bundles on  $W$ :

$$0 \rightarrow N_{W/Z} \rightarrow N_{W/L} \rightarrow N_{Z/L}|_W \rightarrow 0 \quad (\dagger)$$

where  $N_{W/L}$  is the normal bundle of  $W$  in  $L$ . By dualizing  $(\dagger)$ , we obtain an exact sequence of bundles where the extreme terms  $N_{Z/L}^*$  and  $N_{W/Z}^*$  are ample. A result in [3] tells us that  $N_{W/L}^*$  is also ample, i.e.,  $N_{W/L}$  is weakly negative.

*Step 4.* Since  $N_{W/L}$  is weakly negative, following [2],  $W$  is exceptional in  $L$ . From Definition 1, this implies the existence of a  $\mathbb{C}$ -analytic space  $X$  and a birational morphism  $\mathbb{I}: L \rightarrow X$  inducing a biholomorphism

$$L \setminus W \simeq X \setminus \{\omega\} \quad (\dagger)$$

where  $\{\omega\}$  is a point in  $X$ .

Clearly  $X$  is a 3-dimensional normal  $\mathbb{C}$ -analytic space with only one isolated singular point  $\{\omega\}$ . Consequently, Riemann's extension theorem tells us that  $X$  is a holomorphically convex space since  $L$  is the one. Furthermore, in view of  $(\dagger)$ ,  $S := \mathbb{I}(Z)$  is the maximal compact subvariety in  $X$ , in the sense of [2]. From there, one can check that  $(X, S)$  is actually 1-convex in the sense of Definition 2.

*Step 5.*  $S$  is not projective (see [4]).

If it were, there would exist a curve, say  $D$ , on  $S$  with  $\omega \notin D$ . Consequently  $\mathbb{I}^{-1}(D) \subset Z$  would be a curve not intersecting  $W$  and the image  $D_0 := \theta(\mathbb{I}^{-1}(D))$  would be a curve in  $\mathbb{P}_2$  which does not meet  $C$  except at the points  $p_1, \dots, p_{10}$ , where  $\theta: Z \rightarrow \mathbb{P}_2$  is the blowing up map. Let  $d := \deg D_0$ . In view of Bezout's theorem  $D_0 \cdot C = 3d > 0$ . So one can write

$$D_0 \cap C = \sum_{i=1}^{10} n_i p_i \quad \text{on } C$$

with  $n_i \geq 0$  and  $\sum n_i = 3d$ . But  $D_0 \sim dL$  (linear equivalence) where  $L$  is a line in  $\mathbb{P}_2$  and  $L \cdot C \sim 3p_0$ , so one has  $\sum n_i p_i = 0$  in the group law on  $C$ . But this contradicts the linear independency of the points  $p_1, \dots, p_{10}$ .

(B) Construction of 3-dimensional non-kählerian 1-convex manifolds.<sup>2</sup>

*Step 1.* Let  $Z$  be the blowing up of  $\mathbb{C}^3$  at the origin and let  $M \simeq \mathbb{P}_2$  be the exceptional subvariety. Let  $K$  be a singular cubic curve with only one node  $\omega$ , embedded in  $\mathbb{P}_2$ . Certainly  $K \setminus \omega$  is smooth and there exists an open neighborhood  $U$  of  $\omega$  in  $Z$  such that  $K \cap U$  is a union of two smooth irreducible branches, say  $K_1$  and  $K_2$  which intersect at  $\omega$  with distinct tangents.

*Step 2.* We are going to use a basic idea which is due to Hironaka (see [4, Appendix B]). Let  $(\hat{U}, f)$  be the composite of two blowings up over  $U$  in which the first is the blowing up with center  $K_1$  and the second is the blowing up with center  $K'_2$  where  $K'_2$  denotes the proper transform of  $K_2$  by the first blowing up. Let  $(\hat{V}, g)$  be the blowing up over  $V := Z \setminus \omega$  with center  $K \setminus \omega$ . Now let  $\hat{S}_1 := f^{-1}(K_1 \cup K_2)$  and let  $\hat{S}_2 := g^{-1}(K \setminus \omega)$ .

*Step 3.* Notice that  $(\hat{U}, f)$  and  $(\hat{V}, g)$  agree on the inverse image  $W := f^{-1}(U \setminus \omega)$ . Glue  $(\hat{U}, f)$  and  $(\hat{V}, g)$  along  $W$  to obtain a 3-dimensional  $\mathbb{C}$ -analytic manifold, say  $X$  and a proper morphism  $\Pi: X \rightarrow Z$  which induces

<sup>2</sup>The author would like to thank Professor Mumford for his penetrating remark which greatly simplified the construction.

the blowing up  $(\hat{U}, f)$  (resp.,  $(\hat{V}, g)$ ) in the open subset  $U \subset Z$  (resp.,  $V \subset Z$ ). Let  $S$  be the total transform of  $M$  by  $\Pi$ . One has  $S = \hat{M} \cup \hat{S}$  where  $\hat{M}$  (resp.,  $\hat{S}$ ) is the proper transform of  $P_2$  (resp.,  $K$ ) by  $\Pi$ . Notice that  $\hat{S}$  is obtained by glueing  $\hat{S}_1$  and  $\hat{S}_2$ . Furthermore, the inverse image of  $\omega$  by  $\Pi$  is the union of two compact 1-cycles, say  $\bar{\alpha}$  and  $\bar{\beta}$  which intersect transversally (see Figure 2). Clearly  $(X, S)$  is a 3-dimensional 1-convex manifold.




FIGURE 2

*Step 4.* Let  $a$  (resp.,  $b$ ) be a general point on  $K_1 \setminus \omega$  (resp.,  $K_2 \setminus \omega$ ). Their inverse images  $\alpha$  (resp.,  $\beta$ ) are isomorphic to the projective line. Hence  $\alpha \sim \beta$  (homological equivalence). Furthermore, in view of the order of blowing up  $K_1$  and  $K_2$  in  $U$ , one has

$$\alpha \sim \bar{\alpha} + \bar{\beta}, \quad \beta \sim \bar{\beta}.$$

This implies that

$$\bar{\alpha} \sim 0 \quad (††)$$

*Step 5.*  $X$  is not kählerian.

If it were, let  $\Omega$  be the positive, closed  $(1, 1)$  form associated to some kähler metric on  $X$ . Then one has

$$\int_{\bar{\alpha}} \Omega > 0.$$

But this contradicts the existence of the compact 1-cycle  $\bar{\alpha}$  satisfying  $(\dagger\dagger)$ .

REMARKS. (a) Dimensionwise, example  $(\hat{B})$  is sharp. In fact, in a forthcoming paper, the following result will be proved.

THEOREM V. *Let  $X$  be a given 1-convex manifold with its exceptional subvariety  $S$  (singular, in general). If either*

(i)  $\dim X = 3$  and  $\dim S = 1$ , or

(ii)  $\dim X = 2$ ,

*then  $X$  is kählerian.*

(b) In both examples  $(\hat{A})$  and  $(\hat{B})$  above, their exceptional subvariety  $S$  is Moishezon. This is by no means accidental. This fact has been pointed out in [5, Corollary to Theorem 2] namely:

PROPOSITION 2. *Let  $(X, S)$  be a given 1-convex space. Then the exceptional subvariety  $S$  is Moishezon.*

To round off this discussion, we would like to mention a few problems related to the structure of 1-convex spaces.

*Problem 1.* Let  $(X, S)$  be a 1-convex space such that  $S$  is projective algebraic. Is  $X$  embeddable?

*Problem 2.* Let  $X$  be a 1-convex kähler manifold. Is  $X$  embeddable?

However a weaker version than Problems 1 and 2 seems more interesting.

*Problem 3.* Let  $X$  be a 1-convex manifold with its exceptional subvariety  $S$  (singular, in general).

(a) If  $S$  is projective, is  $X$  then kählerian?

(b) If  $X$  is kählerian, is  $S$  then projective?

Finally, in correlation with the previous Proposition 2, one would like to raise the following:

*Problem 4.*<sup>3</sup> Let  $S$  be a given Moishezon space. Is it always possible to construct a 1-convex space  $X$ , admitting  $S$  as its exceptional subvariety?

## REFERENCES

0. A. Andreotti and G. Tomassini, *A remark on the vanishing of certain cohomology groups*, Compositio Math. **21** (1969), 417–430.
1. S. Eto, H. Kazama and K. Watanabe, *On strongly  $q$ -pseudoconvex spaces with positive vector bundles*, Mem. Fac. Sci. Kyushu Univ. Ser. A **28** (1974), 135–146.

---

<sup>3</sup>This problem has recently been settled by the author in the affirmative.

2. H. Grauert, *Über Modifikationen und exzeptionell analytische Mengen*, Math. Ann. 146 (1962), 331–368.
3. R. Hartshorne, *Ample vector bundles*, Inst. Hautes Études Sci. Publ. Math. No. 29 (1966), 63–94.
4. ———, *Algebraic geometry*, Graduate Texts in Mathematics, vol. 52, Springer-Verlag, Berlin and New York, 1977.
5. B. G. Moishezon, *Modifications of complex varieties and Chow lemma*, Classification of Algebraic Varieties and Compact Complex Manifolds, Lecture Notes in Math., vol. 412, Springer-Verlag, Berlin and New York, 1974, pp. 133–139.
- 6a. Vo Van Tan, *On the classification of 1-convex spaces* (preprint).
- 6b. ———, *On the positivity of line bundles on 1-convex spaces* (in preparation)
7. K. W. Wiegmann, *Einbettungen komplexer Raumer in Zahlenraume*, Invent. Math. 1 (1966), 229–242.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MASSACHUSETTS AT BOSTON, BOSTON, MASSACHUSETTS 02125