

ALGEBRAIC FIBER BUNDLES

BY

STEVEN E. LANDSBURG

ABSTRACT. When X is a finite simplicial complex and G is any of a certain class of groups, a classification of G -principal bundles over X in terms of projective modules over a ring $R(G, X)$ is given. This generalizes Swan's classification of vector bundles and uses the results of Mulvey. Often, R can be taken to be noetherian; in this case $\text{Spec}(R)$ is usually reducible with "cohomologically trivial" irreducible components. Information is derived concerning the nature of projective modules over such rings, and some results are obtained indicating how such information reflects information about X .

Introduction. In 1962, Swan [11] demonstrated that the category of real or complex vector bundles over a paracompact space is equivalent via the global section functor to the category of projective modules over the ring of real- or complex-valued functions on the space. Combining a result of Mulvey [8] with some ideas based on those of Swan, we shall investigate the question of what other categories of fiber bundles can be represented "algebraically". In particular, when X is a topological space and G a topological group, we show in certain circumstances how to construct a noetherian scheme \hat{X} and a map $X \rightarrow \hat{X}$ such that the pullback operation converts algebraic vector bundles on \hat{X} into principal G -bundles on X in such a way that some interesting information is retained. The schemes \hat{X} are often affine and reducible with "cohomologically trivial" irreducible components; thus we devote §II to the study of vector bundles over such schemes. We first show (Theorem 2.1) that the vector bundles can be classified in a way analogous to the "Mayer-Vietoris" type classification which is well known in the case in which there are only two components. We then derive several corollaries regarding what the structure of these bundles can be.

In §III we derive stable analogues of some of these results.

Finally, in §IV, we give a few applications to bundles over simplicial complexes and complex varieties and to torsion in Picard Groups.

I. Principal bundles and projective modules.

DEFINITION. A ring is local if it has a unique maximal left (equivalently right) ideal. A ringed space is local if each of its stalks is local.

DEFINITION. Let (X, \mathcal{O}) be a ringed space. Then $\text{GL}_n \mathcal{O}$ will denote the sheaf given by $\Gamma(U, \text{GL}_n \mathcal{O}) = \text{GL}_n(\Gamma(U, \mathcal{O}))$ with the obvious restriction maps.

Received by the editors August 21, 1979 and, in revised form, June 12, 1980.

AMS (MOS) subject classifications (1970). Primary 13C10, 13D15, 55F25; Secondary 16A50, 55F10.

Key words and phrases. Projective module, G -principal bundle, vector bundle, affine scheme.

© 1981 American Mathematical Society
0002-9947/81/0000-0312/\$04.75

THEOREM 1.1 (MULVEY). *Let (X, \mathcal{O}) be a compact local ringed space, and suppose that the center of $\Gamma(X, \mathcal{O})$ separates points (in the sense that for any $x, y \in X$, $\exists s \in \Gamma(X, \mathcal{O})$ with $s_x = 0$ and $s_y = 1$). Then the category of finitely generated projective $\Gamma(X, \mathcal{O})$ -modules is equivalent to the category of locally free, locally finitely generated \mathcal{O} -modules.*

PROOF. See Mulvey [8].

REMARKS. (1) The isomorphism classes of objects in the category of locally free, locally finitely generated \mathcal{O} -modules of rank n form the pointed set $\check{H}^1(X, \mathrm{GL}_n \mathcal{O})$.

(2) Let G be the group of units of a topological ring and let \mathcal{O} be a subsheaf of the sheaf of continuous functions from X to that ring. If the theorem can be applied to \mathcal{O} , then it reduces the study of G -bundles on X (where “ G -bundle” may mean “continuous G -bundle”, “differentiable G -bundle”, “PL- G -bundle”, “holomorphic G -bundle”, etc., depending on \mathcal{O}) to the study of algebraic line bundles on $\hat{X} = \mathrm{Spec}(\Gamma(X, \mathcal{O}))$. \hat{X} is in general a very unwieldy construction (being non-noetherian), and our next theorem will be aimed at correcting this situation. However, \hat{X} can be both noetherian and useful, as the following example shows:

EXAMPLE 1.2. Let X be a finite simplicial complex linearly embedded in general position in \mathbb{R}^N . Let \mathcal{O} be the sheaf given by $\Gamma(U, \mathcal{O}) = \{f: U \rightarrow \mathbb{R} \mid f|_{U \cap \Delta} \text{ is a rational function for each simplex } \Delta \subset X\}$. Then, as in the remark, we construct a noetherian scheme $(\hat{X}, \hat{\mathcal{O}})$ (indeed, $\Gamma(\Delta, \mathcal{O}) \cong \mathbb{R}[X_1, \dots, X_n]$ for each n -simplex Δ and $\Gamma(X, \mathcal{O})$ is a colimit of finitely many of these) such that $H^1(\hat{X}, \hat{\mathcal{O}}^*) \approx H^1(X, \mathcal{O}^*) = H^1(X, \mathbb{R}^*)$ (here \mathbb{R}^* is occurring as a discrete group). Note that $\mathbb{Z}/2$ is a direct summand of \mathbb{R}^* , so that there is a surjection $H^1(X, \mathbb{R}^*) \twoheadrightarrow H^1(X, \mathbb{Z}/2)$, the latter being the group of line bundles on X . Thus we have constructed an affine noetherian scheme \hat{X} and a continuous injection $X \xrightarrow{\mathbb{P}} \hat{X}$ such that every line bundle on X is the pullback of an algebraic line bundle on \hat{X} . Later, this example will allow us to translate theorems about affine schemes into theorems about topology.

EXAMPLE 1.3. It is possible to carry out the above construction using restrictions of analytic functions rather than restrictions of polynomials, as Lønsted has shown [6]. In that case we actually get $H^1(\hat{X}, \hat{\mathcal{O}}^*) \approx H^1(X, \mathbb{Z}/2)$, but it is quite hard to show \hat{X} is noetherian. For another such construction, see Swan [12].

So far, our construction has only applied in cases where it was possible to recognize particular groups as groups of units in local rings. The following theorem will allow us to apply our methods to a much more general class of groups. First we need some definitions:

DEFINITION. Let \mathcal{X} be a category of ringed spaces. Then \mathcal{X} is acceptable if for every pushout diagram

$$\begin{array}{ccc} (A, \mathcal{O}_A) & \rightarrow & (B, \mathcal{O}_B) \\ \downarrow & & \downarrow \\ (C, \mathcal{O}_C) & \rightarrow & (D, \mathcal{O}_D) \end{array}$$

in \mathcal{X} , the diagram

$$\begin{array}{ccc} \Gamma(\theta_A) & \leftarrow & \Gamma(\theta_B) \\ \uparrow & & \uparrow \\ \Gamma(\theta_C) & \leftarrow & \Gamma(\theta_D) \end{array}$$

is a pullback.

EXAMPLE 1.4. If every arrow in \mathcal{X} is an open inclusion, then \mathcal{X} is acceptable. Suppose that X is an affine scheme and that \mathcal{X} is the category whose objects are intersections of irreducible components of X and whose arrows are inclusions. Then if X is the direct limit of \mathcal{X} , it follows that \mathcal{X} is acceptable.

DEFINITION. Let \mathcal{X} be any category and let $\{\Delta_i \rightarrow X\}_{i \in I}$ be some collection of arrows with common codomain. If Δ is the colimit of this collection (i.e., the pullback of the Δ_i over X), we define the union $\bigcup_{i \in I} \Delta_i$ to be the limit of the collection $\{\Delta \rightarrow \Delta_i\}$ (i.e., the pushout of the Δ_i over Δ).

DEFINITION. Let \mathcal{X} be a Grothendieck topology (e.g., a category of spaces) and $\Delta \subset \mathcal{X}$ a subcategory. Then a Δ -covering of an object $X \in \mathcal{X}$ is a covering $\{\Delta_i\}_{i \in I}$ such that:

- (1) $\Delta_i \in \Delta, \forall i$.
- (2) $\Delta_i \times_X (\bigcup_{j \in S} \Delta_j) \in \Delta$ for each subset $S \subset I$ such that $i \notin S$.

THEOREM 1.5. Let \mathcal{X} be an acceptable subcategory of the category of ringed spaces which satisfy the hypotheses of Theorem 1.1. Let Δ be a subcategory of \mathcal{X} such that every object in \mathcal{X} has a finite Δ -covering. Suppose that there exists a constant sheaf $G \subset \theta^*$ such that the inclusion induces isomorphisms

$$H^0(\Delta, G) \approx \Gamma(\Delta, \theta_\Delta^*) \quad \text{for each } (\Delta, \theta) \in \Delta$$

and

$$H^1(\Delta, G) \approx H^1(\Delta, \theta_\Delta^*) \quad \text{for each } (\Delta, \theta) \in \Delta.$$

Then $H^1(X, G) = H^1(X, \theta_X^*) = \text{Pic}(\Gamma(X, \theta_X))$ for each $(X, \theta_X) \in \mathcal{X}$.

PROOF. Given an object $(X, \theta_X) \in \mathcal{X}$ we may write $(X, \theta_X) = (Y, \theta_Y) \cup (\Delta, \theta_\Delta)$ where $(\Delta, \theta_\Delta) \in \Delta$, (Y, θ_Y) has a smaller Δ -covering than (X, θ_X) , and $(Y, \theta_Y) \times_{(X, \theta_X)} (\Delta, \theta_\Delta) \in \Delta$. Writing the pushout diagram

$$\begin{array}{ccc} (Y, \theta_Y) \times_{(X, \theta_X)} (\Delta, \theta_\Delta) & \rightarrow & (\Delta, \theta_\Delta) \\ \downarrow & & \downarrow \\ (Y, \theta_Y) & \rightarrow & (X, \theta_X) \end{array}$$

in \mathcal{X} , we get a pullback diagram

$$\begin{array}{ccc} \Gamma(X, \theta_X) & \rightarrow & \Gamma(\Delta, \theta_\Delta) \\ \downarrow & & \downarrow \\ \Gamma(Y, \theta_Y) & \rightarrow & (Y \times_X \Delta, \theta_{Y \times_X \Delta}) \end{array}$$

in the category of rings. This gives two Mayer-Vietoris sequences:

$$\begin{array}{ccccccc}
 H^0(\Delta, G) \oplus H^0(Y, G) & \rightarrow & H^0(Y \times_X \Delta, G) & \rightarrow & H^1(X, G) & & \\
 \parallel & & \parallel & & & & \\
 \Gamma(\Delta, \mathcal{O}_\Delta^*) \oplus \Gamma(Y, \mathcal{O}_Y^*) & \rightarrow & \Gamma(Y \times_X \Delta, \mathcal{O}_{Y \times_X \Delta}^*) & \rightarrow & \text{Pic } \Gamma(X, \mathcal{O}_X) & & \\
 & & & & \rightarrow & H^1(\Delta, G) \oplus H^1(Y, G) & \rightarrow H^1(Y \times_X \Delta, G) \\
 & & & & & \parallel & \parallel \\
 & & & & \rightarrow & \text{Pic } \Gamma(\Delta, \mathcal{O}_\Delta) \oplus \text{Pic } \Gamma(Y, \mathcal{O}_Y) & \rightarrow \text{Pic}(Y \times_X \Delta, \mathcal{O}_{Y \times_X \Delta})
 \end{array}$$

The first, second, and fifth groups in these sequences are equal by assumption and the fourth by induction. But the latter two maps are induced by the natural transformation $H^1(-, G) \rightarrow H^1(-, \mathcal{O}_*^*) = \text{Pic}(\Gamma(-, \mathcal{O}_*))$ (the latter equality by the second remark following Theorem 1.1), and this gives an arrow $H^1(X, G) \rightarrow \text{Pic}(\Gamma(X, \mathcal{O}_X))$ which makes everything commute. The five-lemma completes the proof. Q.E.D.

REMARK. We have constructed an affine noetherian scheme $\hat{X} = \text{Spec}(\Gamma(X, \mathcal{O}_X))$ and a continuous injection $X \xrightarrow{\psi} \hat{X}$ such that $\psi^* \mathcal{O}_{\hat{X}}^* = G$ and such that the correspondence between G -principal bundles on X and line bundles on \hat{X} is bijective.

EXAMPLE 1.6. Let $T \subset \mathbb{C}$ be any subring such that $\text{Pic}(T[X_1, \dots, X_k]) = 0$ for each k . (By Gauss's Lemma, this is the case when T is a Dedekind ring with class group zero.) Let \mathcal{X} be the category of simplicial complexes X linearly embedded in general position in \mathbb{R}^N and endowed with the sheaf \mathcal{O}_X given by $\Gamma(U, \mathcal{O}_X) = \{f: U \rightarrow \mathbb{C} \mid \text{for each simplex } \Delta, f|_{\Delta \cap U} \text{ is of the form } \varphi/\psi \text{ where } \varphi \text{ and } \psi \text{ are polynomial functions with coefficients in } T\}$. Then the theorem applies with $G = T^*$. Letting $T = \mathbb{Z}[1/2]$, for example, we get $H^1(X, \mathbb{Z}_2 \times \mathbb{Z}) = \text{Pic}(\Gamma(X, \mathcal{O}_X))$ so that we have constructed a map from X to an affine noetherian scheme \hat{X} which induces a bijection between line bundles on \hat{X} and $\mathbb{Z}_2 \times \mathbb{Z}$ -bundles on X . Alternatively, letting $T = \mathbb{Z}$, we get precisely the real line bundles on X .

EXAMPLE 1.7. In the example above we could take $T = \mathbb{Z}[\xi]$ where ξ is a primitive n th root of unity such that $\text{Pic}(T) = 0$ (for example, n could be a prime < 19 —see Milnor [7]). Then the Dirichlet Unit Theorem guarantees that $\mathbb{Z}/n\mathbb{Z}$ will be a direct summand of T^* , so we have maps $\text{Pic}(\hat{X}) \approx H^1(X, T^*) \twoheadrightarrow H^1(X, \mathbb{Z}/n\mathbb{Z})$. Thus we have constructed a map from X to an affine noetherian scheme \hat{X} such that every $\mathbb{Z}/n\mathbb{Z}$ -bundle on X is a pullback from \hat{X} .

In most of the above examples, the schemes we have constructed have been reducible with irreducible components which are cohomologically trivial. Our next goal is to understand the vector bundles over such schemes.

II. Vector bundles over reducible affine schemes. We now turn to the problem of classifying vector bundles over reducible affine schemes in terms of the vector bundles over the subschemes of a closed covering. The solution to this problem (Theorem 2.1) will be analyzed to provide information about the number of sections necessary to generate an arbitrary vector bundle. In §IV we will apply this information to the results of §I.

Notation. In what follows, X will always denote an affine reducible scheme with a closed covering $\{X_1, \dots, X_n\}$. For example, the X_i could be the irreducible components of X . As X is affine, we shall write $X = \text{Spec}(R)$ or $R = \Gamma(X)$, and $X_i = \text{Spec}(R_i)$. $VB_r(X)$ will denote the set of rank- r vector bundles on X . \mathcal{X} will denote the nerve of the covering $\{X_i\}$ of X . $\check{H}^1(\mathcal{X}, \text{GL}_r)$ will denote cohomology with respect to the system of local coefficients which assigns to the simplex $\{i_0, i_1, \dots, i_n\}$ the group $\text{GL}_r(\Gamma(X_{i_0} \cap X_{i_1} \cap \dots \cap X_{i_n}))$.

We need two trivial algebraic lemmas:

LEMMA. *Let R be a ring and let $e, e': R^n \rightarrow R^n$ be idempotent. Then e and e' have isomorphic images if and only if there are maps $f, g: R^n \rightarrow R^n$ satisfying the six (redundant) equations:*

- (1) $fe = f$,
- (2) $eg = g$,
- (3) $e'f = f$,
- (4) $ge' = g$,
- (5) $gf = e$,
- (6) $fg = e'$.

PROOF. Write $P = \text{im}(e)$, $P' = \text{im}(e')$, $Q = \text{im}(1 - e)$ and $Q' = \text{im}(1 - e')$. Supposing that P and P' are isomorphic, let $P \xrightarrow{f} P' \xrightarrow{g} P$ be inverse isomorphisms and extend f and g to be zero on Q and Q' to get a diagram

$$\begin{array}{ccc} R^n & \xrightarrow{e} & P \oplus Q = R^n \\ f \downarrow & & \downarrow f \\ R^n & \xrightarrow{e'} & P' \oplus Q' = R^n \\ g \downarrow & & \downarrow g \\ R^n & \xrightarrow{e} & P \oplus Q = R^n \end{array}$$

Then the facts that f and g are inverse isomorphisms and have been extended by zero are summarized in equations (1) through (6). Conversely, it is easy to see that if f, g satisfy (1)–(6) then $f|_P$ and $g|_{P'}$ are inverse isomorphisms. Q.E.D.

REMARK. In order for P to be isomorphic to P' and Q to be isomorphic to Q' , it is necessary and sufficient that there exist f, g as above where $f = e'ue$ and $g = eu^{-1}e'$ for some isomorphism $u: R^n \rightarrow R^n$.

LEMMA. *Let e be an $m \times m$ idempotent matrix of rank r over a ring R and suppose that $\text{im}(e)$ is free. Then there are matrices*

$$A = \begin{bmatrix} A_1^1 & \cdots & A_m^1 \\ \vdots & & \vdots \\ A_1^r & \cdots & A_m^r \end{bmatrix}$$

and

$$B = \begin{bmatrix} B_1^1 & \cdots & B_m^1 \\ \vdots & & \vdots \\ B_1^r & \cdots & B_m^r \end{bmatrix}$$

such that $e = {}^tA \cdot B$ and $B \cdot {}^tA$ is the $r \times r$ identity matrix. (Here tA denotes the transpose of A .)

PROOF. Let

$$e' = \left(\begin{array}{c|c} I_r & 0 \\ \hline 0 & 0 \end{array} \right)$$

and choose f and g as in the last lemma. Let B consist of the top r rows of f and let tA consist of the left r columns of g . Q.E.D.

REMARK. More generally, in any additive category let $M = M_1 \oplus M_2$, $e =$ the projection of M on M_1 , $N = N_1 \oplus N_2$, and $e' =$ the projection of N on N_1 . Then M_1 and N_1 are isomorphic if and only if there are maps

$$M \begin{array}{c} \xrightarrow{f} \\ \xleftarrow{g} \end{array} N$$

such that $e = gf$ and $e' = fg$. In this case, f and g can be chosen to satisfy equations (1)–(6).

THEOREM 2.1. *If X is the direct limit of the system of X_i 's and their intersections, and the same is true for each finite union $X_{i_1} \cup \cdots \cup X_{i_k}$, then there is a natural long exact sequence of pointed sets*

$$1 \rightarrow \check{H}^1(\mathcal{X}, \text{GL}_r) \rightarrow VB_r(X) \xrightarrow{\varphi} \bigoplus_{i=1}^n VB_r(X_i).$$

These sequences commute with tensor products in the sense indicated by the diagram:

$$\begin{array}{ccccc} & 1 & & 1 & & 1 \\ & \downarrow & & \downarrow & & \downarrow \\ \check{H}^1(\mathcal{X}, \text{GL}_r) & \times & \check{H}^1(\mathcal{X}, \text{GL}_s) & \xrightarrow{\otimes} & \check{H}^1(\mathcal{X}, \text{GL}_{rs}) \\ \downarrow & & \downarrow & & \downarrow \\ VB_r(X) & \times & VB_s(X) & \xrightarrow{\otimes} & VB_{rs}(X) \\ \downarrow & & \downarrow & & \downarrow \\ \bigoplus_{i=1}^n VB_r(X_i) & \times & \bigoplus_{i=1}^n VB_s(X_i) & \xrightarrow{\otimes} & \bigoplus_{i=1}^n VB_{rs}(X_i) \end{array}$$

In particular, when $r = 1$ the maps are group homomorphisms.

PROOF. Letting φ be the restriction map, we begin by defining a map $\text{Ker}(\varphi) \xrightarrow{\alpha} \check{H}^1(\mathcal{X}, \text{GL}_r)$. Given $P \in \text{ker}(\varphi)$, write $P \approx \text{im}(e)$ where $e: R^m \rightarrow R^m$ is idempotent. (We are thinking of P as a projective module over R .) Using the

embedding $R \rightarrow \prod_{i=1}^n R_i$, write $e = (e_1, \dots, e_n)$. Then, since $P \in \ker(\varphi)$, $\text{im}(e_i) \approx R_i'$. Thus we can write $e_i = {}'A_i \cdot B_i$ where $B_i \cdot {}'A_i = \underline{I_r}$ as in the lemma. For each $i, i' \in \{1, \dots, n\}$ define $p_{ii'}$ to be the matrix $\overline{B_i} \cdot {}'A_{i'}$ over $\Gamma(X_i \times_X X_{i'})$. (Here "bar" denotes restriction to $X_i \times_X X_{i'}$.) We claim that $p_{ii'}$ is a well-defined GL_r -cocycle.

First to check that $p_{ii'}$ is invertible, note that $\overline{B_i} \cdot \overline{{}'A_i} = \overline{e_i} = \overline{e_{i'}} = \overline{B_{i'}} \cdot \overline{{}'A_{i'}}$ because e has all of its entries in R . Thus:

$$p_{ii'} \overline{B_{i'}} = \overline{B_i} \cdot \overline{{}'A_{i'}} \cdot \overline{B_{i'}} = \overline{B_i} \cdot \overline{{}'A_i} \cdot \overline{B_i} = \overline{B_i}.$$

But $\overline{B_i} \cdot \overline{{}'A_i} = \underline{I_r}$, so $p_{ii'}$ is invertible.

To see that $(p_{ii'})$ is a cocycle, write

$$p_{ii'} p_{i'i''} = p_{ii'} \cdot \overline{B_{i'}} \cdot \overline{{}'A_{i''}} = \overline{B_i} \cdot \overline{{}'A_{i''}} = p_{ii''}.$$

Next we check that $p_{ii'}$ is well defined: Suppose that e and \tilde{e} have isomorphic images (possibly $e = \tilde{e}$). Clearly, adding rows and columns of zeroes does not affect anything, so we may assume that e and \tilde{e} are of the same size, say $m \times m$. Let $f: \text{im}(e) \rightarrow \text{im}(e')$ be an isomorphism and extend f by zero on the complement of $\text{im}(e)$ in R^n so we have a commutative diagram

$$\begin{array}{ccc} R^m & \xrightarrow{e} & R^m \\ f \downarrow & & \downarrow f \\ R^m & \xrightarrow{\tilde{e}} & R^m. \end{array}$$

(We will write $f = (f_1, \dots, f_n)$.) Suppose that $e = {}'A \cdot B$ and $\tilde{e} = {}'C \cdot D$, and set $p_{ii'} = \overline{B_i} \cdot {}'A_{i'}$ and $\tilde{p}_{ii'} = \overline{D_i} \cdot {}'C_{i'}$. Then

$$\begin{aligned} \overline{D_i} \cdot \overline{f_i} \cdot \overline{{}'A_i} p_{ii'} &= \overline{D_i} \cdot \overline{f_i} \cdot \overline{{}'A_i} \cdot \overline{B_i} \cdot \overline{{}'A_{i'}} \\ &= \overline{D_i} \cdot \overline{f_i} \cdot \overline{e_i} \cdot \overline{{}'A_{i'}} = \overline{D_i} \cdot \overline{\tilde{e}_i} \cdot \overline{f_i} \cdot \overline{{}'A_{i'}} \\ &= \overline{D_i} \cdot \overline{{}'C_i} \cdot \overline{D_i} \cdot \overline{f_i} \cdot \overline{{}'A_{i'}} = \overline{D_i} \cdot \overline{{}'C_{i'}} \cdot \overline{D_{i'}} \cdot \overline{f_{i'}} \cdot \overline{{}'A_{i'}} \\ &= \tilde{p}_{ii'} \cdot \overline{D_{i'}} \cdot \overline{f_{i'}} \cdot \overline{{}'A_{i'}}. \end{aligned}$$

In other words, if we let ξ be the zero-cochain defined by $\xi_i = D_i f_i {}'A_i$, then $p_{ii'}$ and $\tilde{p}_{ii'}$ differ by the coboundary of ξ_i . This shows that $(p_{ii'})$ is well defined as a cohomology class.

To check commutativity with tensor products, note that in general if A, B, C, D are matrices then $AB \otimes CD = (A \otimes C)(B \otimes D)$ so if $e = {}'A \cdot B$ and $\tilde{e} = {}'C \cdot D$, we get

$$\begin{aligned} \text{im}(e) \otimes \text{im}(\tilde{e}) &= \text{im}(e \otimes \tilde{e}) = \text{im}(({}'A \otimes {}'C)(B \otimes D)) \mapsto ((\overline{B_i} \otimes \overline{D_i})(\overline{{}'A_{i'}} \otimes \overline{{}'C_{i'}})) \\ &= ((\overline{B_i} \cdot \overline{{}'A_{i'}}) \otimes (\overline{D_i} \cdot \overline{{}'C_{i'}})). \end{aligned}$$

We now have a well-defined map $\alpha: \text{Ker}(\varphi) \rightarrow \check{H}^1(\mathcal{X}, \text{GL}_r)$. We claim that α is injective. To see this, suppose that $P \approx \text{im}(e)$ and $\tilde{P} \approx \text{im}(\tilde{e})$ have the same image and write $e = {}'A \cdot B$ and $\tilde{e} = {}'C \cdot D$. Then there is a zero-cochain (ξ_i) such that for

each i, i' we have $\overline{\xi_i} \overline{B_i} \overline{{}^t A_i} = \overline{D_i} \overline{{}^t C_i} \overline{\xi_{i'}}$. For each i , let $g_i = {}^t C_i \xi_i B_i: R_i^m \rightarrow R_i^m$. Then

$$\begin{aligned} \overline{g_i} &= \overline{{}^t C_i} \overline{\xi_i} \overline{B_i} \\ &= \overline{{}^t C_i} \overline{D_i} \overline{{}^t C_i} \overline{\xi_i} \overline{B_i} \overline{{}^t A_i} \overline{B_i} \\ &= \overline{{}^t C_{i'}} \overline{D_{i'}} \overline{{}^t C_i} \overline{\xi_i} \overline{B_i} \overline{{}^t A_{i'}} \overline{B_{i'}} \\ &= \overline{{}^t C_{i'}} \overline{\xi_{i'}} \overline{B_{i'}} = \overline{g_{i'}} \end{aligned}$$

so the g_i define a map $g: R^m \rightarrow R^m$. We can likewise define a map h by setting $h_i = {}^t A_i \xi_i^{-1} \cdot D_i$. Then $gh = \tilde{e}$ and $hg = e$ so g and h are inverse isomorphisms between $\text{im}(e)$ and $\text{im}(\tilde{e})$. This proves the claim.

Next we define a map $\beta: \check{H}^1(\mathcal{X}, \text{GL}_r) \rightarrow \text{Ker}(\varphi)$ which is inverse to α . Given $(p_{ii'}) \in \check{H}^1(\mathcal{X}, \text{GL}_r)$, we define $\beta(p_{ii'})$ as the image of an idempotent matrix which we shall construct. We may assume by induction that there is an idempotent matrix $\hat{e} = (\hat{e}_1, \dots, \hat{e}_{n-1})$ over $\Gamma(X_1 \cup \dots \cup X_{n-1}) \subset \prod_{i=1}^{n-1} R_i$ such that each $\text{im}(e_i)$ is free and such that $\alpha(\hat{e}) = (p_{ii'})_{i,i' \leq n-1}$. Now p_{in} is invertible in $\Gamma(X_i \times_X X_n)$; write $q_{in} p_{in} = 1$, and let p_i and q_i be arbitrary liftings of p_{in} and q_{in} to R_i . Write $p_i q_i + u_i = 1$; squaring both sides write $p_i z_i + u_i^2 = 1$ where $z_i = q_i p_i q_i + 2q_i u_i$ (using the fact that $p_i q_i$ commutes with $u_i = 1 - p_i q_i$). Writing $e_i = {}^t \hat{A}_i \hat{B}_i$ as in the lemma, set

$$e_i = \left[\begin{array}{c|c} z_i p_i & z_i u_i \hat{B}_i \\ \hline {}^t \hat{A}_i u_i p_i & {}^t \hat{A}_i u_i^2 \hat{B}_i \end{array} \right] = \left(\frac{z_i}{{}^t \hat{A}_i u_i} \right) (p_i | u_i \hat{B}_i)$$

for $i \leq n-1$ and

$$e_n = \begin{bmatrix} 1 & & & & & 0 \\ & \ddots & & & & \\ & & 1 & & & \\ & & & 0 & & \\ & & & & \ddots & \\ 0 & & & & & 0 \end{bmatrix}.$$

Using the facts that $\hat{B}_i {}^t \hat{A}_i = 1$ and $p_i z_i + u_i^2 = 1$, it is easy to check that e_i is idempotent.

We shall check to see that e really has its entries in R and that $\alpha(\text{im}(e)) = (p_{ii'})$.

To see that e has its entries in R , we need to check that $\overline{e_i} = \overline{e_{i'}}$ where "bar" denotes restriction to $X_i \times_X X_{i'}$. By the Chinese Remainder Theorem and induction, we may assume that the p_i have been chosen to satisfy the equation $p_{i'i} \cdot \overline{p_i} = \overline{p_{i'}}$ and consequently the q_i can be chosen to satisfy $\overline{q_i p_{i'}} = \overline{q_{i'}}$, whence $\overline{u_i} = 1 - \overline{p_i q_i} = \overline{p_{i'i} u_i p_{i'}}$ and $\overline{z_i p_{i'}} = \overline{z_{i'}}$. We also know as before that $\overline{p_{i'i} \hat{B}_{i'}} = \overline{\hat{B}_{i'}}$ and similarly $\overline{{}^t \hat{A}_{i'} p_{i'i}} = \overline{{}^t \hat{A}_{i'}}$. Using this information it is easy to check each of the four blocks of e

to see that $\overline{e_i} = \overline{e_{i'}}$. Specifically

$$\begin{aligned}\overline{z_{i'}} \overline{p_{i'}} &= \overline{z_i p_{ii'} p_{i'i} p_i} = \overline{z_i} \overline{p_i}, \\ \overline{z_{i'}} \overline{u_{i'}} \overline{\widehat{B}_{i'}} &= \overline{z_i p_{ii'} p_{i'i} u_i p_{ii'} p_{i'i} \widehat{B}_i} = \overline{z_i} \overline{u_i} \overline{\widehat{B}_i}, \\ \overline{{}^t \widehat{A}_{i'}} \overline{u_{i'}} \overline{p_{i'}} &= \overline{{}^t \widehat{A}_i p_{ii'} p_{i'i} u_i p_{ii'} p_{i'i} p_i} = \overline{{}^t \widehat{A}_i} \overline{u_i} \overline{p_i}, \\ \overline{{}^t \widehat{A}_{i'}} \overline{u_{i'}^2} \overline{\widehat{B}_{i'}} &= \overline{{}^t \widehat{A}_i p_{ii'} p_{i'i} u_i^2 p_{ii'} p_{i'i} \widehat{B}_i} = \overline{{}^t \widehat{A}_i} \overline{u_i^2} \overline{\widehat{B}_i}.\end{aligned}$$

Next we compute $\alpha(\text{im}(e)) = (\xi_{ii'})$ where

$$\begin{aligned}\xi_{ii'} &= (\overline{p_i} | \overline{u_i} \overline{\widehat{B}_i}) \left(\frac{\overline{z_{i'}}}{\overline{{}^t \widehat{A}_{i'}} \overline{u_{i'}}} \right) \\ &= \overline{p_i} \overline{z_{i'}} + \overline{u_i} \overline{\widehat{B}_i} \overline{{}^t \widehat{A}_{i'}} \overline{u_{i'}} = \overline{p_i z_i p_{ii'}} + \overline{u_i p_{ii'} p_{i'i} u_i p_{ii'}} \\ &= (\overline{p_i z_i} + \overline{u_i}) p_{ii'} = p_{ii'}\end{aligned}$$

when $i, i' \leq n-1$. Also

$$\xi_{in} = (\overline{p_i} | \overline{u_i} \overline{\widehat{B}_i}) \left(\frac{I}{0} \right) = \overline{p_i} = p_{in}.$$

This (combined with the fact that α is one-one) shows that β is well-defined and inverse to α , proving the theorem. Q.E.D.

REMARK. R. G. Swan has pointed out to me via the following example that the condition $X = \text{ind lim } X_i$ is really necessary: Let $X_1 = \text{Spec}(k[X, Y, Z]/XYZ)$ and let $X_2 = \text{Spec}(k[X, Y]/XY(X - Y))$. Then the coverings \mathcal{X}_1 and \mathcal{X}_2 of X_1 and X_2 by their irreducible components are isomorphic, and $\text{Pic}(X_1) = H^1(\mathcal{X}_1, \mathcal{O}_{X_1}^*) = 0$ by the theorem, whereas $\text{Pic}(X_2) = k \neq H^1(\mathcal{X}_2, \mathcal{O}_{X_2}^*)$.

In everything that follows, X will denote an affine scheme.

COROLLARY 2.2. If $\{X_i\}$ has cardinality n and ξ is a vector bundle of rank r such that $\xi|_{X_i}$ is trivial for each i , then ξ can be generated by nr sections.

PROOF. The matrix e constructed in the proof of the theorem is an $nr \times nr$ matrix. Q.E.D.

In some circumstances, the proof of Theorem 2.1 gives much better information about generators:

DEFINITION. A cochain $(p_{ii'})$ is special if there is an index i_0 such that whenever $i \neq i_0$ and $i' \neq i_0$ it follows that $p_{ii'} = 1$.

LEMMA. In the situation of the theorem, if $P \in \ker(\varphi)$ is represented by a special cocycle then P can be generated by $2r$ sections (where r is the rank of P).

PROOF. Let $(p_{ii'})$ represent P and let i_0 be as in the definition of special. For each i , write $p_{ii_0} \cdot q_{ii_0} = 1$ and let p_i and q_i be arbitrary liftings of p_{ii_0} and q_{ii_0} to R_i . Write $p_i q_i + u_i = 1$ so $p_i z_i + u_i^2 = 1$ where $z_i = q_i p_i q_i + 2q_i u_i$. For each $i \neq i_0$ let

$$e_i = \left(\begin{array}{c|c} z_i p_i & z_i u_i \\ \hline u_i p_i & u_i^2 \end{array} \right)$$

and let

$$e_{i_0} = \left(\begin{array}{c|c} I & 0 \\ \hline 0 & 0 \end{array} \right).$$

Having chosen the p_i and q_i carefully as in the proof of the theorem, we may assume $\overline{p_i} = \overline{p_{i'}}$ (and consequently $\overline{q_i} = \overline{q_{i'}}$, $\overline{u_i} = \overline{u_{i'}}$) for each i, i' distinct from i_0 . Then $\alpha(\text{im } e) = (\xi_{ii'})$ where for $i, i' \neq i_0$ we have $\xi_{ii'} = \overline{p_i z_{i'}} + \overline{u_i u_{i'}} = \overline{p_i z_i} + \overline{u_i^2} = 1$, and where $\xi_{i_0 i} = \overline{p_i} = \overline{p_{i_0}}$. That is, $(\xi_{ii'}) = (p_{ii'})$. But $\alpha: \text{Ker } \varphi \rightarrow \check{H}^1(\mathcal{X}, \text{GL}_r)$ is injective so $P \approx \text{im}(e)$. Thus P can be generated by $2r$ sections. Q.E.D.

REMARK. As a consequence of this lemma, any line bundle which becomes trivial on each X_i and whose cocycle is a product of special cocycles is a tensor product of line bundles which can be generated by two sections. In particular, when the special cocycles generate the group of cocycles, every line bundle can be written as such a tensor product. The special *cochains* always generate the group of cochains, so we get

COROLLARY 2.3. *If whenever i, i', i'' are all distinct it follows that $X_i \times_X X_{i'} \times_X X_{i''} = \emptyset$, then every line bundle which becomes trivial on each X_i is a tensor product of line bundles which can be generated by two sections.*

PROOF. In this case the group of cochains is the group of cocycles, so we are done by the remark above. Q.E.D.

We can strengthen this:

COROLLARY 2.4. *Suppose that for every three distinct indices i, i', i'' we have either $X_i \times_X X_{i'} \times_X X_{i''} = \emptyset$ or $X_i \times_X X_{i'} \times_X X_{i''} = X_i \times_X X_{i'}$. Then every line bundle on X which becomes trivial on each X_i is a tensor product $L_1 \otimes \cdots \otimes L_{n-1}$ where each L_i can be generated by two sections.*

PROOF. Let L be a line bundle and set $\alpha(L) = (p_{ii'})$. For each index i_0 let $(p_{ii_0}^{i_0})$ be the cocycle defined by

$$\begin{aligned} p_{ii_0}^{i_0} &= 1, \\ p_{ii_0}^{i_0} &= p_{i_0 i_0} \text{ where } l_0 = l_0(i_0, i') = \max\{l | X_{i_0} \times_X X_{i'} \times_X X_l \neq \emptyset\}, \\ p_{ii_0}^{i_0} &= (p_{i_0 i_0}^{i_0})^{-1}, \\ p_{ii_0}^{i_0} &= 1 \text{ if } i, i' \neq i_0. \end{aligned}$$

To see that this is a cocycle, suppose that $X_{i_0} \times_X X_i \times_X X_{i'} \neq \emptyset$. Then $X_{i_0} \times_X X_i = X_{i_0} \times_X X_{i'}$ so $l_0(i_0, i) = l_0(i_0, i')$, whence $p_{ii_0}^{i_0} p_{i_0 i_0}^{i_0} = p_{i_0 i_0} p_{i_0 i_0} = 1 = p_{ii'}$ (assuming $i, i' \neq i_0$). One checks the other triple intersections similarly.

Now for fixed i, i' ,

$$\prod_{i_0=1}^n p_{ii'}^{i_0} = p_{ii'}^i p_{ii'}^{i'} = p_{ii_0(i,i')} \cdot p_{i_0(i,i')} = p_{ii'},$$

so

$$(p_{ii'}) = \prod_{i=1}^n (p_{ii'}^{i_0}).$$

Thus if L_{i_0} is the line bundle defined by $(p_{ii'}^{i_0})$, we have $L = L_1 \otimes \cdots \otimes L_n$. But each $(p_{ii'}^{i_0})$ is special, so each L_i can be generated by two sections. Moreover, L_n is

trivial since $p_{ii'}^n = 1$ for all i, i' . Therefore $L = L_1 \otimes \cdots \otimes L_{n-1}$ where each L_i can be generated by two sections. Q.E.D.

REMARK. The proof of the above corollary is still valid if we define $p_{ii'}^{i_0}$ separately on each connected component Y of $X_i \times_X X_{i'}$ by

$$p_{ii'}^{i_0}|_Y = p_{i_0 i_0} \quad \text{where } i_0 = i_0(i_0, Y) = \max\{l | Y \times_X X_l \neq \emptyset\}.$$

This gives the stronger result:

COROLLARY 2.5. *Suppose that for every three indices i, i', i'' it is the case that each connected component of $X_i \times_X X_{i'} \times_X X_{i''}$ is a connected component of $X_i \times_X X_{i'}$. Then every line bundle over X which becomes trivial on each X_i can be written as a tensor product $L_1 \otimes \cdots \otimes L_{n-1}$ where each L_i can be generated by two sections.*

An immediate consequence of this is

COROLLARY 2.6. *Suppose that all but at most one of the X_i are one-dimensional. Then every line bundle over X which becomes trivial on each X_i can be written as a tensor product $L_1 \otimes \cdots \otimes L_{n-1}$ where each L_i can be generated by two sections.*

As another corollary to the proof of Theorem 2.1, we get a partial generalization of the well-known fact that if I is any nil ideal in R then any projective module over R/I is extended from R .

COROLLARY 2.7. *Suppose that there exists an index i_0 such that each $X_i \times_X X_{i_0}$ is defined by a nil ideal in $\Gamma(X_i)$. Then any vector bundle which is trivial on each X_i is trivial.*

PROOF. Assume $i_0 = n$. Then, in the proof of Theorem 2.1, u_i is in the ideal defining $\Gamma(X_i \times_X X_n)$ in $\Gamma(X_i)$. Write $u_i^{k_i} = 0$ and let k be greater than $\max_i k_i$. Writing $p_i z_i = u_i^2 = 1$ as in the theorem, raise each side to the k to get $p_i w_i = 1$. Then for each $i, i', p_{ii'} = p_i w_{i'}$, so $(p_{ii'})$ is a coboundary. Q.E.D.

III. **Stability.** We continue to use the notation of the previous section. In addition, E_r will denote the sheaf of $r \times r$ elementary matrices on X and E will denote $\bigcup_{r=1}^{\infty} E_r$ (where E_r is contained in E_{r+1} in the usual way).

THEOREM 3.1. *Let L be the kernel of the restriction map*

$$\tilde{K}_0(X) \rightarrow \bigoplus_{i=1}^n \tilde{K}_0(X_i).$$

Then there is a natural exact sequence

$$H^1(\mathcal{X}, E) \rightarrow L \xrightarrow{\hat{\alpha}} H^1(\mathcal{X}, K_1).$$

PROOF. We use the notation from the proof of Theorem 2.1. Given $([P] - [F]) \in L$ with F free and P of rank r , let $\hat{\alpha}(P)$ be the image of $\alpha(P)$ under the map $H^1(\mathcal{X}, \text{GL}_r) \rightarrow H^1(\mathcal{X}, K_1)$.

This is well defined: if $[P] - [F] = [Q] - [F']$ with F and F' free, then $P \oplus F'' \approx Q \oplus F'''$ for appropriate free modules F'', F''' . The claim that $\hat{\alpha}$ is well defined now follows from the fact that $\hat{\alpha}(P) = \hat{\alpha}(P \oplus F'') = \hat{\alpha}(Q \oplus F''') = \hat{\alpha}(Q)$.

This is a group homomorphism: $\hat{\alpha}([P] - [F] + [Q] - [F'])$ is the image of

$$\alpha(P \oplus Q) = \left(\begin{array}{c|c} \alpha(P) & 0 \\ \hline 0 & \alpha(Q) \end{array} \right),$$

which, by Whitehead's Lemma, is equal in K_1 to $\alpha(P)\alpha(Q)$.

Finally defining a map $H^1(\mathcal{X}, E) \rightarrow L$ by composing the natural map $H^1(\mathcal{X}, E) \rightarrow H^1(\mathcal{X}, GL)$ with the map $H^1(\mathcal{X}, GL) \rightarrow L$ given by Theorem 2.1, we see that if $\hat{\alpha}(P) = 0$, then $\alpha(P)$ is stably elementary and so comes from $H^1(\mathcal{X}, E)$. Q.E.D.

Note. The above proof simply amounts to noting that the natural map $H^1(\mathcal{X}, GL) \rightarrow \tilde{K}_0(X)$ factors through L and that the vertical arrow in the diagram

$$\begin{array}{ccccccc} 1 & \rightarrow & H^1(\mathcal{X}, E) & \rightarrow & H^1(\mathcal{X}, GL) & \rightarrow & H^1(\mathcal{X}, K_1) \\ & & & & \downarrow & \nearrow & \\ & & & & L & & \end{array}$$

is onto.

COROLLARY 3.2. *If $H^1(\mathcal{X}, E) = 0$ then L maps injectively to $H^1(\mathcal{X}, K_1)$.*

As an example, when \mathcal{X} satisfies the hypothesis of Corollary 2.5 an argument using the fact that E_r is "flasque" with respect to the covering \mathcal{X} shows that $H^1(\mathcal{X}, E_r) = 0$ for each r , so that Corollary 3.2 applies.

COROLLARY 3.3. *Suppose that X satisfies the hypotheses of Corollary 2.5 and that $\tilde{K}_0(X_i) = 0$ for each i . Suppose also that $H^1(\mathcal{X}, SK_1) = 0$. Then $\tilde{K}_0(X) = \text{Pic}(X)$.*

PROOF. We have $\tilde{K}_0(X) \hookrightarrow H^1(\mathcal{X}, K_1) = H^1(\mathcal{X}, GL_1) \oplus H^1(\mathcal{X}, SK_1) = \text{Pic}(X)$; it is easy to check that this is just the determinant map, which is onto. Q.E.D.

EXAMPLE 3.4. Let R be a regular ring such that $\tilde{K}_0(R) = SK_1(R) = 0$. Suppose that each X_i and each $X_i \times_X X_j$ is an affine space over R , and that the hypothesis of Corollary 2.5 is satisfied. Then $\tilde{K}_0(X) = \text{Pic}(X)$.

IV. Applications.

A. Coverings and triangulations. Suppose that either $K = \mathbf{R}$, $k = 1$, $R = \mathbf{Z}_2$, or $K = \mathbf{C}$, $k = 2$, and $R = \mathbf{Z}$. Let X be a compact space, and ξ a K -line bundle on X . Let $c \in H^k(X, R)$ be the first Stiefel-Whitney class of ξ if $K = \mathbf{R}$, or the first Chern class of ξ if $K = \mathbf{C}$. The following argument must be well known, but I have not been able to find a specific reference:

LEMMA. *If ξ can be generated by n sections then the n -fold cup product $c^n \in H^{nk}(X, R)$ is zero.*

PROOF. Let $\varphi_\xi: X \rightarrow K\mathbf{P}^\infty$ be the classifying map for ξ . The n sections define a factorization

$$\begin{array}{ccc} X & \xrightarrow{\varphi_\xi} & K\mathbf{P}^\infty \\ \psi \searrow & & \nearrow i \\ & K\mathbf{P}^{n-1} & \end{array}$$

so that $c = \psi^* \hat{c}$ for some $\hat{c} \in H^k(K\mathbf{P}^\infty, R)$. But $H^{nk}(K\mathbf{P}^{n-1}, R) = 0$. Q.E.D.

THEOREM 4.1. *Let X be any compact Hausdorff space which can be covered by finitely many closed subspaces X_1, \dots, X_n such that $H^k(X_i, R) = 0$ for each i . Suppose that for any distinct i, i', i'' , every connected component of $X_i \cap X_{i'} \cap X_{i''}$ is a connected component of $X_i \cap X_{i'}$. Then every K -line bundle over X can be written as a tensor product $L_1 \otimes \dots \otimes L_{n-1}$ where each L_i can be generated by two sections.*

PROOF. By Swan's Theorem [11], every vector bundle on X is a pullback of a vector bundle over $\hat{X} = \text{Spec}(C(X, K))$ under the natural inclusion $X \hookrightarrow \hat{X}$. Because X is normal, each X_i is the zero set of some continuous function, so $\{\hat{X}_i\}_{i=1, \dots, n}$ is a closed covering of \hat{X} . Each $\text{Pic}(\hat{X}_i)$ is trivial because we assumed $H^k(X_i, R) = 0$, so, by Corollary 2.5, every line bundle over \hat{X} is of the required form, whence every line bundle over X is also. Q.E.D.

COROLLARY 4.2. *If X is a space which can be covered as in the statement of the theorem, then $H^k(X, R)$ is generated as a group by elements c satisfying $c \cap c = 0$.*

COROLLARY 4.3. *Neither \mathbf{RP}^n nor \mathbf{CP}^n can be covered as in the statement of the theorem, for $n \geq 2$.*

COROLLARY 4.4. *For $n \geq 2$, any triangulation of \mathbf{RP}^n or \mathbf{CP}^n includes three n -simplices A, B, C such that $\emptyset \neq A \cap B \cap C \neq A \cap B$.*

PROOF. By "fattening things up" where necessary, we may assume that every simplex is a face of an n -simplex, so the n -simplices cover. Q.E.D.

REMARK. Let X be any space for which Swan's Theorem holds. Set

$L(X)$ = the Lusternik-Schnirelman category of X (i.e., the smallest number of closed contractible sets needed to cover X).

$V(X)$ = the maximum, as ξ varies over all K -line bundles over X , of the minimal number of sections necessary to generate ξ .

$C(X) = \text{cuplong}(X)$ = the largest integer n such that for some field F and elements $c_i \in H^{k_i}(X, F)$, $k_i > 0$, $1 \leq i \leq n$, we have $c_1 \cup \dots \cup c_n \neq 0$.

$D(X)$ = the largest integer n such that there exists $c \in H^k(X, R)$ with $c^n \neq 0$.

Then the sorts of arguments made above show that $L(X) \geq V(X) \geq D(X) + 1$. It is known [9] that $L(X) \geq C(X) + 1$ and obvious that $C(X) \geq D(X)$, so we have an array

$$\begin{array}{ccc} & L(X) & \\ & / \quad \backslash & \\ V(X) & & C(X) + 1 \\ & \backslash \quad / & \\ & D(X) + 1 & \end{array}$$

where each number is greater than those displayed below it.

We also have

THEOREM 4.5. *Let X be any compact Hausdorff space which can be covered by finitely many closed subspaces X_1, \dots, X_n such that $\tilde{K}_0(X_i) = 0$ for each i . Suppose that for any distinct i, i', i'' , every connected component of $X_i \cap X_{i'} \cap X_{i''}$ is a connected component of $X_i \cap X_{i'}$. Then $\tilde{K}_0(X) = \text{Pic}(X)$ (where K_0 and Pic refer to the topological K_0 and Picard groups).*

PROOF. This can be proved just as Theorem 4.1 is proved, invoking Corollary 3.3 instead of Corollary 2.5.

B. Cohomology of complex varieties.

THEOREM 4.6. *Let X be a complex affine variety with irreducible components X_1, \dots, X_n satisfying the hypotheses of Theorem 2.1. Suppose that there is a group G such that for each i, j the group of units on $X_i \cap X_j$ is isomorphic to G . Suppose also that for each i , $\text{Pic}(X_i) = 0 = H^1(X_i, G)$. (For example, the X_i could all be nonsingular and contractible—see Gurjar [4].) Then $\text{Pic}(X) = H^1(X, G)$.*

PROOF. Consider the sequences

$$\begin{array}{ccccccc} 0 & \rightarrow & \check{H}^1(\mathcal{X}, \mathcal{O}_X^*) & \rightarrow & \text{Pic}(X) & \rightarrow & \bigoplus_{i=1}^n \text{Pic}(X_i) = 0 \\ & & \parallel & & & & \\ 0 & \rightarrow & \check{H}^1(\mathcal{X}, G) & \rightarrow & H^1(X, G) & \rightarrow & \bigoplus_{i=1}^n H^1(X_i, G) = 0 \end{array}$$

Here the top row is the exact sequence from Theorem 2.1, and the bottom row is the standard exact sequence from topology [1] (actually, this sequence requires the X_i to be open, but this can be arranged by embedding X in \mathbb{C}^N and then replacing X and the X_i by sufficiently small open neighborhoods of themselves, which does not affect the nerve of \mathcal{X} or the singular cohomology groups). The indicated equalities hold by assumption and the theorem follows immediately. Q.E.D.

REMARK. This does not follow directly from Theorem 1.5, as that would require an assumption about $X_i \cap (\bigcup_{j \in S} X_j)$ for various subsets $S \subset \{1, \dots, n\}$.

THEOREM 4.7. *Suppose that X is a complex affine variety and that there is a group G such that for each i, j we have $K_1(X_i \times_X X_j) = G$. Suppose also that for each i there is an injection $\check{K}_0(X_i) \hookrightarrow H^1(X_i, G)$, and that the hypothesis of Corollary 2.5 is satisfied. Then there is an injection $\check{K}_0(X) \hookrightarrow H^1(X, G)$.*

PROOF. Invoke Corollary 3.3.

C. Torsion in Picard groups.

THEOREM 4.8. *Let K be an affine scheme with irreducible components X_1, \dots, X_n . Suppose that there is a group $G \subset \mathcal{O}^*$ such that for each i, j the units on $X_i \times_X X_j$ are isomorphic to G , and suppose also that $\text{Pic}(X_i) = 0$ for each i . Let $k > 0$ be an integer. Then:*

- (1) *If G has no k -torsion, then neither does $\text{Pic}(X)$.*
- (2) *If G is a k -torsion group, then so is $\text{Pic}(X)$.*

PROOF. Let $\bar{X} = X \otimes_{\mathbb{Z}} \mathbb{C}$ and let $\bar{\mathcal{X}} = \{\bar{X}_1, \dots, \bar{X}_n\}$. Then by Theorem 2.1, $\text{Pic}(X) = \check{H}^1(\mathcal{X}, \mathcal{O}^*)$. By assumption this is the same as $\check{H}^1(\mathcal{X}, G)$, which it is easy to see is the same as $\check{H}^1(\bar{\mathcal{X}}, G)$. But this is a subgroup of $H^1(\bar{X}, G)$ as in the proof of Theorem 4.6, and it is well known [1] that $H^1(\bar{X}, G) = \text{Hom}_{\mathbb{Z}}(H_1(\bar{X}, \mathbb{Z}), G)$ which completes the proof. Q.E.D.

REFERENCES

1. A. Dold, *Lectures on algebraic topology*, Springer-Verlag, Berlin, 1972.
2. E. G. Evans, Jr., *Projective modules as fiber bundles*, Proc. Amer. Math. Soc. **27** (1971), 623–626.
3. R. Fossum, *Vector bundles over spheres are algebraic*, Invent. Math. **8** (1969), 222–225.
4. R. Gurjar, Thesis, University of Chicago, 1979.
5. K. Lønsted, *An algebraization of vector bundles on compact manifolds*, J. Pure Appl. Algebra **2** (1972), 193–207.
6. ———, *Vector bundles over finite CW complexes are algebraic*, Proc. Amer. Math. Soc. **105** (1962), 264–277.
7. J. Milnor, *Introduction to algebraic K-theory*, Princeton Univ. Press, Princeton, N. J., 1971.
8. C. Mulvey, *A generalization of Swan's Theorem*, Math. Z. **151** (1976), 57–70.
9. R. S. Palais, *Lusternik-Schnirelman theory on Banach manifolds*, Topology **5** (1966), 115–132.
10. J.-P. Serre, *Faisceaux algébriques cohérents*, Ann. of Math. (2) **6** (1955), 197–278.
11. R. G. Swan, *Vector bundles and projective modules*, Trans. Amer. Math. Soc. **105** (1962), 264–277.
12. ———, *Topological examples of projective modules*, Trans. Amer. Math. Soc. **230** (1977), 201–234.
13. R. G. Swan and J. Towber, *A class of projective modules which are nearly free*, J. Algebra **36** (1975), 427–434.

DEPARTMENT OF ECONOMICS, UNIVERSITY OF CHICAGO, CHICAGO, ILLINOIS 60637