

A TWO CARDINAL THEOREM FOR HOMOGENEOUS SETS AND THE ELIMINATION OF MALITZ QUANTIFIERS

BY

PHILIPP ROTHMALER AND PETER TUSCHIK

ABSTRACT. Sufficient conditions for the eliminability of Malitz quantifiers in a complete first order theory are given. Proving that certain superstable and not ω -stable theories satisfy these conditions, a question of Baldwin and Kueker is answered negatively.

Introduction. The aim of the present paper is to give a first approximation to the problem of finding natural conditions for a theory in the first order language to admit the elimination of certain Malitz quantifiers in the same sense as the substructure completeness theorem does for elimination of elementary quantifiers. In other words, we look for the classes of first order theories which remain essentially the same when we add certain Malitz quantifiers. Thus, the present paper is a contribution to first order model theory, and not to logics with additional quantifiers. Our investigations were inspired by the second author's mostly unpublished work (cf. [TU1] and [TU2]) concerning the eliminability of the quantifiers "there are \aleph_α many" (= Malitz quantifiers for 1-tuples) which we intended to extend to the general case. Although we did not completely succeed in our design, we obtained some partial results which we present in the following order.

§1 contains definitions, conventions, central properties related to the quantifiers, and an eliminability condition for Ramsey quantifiers (= Malitz quantifiers in the \aleph_0 -interpretation). §2 is devoted to a two cardinal theorem for maximally homogeneous sets (without any reference to quantifiers). §3 is an application of §2 to the eliminability problem. §4 provides a negative answer to a question of Baldwin and Kueker [BK, Question 4]. Open questions are scattered about the paper. Our results were obtained independently of [BK] in September–October 1979. For further historical remarks, see below.

1. Preliminaries. In our notation we follow primarily [SH] with the following exceptions.

A, B, \dots, M, N are models and we do not distinguish between a model and its universe. X, Y are sets; u, v, x, z are variables; $\bar{u}, \bar{v}, \bar{x}$ are finite sequences (tuples) of variables. y is reserved for an element of a homogeneous set. Denote by $l(\bar{z})$ the

Received by the editors January 2, 1980.

AMS (MOS) subject classifications (1970). Primary 02H05, 02B20, 02G15, 02G20.

Key words and phrases. Elimination of generalized quantifiers, first order theories, f.c.p., two cardinal theorem, stability, categoricity, finite axiomatizability.

© 1982 American Mathematical Society
 0002-9947/82/0000-0318/\$03.75

length of the sequence \bar{z} . Throughout the paper T denotes a complete theory with infinite models in a countable first order language L . In §4, T has a more specific meaning. We do not distinguish between symbols of L and their interpretations in a given model, nor between elements and their names with one exception: For $X \subseteq A$ we write $\text{Th}(\langle A, \underline{c} \rangle_{c \in X})$ (in short $\text{Th}(\langle A, X \rangle)$) to mark out that \underline{c} is a new constant symbol which is the name of c . $A \models F(\bar{a})$ simply denotes that $\bar{a} \in A$ satisfies the formula $F(\bar{x})$ in A , where it is always assumed that $l(\bar{a}) = l(\bar{x})$. We write $F(A)$ for the set $\{a \in A : A \models F(a)\}$. For $F(x)$ a 1-placed formula $\bar{F}(\bar{x})$ is an abbreviation of $\&_{i=1}^m F(x_i)$, where $\bar{x} = (x_1, \dots, x_m)$.

We use also the following notion of restriction which can be found in the literature. If $F(\bar{x})$, $K(x)$ are formulae of L , $F^K(\bar{x})$, the restriction of F to K , and $A \upharpoonright K(A)$, the restriction of A to $K(A)$, are defined in such a way that for all $\bar{a} \in K(A)$, $A \upharpoonright K(A) \models F(\bar{a})$ iff $A \models F^K(\bar{a})$. For X a set of L -formulae, X^K denotes the set $\{F^K : F \in X\}$.

A subset Y of a model A is said to be *homogeneous* for a formula $\varphi(x_1, \dots, x_m)$ if $A \models \varphi(y_1, \dots, y_m)$ for all (y_1, \dots, y_m) of Y^m . Y is called *maximally homogeneous* for φ in A if Y is homogeneous for φ in A and no proper superset of Y , which is also a subset of A , is homogeneous for φ in A .

Let us now introduce the interpretations of the *Malitz quantifiers* (for syntactical and other details we refer to [MM]). We define $A \models Q_\alpha^m x_1 \dots x_m \varphi(x_1, \dots, x_m)$ if there is a subset Y in A of cardinality \aleph_α which is homogeneous for φ in A . For $\alpha = 0$ the quantifiers are also called *Ramsey* quantifiers.

In [MM] there is considered in fact another interpretation connected with the following weaker notion of homogeneity. Y is called *weakly homogeneous* for φ in A if $A \models (\&_{1 \leq i < j \leq m} y_i \neq y_j) \rightarrow \varphi(y_1, \dots, y_m)$ for all $(y_1, \dots, y_m) \in Y^m$. Denote $+Q^m$ the appropriate quantifier. Then it is easily seen that the following hold.

$$\begin{aligned} A \models +Q_\alpha^m x_1 \dots x_m \varphi(x_1, \dots, x_m) \\ \Leftrightarrow Q_\alpha^m x_1 \dots x_m \left(\left(\&_{1 \leq i < j \leq m} x_i \neq x_j \right) \rightarrow \varphi(x_1, \dots, x_m) \right), \\ A \models Q_\alpha^m x_1 \dots x_m \varphi(x_1, \dots, x_m) \Leftrightarrow +Q_\alpha^m x_1 \dots x_m \left(\&_{\bar{u} \in J} \varphi(\bar{u}) \right), \end{aligned}$$

where J is the set of all sequences of elements from $\{x_1, \dots, x_m\}$ having length m . Thus, Q_α^m is eliminable iff $+Q_\alpha^m$ is eliminable. Notice, Q_α^1 is the usual quantifier “there are \aleph_α many”.

If $\varphi(\bar{x}, \bar{v}) \in L$, call the quantifier $Q_\alpha^m \varphi$ -definable in T if there is an n such that $A \models \forall \bar{v} (H_n^m \bar{x} \varphi(\bar{x}, \bar{v}) \rightarrow Q_\alpha^m \bar{x} \varphi(\bar{x}, \bar{v}))$ for all models A of T of cardinality $\geq \aleph_\alpha$, where $H_n^m \bar{x} \varphi(\bar{x}, \bar{v})$ is the first order expression of “there is a set of cardinality $\geq n$ which is homogeneous for φ ”. Call the quantifier $Q_\alpha^m \varphi$ -eliminable in T if there is a first order formula $\psi(\bar{v})$ such that $A \models \forall \bar{v} (Q_\alpha^m \bar{x} \varphi(\bar{x}, \bar{v}) \leftrightarrow \psi(\bar{v}))$ for all models A of T of cardinality $\geq \aleph_\alpha$.

The following observation is basic for elimination results.

LEMMA. *The quantifier Q_α^m is φ -eliminable in T iff it is φ -definable in T .*

For $\alpha = 0$ this is Theorem 1 of [BK]. However, the second author was familiar with various versions of it. In fact the lemma holds for every quantifier we deal with in the present paper (mutatis mutandis in the sentence $H_n^m \bar{x} \varphi(\bar{x}, \bar{v})$) and we will use it without making mention of it. This observation is true also for the *equicardinality interpretation* of Q^m which is defined by $A \models Q_c^m \bar{x} \varphi(\bar{x})$ iff $A \models Q_\alpha^m \bar{x} \varphi(\bar{x})$, where $|A| = \aleph_\alpha$ (cf. [BK]).

Following [TU2], a formula $\varphi(x, \bar{v})$ is called *graduated* in T if Q_0^1 is φ -definable in T (i.e. if $\varphi(x, \bar{a})$ defines a finite set then this set has cardinality $< n$, where n can be chosen universally for all parameters \bar{a}). A theory is *graduated* if all formulae are graduated in T . Now it is easily seen that a theory admits the elimination of Q_0^1 iff it is graduated [TU2].

It turns out that the following property introduced by Keisler is a good generalization of nongraduatedness. $\varphi(\bar{x}, \bar{v})$ has the *finite cover property* (f.c.p.) if in some model A of T for arbitrarily large n there exist $\bar{a}_0, \dots, \bar{a}_{n-1}$ such that

$$A \models \bigwedge_{j < n} \left(\bigwedge_{i \neq j} \varphi(\bar{x}, \bar{a}_i) \right) \& \neg \exists \bar{x} \bigwedge_{i < n} \varphi(\bar{x}, \bar{a}_i).$$

Note that $\psi(x, \bar{u}) \leftarrow (\varphi(x, \bar{v}) \& x \neq u)$, where $\bar{u} = \bar{v} \cap u$, has f.c.p. if $\varphi(x, \bar{v})$ is not graduated. By definition, T has the f.c.p. if some formula $\varphi(x, \bar{v})$ has the f.c.p.

Feferman introduced another quantifier which is closely related to the f.c.p. Let $\varphi(x, z)$ be an L -formula. Then $\forall z (\varphi(x_1, z) \leftrightarrow \varphi(x_2, z))$ determines an equivalence relation. We define $A \models Q_\alpha^E xz \varphi(x, z)$ if this equivalence relation has at least \aleph_α distinct equivalence classes on A (cf. [FE, p. 129]).

We will use another version of this quantifier defined by $A \models E_\alpha xz \varphi(x, z)$ if $\varphi(x, z)$ is an equivalence relation with $\geq \aleph_\alpha$ many equivalence classes on A .

Eliminability of Q_α^E is equivalent to eliminability of E_α , since they are definable by each other:

$$\begin{aligned} A \models E_\alpha xz \varphi(x, z) &\leftrightarrow (Q_\alpha^E xz \varphi(x, z) \& el(\varphi)), \\ A \models Q_\alpha^E xz \varphi(x, z) &\leftrightarrow E_\alpha xz (\forall v (\varphi(x, v) \leftrightarrow \varphi(z, v))), \end{aligned}$$

where $el(\varphi)$ is a first order sentence which says " $\varphi(x, z)$ is an equivalence relation". Note that Q_α^1 is definable using E_α .

Having in mind the equivalence of definability and eliminability of the quantifier E_α , we are able to give the following restatement of Shelah's f.c.p. theorem (cf. [SH, II, 4.4]):

THEOREM (THE F.C.P. THEOREM). *Let T be stable. Then the following are equivalent:*

- (i) T has the f.c.p., i.e. some $\varphi(x, \bar{v})$ has the f.c.p.
- (ii)_m Some $\varphi(\bar{x}, \bar{v})$ has the f.c.p., $l(\bar{x}) = m$.
- (iii) E_0 is not eliminable in T .¹

Obviously, E_α is definable by Q_α^2 . Thus eliminability of Q_α^2 implies eliminability of E_α . Further, one can directly prove that a theory without the f.c.p. admits the

¹One can introduce E_α^m also for m -tuples. Then (iii)_m is equivalent to the other conditions.

elimination of all the Q_0^m . So, by the f.c.p. theorem, we get for T a stable theory, T does not have the f.c.p. iff Q_0^m is eliminable in T ($m > 1$) iff E_0 is eliminable in T .² This is Theorem 6 of [BK]. We proved it independently.

Note that an \aleph_1 -categorical theory does not have the f.c.p. (Keisler's theorem), and all unstable theories do have it.

There are examples of unstable theories admitting the elimination of all Q_0^m (e.g. \aleph_0 -categorical theories; see also [BK]), but at present it is not known what class of unstable theories is determined by this property.

2. The two cardinal theorem. This section partially resembles §22 of [SA]. So we will omit some proofs. First of all, let us prove some technical lemmas. Let R, K be new unary predicate symbols, \underline{c} a new constant symbol. Then $S(\underline{c}, \varphi(\bar{x}, \bar{a}))$ denotes the union of the following sets:

- (1) $\{\forall \bar{v}(\bar{K}(\bar{v}) \rightarrow (F(\bar{v}) \leftrightarrow F^K(\bar{v}))) : F(\bar{v}) \in L\} \cup T$;
- (2) $\{\neg K(\underline{c})\}$;
- (3) $\{\forall x(R(x) \rightarrow K(x)) \& \bar{K}(\bar{a})\}$;
- (4) $\{\forall \bar{x}(\bar{R}(\bar{x}) \rightarrow \varphi(\bar{x}, \bar{a}))\}$;
- (5) $\{\forall x_0(\forall x_1 \dots x_{m-1}(\&_{i=1}^{m-1} R(x_i) \rightarrow \&_{\bar{u} \in J} \varphi(\bar{u}, \bar{a})) \rightarrow R(x_0))\}$,

where $J = \{x_0, \dots, x_{m-1}\}^m$.

LEMMA 2.1. *B is a model of $S(\underline{c}, \varphi(\bar{x}, \bar{a}))$ iff B is a model of T , there is $A \prec B$ with a subset Y (of A) such that $A \neq B$, and Y is maximally homogeneous for $\varphi(\bar{x}, \bar{a})$ in B , and $\bar{a} \in A$.*

PROOF. Let B be a model of that set, A the restriction of B to $K(B)$, Y the set $R(B)$. Then $A \prec B$, $A \neq B$, $\bar{a} \in A \supseteq Y$ by (1)–(3) resp. (4) guarantees homogeneity, and (5) maximality of Y . Similarly the other direction. Q.E.D.

Now we will prove a series of analoga of lemmata from [SA, §22]. Notice, a maximal homogeneous set for a formula $\varphi(x)$ (in one free variable) in A is $\varphi(A)$ itself. Thus the maximal homogeneous sets for formulae in one free variable are exactly the definable sets. This is the point in the analogy mentioned above: We will extend the two cardinal theorem to formulae in m free variables.

The next lemma is the appropriate analogon of [SA, 22.2].

LEMMA 2.2. *If $A \prec B$, $A \neq B$, $\bar{a} \cup Y \subseteq A$, B countable, $b \in B - A$, p a complete n -type over $\{a_i : i < k\} \subseteq A$, and Y maximally homogeneous for $\varphi(\bar{x}, \bar{a})$ in B , then there are $A' \succ A$, $B' \succ B$, Y' with the same properties, but p realized in A' .*

PROOF. Let S be the union of the following sets:

- (6) $\text{Th}(\langle B, B \rangle) \cup \{K(a) : a \in A\}$;
- (7) $(\text{Th}(\langle A, A \rangle))^K$;
- (8) $S(b, \varphi(\bar{x}, \bar{a}))$, where b and \bar{a} are as in the hypothesis (notice, b occurs also in (6));

² E_α^m is definable using $Q_\alpha^{*,2,m}$ of [BK, p. 7]. The eliminability of E_0^m is also equivalent to \neg f.c.p. in the stable case by Theorem 7 of [BK].

(9) $\{F(\bar{c}) \& \bar{K}(\bar{c}): F(\bar{x}) \in p\}$, where \bar{c} is an n -tuple of new constant symbols not occurring in $\{\underline{b}': b' \in B\}$.

S is consistent, as every finite subset is realized in B for the interpretation $K(B) = A$, $R(B) = Y$. Take a countable model B' of S , A' the restriction of B' to $K(B')$, $Y' = R(B')$. Then $B' \succ B$ by (6), $A' \succ A$ by (6) and (7), p is realized in A' by (9), $b \in B' - A'$ by (2) and (8). The rest follows by (8) and Lemma 2.1. Q.E.D.

The next lemmata are the analoga of 22.3, 22.4 of [SA], resp. We omit the proofs, since one only has to put " $Y \subseteq N$ and Y is maximally homogeneous for $\varphi(\bar{x}, \bar{a})$ in M " instead of " $R(N) = R(M)$ " whenever $N < M$ and $N \neq M$ in the original proofs.

LEMMA 2.3. *Let A , B , and Y be as in the hypothesis of Lemma 2.2. Then there are A' , B' , Y' with the same properties such that $A' \succ A$, $B' \succ B$, $A' \cong B'$, A' and B' are countable homogeneous models.*

This lemma tells us that w.l.o.g. we can assume A and B to be isomorphic and homogeneous models in the situation of the next lemma.

LEMMA 2.4. *Let A , B , Y be as in Lemma 2.3. Then there are a model $C \succ B$ and a maximal homogeneous set Y' for $\varphi(\bar{x}, \bar{a})$ in C such that $|C| = \aleph_1$ and $|Y'| \leq \aleph_0$. If $|Y| = \aleph_0$, then one can assume $|Y'| = \aleph_0$.*

Now we introduce a generalization of the notion of a Vaughtian pair.

DEFINITION 2.5. (i) (A, B) is called a *generalized Vaughtian pair* for $\varphi(\bar{x}, \bar{a})$ if $A < B$, $A \neq B$, $\bar{a} \in A$, and there is an infinite subset Y of A which is maximally homogeneous for $\varphi(\bar{x}, \bar{a})$ in B .

(ii) (A, B) is called a *generalized Vaughtian pair (of index m)* if (A, B) is a generalized Vaughtian pair for a formula $\varphi(\bar{x}, \bar{a})$ with $\bar{a} \in A$ (and $l(\bar{x}) = m$).

(iii) Let $\varphi(\bar{x})$ be a fixed formula of L . T is said to *admit a pair of cardinals* (μ, χ) if T has a model B of cardinality μ and there is a subset Y of B of cardinality χ such that Y is maximally homogeneous for φ in B .

Notice, a generalized Vaughtian pair of index 1 is a usual Vaughtian pair.

LEMMA 2.6. *If there is a generalized Vaughtian pair for a certain formula, then there is a countable one for that formula.*

PROOF. By the hypothesis and Lemma 2.1 the set $S(\underline{c}, \varphi(\bar{x}, \bar{a})) \cup \{R(\underline{c}_i) \& \underline{c}_i \neq \underline{c}_j: i < j < \omega\}$ is consistent. Take a countable model and define a generalized Vaughtian pair as in Lemma 2.1. Q.E.D.

DEFINITION 2.7. (i) A model A is called φ -*regular* if any infinite set which is maximally homogeneous for φ in A has the cardinality $|A|$. A is called φ -*singular* if it is not φ -regular.

(ii) A model A is called m -*regular* if it is regular for all formulae in m free variables.

(iii) A model A is called *regular* if it is m -regular for all m .

(iv) A theory is called φ -regular if all models of T are φ -regular. Similarly for m -regularity and regularity of T .

Now we are able to prove the main theorem of this section.

THEOREM 2.8 (THE TWO CARDINAL THEOREM FOR HOMOGENEOUS SETS). (i) T has no Vaughtian pairs for $\varphi(\bar{x}, \bar{a})$ iff any model containing \bar{a} is φ -regular.

(ii) T has no Vaughtian pairs (of index m) iff any model is φ -regular for all formulae φ with (m free variables and) parameters from that model.

(iii) If T admits (μ, χ) for $\mu > \chi \geq \aleph_0$, T admits (\aleph_1, \aleph_0) .

PROOF. (i) \Rightarrow . Suppose to the contrary there is an infinite set $Y \subseteq A$ of cardinality less than $|A|$, such that Y is maximally homogeneous for $\varphi(\bar{x}, \bar{a})$ in A . Let B be an elementary submodel of A extending Y which has the same cardinality as Y . Then $B \neq A$ and thus (B, A) is a generalized Vaughtian pair for $\varphi(\bar{x}, \bar{a})$.

\Leftarrow . Suppose (A, B) is a generalized Vaughtian pair for $\varphi(\bar{x}, \bar{a})$. By Lemma 2.6, w.l.o.g. B is countable. By Lemma 2.4, there is a model $C \succ B$ and a maximal homogeneous set Y for $\varphi(\bar{x}, \bar{a})$ in C such that $|C| = \aleph_1$ and $|Y| = \aleph_0$, contradicting the hypothesis.

(ii) follows from (i).

(iii) Suppose $B \models T$, $|B| = \mu$, $Y \subseteq B$ is a maximal homogeneous set for $\varphi(\bar{x}, \bar{a})$ in B , $|Y| = \chi$. By the first half of (i), there is a generalized Vaughtian pair (A, B) for φ which is countable by Lemma 2.6. Now, by Lemma 2.4, T admits (\aleph_1, \aleph_0) . Q.E.D.

The reader will not be very surprised by the following.

THEOREM 2.9. Any model of an \aleph_1 -categorical theory is regular (or, equivalently, an \aleph_1 -categorical theory has no generalized Vaughtian pairs).

PROOF. By definition, a singular model is uncountable. Now the assertion follows from the next lemma. Q.E.D.

LEMMA 2.10. Any saturated model is regular.

PROOF. Let A be a singular model which is saturated, $Y \subseteq A$ a maximal homogeneous set for a formula in A which has infinite cardinality less than $|A|$. Let $p(x)$ be the type saying " $x \notin Y$ and $\{x\} \cup Y$ is homogeneous for that formula". Then p is realized in A , contradicting the maximality of Y . Q.E.D.

The class of regular theories seems to us to be of interest for further investigation. We close this section with the following observation.

LEMMA 2.11. If all models of T of power \aleph_1 are φ -regular, then T is φ -regular.

PROOF. By Theorem 2.8(iii). Q.E.D.

COROLLARY 2.12. If all models of T of power \aleph_1 are regular, T is regular.

3. The elimination of Malitz quantifiers. Now we are able to give sufficient conditions for the eliminability of the quantifiers in terms of regularity introduced in the preceding section.

THEOREM 3.1. (i) If every model of T is φ -regular ($\varphi = \varphi(\bar{x}, \bar{v})$, $l(\bar{x}) = m$), then the quantifier Q_c^m , hence also Q_α^m for $\alpha \geq 0$, is φ -eliminable in T .

(ii) If T is m -regular, or, equivalently, T has no generalized Vaughtian pairs of index m , all the quantifiers Q_c^m and Q_α^m are eliminable in T ($\alpha \geq 0$).

(iii) If T is regular, or, equivalently, T has no generalized Vaughtian pairs, the quantifiers Q_c^m and Q_α^m are eliminable in T for all $m \geq 1$ and $\alpha \geq 0$.

PROOF. (i) implies (ii) and (iii). First of all we prove

Claim. If every model of T is φ -regular, then Q_0^m is φ -eliminable in T ($\varphi = \varphi(\bar{x}, \bar{v})$, $l(\bar{x}) = m$).

PROOF OF THE CLAIM. For the contrary, suppose Q_0^m is not φ -eliminable in T . Then for all n there is a model A_n of T and $\bar{a}_n \in A_n$ such that A_n has a maximal homogeneous set for $\varphi(\bar{x}, \bar{a})$ which has finite cardinality $> n$. Then every finite subset of $S(\bar{c}, \varphi(\bar{x}, \bar{a})) \cup \{R(\bar{c}_i) \& \bar{c}_i \neq \bar{c}_j : i < j < \omega\}$ (cf. §2) is consistent (w.l.o.g. A_n is uncountable; put $K(A_n) = B$, where B is a countable elementary submodel of A_n containing the finite maximal homogeneous set).

By Lemma 2.1, every model yields a Vaughtian pair for $\varphi(\bar{x}, \bar{a})$. Thus, by the two cardinal theorem, there is a model which is not φ -regular, whence the claim is proved.

Now suppose Q_c^m is not φ -eliminable in T . Then there is a model B of T with $\bar{a} \in B$ and

$$B \models H_n^m \bar{x} \varphi(\bar{x}, \bar{a}) \& \neg Q_c^m \bar{x} \varphi(\bar{x}, \bar{a}),$$

where n is chosen to satisfy

$$A \models \forall \bar{v} (H_n^m \bar{x} \varphi(\bar{x}, \bar{v}) \rightarrow Q_0^m \bar{x} \varphi(\bar{x}, \bar{v}))$$

for all $A \models T$ using the claim. So

$$B \models Q_0^m \bar{x} \varphi(\bar{x}, \bar{a}) \& \neg Q_c^m \bar{x} \varphi(\bar{x}, \bar{a}),$$

whence B is a φ -singular model, contradicting the hypothesis. Q.E.D.

The converse seems hard to come by, since a formula with more than one free variable ($m > 1$) can have maximal homogeneous sets of different cardinality. However, for $m = 1$ the following were already proved in [TU2].

COROLLARY 3.2. The following properties are equivalent for a theory T :

- (i) Q_1^1 is eliminable;
- (ii) Q_c^1 is eliminable;
- (iii) for all $\alpha \geq 0$, Q_α^1 is eliminable;
- (iv) T has no (usual) Vaughtian pairs, or, equivalently, T is 1-regular.

PROOF. Remember, the usual Vaughtian pairs are exactly the generalized Vaughtian pairs of index 1. Now one direction is Theorem 3.1(ii). For the other, note that a formula in one free variable has only one maximal homogeneous set. Thus, if Q_1^1 is eliminable, every model of power \aleph_1 is 1-regular. By Lemma 2.11, T is regular. Q.E.D.

Similarly one can prove

COROLLARY 3.3. If Q_α^1 is eliminable in T for some $\alpha \geq 0$, Q_0^1 is eliminable in T .

So, eliminability of Q_1^1 is the strongest notion, that of Q_0^1 the weakest, whereas eliminability of the Q_α^1 's is between them.

In case of stability an analogon of Morley's categoricity theorem is true (cf. [TU2]):

PROPOSITION 3.4. *For T a stable theory, T admits the elimination of Q_1^1 iff T admits the elimination of Q_α^1 for some $\alpha \geq 1$.*

We do not know what can happen if T is unstable. The general case presumably depends on set theory. Also we do not know what is the matter in case $m > 1$. Generally, there are two kinds of questions concerning the relative strength of eliminability of Malitz quantifiers:

What is the relation between the eliminability of

(a) Q_α^m and Q_α^n (Q_c^m and Q_c^n) and of

(b) Q_α^m and Q_β^m .

For $\alpha = 0$, there is an answer to (a) for stable theories:

Eliminability of Q_0^2 is equivalent to that of Q_0^m ($m > 1$); cf. §1 or [BK].

For $m = 1$, the present section provides some answers to (b) which are complete when stability is assumed.

The theory RCF of real closed fields is an example of an unstable complete theory which admits elimination of Q_1^1 , but not of Q_1^2 . (RCF has no (usual) Vaughtian pairs; using Q_1^2 one can define the sentence "there is an uncountable discrete subset cofinal in the field" which is consistent with RCF, but not true in \mathbf{R} ; similarly, RCF admits elimination of Q_0^1 , but not of Q_0^2 .)

Perhaps from [GA] one can obtain examples of theories admitting the elimination of Q_α^m , but not of Q_α^{m+1} for all $\alpha > 0$ and $m \geq 1$ (for this one has to show that not only the structures \mathfrak{A} and \mathfrak{B} , but the theory $\text{Th}(\mathfrak{A})$ admits elimination of Q_α^m ; cf. [GA, Lemma 2]).

Let us conclude this section with a theorem which was also proved by Baldwin and Kueker [BK, Theorem 9].

THEOREM 3.5. *An \aleph_1 -categorical theory admits the elimination of Q_c^m and Q_α^m for all $m \geq 1$ and $\alpha \geq 0$.*

PROOF. By Theorem 3.1(iii) and 2.9. Q.E.D.

4. On a question of Baldwin and Kueker. In the present section we will show that an example of Makowsky, which goes back to R. Robinson, is in fact a counterexample to the following question of Baldwin and Kueker [BK, Question 4]:

For T a theory in a finite language, if T admits the elimination of Q_c^m for all $m \geq 1$, must T be \aleph_1 -categorical (or, equivalently, ω -stable)?

While investigating the elimination of Q_1^m the second author had the same question (however for Q_1^m), whereas the first author suggested Makowsky's example for a negative answer.

We would like to thank H. Herre who brought to our attention that [MA] provides an example of a superstable theory without (usual) Vaughtian pairs which is finitely axiomatizable.

Now let us introduce the example. For details and proofs we refer to [MA]. Let $G = F/R$ be a group with F, R free countably generated groups. Let A_G be the graph of the group G , i.e. $A_G = \langle A, f_i: i \in I \rangle$, where $\{f_i: i \in I\}$ is a set of generators of F , f_i is a unary function symbol ($i \in I$), and for all $a \in A$, $f_{i_1} \cdots f_{i_k}(a) = a$ iff $f_{i_1} \cdots f_{i_k} \in R$.

We assume that for all $i \in I$ there is a $j \in I$ such that $f_i f_j(a) = f_j f_i(a) = a$ for all $a \in A$. In other words, we assume that the set of generators of F is closed under taking the inverse.

Let P_1, \dots, P_n be unary predicates on A_G . From now on, $T = \text{Th}(A_G, P_1, \dots, P_n)$, and $T_G = \text{Th}(A_G)$. T is model complete and superstable. Following [MA], call a substructure of a model of T which is generated by a single element a *component* of that model. Denote by $C(a)$ the component generated by a . Then every component is countable and every model of T splits into disjoint components which are all isomorphic relative to L_G , the language of T_G .

Makowsky proved for universal T that T is \aleph_1 -categorical iff it is ω -stable. This follows also from Corollary 4.4. Our aim is to show that T has no generalized Vaughtian pairs if T is universal.

From the model completeness of T we obtain the following.

LEMMA 4.1. *If T is universal, T admits the elimination of (usual) quantifiers.*

Considering various examples of graphs of groups the next lemma becomes evident.

LEMMA 4.2. *A_G is totally homogeneous, i.e. for all $a, b \in A_G$ there is an L_G -automorphism F of A_G mapping a onto b .*

PROOF. Let $1 \in A$ correspond to the unity of the group G . There are representations of a and b through 1 , since A_G is connected: $a = f_{i_1} \cdots f_{i_m}(1)$, $b = f_{j_1} \cdots f_{j_n}(1)$. Now we define F as follows: for $c = f_{k_1} \cdots f_{k_l}(1)$ an arbitrary element of A , set $F(c) = f_{k_1} \cdots f_{k_l} f_{i_m}^{-1} \cdots f_{i_1}^{-1} f_{j_1} \cdots f_{j_n}(1)$. Clearly, $F(a) = b$ and $f(F(c)) = F(f(c))$ for $f \in \{f_i: i \in I\}$, whence F is a homomorphism. Also it is not hard to verify that F is a 1-1 map, since

$$A_g \models \forall x (f_{k_1} \cdots f_{k_l}(x) = x) \leftrightarrow \exists x (f_{k_1} \cdots f_{k_l}(x) = x). \quad \text{Q.E.D.}$$

The following lemma is the main one.

LEMMA 4.3. *Let $B \models T$, $A < B$, $A \neq B$, $Y \subseteq A$, Y homogeneous for $\varphi(\bar{x}, \bar{a})$, $\bar{a} \in A$. If T is universal, Y is not maximally homogeneous for φ in B .*

PROOF. $Y \cup \{c\}$ is homogeneous for φ in C for some proper extension $C \succ A$ and some $c \in C - A$. W.l.o.g. $C = A \cup C(c)$.

By Lemma 4.1, there are $G_i(\bar{x}, \bar{v})$, $R_i(\bar{x})$, and $F_i(\bar{v})$ ($i = 1, \dots, k$) such that

$$T \vdash \varphi(\bar{x}, \bar{v}) \leftrightarrow \bigvee_{i=1}^k (G_i(\bar{x}, \bar{v}) \& R_i(\bar{x}) \& F_i(\bar{v})),$$

and $G_i \in L_G$, $R_i(\bar{x})$ and $F_i(\bar{v})$ are conjunctions of the P_j 's and their negations ($i = 1, \dots, k$). (Generally, there can be L_G -terms in R_i, F_i ; the proof remains the

same.) In B there is a component $C(b)$ elementarily equivalent to $C(c)$, since both are models of T for T is universal. W.l.o.g. $B = A \cup C(b)$. In $C(b)$ there is an element which has the same " $\{P_1, \dots, P_n\}$ -type" as has c . Therefore without loss we can assume that b has that property.

We claim $Y \cup \{b\}$ is homogeneous for φ . For this, let $\bar{y} \in (Y \cup \{b\})^m$ and \bar{y}' be the sequence obtained from \bar{y} by substituting b by c .

We have to show $B \models \varphi(\bar{y}, \bar{a})$. $C(c)$ and $C(b)$ are models of T , in particular, both are models of T_G , whence they are isomorphic to A_G as L_G -structures. Using Lemma 4.2 we choose an L_G -isomorphism $F: C \cong B$ which maps c onto b and is the identity map on A . Thus, $B \models G_i(\bar{y}, \bar{a})$ iff $C \models G_i(\bar{y}', \bar{a})$ for all i . Also, $B \models R_i(\bar{y})$ iff $C \models R_i(\bar{y}')$, since by construction \bar{y} and \bar{y}' have the same $\{P_1, \dots, P_n\}$ -type. So we have $C \models \varphi(\bar{y}', \bar{a})$ iff $B \models \varphi(\bar{y}, \bar{a})$. But $C \models \varphi(\bar{y}', \bar{a})$ for $Y \cup \{c\}$ is homogeneous for φ , whence the result. Q.E.D.

COROLLARY 4.4. *Let T be universal. Then:*

- (i) *T is regular, or, equivalently, T has no generalized Vaughtian pairs.*
- (ii) *T admits the elimination of Q_c^m and Q_α^m for all $m \geq 1$ and $\alpha \geq 0$.*

PROOF. (i) follows from Lemma 4.3, (ii) from (i) and Theorem 3.1(iii). Q.E.D.

Using a domino of R. Robinson, Makowsky showed that there is a universal theory $T = \text{Th}(\langle A_G, P_1, \dots, P_n \rangle)$ in a finite language which is finitely axiomatizable and not ω -stable (cf. [MA, p. 200]). Thus we obtain the following theorem which yields the desired answer to the question mentioned in the beginning of this section.

THEOREM 4.5. *There is a theory T in a finite language such that:*

- (i) *T is superstable, but not ω -stable;*
- (ii) *T has no generalized Vaughtian pairs;*
- (iii) *T admits the elimination of Q_c^m and Q_α^m for all $m \geq 1$ and $\alpha \geq 0$.*
- (iv) *T is finitely axiomatizable.*

APPENDIX. One can define homogeneity and regularity for $\psi(\bar{x}_1, \dots, \bar{x}_m, \bar{v})$ and extend 2.8 to that case. Then 3.1 extends to $Q_\alpha^{*,m,n}$ of [BK]. The converse of 3.1 holds if all infinite maximal homogeneous sets have the same cardinality. Of course, this is true for an equivalence relation (a maximal homogeneous set is a set of representatives of each class). Thus, eliminability of E_1 implies that of E_0 . Now, by the f.c.p. theorem, for T a stable theory, if T admits elimination of Q_1^2 , T does not have the f.c.p., whence all Q_0^m are eliminable.

ADDED IN PROOF. We have proved the converse of Theorem 3.1(iii) for stable theories T . This will appear in a subsequent paper.

REFERENCES

- [BK] J. T. Baldwin and D. W. Kueker, *Ramsey quantifiers and the finite cover property*, Pacific J. Math. **90** (1980), 11–19.
- [FE] S. Feferman, *Two notes on abstract model theory*. II, Fund. Math. **89** (1975), 111–130.
- [GA] S. Garavaglia, *Relative strength of Malitz quantifiers* (preprint).
- [MA] J. A. Makowsky, *On some conjectures connected with complete sentences*, Fund. Math. **81** (1974), 193–202.

[MM] M. Magidor and J. Malitz, *Compact extensions of $L(Q)$* , Ann. Math. Logic **11** (1977), 217–261.

[SA] G. E. Sacks, *Saturated model theory*, Benjamin, New York, 1972.

[SH] S. Shelah, *Classification theory and the number of non-isomorphic models*, North-Holland, Amsterdam, 1978.

[TU1] P. Tuschik, \aleph_1 -Kategorizität, Dissertation (A), Humboldt-Univ., Berlin, 1975.

[TU2] ———, *Elimination of cardinality quantifiers* (contributed paper at the 4th Bierutowice Conf., Karpacz, Poland, September 1979).

AKADEMIE DER WISSENSCHAFTEN DER DDR, INSTITUT FÜR MATHEMATIK, 1080 BERLIN, GERMAN DEMOCRATIC REPUBLIC

SEKTION MATHEMATIK, HUMBOLDT-UNIVERSITÄT ZU BERLIN, 1086 BERLIN, GERMAN DEMOKRATIK REPUBLIK