

## ITERATING THE BASIC CONSTRUCTION

MIHAI PIMSNER AND SORIN POPA

**ABSTRACT.** Let  $N \subset M$  be a pair of type  $\text{II}_1$  factors with finite Jones' index and  $N \subset M \subset M_1 \subset M_2 \subset \cdots \subset M_n \subset \cdots \subset M_{2n+1}$  be the associated tower of type  $\text{II}_1$  factors obtained by iterating Jones' basic construction. We give an explicit formula of a projection in  $M_{2n+1}$  which implements the conditional expectation of  $M_n$  onto  $N$ , thus showing that  $M_{2n+1}$  comes naturally from the basic construction associated to the pair  $N \subset M_n$ . From this we deduce several properties of the relative commutant  $N' \cap M_n$ .

**Introduction.** Let  $N \subset M$  be a pair of finite factors. Jones defined in [1] the index  $[M : N]$  of  $N$  in  $M$  to be the coupling constant of  $N$  in its representation on  $L^2(M)$ . If this index is finite, then the trace preserving conditional expectation of  $M$  onto  $N$ , regarded as an operator on  $L^2(M)$ , generates together with  $M$  a finite factor  $M_1$ . This factor is called in Jones' terminology the extension of  $M$  by  $N$  and the construction of  $M_1$  from  $M$  and  $N$ , the basic construction. The pair  $M \subset M_1$  has the remarkable property that  $[M_1 : M] = [M : N]$ , so this procedure may be iterated to get an increasing sequence of finite factors  $N \subset M \subset M_1 \subset M_2 \subset \cdots$  and together with it a sequence of projections  $e_i \in M_{i+1}$ ,  $i \geq 0$ , implementing the conditional expectations at consecutive steps.

We prove in this paper that in this sequence of factors the basic construction arises periodically from  $n$  to  $n$  steps, for any  $n$ . In fact we give a formula for a projection  $f_n$  in  $M_{2n+1}$  that implements the conditional expectation of  $M_n$  onto  $N$ :  $f_n$  is a scalar multiple of the word of maximal length in  $\{e_i\}_{0 \leq i \leq 2n}$ , namely

$$f_n = [M : N]^{n(n+1)/2} (e_n e_{n-1} \cdots e_0) (e_{n+1} e_n \cdots e_1) \cdots (e_{2n} \cdots e_n).$$

We mention that this result was independently obtained by A. Ocneanu [2]. We apply this theorem to show that if the logarithm of the index  $[M : N]$  equals the relative entropy  $H(M|N)$  considered in [3], then one also has

$$H(M_n|N) = \ln[M_n : N] \quad \text{for every } n.$$

Since this equality characterizes an extremal case for an inclusion of factors, from the analysis of a similar situation in [3] we deduce several properties of the inclusion  $N \subset M_n$  and of the relative commutant  $N' \cap M_n$ .

**1. Preliminaries.** Throughout this paper  $M$  will be a finite factor with normalized trace  $\tau$ ,  $\tau(1) = 1$ . We denote by  $\|x\|_2 = \tau(x^*x)^{1/2}$ ,  $x \in M$ , the Hilbert norm given by  $\tau$  and by  $L^2(M, \tau)$  the Hilbert space completion of  $M$  in this norm. The canonical conjugation of  $L^2(M, \tau)$  is denoted by  $J$ . It acts on  $M \subset L^2(M, \tau)$

---

Received by the editors February 23, 1987.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 46L35; Secondary 46L10.

©1988 American Mathematical Society  
0002-9947/88 \$1.00 + \$.25 per page

by  $Jx = x^*$  and satisfies  $JMJ = M'$ . In fact, if we regard  $M$  as acting by left multiplication on  $L^2(M, \tau)$  then for  $x \in M$ ,  $JxJ$  is the operator of right multiplication by  $x^*$ .

$N \subset M$  will denote a subfactor of  $M$  with  $1_N = 1_M$  and  $E_N$  will be the unique normal trace preserving conditional expectation of  $M$  onto  $N$ . Note that  $E_N$  is just the restriction to  $M \subset L^2(M, \tau)$  of the orthogonal projection  $e_N$  of  $L^2(M, \tau)$  onto  $L^2(N, \tau)$  (the closure of  $N$  in  $L^2(M, \tau)$ ). The conditional expectation  $E_N$ , the projection  $e_N$  and the conjugation  $J$  are related by the properties

- (i) If  $x \in M$  then  $x \in N$  iff  $e_N x = x e_N$ .
- (ii)  $e_N x e_N = E_N(x) e_N$ ,  $x \in M$ .
- (iii)  $J$  commutes with  $e_N$ .

If the index of  $N$  in  $M$  is finite then from the pair  $N \subset M$  one can construct a new pair of finite factors  $M \subset M_1$  with the same index  $[M_1 : M] = [M : N]$ . The construction of  $M_1$  is called the basic construction and the factor  $M_1$  is called the extension of  $M$  by  $N$ .

We recall from [1] the definition and main properties of  $M_1$ :

1.1 PROPOSITION. *Define  $M_1 = JN'J$ . Then we have*

1°  $M_1 = (M \cup \{e_N\})''$ ,

2°  $[M_1 : M] = [M : N]$  and if  $\tau$  denotes the unique normalized trace on  $M_1$  and  $E_M$  the  $\tau$  preserving conditional expectation of  $M_1$  onto  $M$ , then  $E_M(e_N) = [M : N]^{-1} 1_M$  or equivalently  $\tau(e_N x) = [M : N]^{-1} \tau(x)$  for every  $x \in M$ .

Part 1° of this proposition can be made more precise: by [3], if  $n + 1 \geq [M : N]$  then any element in  $M_1$  is a sum of at most  $(n + 1)^2$  monomials of the form  $x e_N y$ ,  $x, y \in M$ . Note that  $M_1$  can also be described abstractly as the unique (up to isomorphism) finite factor  $M_1$  which contains  $M$  and a projection  $e$  so that  $[M_1 : M] = [M : N]$ ,  $[e, y] = 0$  for  $y \in N$ ,  $e x e = E_N(x) e$  for  $x \in M$ , and with the trace  $\tau$  satisfying  $\tau(e x) = [M_1 : M]^{-1} \tau(x)$ ,  $x \in M$ . In fact one of the conditions is redundant: the next proposition gives two equivalent ways of characterizing  $M_1$ .

1.2 PROPOSITION. *Let  $N \subset M$  be a pair of finite factors with finite index and  $M_1$  the extension of  $M$  by  $N$ . Let  $\tilde{M}$  be a finite factor that contains  $M$  and with normalized trace  $\tilde{\tau}$ ,  $E_M$  the  $\tilde{\tau}$ -preserving conditional expectation of  $\tilde{M}$  onto  $M$  and  $e \in \tilde{M}$  an orthogonal projection. Then the following conditions are equivalent:*

1° *There exists an isomorphism  $\phi$  of  $M_1$  onto  $\tilde{M}$  such that  $\phi(x) = x$  for  $x \in M$  and  $\phi(e_N) = e$ .*

2° (i)  $[e, y] = 0$ ,  $y \in N$ ;

(ii)  $E_M(e) = [\tilde{M} : M]^{-1} 1_M = [M : N]^{-1} 1_M$ .

3° (i)  $e x e = E_N(x) e$ ,  $x \in M$ , and  $e \neq 0$ ;

(ii)  $e$  and  $M$  generate  $\tilde{M}$  as a von Neumann algebra.

PROOF. 1° implies 2° by the known properties of  $e_N$ .

Suppose 2° holds. Then by 1.8 of [3] we get that  $\tilde{M}$  is the extension of  $M$  by  $P$  where  $P = \{e\}' \cap M$ . But (i) implies that  $N \subset P$  and since  $[M : P] = [\tilde{M} : M] = [M : N]$  we conclude that  $N = P$ . Thus  $e$  and  $M$  generate  $\tilde{M}$  as a von Neumann algebra and again by 1.8 of [3] we get  $E_N(x) e = e x e$ , for every  $x \in M$ .

Assume that  $3^\circ$  holds. Using the “orthonormal basis” of [3] it is easy to see that the map  $\phi: M_1 \rightarrow \tilde{M}$  that sends  $\sum x_i e_N y_i$  to  $\sum x_i e y_i$  is a well-defined  $*$ -homomorphism. Moreover  $\phi$  satisfies  $m\phi(x) = \phi(mx)$  for every  $m \in M$  and  $x \in M_1$ . This shows that  $\phi(1)$  is a projection that commutes with  $e$  and with every  $m \in M$ . By (ii) we conclude that  $\phi(1)$  is central and since  $e \neq 0$  and  $M$  is a factor  $\phi(1) = 1$ . This implies now that  $\phi(m) = \phi(m1) = m\phi(1) = m$  and since obviously  $\phi(e_N) = e$  we get  $1^\circ$ . Q.E.D.

The pair  $M \subset M_1$  having finite index one can construct its extension  $M_1 \subset M_2$  and in fact the whole procedure may be iterated to get an increasing sequence of finite factors  $N \subset M \subset M_1 \subset M_2 \subset \dots$ , and orthogonal projections  $e_i \in M_{i+1}$ ,  $i \geq 0$  ( $N = M_{-1}$ ,  $M = M_0$ ) in which  $M_{i+1}$  is the extension of  $M_i$  by  $M_{i-1}$  or in other words  $M_{i+1}$  and  $e_i$  are obtained by the basic construction from the pair  $M_{i-1} \subset M_i$ . Thus if  $\tau$  denotes the unique normalized trace on  $\bigcup_i M_i$  and  $E_{M_{i-1}}$  the  $\tau$ -preserving conditional expectation of  $M_i$  onto  $M_{i-1}$ ,  $i \geq 0$ , then:

- (a)  $[e_i, y] = 0$  for  $y \in M_{i-1}$ ;
- (b)  $e_i x e_i = E_{M_{i-1}}(x) e_i$ ,  $x \in M_i$ ;
- (c)  $[M_{i+1} : M_i] = [M : N]$  and  $E_{M_i}(e_i) = [M : N]^{-1} 1$ .

In particular it follows that the sequence of projections  $e_i$  satisfies  $[e_i, e_j] = 0$ ,  $|i - j| \geq 2$ ,  $e_i e_{i\pm 1} e_i = [M : N]^{-1} e_i$  and  $\tau(e_i w) = [M : N]^{-1} \tau(w)$  for every word in  $1, e_0, e_1, \dots, e_{i-1}$ .

**2.  $n$ -step extensions.** In this section we prove the main result of the paper: we show that if  $N \subset M \subset M_1 \subset \dots$  is the sequence of finite factors obtained by iterating the basic construction as in §1, then, for each  $n > 0$ ,  $M_{2n+1}$  is the extension of  $M_n$  by  $N$ . In fact we give an explicit formula for a projection  $f_n \in M_{2n+1}$  which implements the conditional expectation of  $M_n$  onto  $N$  and generates with  $M_n$  the factor  $M_{2n+1}$ :  $f_n$  will be a scalar multiple of the word of maximal length in  $e_0, e_1, \dots, e_{2n}$  where  $e_i \in M_{i+1}$  are as in §1.

We define for each  $n, k \geq 0$  the element

$$g_n^k = (e_{n+k} e_{n+k-1} \dots e_k) (e_{n+k+1} e_{n+k} \dots e_{k+1}) \dots (e_{2n+k} e_{2n+k-1} \dots e_{n+k})$$

(there are  $n+1$  products of parentheses and in each parentheses the product of  $n+1$  consecutive projections  $e_i$  in decreasing order). We put  $f_n^k = [M : N]^{n(n+1)/2} g_n^k \in M_{2n+k+1}$  and  $f_n = f_n^0 \in M_{2n+1}$ .

To prove that the above defined  $f_n$  implements the basic construction in the extension of  $M_n$  by  $N$ , we only have to show that  $f_n$  is an orthogonal projection, that  $f_n \in N' \cap M_{2n+1}$  and that  $E_{M_n}(f_n) = [M_n : N]^{-1} = [M_{2n+1} : M_n]^{-1}$ . (See Proposition 1.2.) Note that since  $[M_{i+1} : M_i] = [M : N]$ , by the multiplicative property of the index we do have  $[M_n : N] = [M : N]^{n+1} = [M_{2n+1} : M_n]$ . To prove the other properties, let us first recall some facts about the algebra generated by  $\{e_i\}_{i \geq 0}$  (cf. [1]).

A finite product of  $e_i$ 's is called a word. It is called a reduced word if it is of minimal length for the grammatical rules  $e_i e_{i\pm 1} e_i \leftrightarrow e_i$ ,  $e_i^2 \leftrightarrow e_i$  and  $e_i e_j \leftrightarrow e_j e_i$  for  $|i - j| \geq 2$ . Note that any word is a scalar multiple of a reduced word. Jones pointed out (in [1, 4.1.4]) that reduced words can be uniquely written in the ordered form

$$(*) \quad w = (e_{j_1} e_{j_1-1} \dots e_{k_1}) (e_{j_2} e_{j_2-1} \dots e_{k_2}) \dots (e_{j_p} e_{j_p-1} \dots e_{k_p})$$

where  $j_i \geq k_i$ ,  $j_{i+1} > j_i$ ,  $k_{i+1} > k_i$ .

From this description of reduced words it follows that if a reduced word  $w$  is written with the letters  $e_r, e_{r+1}, \dots, e_s$  ( $s \geq r$ ) then  $e_{r+i}$  and  $e_{s-i}$  appear at most  $i + 1$  times in  $w$ .

To prove the theorem we first show that  $g_n^0$  are selfadjoint elements. This will be an easy consequence of the next two lemmas.

**2.1 LEMMA.**  $g_n^0$  is the unique reduced word of maximal length in  $e_0, e_1, \dots, e_{2n}$ .

**PROOF.** Since by definition  $g_n^0$  is of the form  $(*)$  it is a reduced word. As noted before if  $w$  is an arbitrary reduced word in  $e_0, e_1, \dots, e_{2n}$  then  $e_0, e_{2n}$  appear at most once in  $w$ ,  $e_1, e_{2n-1}$  at most twice and more generally  $e_k, e_{2n-k}$  at most  $k + 1$  times. Thus the length of  $w$  is at most equal to  $1 + 2 + \dots + n + (n + 1) + n + \dots + 2 + 1$  and by inspecting the conditions  $j_i \geq k_i$ ,  $j_{i+1} > j_i$ ,  $k_{i+1} > k_i$  of  $(*)$  it follows that the only reduced word  $w$  with this length is obtained when  $j_i = n + i$ ,  $k_i = i$ , i.e.  $w = g_n^0$ . Q.E.D.

**2.2 LEMMA.** If  $w$  is a reduced word in  $e_0, e_1, \dots, e_{2n}$  then the reduced form of  $w^*$  has the same length as  $w$ .

**PROOF.** Indeed,  $w^*$  has length at most equal to that of  $w$  and since  $(w^*)^* = w$ , the statement follows. Q.E.D.

To prove that  $g_n^0$  are scalar multiples of projections we have to compute  $(g_n^0)^2$ . To do this we use an induction argument based on the formula

**2.3 LEMMA.**  $g_n^0 = (e_n e_{n+1} \dots e_{2n}) g_{n-1}^0 (e_{2n-1} \dots e_n)$ .

**PROOF.** The equality follows by pushing  $e_{2n}$  to the left as much as possible in the formula giving  $g_n^0$ . Q.E.D.

**2.4 REMARK.** Two other equalities that can be obtained in a similar fashion and seem to be of interest are

$$g_n^0 = g_{n-1}^1 (e_{2n} \dots e_{n+1}) (e_0 \dots e_n) = (e_n e_{n-1} \dots e_0) g_{n-1}^2 (e_1 e_2 \dots e_n).$$

To show that  $g_n^0$  projects on a scalar in  $M_n$  we prove

**2.5 LEMMA.**  $E_{M_{2n}}(g_n^0) = [M : N]^{-(n+1)} g_{n-1}^1$ . More generally

$$E_{M_{2n+k}}(g_n^k) = [M : N]^{-(n+1)} g_{n-1}^{k+1}.$$

**PROOF.** It is enough to prove that  $E_{M_{2n}}(g_n^0) = \lambda^{n+1} g_{n-1}^1$ , where  $\lambda = [M : N]^{-1}$ , because the rest of the statement follows by starting the sequence of factors from  $M_{k-1} \subset M_k$ , instead of  $N = M_{-1} \subset M_0 = M$ .

We first show that for  $j \geq p \geq k + 1$  we have

$$(**) \quad (e_j e_{j-1} \dots e_k) (e_p e_{p-1} \dots e_{k+1}) = \lambda (e_{p-2} \dots e_k) (e_j \dots e_{k+1}).$$

Indeed we have

$$\begin{aligned} (e_j e_{j-1} \dots e_p e_{p-1} \dots e_k) e_p &= \lambda (e_j e_{j-1} \dots e_p) (e_{p-2} e_{p-3} \dots e_k) \\ &= \lambda (e_{p-2} \dots e_k) (e_j e_{j-1} \dots e_p), \end{aligned}$$

which easily implies (\*\*). Applying recursively (\*\*) we get

$$\begin{aligned}
 E_{M_{2n}}(g_n^0) &= (e_n e_{n-1} \cdots e_0) \cdots (e_{2n-1} \cdots e_{n-1}) E_{M_{2n}}(e_{2n})(e_{2n-1} \cdots e_n) \\
 &= \lambda(e_n \cdots e_0) \cdots (e_{2n-1} \cdots e_{n-1})(e_{2n-1} \cdots e_n) \\
 &= \lambda^2(e_n \cdots e_0) \cdots (e_{2n-2} e_{2n-3} \cdots e_{n-2})(e_{2n-3} \cdots e_{n-1})(e_{2n-1} \cdots e_n) \\
 &= \lambda^3(e_n \cdots e_0) \cdots (e_{2n-5} \cdots e_{n-2})(e_{2n-2} \cdots e_{n-1})(e_{2n-1} \cdots e_n) \\
 &= \cdots = \lambda^n(e_n \cdots e_0) e_1(e_{n+1} \cdots e_2) \cdots (e_{2n-2} \cdots e_{n-1})(e_{2n-1} \cdots e_n) \\
 &= \lambda^{n+1}(e_n \cdots e_1)(e_{n+1} \cdots e_2) \cdots (e_{2n-1} \cdots e_n) = \lambda^{n+1} g_{n-1}^1. \quad \text{Q.E.D.}
 \end{aligned}$$

We can now prove the theorem.

**2.6 THEOREM.** *Let  $N \subset M$  be a pair of finite factors with  $[M : N] < \infty$ . Let  $N \subset M \subset M_1 \subset \cdots$  be the sequence of finite factors obtained by iterating the basic construction and  $e_i \in M_{i+1}$  the projection implementing the conditional expectation of  $M_i$  onto  $M_{i-1}$  at each step of the basic construction as in §1, for  $i \geq 0$  ( $M_{-1} = N$ ,  $M_0 = M$ ). Let*

$$f_n = [M : N]^{n(n+1)/2} (e_n e_{n-1} \cdots e_0)(e_{n+1} e_n \cdots e_1) \cdots (e_{2n} e_{2n-1} \cdots e_n) \in M_{2n+1}.$$

*Then  $M_{2n+1}$  is the extension of  $M_n$  by  $N$  and  $f_n \in M_{2n+1}$  is the projection that implements the conditional expectation of  $M_n$  onto  $N$ , i.e.  $f_n \in N' \cap M_{2n+1}$ ,  $f_n x f_n = E_N(x) f_n$ ,  $x \in M_n$ ,  $E_{M_n}(f_n) = [M_n : N]^{-1}$  and  $M_{2n+1} = (M_n \cup \{f_n\})''$ .*

**PROOF.** We will prove the theorem by induction over  $n \geq 0$ . If  $n = 0$  then  $f_0 = e_0$  and we have nothing to prove. Assume the statement is true up to  $n - 1$ . Let  $\lambda = [M : N]^{-1}$  and  $c_n = \lambda^{-n(n+1)/2}$ . Since  $f_n = c_n g_n^0$  and  $g_n^0$  is a word in  $e_0, e_1, \dots, e_{2n}$ , which all commute with  $N$ , it follows that  $f_n \in N' \cap M_{2n+1}$ . Note also that since  $e_{2n} \in M_{2n-1}' \cap M_{2n+1}$ ,  $e_{2n}$  commutes with  $g_{n-1}^0 \in M_{2n-1}$ . To see that  $g_n^0$  is selfadjoint we use Lemma 2.2 to obtain that  $g_n^{0*}$  has the same length as  $g_n^0$  and thus by Lemma 2.1  $g_n^0 = (g_n^0)^*$ . Further, Lemma 2.3 implies that

$$\begin{aligned}
 (g_n^0)^2 &= g_n^{0*} g_n^0 \\
 &= (e_n e_{n+1} \cdots e_{2n-1}) g_{n-1}^0 (e_{2n} e_{2n-1} \cdots e_{n+1} e_n e_{n+1} \\
 &\quad \cdots e_{2n-1} e_{2n}) g_{n-1}^0 (e_{2n-1} \cdots e_n) \\
 &= \lambda^n (e_n e_{n+1} \cdots e_{2n-1}) g_{n-1}^0 e_{2n} g_{n-1}^0 (e_{2n-1} \cdots e_n) \\
 &= \lambda^n (e_n e_{n+1} \cdots e_{2n}) (g_{n-1}^0)^2 (e_{2n-1} \cdots e_n) \\
 &= \lambda^n c_{n-1}^{-1} (e_n e_{n+1} \cdots e_{2n}) g_{n-1}^0 (e_{2n-1} \cdots e_n) \\
 &= \lambda^n c_{n-1}^{-1} g_n^0 = c_n^{-1} g_n^0.
 \end{aligned}$$

Thus  $f_n = c_n g_n^0$  is a selfadjoint projection in  $N' \cap M_{2n+1}$ . Next we apply recursively Lemma 2.5 to get

$$\begin{aligned}
 E_{M_n}(f_n) &= c_n E_{M_n}(g_n^0) = c_n E_{M_n} E_{M_{2n}}(g_n^0) = c_n \lambda^{n+1} E_{M_n}(g_{n-1}^1) \\
 &= c_n \lambda^{n+1} E_{M_n} E_{M_{2n-1}}(g_{n-1}^1) = c_n \lambda^{(n+1)+n} E_{M_n}(g_{n-2}^2) \\
 &= \cdots = c_n \lambda^{(n+1)+n+\cdots+2} E_{M_n}(g_0^n) = c_n \lambda^{(n+1)+n+\cdots+2} E_{M_n}(e_n) \\
 &= c_n \lambda^{(n+2)(n+1)/2} 1_{M_n} = \lambda^{n+1} 1_{M_n}
 \end{aligned}$$

(we used  $g_0^n = e_n$ ).

Moreover by [1],

$$\begin{aligned} [M_{2n+1} : M_n] &= \prod_{n \leq i \leq 2n} [M_{i+1} : M_i] = [M : N]^{n+1} \\ &= \prod_{0 \leq i \leq n} [M_{i+1} : M_i] = [M_n : N]. \end{aligned}$$

By Proposition 1.2 the rest of the properties of  $f_n$  follow automatically. Q.E.D.

**2.7 REMARK.** We could include the proof of  $g_n^0 = g_n^{0*}$  in the induction argument. Indeed by Lemma 2.3 and using  $g_{n-1}^0 = (g_{n-1}^0)^*$  and  $[e_{2n}, g_{n-1}^0] = 0$  we get

$$\begin{aligned} (g_n^0)^* &= e_n e_{n+1} \cdots e_{2n} (g_{n-1}^0)^* e_{2n-1} e_{2n-2} \cdots e_n \\ &= e_n e_{n+1} \cdots e_{2n} g_{n-1}^0 e_{2n-1} \cdots e_n = g_n^0. \end{aligned}$$

We preferred however the deductive argument of Lemmas 2.1 and 2.2 as it points out some properties of  $f_n$ .

**3. Some applications.** In this section we derive some consequences on the inclusion  $N \subset M_n$ . We consider the case when the relative entropy  $H(M|N)$  considered in [3] satisfies  $H(M|N) = \ln[M : N]$ . An important case when this equality occurs is when  $N' \cap M = \mathbb{C}$  (cf. [3]). First we compute the relative entropy from  $n$  to  $n$  steps.

**3.1 THEOREM.** *If  $H(M|N) = \ln[M : N]$  then*

$$H(M_{n+k}|M_{k-1}) = \ln[M_{n+k} : M_{k-1}], \quad \text{for every } n, k \geq 0.$$

*In particular  $H(M_n|N) = \ln[M_n : N]$  and  $H(M_k|M_{k-1}) = \ln[M_k : M_{k-1}]$ , for every  $k \geq 0$ .*

**PROOF.** Since  $H(M, N) = \ln[M : N]$ ,  $E_{N' \cap M}(e_0) = \lambda 1$  and the anti-isomorphism  $N' \cap M \ni x' \mapsto \theta_0(x') = J_M x' J_M \in M' \cap M_1$  is trace preserving (cf. 4.5 in [3]). To show that  $E_{M' \cap M_1}(e_1) = \lambda 1$  it suffices to prove that  $M' \cap M_1 \ni y' \mapsto \theta_1(y') = J_{M_1} y' J_{M_1} \in M' \cap M_1$  is also trace preserving (cf. [3]). But  $\theta_1 \theta_0 = \sigma'$ , where  $\sigma'$  is the restriction to  $N' \cap M$  of the isomorphism  $\sigma$  defined in [3, 1.3],  $\sigma'(x') = \lambda^{-1} \sum_i m_i e_0 e_1 x' e_0 m_i^*$ , with  $\{m_i\}$  an orthonormal basis of  $M$  over  $N$ . Indeed if  $x' = \sum_i m_i n_i \in N' \cap M$ , with  $n_i \in N$ , then  $\theta_0(x') \in M_1$  implies  $\theta_0(x') = \sum_{i,j} m_i E_N(m_j^* x'^* e_N m_j^*) e_N m_j^*$  and thus in  $L^2(M_1, \tau)$  we have

$$\begin{aligned} \theta_1(\theta_0(x'))(m_p n e_0 m_\tau^*) &= \sum_{i,j} m_p n e_0 m_\tau^* m_j E_N(x' m_j^* m_i) e_0 m_i^* \\ &= \sum_i m_p n e_0 E_N(x' m_\tau^* m_i) m_i^* = m_p n e_0 x' m_\tau^* \\ &= m_p e_0 x' n m_\tau^* = \sigma'(x')(m_p n e_0 m_\tau^*), \end{aligned}$$

for all  $n \in N$ . Thus, since  $\sigma', \theta_0$  are trace preserving,  $\theta_1$  is also trace preserving. Induction now shows that  $E_{M'_k \cap M_{k+1}}(e_k) = \lambda 1$ ,  $k \geq -1$ , and thus  $H(M_{k+1}, M_k) = \ln[M_{k+1} : M_k]$ .

To prove that  $H(M_{n+k}|M_{k-1}) = \ln[M_{n+k} : M_{k-1}]$  it now suffices to prove that  $H(M_n|N) = \ln[M_n : M]$  or, by [3],  $E_{M'_n \cap M_{2n+1}}(f_n) = \lambda^{n+1} 1_{M_{2n+1}}$ . Since

$M'_n \cap M_{2n+1} \subset M'_{n-1} \cap M_{2n+1} \subset \cdots \subset M' \cap M_{2n+1}$  we have  $E_{M'_n \cap M_{2n+1}} = E_{M'_n \cap M_{2n+1}} E_{M'_{n-1} \cap M_{2n+1}} \cdots E_{M' \cap M_{2n+1}}$ . Since  $e_0$  appears only once in  $g_n^0$  and  $E_{M' \cap M_{2n+1}}(e_0) = \lambda 1$  and  $e_i \in M'_{i-1}$ , it follows that

$$E_{M' \cap M_{2n+1}}(g_n^0) = (e_n \cdots e_1 E_{M' \cap M_{2n+1}}(e_0))(e_{n+1} \cdots e_1) \cdots (e_{2n} e_{2n-1} \cdots e_n).$$

Using now the same computations as in the proof of 2.6 it follows that

$$E_{M' \cap M_{2n+1}}(g_n^0) = \lambda^{n+1} g_{n-1}^1.$$

By induction it follows that

$$\begin{aligned} E_{M'_n \cap M_{2n+1}}(g_n^0) &= \lambda^{n+1} E_{M'_n \cap M_{2n+1}}(g_{n-1}^1) = \lambda^{n+1} E_{M'_n \cap M_{2n+1}} E_{M'_1 \cap M_{2n+1}}(g_{n-1}^1) \\ &= \lambda^{n+1} \lambda^n E_{M'_n \cap M_{2n+1}}(g_{n-2}^2) = \cdots = \lambda^{(n+1)+n+\cdots+1} I \end{aligned}$$

and thus  $E_{M'_n \cap M_{2n+1}}(f_n) = \lambda^{n+1} I$ . Q.E.D.

**3.2 COROLLARY.** *Let  $N \subset M$  be as in Theorem 3.1. Let  $J_n$  be the canonical conjugation on  $L^2(M_n, \tau)$ . Suppose  $M_{2n+1}$  is represented on  $L^2(M_n, \tau)$  so that to coincide with the basic construction of  $N \subset M_n$ . Then we have*

- (i) *For every projection  $f \in N' \cap M_n$ ,  $[(M_n)_f : N_f] = [M_n : N] \tau(f)^2$ .*
- (ii) *The anti-isomorphism  $N' \cap M_n \ni x \mapsto J_n x J_n \in M'_n \cap M_{2n+1}$  is trace preserving.*
- (iii) *For every  $k \geq 0$  there exists a trace preserving isomorphism  $N' \cap M \ni x \mapsto x' \in M'_{k-1} \cap M_k$  so that for every minimal projection  $f \in N' \cap M$ ,  $[M_f : N_f] = [(M_k)_f : (M_{k-1})_{f'}]$ .*

**PROOF.** By 4.5 in [3] the condition  $H(M_n|N) = \ln[M_n : N]$  is equivalent to the above conditions (i) and (ii). Then (iii) follows by (i), (ii) and by the fact that given any trace preserving anti-isomorphism between two finite-dimensional algebras there exists a trace preserving isomorphism between them which acts on the centers in the same way the anti-isomorphism does. Q.E.D.

## REFERENCES

1. V. Jones, *Index for subfactors*, Invent. Math. **75** (1983), 1-25.
2. A. Ocneanu, in preparation.
3. M. Pimsner and S. Popa, *Entropy and index for subfactors*, Ann. Sci. Ecole Norm. Sup. **19** (1986), 57-106.

THE NATIONAL INSTITUTE FOR SCIENTIFIC AND TECHNICAL CREATION, DEPARTMENT OF MATHEMATICS, BD. PACII 220, 79622 BUCHAREST, ROMANIA

*Current address* (M. Pimsner): Culmea Veche No 2, Sector 3, Bucharest, Romania

*Current address* (S. Popa): Department of Mathematics, University of California, Los Angeles, California 90024