

## ON A CLASS OF FUNCTIONALS INVARIANT UNDER A $\mathbf{Z}^n$ ACTION

PAUL H. RABINOWITZ

**ABSTRACT.** Consider a system of ordinary differential equations of the form

$$(*) \quad \ddot{q} + V_q(t, q) = f(t)$$

where  $f$  and  $V$  are periodic in  $t$ ,  $V$  is periodic in the components of  $q = (q_1, \dots, q_n)$ , and the mean value of  $f$  vanishes. By showing that a corresponding functional is invariant under a natural  $\mathbf{Z}^n$  action, a simple variational argument yields at least  $n + 1$  distinct periodic solutions of  $(*)$ . More general versions of  $(*)$  are also treated as a class of Neumann problems for semilinear elliptic partial differential equations.

**Introduction.** Consider the system of ordinary differential equations

$$(0.1) \quad \ddot{q} + V_q(t, q) = f(t),$$

where  $q = (q_1, \dots, q_n) \in \mathbf{R}^n$  and  $V$  satisfies

$$(V_1) \quad V \in C^1(\mathbf{R} \times \mathbf{R}^n, \mathbf{R}) \text{ and is } T \text{ periodic in } t \text{ and } T_i \text{ periodic in } q_i, \\ 1 \leq i \leq n.$$

Suppose further that  $f$  satisfies

$$(f_1) \quad f \in C(\mathbf{R}, \mathbf{R}^n) \text{ and is } T \text{ periodic in } t$$

and

$$(f_2) \quad [f] \equiv \frac{1}{T} \int_0^T f(t) dt = 0.$$

Note that if  $q(t)$  is a solution of (0.1), so is  $q(t) + (k_1 T_1, \dots, k_n T_n)$  for all  $k = (k_1, \dots, k_n) \in \mathbf{Z}^n$ . It was shown by Mawhin and Willem [1], Serrin and Pucci [2–3], Li [4], and Franks [5] that if  $n = 1$  and  $V$  is independent of  $t$ , (0.1) possesses at least two  $T$  periodic solutions which do not differ by a multiple of  $T_1$ . In [5], the proof relies on a generalized version of the Poincaré-Birkhoff Theorem while [1–4] use variational arguments. Part of the difficulty in treating (0.1) in [1–4] is caused by the fact that the corresponding functional

$$(0.2) \quad I(q) = \int_0^T \left[ \frac{1}{2} |\dot{q}|^2 - V(t, q) + f \cdot q \right] dt$$

---

Received by the editors August 21, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 34C25, 35J60, 58E05, 58F05, 58F22.

*Key words and phrases.*  $\mathbf{Z}^n$  action periodic solution, critical point, minimax argument, Ljusternik-Schnirelmann category, Neumann problem.

This research was supported in part by the United States Army under Contract No. DAAL03-87-K-0043, the Air Force Office of Scientific Research under Grant No. AFOSR-87-0202 and the National Science Foundation under Grant No. MCS-8110556.

defined on the natural Sobolev space associated with (0.2) does not satisfy the Palais-Smale condition, a compactness criterion very useful for variational problems and henceforth denoted by (PS).

The purpose of this note is to show that in fact if appropriately interpreted, the variational problem does satisfy the (PS) condition. The simple observation that makes this statement precise together with standard techniques leads to a generalization of the above results.

**THEOREM 0.3.** *Under the above hypotheses on  $V$  and  $f$ , (0.1) possesses at least  $n + 1$  "distinct" solutions.*

What is meant by distinct will be explained in §1. Theorem 0.3 will be obtained from a more general result involving a Lagrangian of the form

$$\mathcal{L}(q, \dot{q}) = \frac{1}{2}L(t, q)\dot{q} \cdot \dot{q} - V(t, q) + f(t) \cdot q,$$

where  $L(t, q)$  is a symmetric positive definite matrix possessing the same periodicity properties as does  $V$ . Actually as will be shown in §1, when  $L$  is independent of  $t$  and  $q$  and  $f \equiv 0$ , this more general result can be obtained from a theorem of Conley and Zehnder [6, Theorem 3]. They were mainly interested in the much more difficult case of indefinite  $L$ .

In §1, the generalizations of Theorem 0.3 will be carried out. Some of the technical details will be given in §2. The ideas used in §§1 and 2 can also be applied to a class of Neumann problems for semilinear elliptic partial differential equations. Consider

$$(0.4) \quad \begin{aligned} -\Delta u &= p(x, u) + h(x), & x \in \Omega, \\ \partial u / \partial \nu &= 0, & x \in \partial \Omega. \end{aligned}$$

Here  $\Omega$  denotes a bounded domain in  $\mathbf{R}^n$  with a smooth boundary and outward pointing normal  $\nu(x)$  and  $\partial u / \partial \nu = \nu(x) \cdot \nabla u$ . Suppose  $p(x, \xi) = \partial P(x, \xi) / \partial \xi$  and  $|\Omega|$  denotes the volume of  $\Omega$ . Then we have

**THEOREM 0.5.** *Suppose  $P$  satisfies*

$$(p_1) \quad P \in C^2(\overline{\Omega} \times \mathbf{R}, \mathbf{R}) \text{ and } P(x, \xi + r) = P(x, \xi) \text{ for all } (x, \xi) \in \overline{\Omega} \times \mathbf{R}$$

*and  $h$  satisfies*

$$(h_1) \quad h \in C^1(\Omega, \mathbf{R}),$$

*and*

$$(h_2) \quad [h] \equiv \frac{1}{|\Omega|} \int_{\Omega} h(x) dx = 0.$$

*Then (0.4) possesses at least two classical solutions which do not differ by an integer multiple of  $r$ .*

The proof of Theorem 0.5 will be carried out in §3. We thank Ed Fadell and Sufian Hussein for helpful conversations.

**1. A generalized version of Theorem 0.3.** Let  $E = W_T^{1,2}(\mathbf{R}, \mathbf{R}^n)$  denote the Sobolev space of  $T$  periodic functions with values in  $\mathbf{R}^n$  under the norm

$$\|q\| = \left( \int_0^T |\dot{q}|^2 dt + [q]^2 \right)^{1/2}.$$

Suppose  $L$  satisfies

- (L<sub>1</sub>)  $L \in C^1(\mathbf{R} \times \mathbf{R}^n, \mathbf{R}^{n^2})$  is a symmetric matrix with  $L(t, q)$  being  $T$  periodic in  $t$  and  $T_i$  periodic in  $q_i$ ,  $1 \leq i \leq n$ ,

and

- (L<sub>2</sub>) there is an  $\alpha > 0$  such that  $L(t, q)\xi \cdot \xi \geq \alpha|\xi|^2$  for all  $t \in \mathbf{R}$  and  $q, \xi \in \mathbf{R}^n$ .

Suppose  $V$  and  $f$  satisfy (V<sub>1</sub>) and (f<sub>1</sub>)–(f<sub>2</sub>) respectively. Let  $q \in E$ . Then

$$(1.1) \quad I(q) \equiv \int_0^T \left[ \frac{1}{2} L(t, q) \dot{q} \cdot \dot{q} - V(t, q) + f \cdot q \right] dt$$

is well defined and the argument of [7, Proposition B10] shows that  $I \in C^1(E, \mathbf{R})$  and critical points of  $I$  are classical solutions of

$$(1.2) \quad \frac{d}{dt}(L(t, q)\dot{q}) - \frac{1}{2} \frac{\partial L}{\partial q}(t, q)\dot{q} \cdot \dot{q} + V_q(t, q) = f(t).$$

Suppose  $Q, q \in E$  and

$$(1.3) \quad Q - q = (k_1 T_1, \dots, k_n T_n),$$

where  $k = (k_1, \dots, k_n) \in \mathbf{Z}^n$ . The properties of  $L, V$ , and  $f$  then imply

$$(1.4) \quad I(Q) = I(q).$$

A functional  $J$  on  $E$  satisfies (PS) if any sequence  $q_m$  such that  $I(q_m)$  is bounded and  $I'(q_m) \rightarrow 0$  possesses a convergent subsequence. Equation (1.4) shows that  $I$  does not satisfy (PS) on  $E$ . However, (1.4) implies  $I$  possesses a free  $\mathbf{Z}^n$  action on  $E$  and this will provide the topological basis for an existence result for (1.2). Before stating it some further preliminaries are required.

We introduce an equivalence relation  $\sim$  on  $E$  via  $Q \sim q$  if (1.3) holds for some  $k \in \mathbf{Z}^n$ . Let  $\tilde{E} = E/\sim$  with the corresponding quotient topology. One could now consider  $I$  on  $\tilde{E}$ . However, we find it more convenient technically to work on  $E$  itself and proceed as follows. For  $k \in \mathbf{Z}^n$ , set  $\theta_k = (k_1 T_1, \dots, k_n T_n)$ . For  $q \in \mathbf{R}^n$ , let  $g_k(q) = q + \theta_k$ . Then  $\mathcal{G} \equiv \{g_k \mid k \in \mathbf{Z}^n\}$  is a group of mappings of  $E$  onto  $E$  and  $I$  is invariant under  $\mathcal{G}$ , i.e.  $I(g(q)) = I(q)$  for all  $q \in E$  and  $g \in \mathcal{G}$ . A set  $A \subset E$  is called *invariant* (with respect to  $\mathcal{G}$ ) if  $g(A) \subset A$  for all  $g \in \mathcal{G}$ .  $A$  is an invariant set if and only if there is a set  $\tilde{A} \subset \tilde{E}$  such that  $A/\sim = \tilde{A}$ .

With the aid of the above notions, a generalized version of Theorem 0.3 can be stated.

**THEOREM 1.5.** *Let  $V$  satisfy  $(V_1)$ ,  $f$  satisfy  $(f_1)$ – $(f_2)$  and  $L$  satisfy  $(L_1)$ – $(L_2)$ . Then equation (1.2) has at least  $n+1$  distinct solutions  $Q_1, \dots, Q_{n+1}$ , i.e. for  $i \neq j$ ,  $Q_i \neq g(Q_j)$  for all  $g \in G$ .*

These solutions will be determined as critical points of  $I$  by means of a minimax argument. To characterize the corresponding critical values of  $I$ , let  $A \subset E$  be closed and define

$$(1.6) \quad \Gamma_k = \{A \subset E | A \text{ is invariant and } \text{cat}_{\tilde{E}} \tilde{A} \geq k\}.$$

In (1.6),  $\text{cat}_X Y$  denotes the Ljusternik-Schnirelmann category of the closed set  $Y$  in the topological space  $X$ . (See e.g. [8].)

**LEMMA 1.7.**  $\Gamma_k \neq \emptyset$ ,  $1 \leq k \leq n+1$ .

**PROOF.** It suffices to show  $\Gamma_{n+1} \neq \emptyset$ . Let  $A = \{q \in E | q = [q]\}$ . Then  $A$  is isomorphic to  $\mathbf{R}^n$  and  $A/\sim$  to  $T^n$ . Therefore  $\tilde{E}/\sim = T^n \oplus F$  where  $F = A^\perp$ , the orthogonal complement of  $A$ . Since  $F$  is a linear space, it is contractible to a point in itself. Therefore by standard results on Ljusternik-Schnirelmann category—see pp. 459–460 of [9]—

$$(1.8) \quad \text{cat}_{\tilde{E}} \tilde{E} = \text{cat}_{\tilde{E}} T^n \oplus \{0\} = \text{cat}_{T^n} T^n = n+1.$$

Now  $n+1$  critical values  $c_k$  of  $I$  can be defined as

$$(1.9) \quad c_k = \inf_{B \in \Gamma_k} \max_{q \in B} I(q), \quad 1 \leq k \leq n+1.$$

To prove Theorem 1.5, it will be shown that each of the numbers  $c_k$  is a critical value of  $I$  and there are at least  $n+1$  corresponding critical points. To carry out the proof some further preliminaries are required. A mapping  $\Psi: E \rightarrow E$  is called *equivariant* with respect to  $\mathcal{G}$  if  $\Psi \circ g = g \circ \Psi$  for all  $g \in \mathcal{G}$ . For  $s \in \mathbf{R}$ , let  $A_s \equiv \{q \in E | I(q) \leq s\}$  and  $K_s = \{q \in E | I(q) = s \text{ and } I'(q) = 0\}$ . The following version of a standard “Deformation Theorem” is needed:

**PROPOSITION 1.10.** *For any  $c \in \mathbf{R}$  and invariant neighborhood  $\mathcal{O}$  of  $K_c$ , there is an  $\varepsilon > 0$  and  $\eta \in C([0, 1] \times E, E)$  such that*

- 1°  $\eta(0, q) = q$  for all  $q \in E$ ,
- 2°  $\eta(t, q)$  is equivariant for each  $t \in [0, 1]$ ,
- 3°  $\eta(q, A_{c+\varepsilon} \setminus \mathcal{O}) \subset A_{c-\varepsilon}$ ,
- 4° If  $K_c = \emptyset$ ,  $\eta(1, A_{c+\varepsilon}) \subset A_{c-\varepsilon}$ .

The proof of Proposition 1.10 will be given in §2.

*Completion of proof of Theorem 1.5.* By 1° and 2° of Proposition 1.10,  $\eta(1, \cdot)$  is homotopic to the identity map and is equivariant. Hence by Lemma 5.3 of [8],  $\eta(1, \cdot): \Gamma_k \rightarrow \Gamma_k$ ,  $1 \leq k \leq n+1$ . Now the conclusions of Theorem 1.5 follow in a standard way: suppose  $c_k = \dots = c_{k+p} \equiv c$ . Then

$$(1.11) \quad \text{cat}_{\tilde{E}} \tilde{K}_c \geq p+1$$

for if not, by the proof of Proposition 1.10 and Lemma 5.6 of [8], we can find a uniform neighborhood  $\mathcal{O}$  of  $K_c$  such that  $\text{cat}_{E^*} \mathcal{O}^* < p+1$ . Choose  $B \in \Gamma_{k+p}$  such that

$$(1.12) \quad \max_B I \leq c + \varepsilon.$$

By the subadditivity property of category [8, Lemma 5.3],  $\text{cat}_{\widetilde{B} \setminus \widetilde{\mathcal{O}}} \geq k + p - p = k$  so  $\widetilde{B} \setminus \widetilde{\mathcal{O}} \in \Gamma_k$  as is  $\eta(1, \widetilde{B} \setminus \widetilde{\mathcal{O}}) \equiv B_1$ . But by (1.12) and  $3^\circ$  (or if  $p = 0$ ,  $4^\circ$ ) of Proposition 1.10,

$$(1.13) \quad \max_{B_1} I \leq c - \varepsilon,$$

contrary to the definition of  $c = c_k$ . Finally, if  $p \geq 1$ , the definition of category implies  $\widetilde{K}_c$  contains infinitely many distinct points completing the proof.

REMARK 1.14. In [4], Li proved an abstract theorem about critical points of periodic functionals which the above arguments can be used to generalize and put in a more natural form.

REMARK 1.15. If e.g.  $L$  and  $V$  are independent of  $t$  and  $f = 0$ , the hypotheses and therefore conclusions of Theorem 1.5 hold for all  $T \in \mathbf{R}$ . However, by the periodicity properties of  $V$ , it belongs to  $C^1(T^n, \mathbf{R})$ . Therefore  $V$  possesses at least  $\text{cat}_{T^n} T^n = n + 1$  distinct critical points which will be equilibrium solutions of (1.2). Thus it is not clear that there need exist any time dependent solutions for this special case without more structure for  $V$ . In this regard, see Theorem 5.2 of [4] when  $L = \text{id}$  and  $n = 1$ .

REMARK 1.16. In [6], Conley and Zehnder proved that if (i)  $H(t, p, q) \in C^2(\mathbf{R} \times \mathbf{R}^n \times \mathbf{R}^n, \mathbf{R})$ , where  $H$  is 1-periodic in  $t$  and the components of  $q$ , and (ii) there is an  $R > 0$  such that for  $|p| \geq R$ ,

$$H(t, p, q) = \frac{1}{2}Mp \cdot p + a \cdot p$$

where  $a \in \mathbf{R}^n$  and  $M$  is a symmetric nonsingular time independent matrix, then the corresponding Hamiltonian system possesses at least  $n + 1$  distinct periodic solutions. This theorem applies to a subclass of the problems treated here including Theorem 0.3. Note however, that the Conley-Zehnder result permits *indefinite* matrices  $M$ . Assuming for the moment its applicability here to Hamiltonians of the form

$$(1.17) \quad H = \frac{1}{2}Mp \cdot p + V(t, q),$$

observe that the corresponding Hamiltonian system is

$$(1.18) \quad \dot{p} = -V_q, \quad \dot{q} = Mp.$$

Therefore

$$(1.19) \quad \dot{p} = \frac{d}{dt}M^{-1}\dot{q} = -V_q.$$

Choosing  $M = L^{-1}$ , (1.19) now gives (1.2).

To applicability of the Conley-Zehnder result is not quite immediate here since (1.17) does not satisfy condition (ii). However, using a trick, a modified Hamiltonian can be constructed which satisfies (ii) and whose solutions satisfy (1.2). For simplicity we will just verify this for (0.3) where  $M = \text{id}$ . For this case by (1.2) or (0.1), if  $q$  is a  $T$  periodic solution,

$$(1.20) \quad \|\ddot{q}\|_{L^\infty} \leq \max_{t \in \mathbf{R}, \xi \in \mathbf{R}^n} |V_q(t, \xi)| \equiv K_1 \leq K_1 + \max_{t \in \mathbf{R}, \xi \in \mathbf{R}^n} |V(t, \xi)| \equiv K.$$

Simple estimates then show

$$(1.21) \quad \|\dot{q}\|_{L^\infty} \leq T\|\ddot{q}\|_{L^\infty} < 2TK.$$

Let  $\varphi \in C^\infty(\mathbf{R}, \mathbf{R})$  such that  $\varphi(s) = 1$  if  $s \leq (1+T)K \equiv K_2$ ,  $\varphi(s) = 0$  if  $s \geq 2K_2$ , and  $-1 < \varphi'(s) < 0$  if  $s \in (K_2, 2K_2)$ . The Hamiltonian

$$(1.22) \quad \widehat{H}(t, p, q) = \frac{1}{2}|p|^2 + \varphi(|p|)V(t, q)$$

satisfies (i) and (ii). The corresponding Hamiltonian system is

$$(1.23) \quad \dot{p} = -V_q \varphi, \quad \dot{q} = p + V \varphi_p.$$

If  $(p, q)$  is a  $T$ -periodic solution of (1.23), the second equation implies

$$(1.24) \quad [p] = -[V \varphi_p].$$

By the definition of  $K_2$ ,  $|\varphi_p| \leq 1$  and therefore

$$(1.25) \quad |[p]| \leq K.$$

Since

$$(1.26) \quad p(t) = [p] + \frac{1}{T} \int_0^T \left( \int_\tau^T \dot{p}(s) ds \right) d\tau,$$

(1.26), (1.23), and (1.25) imply

$$(1.27) \quad \|p\|_{L^\infty} \leq K + TK \leq K_2.$$

Consequently  $\varphi(|p|) = 1$ ,  $\varphi_p(|p|) = 0$ , and any  $T$ -periodic solution of the modified equation satisfies (0.3).

**2. The proof of Proposition 1.10.** A mapping  $\eta \in C([0, 1] \times E, E)$  and  $\varepsilon > 0$  must be found satisfying 1°–4° of Proposition 1.10. Much of this construction is standard and therefore we will only indicate the modifications that must be made in the proof the “usual case” that can be found in [7, Appendix A].

The mapping  $\eta$  is determined as the solution of an ordinary differential equation of the form

$$(2.1) \quad d\eta/dt = \omega(\eta)\Psi(\eta)$$

where  $\omega$  is a cut-off function with  $0 \leq \omega \leq 1$  and  $\Psi$  is a pseudogradient vector field for  $I'$ . In [7], the choice of  $\varepsilon$  and construction of  $\omega$  depend on the fact that  $I$  satisfies (PS). As was already noted, this is not the case here. Therefore we must show that because of the invariance of  $I$  under  $\mathcal{S}$ , the proof works nevertheless. Heuristically, we will show that  $I|_{\widetilde{E}}$  satisfies (PS). First the form of  $I'$  must be studied. Let  $D$  denote the duality map from  $E$  to  $E^*$ .

PROPOSITION 2.2. *Under the hypotheses of Theorem 1.5,*

$$I'(q) = D(\mathcal{P}_1(q) + \mathcal{P}(q))$$

where  $\mathcal{P}$  is compact and

$$\frac{d}{dt}\mathcal{P}_1(q) = L(t, q)\dot{q} - [L(t, q)\dot{q}].$$

PROOF. By (1.1),

$$(2.3) \quad I'(q)\varphi = \int_0^T \left[ L\dot{q} \cdot \dot{\varphi} + \frac{1}{2} \sum_{i=1}^n L_{q_i} \dot{q} \cdot \dot{q} \varphi_i - V_q \cdot \varphi + f \cdot \varphi \right] dt.$$

Let

$$(2.4) \quad \begin{cases} I_1(q)\varphi \equiv \int_0^T L\dot{q} \cdot \dot{\varphi} dt, \\ I_2(q)\varphi = \frac{1}{2} \int_0^T \sum_{i=1}^n L_{q_i} \dot{q} \cdot \dot{\varphi} \varphi_i dt, \\ I_3(q)\varphi = \int_0^T (-V_q + f) \cdot \varphi dt. \end{cases}$$

By the argument of Proposition B.10 of [7],  $I_3(q) = D\mathcal{P}_3(q)$  where  $\mathcal{P}_3$  is compact. Next consider  $I_2(q)\varphi$ . It has the form

$$(2.5) \quad I_2(q)\varphi = \int_0^T b \cdot \varphi dt$$

where  $b = b(t, q, \dot{q}) \in L^1$ . Since  $I_2(q) \in E^*$ , there is a unique  $x = \xi + y \in \mathbf{R}^n \oplus F$  such that

$$(2.6) \quad I_2(q)\varphi = \int_0^T \dot{y} \cdot \dot{\varphi} dt + \xi \cdot [\varphi].$$

Comparing (2.5)–(2.6) shows  $y$  is a weak solution of

$$(2.7) \quad -\ddot{y} = b - [b].$$

Hence  $y \in W^{2,1}$  and

$$(2.8) \quad \|y\|_{W^{2,1}} \leq 2\|b\|_{L^1}.$$

Similarly  $\xi = T[b]$  and

$$(2.9) \quad |\xi| \leq T\|b\|_{L^1}.$$

Let  $\mathcal{P}_2(q) = y(q) + \xi(q)$ . The form of  $b$  shows that for  $q$  belonging to a bounded set in  $E$ ,  $\mathcal{P}_2(q)$  will be bounded in  $W^{2,1}$ . Hence  $\mathcal{P}_2(q)$  and  $I_2(q) = D\mathcal{P}_2$  are compact on  $E$ .

Finally, consider  $I_1(q)\varphi$ . As was the case with  $I_2(q)$ ,

$$(2.10) \quad I_1(q)\varphi = \int_0^T \dot{w} \cdot \dot{\varphi} dt$$

where  $w \in F$  and there is no mean value term due to the form of  $I_1(q)$ . Comparing with (2.4) yields

$$(2.11) \quad \dot{w} = L\dot{q} - [L\dot{q}].$$

Now (2.11) together with  $[w] = 0$  determines  $w = \mathcal{P}_1(q)$ . Setting  $\mathcal{P} = \mathcal{P}_1 + \mathcal{P}_2$ , the result follows.

**LEMMA 2.12.** *Let  $\hat{K}_c = \{q \in K_c | 0 \leq [q_i] \leq T_i, 1 \leq i \leq n\}$ . Then  $\hat{K}_c$  is compact.*

**PROOF.** Suppose  $(Q_m) \subset \hat{K}_c$ . Let  $Q_m = \xi_m + Y_m$ , where  $\xi_m \in \mathbf{R}^n$  and  $Y_m \in F$ . The form of  $I$  and  $(L_2)$  imply that  $(Y_m)$  is bounded and the definition of  $\hat{K}_c$  shows that  $(\xi_m)$  is bounded. Since  $(Q_m)$  is bounded, along a subsequence,  $Q_m$  converges weakly in  $E$  and strongly in  $L^\infty$  to  $Q$  and  $\mathcal{P}(Q_m)$  converges in  $E$ . Therefore from Proposition 2.2,

$$(2.13) \quad \dot{Q}_m = L^{-1}(t, Q_m) \left[ T^{-1} \int_0^T L(\tau, Q_m) \dot{Q}_m d\tau - \frac{d}{dt} \mathcal{P}(Q_m) \right].$$

Now equation (2.13) shows that along this subsequence  $\dot{Q}_m$  converges in  $L^2$  and  $Q_m$  converges in  $E$  and  $I'(Q) = 0$ . Hence  $Q \in \hat{K}_c$ .

*Completion of the proof of Proposition 1.10.* In the proof of the version of Proposition 1.10 given in [7], (PS) is used in two places. The first is in showing that  $K_c$  is compact and therefore for any neighborhood  $\mathcal{O}$  of  $K_c$ , there exists a uniform neighborhood  $N_\delta \equiv N_\delta(K_c) = \{q \in E \mid \|q - K_c\| \leq \delta\} \subset \mathcal{O}$ . For the current setting, if  $\mathcal{O}$  is an invariant neighborhood of  $K_c$ , since  $\hat{K}_c \subset K_c$  and is compact, there is a  $\delta > 0$  such that  $\mathcal{O} \supset N_\delta(\hat{K}_c)$ . Therefore by the invariance of  $\mathcal{O}$ ,

$$(2.14) \quad \mathcal{O} \supset \bigcup_{g \in \mathcal{G}} g(N_\delta(\hat{K}_c)) = N_\delta(K_c).$$

The second place at which (PS) is required in [7] is in showing that there exist constants  $b, \hat{\varepsilon} > 0$  such that if  $q \in \mathcal{A} \equiv A_{c+\hat{\varepsilon}} \setminus (A_{c-\hat{\varepsilon}} \cup N_{\delta/8})$ ,

$$(2.15) \quad \|I'(q)\| \geq b.$$

Arguing indirectly, (2.15) follows for the subclass of  $q \in \mathcal{A}$  such that  $0 \leq [q_i] \leq T_i$ ,  $1 \leq i \leq n$ , by an argument paralleling the proof of Lemma 2.12, which will be omitted. Since  $A_s$  and  $N_{\delta/8}$  are invariant sets, (2.15) then holds for all  $q \in \mathcal{A}$ .

Given (2.14)–(2.15), the argument of Theorem A.4 of [7] yields 1°, 3°–4° of Proposition 1.10. It remains only to show that by an appropriate construction of  $\eta(t, \cdot)$ , 2° also holds. Since  $I$  is invariant under  $\mathcal{G}$ , i.e.  $I(g_k) = I(q + \theta_k) = I(q)$  for all  $g_k \in \mathcal{G}$ ,

$$(2.16) \quad I'(q + \theta_k) = I'(q) = I'(g_k q)$$

for all  $g_k \in \mathcal{G}$ . The invariance of  $I'$  under  $\mathcal{G}$  allows us to slightly modify the construction of Lemma A.2 of [7] to obtain a pseudogradient vector field  $\Psi$  for  $I'$  which is also invariant under  $\mathcal{G}$ . Also the definition of  $\omega$  in [7] shows that if  $I$  is invariant under  $\mathcal{G}$ , so is  $\omega$ . Thus the right-hand side of (2.1) is invariant under  $\mathcal{G}$ . We claim  $\eta(t, \cdot)$  satisfies 2°, i.e.  $\eta(t, gq) = g\eta(t, q)$  for all  $t \in [0, 1]$ ,  $g \in \mathcal{G}$ ,  $q \in E$ . Indeed set  $w = g\eta(t, q)$ . Then  $w = \eta(t, q) + \theta_k$  for some  $k \in \mathbf{Z}^n$  and

$$(2.17) \quad \begin{cases} dw/dt = d\eta/dt = \omega(\eta)\Psi(\eta) = \omega(g\eta)\Psi(g\eta) = \omega(w)\Psi(w), \\ w(0, g) = gq. \end{cases}$$

Therefore  $w(t, q) = \eta(t, gq) = g\eta(t, q)$  so 2° holds. The proof is complete.

**3. The Neumann problem.** In this section, the Neumann problem

$$(3.1) \quad \begin{cases} -\Delta u = p(x, u) + h(x), & x \in \Omega, \\ u = 0, & x \in \partial\Omega, \end{cases}$$

will be studied and Theorem 0.5 will be proved. The ideas used here are so close to those of §§1–2 that we will be sketchy.

Let  $E = W^{1,2}(\Omega)$ . For  $u \in E$ , define

$$(3.2) \quad I(u) = \int_{\Omega} \left[ \frac{1}{2} |\nabla u|^2 - P(x, u) - h(x)u \right] dx,$$

where  $P, h, \Omega$  satisfy the hypotheses of Theorem 0.5. The proof of Proposition B.10 of [7] shows  $I \in C^1(E, \mathbf{R})$ . By e.g. [10, Chapter 2] critical points of  $I$  are classical solutions of (3.1). Note that  $I(u + kr) = I(u)$  for all  $u \in E$  and  $k \in \mathbf{Z}$  via  $(p_1)$  and



( $h_2$ ). Therefore  $I$  does not satisfy (PS) on  $E$ . For  $u, v \in E$ , we say  $u$  is equivalent to  $v$ ,  $u \sim v$ , if  $u - v = kr$  for some  $k \in \mathbf{Z}$ . Let  $\tilde{E} = E/\sim$ . As in §1,  $A \subset E$  is an invariant set if and only if there is an  $\tilde{A} \subset \tilde{E}$  such that  $\tilde{A} = A/\sim$ . Again  $E$  can be identified with  $\mathbf{R} \oplus F$  where  $F$  is the orthogonal complement of  $\text{span}\{1\}$  in  $E$  and  $\tilde{E}$  can be identified with  $S^1 \oplus F$ .

The argument of Lemma 1.7 shows that

$$\Gamma_k = \{A \subset E \mid \text{cat}_{\tilde{E}} \tilde{A} \geq k\} \neq \emptyset$$

for  $k = 1, 2$ . Noting that  $(\int_{\Omega} |\nabla u|^2 dx)^{1/2}$  is a norm on  $F$  and that

$$(3.3) \quad I'(u) = D(u - [u]) + P(u),$$

where  $D$  again denotes the duality map and  $P$  is compact (see Proposition B.10 of [7]). Therefore there is an analogue of Proposition 1.10 here which implies that

$$(3.4) \quad c_k = \inf_{A \in \Gamma_k} \max_{u \in A} I(u), \quad k = 1, 2,$$

is a critical value of  $I$ ,  $k = 1, 2$  with a multiplicity result if  $c_1 = c_2$ . This completes the proof of Theorem 0.5.

REMARK 3.5. If in (0.6),  $u$  is an  $n$  vector,  $p(x, u) = \nabla_u P(x, u)$ , and  $P$  is periodic in the components of  $u$ , a version of Theorem 0.5 analogous to Theorem 0.3 obtains.

REMARK 3.6. An alternate way to prove Theorem 0.5 is to obtain a first solution as a minimum and a second using a variant of the mountain pass theorem as in [1–4] for (0.1) with  $n = 1$ . Such an approach, however, does not extend to cover the vector case mentioned in Remark 3.5.

REMARK 3.7. Theorem 0.5 can also be extended by replacing  $-\Delta$  by a more general second order divergence structure uniformly elliptic operator with appropriate changes in the boundary conditions. A similar result also obtains if  $\Omega$  is replaced by a rectangular domain and the Neumann boundary conditions by periodic ones.

## REFERENCES

1. J. Mawhin and M. Willem, *Multiple solutions of the periodic boundary value problem for some forced pendulum-type equations*, J. Differential Equations **52** (1984), 264–287.
2. P. Pucci and J. Serrin, *A mountain pass theorem*, J. Differential Equations **60** (1985), 142–149.
3. ———, *Extensions of the mountain pass theorem*, Univ. of Minnesota Math. Rep. 83–150.
4. Shujie Li, *Multiple critical points of periodic functional and some applications*, International Center for Theoretical Physics Tech. Rep. IC-86-191.
5. J. Franks, *Generalizations of the Poincaré-Birkhoff theorem*, preprint.
6. C. C. Conley and E. Zehnder, *The Birkhoff-Lewis fixed point theorem and a conjecture of V. I. Arnold*, Invent. Math. **73** (1983), 33–49.
7. P. H. Rabinowitz, *Minimax methods in critical point theory with applications to differential equations*, CBMS Regional Conf. Ser. in Math., no. 65, Amer. Math. Soc., Providence, R. I., 1986.
8. J. T. Schwartz, *Nonlinear functional analysis*, Gordon & Breach, New York, 1969.
9. G. W. Whitehead, *Elements of homotopy theory*, Springer-Verlag, 1978.
10. P. Grisvard, *Elliptic problems in nonsmooth domains*, Pitman, 1985.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, MADISON, WISCONSIN 53706