

BIFURCATION PHENOMENA ASSOCIATED TO THE p -LAPLACE OPERATOR

MOHAMMED GUEDDA AND LAURENT VERON

ABSTRACT. We determine the structure of the set of the solutions u of $-(|u_x|^{p-2}u_x)_x + f(u) = \lambda|u|^{p-2}u$ on $(0, 1)$ such that $u(0) = u(1) = 0$, where $p > 1$ and $\lambda \in \mathbf{R}$. We prove that the solutions with k zeros are unique when $1 < p \leq 2$ but may not be so when $p > 2$.

0. Introduction. In this article we study the structure of the set E_λ of the solutions of the following nonlinear eigenvalue problem

$$(0.1) \quad \begin{cases} -(|u_x|^{p-2}u_x)_x + f(u) = \lambda|u|^{p-2}u & \text{in } (0, 1), \\ u(0) = u(1) = 0, \end{cases}$$

where $p > 1$, λ is a real number and f is a C^1 real-valued odd function such that

$$(0.2) \quad r \mapsto g(r) = f(r)/(|r|^{p-2}r)$$

is increasing on $(0, +\infty)$ with limits 0 at 0 and $+\infty$ at infinity. We first investigate the unperturbed eigenvalue problem

$$(0.3) \quad \begin{cases} -(|v_x|^{p-2}v_x)_x = \lambda|v|^{p-2}v & \text{in } (0, 1), \\ v(0) = v(1) = 0. \end{cases}$$

By means of an elementary integration process we prove that (0.3) admits a non-trivial solution if and only if

$$(0.4) \quad \lambda = \lambda_k = k^p(p-1) \left[2 \int_0^1 \frac{dt}{(1-t^p)^{1/p}} \right]^p, \quad k \in \mathbf{N}^*.$$

Moreover to each λ_k is associated a one-dimensional eigenspace generated by a function ω_k with exactly $k-1$ zeros in $(0,1)$. Concerning the equation (0.1) we prove that each λ_k is a point of bifurcation as in the semilinear case ($p=2$). More precisely we define for $k \in \mathbf{N}^*$

$$(0.5) \quad S_k = \{\varphi \in C: \varphi \text{ has exactly } k-1 \text{ simple zeros in } (0, 1)\},$$

where $C = \{\varphi \in C^1([0, 1]): \varphi(0) = \varphi(1) = 0\}$ and

$$(0.6) \quad S_k^+ = \{\varphi \in S_k: \varphi_x(0) > 0\}, \quad S_k^- = -S_k^+.$$

As λ_1 is defined as the best Poincaré constant in $W_0^{1,p}(0, 1)$, that is,

$$(0.7) \quad \inf \left\{ \int_0^1 |v_x|^p dx : v \in W_0^{1,p}(0, 1), \int_0^1 |v|^p dx = 1 \right\},$$

Received by the editors July 4, 1987 and, in revised form, October 12, 1987.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 35J60; Secondary 34C30.

©1988 American Mathematical Society
0002-9947/88 \$1.00 + \$.25 per page

it is clear that E_λ is reduced to the zero function when $\lambda \leq \lambda_1$.

When $1 < p \leq 2$ we prove that the configuration of E_λ is exactly the same as in the case $p = 2$ [1], that is,

$$(0.8) \quad E_\lambda = \{0, \pm u_l, l = 1, \dots, k: u_l \in S_l^+\}.$$

When $p > 2$ the structure of E_λ can be quite a bit more complicated for large values of λ . Let h be the inverse function of g and $F(r) = \int_0^r f(s) ds$; we define

$$(0.9) \quad \alpha(\lambda) = \left(\frac{\lambda}{p-1} h^p(\lambda) - \frac{p}{p-1} F(h(\lambda)) \right)^{1/p}$$

and

$$(0.10) \quad x(\lambda) = \int_0^{h(\lambda)} \frac{ds}{(\alpha^p(\lambda) + pF(s)/(p-1) - \lambda s^p/(p-1))^{1/p}};$$

and $\lambda \mapsto x(\lambda)$ is a decreasing positive function defined on $(0, +\infty)$. If $\lambda_k < \lambda \leq \lambda_{k+1}$ we then have

$$(0.11) \quad E_\lambda = \{0\} \cup \{\pm u_1\} \bigcup_{p=2}^k \{\pm E_\lambda^l\},$$

where $u_1 \in S_1^+$ and $E_\lambda^l \subset S_l^+$ such that

- (i) E_λ^l contains only one element if $2lx(\lambda) \geq 1$,
- (ii) E_λ^l is diffeomorphic to $[0, 1]^{l-1}$ if $0 < 2lx(\lambda) < 1$. In case (ii) the elements of E_λ^l are constant with value $(-1)^{j+1}h(\lambda)$ on l closed and disconnected subintervals $I_j \subset (0, 1)$, $j = 1, \dots, l$, with total length $1 - 2lx(\lambda)$.

1. The eigenvalue problem. For $p > 1$ we consider the following eigenvalue problem

$$(1.1) \quad \begin{cases} -(|v_x|^{p-2}v_x)_x = \lambda|v|^{p-2}v & \text{in } (0, 1), \\ v(0) = v(1) = 0 \end{cases}$$

and let S be the subset of $W_0^{1,p}(0, 1) \times \mathbf{R}$ of all the (v, λ) , $v \neq 0$, satisfying (1.1).

THEOREM 1.1. *There exists a unique sequence of functions $v_k \in S_k^+$, $k \in \mathbf{N}^*$, with maximal value 1 on $(0, 1)$ such that*

$$(1.2) \quad S = \{(\mu v_k, \lambda_k): k \in \mathbf{N}^*\},$$

where μ is any nonzero real number and

$$(1.3) \quad \lambda_k = k^p \lambda_1 = k^p(p-1) \left[2 \int_0^1 \frac{dt}{(1-t^p)^{1/p}} \right]^p.$$

Moreover the following holds for $m = 0, \dots, k-1$:

$$(1.4) \quad v_k(x) = (-1)^m v_1(kx - m), \quad m/k \leq x \leq (m+1)/k.$$

Before giving the proof it must be noticed that this result is partially contained in [5], in particular formula (1.4).

PROOF. It is clear from (1.1) and $v \in C^0([0, 1])$ and then $v \in C^1([0, 1])$ when $p > 2$ or $v \in C^2([0, 1])$ when $1 < p \leq 2$ (the complete regularity, due to Otani [5], will be given in Remark 1.1).

Step 1. If $(v, \lambda) \in S$ then $v_x(0) \neq 0$ and $\lambda > 0$. Multiplying (1.1) by v and integrating over $(0, 1)$ yields

$$(1.5) \quad \int_0^1 |v_x|^p dx = \lambda \int_0^1 v^p dx.$$

Hence necessarily $\lambda > 0$. Multiplying (1.1) by v_x and integrating over $(0, x)$, $0 < x < 1$, yields the energy estimate

$$(1.6) \quad (p-1)|v_x(x)|^p + \lambda|v(x)|^p = (p-1)|v_x(0)|^p + \lambda|v(0)|^p.$$

As $v(0) = 0$ we need $v_x(0) \neq 0$ in order to have a nonzero v .

Step 2. The explicit construction. Assume v is a nonzero solution with $v_x(0) = \alpha > 0$ for example. Then $v_x > 0$ on $[0, x_0]$ for some $x_0 \in (0, 1)$ and

$$(1.7) \quad v_x(x) = \left(\alpha^p - \frac{\lambda}{p-1} (v(x))^p \right)^{1/p}$$

on $[0, x_0]$, from (1.6), which gives

$$(1.8) \quad x = \int_0^{v(x)} \frac{dt}{(\alpha^p - \lambda t^p/(p-1))^{1/p}}.$$

Moreover this formula remains valid as long as $v(x)$ remains smaller than the first positive zero of the function

$$(1.9) \quad r \mapsto \varphi(\alpha, r) = \alpha^p - \lambda r^p/(p-1)$$

which is $S(\alpha) = ((p-1)/\lambda)^{1/p} \alpha$. As $S(\alpha)$ is simple we define $\theta(\alpha)$ by

$$(1.10) \quad \theta(\alpha) = \int_0^{S(\alpha)} \frac{dt}{(\alpha^p - \lambda t^p/(p-1))^{1/p}}.$$

Moreover $v(\theta(\alpha)) = S(\alpha)$ and $v_x(\theta(\alpha)) = 0$. As $\alpha^p = \lambda S^p(\alpha)/(p-1)$ we get

$$(1.11) \quad \theta(\alpha) = \theta_\lambda = C \left(\frac{p-1}{\lambda} \right)^{1/p}, \quad C = \int_0^1 \frac{ds}{(1-s^p)^{1/p}}.$$

From (1.6) the function v is decreasing on some interval $[\theta_\lambda, \Theta]$, so we get

$$(1.12) \quad x - \theta_\lambda = - \int_{v(x)}^{S(\alpha)} \frac{dt}{[(\lambda/(p-1))(S^p(\alpha) - t^p)]^{1/p}},$$

or

$$x - \theta_\lambda = - \int_{v(x)}^{S(\alpha)} \frac{dt}{(\alpha^p - \lambda t^p/(p-1))^{1/p}};$$

and this formula remains valid as long as v is decreasing, in particular as long as v is positive. If $x_1 \in (0, \theta_\lambda)$ and $x_2 = 2\theta_\lambda - x_1$ then

$$x_1 = \int_0^{v(x_1)} \frac{dt}{(\varphi(\alpha, t))^{1/p}}, \quad \theta_\lambda - x_1 = - \int_{v(x_2)}^{S(\alpha)} \frac{dt}{(\varphi(\alpha, t))^{1/p}}$$

and $v(x_1) = v(x_2)$. As a consequence $x = \theta_\lambda$ is an axis of symmetry for the restriction of v to $[0, 2\theta_\lambda]$ and $x = 2\theta_\lambda$ is a center of symmetry for the restriction of

v to $[0, 4\theta_\lambda]$. Hence the function v is $4\theta_\lambda$ -periodic on $[0, +\infty)$. The necessary and sufficient condition for the restriction of v to $[0, 1]$ to be a solution of (1.1) is then

$$(1.13) \quad 1/2\theta_\lambda \in \mathbf{N}^*,$$

which means (1.3). As for the number of zeros of v in $(0, 1)$ it is given by $1/2\theta_\lambda - 1$. Using the homogeneity of (1.1) we get the desired result as the uniqueness is a consequence of the construction of v .

REMARK 1.1. Existence and uniqueness of the first positive normalized eigenfunction of $-\operatorname{div}(|D \cdot|^{p-2} D \cdot)$ in $W_0^{1,p}(\Omega)$ have been obtained by De Thelin in the radial case when Ω is a ball [7] and Guedda-Veron for general Ω with a connected C^2 boundary [4].

As for the regularity of v we have

$$(1.14) \quad v \in C^\alpha([0, 1]) \cap C^{\langle p \rangle}([0, 1] \setminus Z)$$

where $Z = \{x \in (0, 1) : v_x(x) = 0\}$, $\alpha = \min(\langle (2-p)/(p-1) \rangle + 1, \langle p \rangle)$ and $\langle r \rangle = +\infty$ if $r \in 2\mathbf{N}^*$ or $\langle r \rangle = \min\{n : n \in \mathbf{N}^*, n \geq r\}$ if not.

REMARK 1.2. We have the following Poincaré type relation

$$(1.15) \quad \lambda_1 = \inf \left\{ \int_0^1 |u_x|^p dx / \int_0^1 |u|^p dx : u \in W_0^{1,p}(0, 1) \setminus \{0\} \right\}$$

and the infimum is achieved for $u = v_1$.

2. The bifurcation phenomena. In this section we consider the following equation

$$(2.1) \quad \begin{cases} -(|u_x|^{p-2} u_x)_x + f(u) = \lambda |u|^{p-2} u & \text{in } (0, 1), \\ u(0) = u(1) = 0, \end{cases}$$

where $p > 1$ and $\lambda \in \mathbf{R}$. As for f we first assume that

$$(2.2) \quad f \text{ is a } C^1 \text{ odd function,}$$

$$(2.3) \quad s \mapsto f(s)/s^{p-1} \text{ is strictly increasing on } (0, +\infty) \text{ with limit } 0 \text{ at } 0,$$

$$(2.4) \quad \lim_{s \rightarrow +\infty} f(s)/s^{p-1} = +\infty.$$

We then define

$$(2.5) \quad h \text{ is the inverse function of the restriction of } f(s)/s^{p-1} \text{ to } (0, +\infty),$$

$$(2.6) \quad H(s) = \lambda s^p - pF(s),$$

where $F(s) = \int_0^s f(t) dt$. For $\lambda > 0$ we shall also consider the following hypothesis:

$$(2.7) \quad (p-1)(H'(s))^2 - pH(s)H''(s) \geq 0 \quad \text{for any } s \in [0, h(\lambda)].$$

Let E_λ be the set of all the solutions of (2.1) in $W_0^{1,p}(0, 1)$ and λ_k be defined by (1.3). When $1 < p \leq 2$ the structure of E_λ is exactly the same as in the case $p = 2$.

THEOREM 2.1. Assume $1 < p \leq 2$ and (2.2)–(2.7). Then

- (i) if $\lambda \leq \lambda_1$, $E_\lambda = \{0\}$, and
- (ii) if $\lambda_k < \lambda \leq \lambda_{k+1}$ for some $k \in \mathbb{N}^*$

$$(2.8) \quad E_\lambda = \{0, \pm u_1, \dots, \pm u_k\},$$

where $u_l \in S_l^+$ for $l = 1, \dots, k$.

REMARK 2.1. The assumption (2.7), which is equivalent to the fact that $s \mapsto H^{p-1}(s)/H^p(s)$ is nondecreasing on $[0, h(\lambda)]$, is essential for uniqueness but not for existence. In the particular case where $f(r) = |r|^{q-1}r$ with $q > p - 1$ then $h(\lambda) = \lambda^{1/(q+1-p)}$, $H(s) = \lambda s^p - p s^{q+1}/(q+1)$ and (2.7) is satisfied.

PROOF OF THEOREM 2.1. As in Theorem 1.1 it is clear that any solution of (2.1) in $W_0^{1,p}(0,1)$ is continuous and at least C^2 (remember that $1 < p \leq 2$). Multiplying the equation by u yields

$$(2.9) \quad \int_0^1 |u_x|^p dx + \int_0^1 u f(u) dx = \lambda \int_0^1 |u|^p dx.$$

From Remark 1.2 a nonzero solution of (2.1) can exist only if $\lambda > \lambda_1$, which will be assumed in the sequel.

Step 1. If u is a nonzero solution of (2.1) then $u_x(0) \neq 0$. Although it is a consequence of a general result due to Franchi, Lanconelli and Serrin, we give here a direct proof which also works when $p > 2$. Multiplying (2.1) by u_x yields the energy relation

$$(2.10) \quad \begin{aligned} & -\frac{p-1}{p} |u_x(x)|^p + F(u(x)) - \frac{\lambda}{p} |u(x)|^p \\ & = -\frac{p-1}{p} |u_x(0)|^p + F(u(0)) - \frac{\lambda}{p} |u(0)|^p. \end{aligned}$$

If we assume that $u_x(0) = 0$ we get

$$(2.11) \quad |u_x(x)|^p = \frac{1}{p-1} (pF(u(x)) - \lambda |u(x)|^p).$$

As the function $x \mapsto pF(x) - \lambda |x|^p$ is negative on $(-\rho, \rho) \setminus \{0\}$, u_x is always 0 and $u \equiv 0$.

Step 2. The explicit construction. Without any loss of generality we assume $u_x(0) = \alpha > 0$. Hence u is increasing on some interval $[0, x_0]$ and from (2.10) we get

$$(2.12) \quad u_x^p(x) = \alpha^p + \frac{p}{p-1} F(u(x)) - \frac{\lambda}{p-1} u^p(x)$$

which gives u as the inverse function of a p -elliptic integral

$$(2.13) \quad x = \int_0^{u(x)} \frac{dt}{(\alpha^p + pF(t)/(p-1) - \lambda t^p/(p-1))^{1/p}}$$

on $[0, x_0]$. Moreover this formula remains valid as long as $u(x)$ is smaller than the first positive zero of

$$(2.14) \quad r \mapsto \Psi(\alpha, r) = \alpha^p + \frac{p}{p-1} F(r) - \frac{\lambda}{p-1} |r|^p.$$

But the function $\Psi(\alpha, \cdot)$ is decreasing in $[0, h(\lambda)]$ and increasing on $[h(\lambda), +\infty)$; hence there are three possibilities.

Case 1. $\alpha^p > \lambda h^p(\lambda)/(p-1) - pF(h(\lambda))/(p-1) = \alpha^p(\lambda)$.

In that case the function $r \mapsto \int_0^r ds/(\Psi(\alpha, s))^{1/p}$ is an increasing C^2 diffeomorphism from \mathbf{R}^+ onto \mathbf{R}^+ and it is the same with u defined by (2.13) which cannot belong to E_λ .

Case 2. $\alpha^p = \alpha^p(\lambda)$.

In that case $h(\lambda)$ is a double zero for $\Psi(\alpha, \cdot)$, and as $1 < p \leq 2$

$$\int_0^{h(\lambda)} ds/(\Psi(\alpha, s))^{1/p} = +\infty.$$

As in Case 1 the function $r \mapsto \int_0^r ds/(\Psi(\alpha, s))^{1/p}$ is a C^2 diffeomorphism from $[0, h(\lambda))$ onto \mathbf{R}^+ and u cannot belong to E_λ .

Case 3. $\alpha^p < \alpha^p(\lambda)$.

In that case $\Psi(\alpha, \cdot)$ admits a simple zero $S(\alpha)$ in $(0, h(\lambda))$. As $(\partial\Psi/\partial r)(\alpha, S(\alpha)) \neq 0$, $r \mapsto (\Psi(\alpha, r))^{-1/p}$ is integrable on $(0, S(\alpha))$ and we define

$$(2.15) \quad \theta(\alpha) = \int_0^{S(\alpha)} \frac{ds}{(\Psi(\alpha, s))^{1/p}}.$$

Relation (2.13) remains valid on $[0, \theta(\alpha)]$ and we have

$$(2.16) \quad u(\theta(\alpha)) = S(\alpha), \quad u_x(\theta(\alpha)) = 0.$$

Using the energy relation at $\theta(\alpha)$ we have

$$(2.17) \quad \frac{p-1}{p} |u_x(x)|^p = \frac{\lambda}{p} S^p(\alpha) - F(S(\alpha)) - \left(\frac{\lambda}{p} u^p(x) - F(u(x)) \right)$$

or

$$|u_x(x)|^p = \alpha^p + \frac{p}{p-1} F(u(x)) - \frac{\lambda}{p-1} u^p(x).$$

Hence u is decreasing on some interval $[\theta(\alpha), \Theta]$ and we have

$$(2.18) \quad x - \theta(\alpha) = - \int_{u(x)}^{S(\alpha)} \frac{ds}{(\Psi(\alpha, s))^{1/p}}.$$

This formula remains valid as long as u is decreasing, and as in §1 $x = \theta(\alpha)$ is an axis of symmetry for the restriction of u to $[0, 2\theta(\alpha)]$ and $x = 2\theta(\alpha)$ is a center of symmetry for the restriction of u to $[0, 4\theta(\alpha)]$; the necessary and sufficient condition for u to be a solution of (2.1) is that

$$(2.19) \quad 1/2\theta(\alpha) \in \mathbf{N}^*.$$

Step 3. The function $\alpha \mapsto S(\alpha)$ is convex, increasing on $[0, \alpha(\lambda))$. We have $\Psi(\alpha, S(\alpha)) = 0$ and $(\partial\Psi/\partial r)(\alpha, S(\alpha)) \neq 0$. By the implicit function theorem $\alpha \mapsto S(\alpha)$ is C^2 . We also have

$$\frac{d}{d\alpha}(\Psi(\alpha, S(\alpha))) = \frac{\partial\Psi}{\partial\alpha}(\alpha, S(\alpha)) + \frac{\partial\Psi}{\partial r}(\alpha, S(\alpha)) \frac{dS}{d\alpha}(\alpha)$$

which gives

$$(2.20) \quad \frac{dS}{d\alpha}(\alpha) = \frac{(p-1)\alpha^{p-1}}{\lambda S^{p-1}(\alpha) - f(S(\alpha))} = \frac{p(p-1)\alpha^{p-1}}{H'(S(\alpha))}.$$

As $S(\alpha) < h(\lambda)$, $\alpha \mapsto S(\alpha)$ is increasing on $[0, \alpha(\lambda))$. Moreover

$$\frac{d^2 S}{d\alpha^2}(\alpha) = p(p-1) \frac{(p-1)\alpha^{p-2}H'(S(\alpha)) - \alpha^{p-1}H''(S(\alpha))dS/d\alpha}{(H'(S(\alpha)))^2}.$$

Using (2.20) and the definition of $S(\alpha)$ and H we get

$$(2.21) \quad \frac{d^2 S}{d\alpha^2}(\alpha) = p(p-1)\alpha^{p-2} \frac{(p-1)(H'(S(\alpha)))^2 - pH(S(\alpha))H''(S(\alpha))}{(H'(S(\alpha)))^3}.$$

From (2.7) we deduce $d^2 S(\alpha)/d\alpha^2 \geq 0$.

Step 4. The function $\alpha \mapsto \theta(\alpha)$ is continuous increasing on $[0, \alpha(\lambda))$. For $t \in [0, \alpha]$ the function $s \mapsto \Psi(t, s)$ admits a first positive zero at $S(t)$ which means

$$t^p + \frac{p}{p-1}F(S(t)) - \frac{\lambda}{p-1}S^p(t) = 0 \quad \text{and} \quad \Psi(\alpha, S(t)) = \alpha^p - t^p.$$

Taking t as a new variable in (2.15) we get

$$(2.22) \quad \theta(\alpha) = \int_0^\alpha \frac{dS}{dt}(t) \frac{dt}{(\alpha^p - t^p)^{1/p}}$$

or

$$(2.23) \quad \theta(\alpha) = \int_0^1 \frac{dS}{dt}(\alpha\sigma) \frac{d\sigma}{(1 - \sigma^p)^{1/p}}.$$

As ds/dt is increasing and C^1 on $[0, \alpha(\lambda))$, it is the same with $\alpha \mapsto \theta(\alpha)$.

Step 5. End of the proof. As $\lim_{\alpha \downarrow 0} S(\alpha) = 0$ and $\lim_{\alpha \downarrow 0} F(S(\alpha))/S^p(\alpha) = 0$ we get

$$(2.24) \quad S(\alpha) \underset{\alpha \downarrow 0}{\sim} \alpha \left(\frac{p-1}{\lambda} \right)^{1/p}$$

which implies

$$\lim_{\alpha \downarrow 0} \frac{dS}{d\alpha}(\alpha) = \left(\frac{p-1}{\lambda} \right)^{1/p}$$

and

$$(2.25) \quad \lim_{\alpha \downarrow 0} \theta(\alpha) = \left(\frac{p-1}{\lambda} \right)^{1/p} \int_0^1 \frac{d\sigma}{(1 - \sigma^p)^{1/p}} = \frac{1}{2} \left(\frac{\lambda_1}{\lambda} \right)^{1/p}.$$

For the other bound we have $\lim_{\alpha \downarrow \alpha(\lambda)} S(\alpha) = h(\lambda)$. As $h(\lambda)$ is just a double zero for $\Psi(\alpha(\lambda), r)$, there exists a continuous and bounded function φ on $[0, \alpha(\lambda)]$ such that

$$\Psi(\alpha(\lambda), r) = (h(\lambda) - r)^2 \varphi(r).$$

Moreover

$$\begin{aligned} \int_0^{S(\alpha)} (\Psi(\alpha, t))^{-1/p} dt &> \int_0^{S(\alpha)} (\Psi(\alpha(\lambda), t))^{-1/p} dt \\ &= \int_0^{S(\alpha)} (h(\lambda) - t)^{-2/p} (\varphi(t))^{-1/p} dt. \end{aligned}$$

As $1 < p \leq 2$ we get

$$(2.26) \quad \lim_{\alpha \uparrow \alpha(\lambda)} \theta(\alpha) = \int_0^{h(\lambda)} (h(\lambda) - t)^{-2/p} (\varphi(t))^{-1/p} dt = +\infty.$$

As a consequence $\alpha \mapsto \theta(\alpha)$ is an increasing diffeomorphism from $(0, \alpha(\lambda))$ onto $(\frac{1}{2}(\lambda_1/\lambda)^{1/p}, +\infty)$ and $1/2\theta(\alpha)$ a decreasing diffeomorphism from $(0, \alpha(\lambda))$ onto $(0, (\lambda/\lambda_1)^{1/p})$. If we assume that $\lambda_k < \lambda \leq \lambda_{k+1}$ for some $k \in \mathbf{N}^*$ there exist exactly k integers $l = 1, \dots, k$ and k positive real numbers α_l such that $1/2\theta(\alpha_l) = l$. If u_l is the solution of the initial value problem

$$(2.27) \quad \begin{cases} -(|u_{lx}|^{p-2}u_{lx})_x + f(u_l) = \lambda|u_l|^{p-2}u_l & \text{on } (0, 1), \\ u_l(0) = 0, \quad u_{lx}(0) = \alpha_l, \end{cases}$$

then $u_l(1) = 0$, $u_l \in S_l^+$. We get the result in considering $-u_l$, $l = 1, \dots, k$.

REMARK 2.2. If we represent the bifurcation diagram (λ, u_λ) then there exists no secondary bifurcation along the branches of solutions in S_k^\pm issuing from λ_k .




FIGURE 1

In the case $p > 2$ the main difference will come from the fact that the following integral

$$(2.28) \quad x(\lambda) = \int_0^{h(\lambda)} \frac{ds}{(\alpha^p(\lambda) + \frac{p}{p-1}F(s) - \frac{\lambda}{p-1}s^p)^{1/p}}$$

is finite as $h(\lambda)$ is a double zero of $\Psi(\alpha(\lambda), r)$.

THEOREM 2.2. Assume $p > 2$ and (2.2)–(2.7). Then

- (i) if $\lambda \leq \lambda_1$ $E_\lambda = \{0\}$,
- (ii) if $\lambda_k < \lambda \leq \lambda_{k+1}$ for some $k \in \mathbf{N}^*$

$$(2.29) \quad E_\lambda = \{0\} \cup \{\pm u_1\} \bigcup_{l=2}^k \{\pm E_\lambda^l\},$$

where $u_1 \in S_1^+$ and $E_\lambda^l \subset S_l^+$, $l = 2, \dots, k$, and

E_λ^l is reduced to a single element if $2lx(\lambda) \geq 1$,

E_λ^l is diffeomorphic to $[0, 1]^{l-1}$ if $0 < 2px(\lambda) < 1$.¹

PROOF. The idea is essentially the same as in Theorem 2.1 except that in Step 2, Case 2 (that is, if $\alpha^p = \alpha^p(\lambda)$) gives rise to solutions of (2.1) with maximum value $h(\lambda)$, and in that case Serrin and Veron's existence and uniqueness result does not apply; moreover the value $u = h(\lambda)$ is a bifurcation value for (2.1).

Step 1. Assume $2x(\lambda) \geq 1$. Then the construction of Theorem 2.1 works: the function $\alpha \mapsto 1/2\theta(\alpha)$ is a decreasing diffeomorphism from $(0, \alpha(\lambda)]$ onto $[1/2x(\lambda), (\lambda/\lambda_1)^{1/p}]$. As $\lambda_k < \lambda \leq \lambda_{k+1}$ there exist exactly k integers $1, 2, \dots, k$ and k positive real numbers $\alpha_1, \dots, \alpha_k$ such that $1/2\theta(\alpha_l) = l \in [1/2x(\lambda), (\lambda/\lambda_1)^{1/p}]$, $l = 1, \dots, k$. and we get the corresponding solutions $u_l \in S_l^+$ by (2.26).

Step 2. Assume $4x(\lambda) \geq 1 > 2x(\lambda)$. All the elements $u_l = 2, \dots, k$ in S_l^+ are constructed as in Step 1. As for the element $u_1 \in S_1^+$ it has necessarily the following form as the initial slope must be $\alpha(\lambda)$:

$$(2.30) \quad \begin{aligned} &\text{for } 0 \leq x \leq x(\lambda) \\ x &= \int_0^{u_1(x)} \frac{dt}{(\Psi(\alpha(\lambda), t))^{1/p}}, \end{aligned}$$

$$(2.31) \quad \begin{aligned} &\text{for } x(\lambda) \leq x \leq 1 - x(\lambda) \\ u_1(x) &= h(\lambda), \\ &\text{for } 1 - x(\lambda) \leq x \leq 1 \end{aligned}$$

$$(2.32) \quad x - (1 - x(\lambda)) = - \int_{u_1(x)}^{h(\lambda)} \frac{dt}{(\Psi(\alpha(\lambda), t))^{1/p}}.$$

Step 3. Assume $0 < 2lx(\lambda) < 1$ for some $l \in \{2, \dots, k\}$. We can construct all the elements of $E_\lambda \cap S_l^+$ in the following way as their initial slope is necessarily $\alpha(\lambda)$:

$$(2.33) \quad \begin{aligned} &\text{for } 0 \leq x \leq x(\lambda) \\ x &= \int_0^{u_l(x)} \frac{dt}{(\Psi(\alpha(\lambda), t))^{1/p}}, \end{aligned}$$

$$(2.34) \quad \begin{aligned} &\text{for } x(\lambda) \leq x \leq x_1 \text{ where } x_1 \in (x(\lambda), 1) \text{ and} \\ x_1 - x(\lambda) &\leq 1 - 2lx(\lambda) \\ &\text{then } u_l(x) = h(\lambda), \end{aligned}$$

$$(2.35) \quad \begin{aligned} &\text{for } x_1 \leq x \leq 2x(\lambda) + x_1 \\ x - x_1 &= - \int_{u_l(x)}^{h(\lambda)} \frac{dt}{(\Psi(\alpha(\lambda), t))^{1/p}}, \end{aligned}$$

$$(2.36) \quad \begin{aligned} &\text{for } x_1 + 2x(\lambda) \leq x \leq x_2 \text{ where } x_2 \in (x_1 + 2x(\lambda), 1) \text{ and} \\ x_2 - (x_1 + 2x(\lambda)) + x_1 - x(\lambda) &\leq 1 - 2lx(\lambda) \\ &\text{then } u_l(x) = -h(\lambda). \end{aligned}$$

¹And more naturally to the set $K_l = \{x = (x^1, \dots, x^l), x^j \geq 0, \sum_{j=1}^l x^j = 1 - 2lx(\lambda)\}$.

Continuing this procedure any solution $u_l \in S_l^+$ is defined by the intervals $I_j = [x_{j-1} + 2x(\lambda), x_j]$, $j = 1, \dots, l$, and $x_0 = -x(\lambda)$ where it takes the constant value $(-1)^{j+1}h(\lambda)$ and the intervals $[x_{j-1}, x_{j-1} + 2x(\lambda)]$ where it is defined by

$$(2.37) \quad x - x_{j-1} = - \int_{u_l(x)}^{h(\lambda)} \frac{dt}{(\Psi(\alpha(\lambda), t))^{1/p}}$$

if j is even or

$$(2.38) \quad x - x_{j-1} = \int_{h(\lambda)}^{u_l(x)} \frac{dt}{(\Psi(\alpha(\lambda), t))^{1/p}}$$

if j is odd.

From the above construction the total length of the I_j is $1 - 2lx(\lambda)$ and the set E_λ^l of the u_l is diffeomorphic to the $(l-1)$ -dimensional cube.




FIGURE 2. Example of construction of E_λ^3

REMARK 2.3. It is important to notice that this type of secondary bifurcation along the branch of solutions issuing from λ_k , $k \geq 2$, always appears if we have

$$(2.39) \quad \lim_{\lambda \rightarrow +\infty} x(\lambda) = 0.$$

This is in particular the case if $f(r) \sim_{r \rightarrow +\infty} |r|^{q-1}r$ which implies

$$(2.40) \quad x(\lambda) \sim_{\lambda \rightarrow +\infty} \lambda^{-1/p} \int_0^1 \left(\frac{q+1-p}{(p-1)(q+1)} + \frac{p}{(p-1)(q+1)} \sigma^{q+1} - \frac{\sigma^p}{p-1} \right)^{-1/p} d\sigma.$$

However this is not always the case under conditions (2.2)–(2.7), for example, with $f(r) = (|r|^{p-2} \text{Log } |r|)r$ for $|r| \geq 2$, where we get

$$(2.41) \quad \lim_{\lambda \rightarrow +\infty} x(\lambda) = \int_0^1 \left(\frac{1}{p(p-1)} (1 - \sigma^p) + \frac{1}{p-1} \sigma^p \text{Log } \sigma \right)^{-1/p} d\sigma.$$

We finally have the following exclusion principle.

THEOREM 2.3. Assume $p > 1$, (2.2)–(2.7), g is a continuous even function increasing on \mathbf{R}^+ and u_1 and u_2 are two solutions of (2.1); then

(i) if u_1 and u_2 have the same number of zeros

$$(2.42) \quad \int_0^1 g(u_1(x)) dx = \int_0^1 g(u_2(x)) dx;$$

(ii) if u_1 and u_2 do not have the same number of zeros

$$(2.43) \quad \int_0^1 g(u_1(x)) dx \neq \int_0^1 g(u_2(x)) dx.$$

PROOF. It is clear that for any function $\int_0^1 g(u(x)) dx$ is equal to $\int_0^1 g(-u(x)) dx$. When $p > 2$ we have only to consider two solutions of E_λ with the same number of zeros and belonging to some E_λ^l , $l \geq 2$, in the case $2lx(\lambda) < 1$. In that case u_1 and u_2 take the value $\pm h(\lambda)$ on l intervals I_j^1 and I_j^2 , $j = 1, \dots, l$, which are disconnected and have the same total length which gives

$$(2.44) \quad \int_{\bigcup_j I_j^1} g(u_1(x)) dx = \int_{\bigcup_j I_j^2} g(u_2(x)) dx = (1 - 2lx(\lambda))g(h(\lambda)).$$

On $(0, 1) \setminus \{\bigcup_j I_j^1\}$ or $(0, 1) \setminus \{\bigcup_j I_j^2\}$ u_1 and u_2 are defined by the same types of formula ((2.32) or (2.30)) and the integral of $g(u_i)$ over these sets is

$$2l \int_0^{x(\lambda)} g(u_1(x)) dx.$$

Hence, for $i = 1, 2$, we get

$$(2.45) \quad \int_0^1 g(u_i(x)) dx = (1 - 2lx(\lambda))g(h(\lambda)) + 2l \int_0^{x(\lambda)} g(u_i(x)) dx$$

which proves (i).

For proving (ii) we shall assume either $1 < p \leq 2$ or $p > 2$ but u_1 and u_2 are not constant on any subinterval of $(0, 1)$ (the other case is essentially the same). If u_1 and u_2 do not have the same number of zeros in $(0, 1)$ we can assume $u_{1x}(0) = \alpha$, $u_{2x}(0) = \beta$, $0 < \alpha < \beta$; u_1 is $4\theta(\alpha)$ -periodic, u_2 is $4\theta(\beta)$ -periodic and $0 < \theta(\alpha) < \theta(\beta)$. Moreover

$$(2.46) \quad \frac{1}{2\theta(\alpha)} = k_1, \quad \frac{1}{2\theta(\beta)} = k_2, \quad k_1, k_2 \in \mathbf{N}^*, \quad k_1 > k_2.$$

Step 1. For $0 < x < \theta(\alpha)$ we have $0 < u_1(x) < u_2(x)$. On a right neighbourhood of 0 we have $u_1 < u_2$, and u_1 and u_2 are increasing on $[0, \theta(\alpha)]$. If we assume the existence of some $x_0 \in [0, \theta(\alpha)]$ such that $u_1(x_0) = u_2(x_0)$, we can always suppose that $u_1 < u_2$ in $(0, x_0)$ and then $u_{1x}(x_0) \geq u_{2x}(x_0)$. The energy relation implies

$$(2.47) \quad \begin{aligned} \alpha^p + \frac{p}{p-1} F(u_1(x_0)) - \frac{\lambda}{p-1} u_1^p(x_0) \\ \geq \beta^p + \frac{p}{p-1} F(u_2(x_0)) - \frac{\lambda}{p-1} u_2^p(x_0) \end{aligned}$$

and $\alpha \geq \beta$ which is impossible.

Step 2. End of the proof. From Step 1: $0 < u_1(x) < u_2(x')$ for $0 < x < \theta(\alpha)$ and $0 < x \leq x' < \theta(\beta)$. Set φ the lowest common multiple to k_1 and k_2 . There exist n_1 and $n_2 \in \mathbf{N}^*$ such that $n_1 k_1 = n_2 k_2 = \varphi$ and

$$(2.48) \quad n_1/\theta(\alpha) = n_2/\theta(\beta), \quad 0 < n_1 < n_2.$$

Then

$$(2.49) \quad \int_0^1 g(u_1(x)) dx = \frac{1}{\theta(\alpha)} \int_0^{\theta(\alpha)} g(u_1(x)) dx,$$

$$(2.50) \quad \int_0^1 g(u_2(x)) dx = \frac{1}{\theta(\beta)} \int_0^{\theta(\beta)} g(u_2(x)) dx.$$

Setting $T = n_2\theta(\alpha) = n_1\theta(\beta)$, we have

$$\begin{aligned} \frac{1}{\theta(\alpha)} \int_0^{\theta(\alpha)} g(u_1(x)) dx &= \frac{1}{n_2\theta(\alpha)} \int_0^{n_2\theta(\alpha)} g\left(u_1\left(\frac{\sigma}{n_2}\right)\right) d\sigma \\ &= \frac{1}{T} \int_0^T g\left(u_1\left(\frac{\sigma}{n_2}\right)\right) d\sigma \end{aligned}$$

and

$$\frac{1}{\theta(\beta)} \int_0^{\theta(\beta)} g(u_2(x)) dx = \frac{1}{T} \int_0^T g\left(u_2\left(\frac{\sigma}{n_1}\right)\right) d\sigma,$$

which implies

$$(2.51) \quad \int_0^1 g(u_1(x)) dx < \int_0^1 g(u_2(x)) dx.$$

REMARK 2.4. As a consequence there exist $k+1$ different critical values for the energy functional

$$(2.52) \quad J(\omega) = \frac{1}{p} \int_0^1 |\omega_x|^p dx + \int_0^1 F(\omega) dx - \frac{\lambda}{p} \int_0^p |\omega|^p dx$$

defined in $W_0^{1,p}(0,1)$, for $\lambda_k < \lambda \leq \lambda_{k+1}$; those critical values only depend on the set S_l , $l = 1, \dots, k$, the critical points of (2.52) belong to. This is an immediate consequence of Theorem 2.3 and the fact that

$$(2.53) \quad J(u) = \int_0^1 \left(F(u) - \frac{1}{p} u f(u) \right) dx$$

for $u \in E_\lambda$.

REFERENCES

1. H. Berestycki, *Le nombre de solutions de certains problèmes semi-linéaires elliptiques*, J. Funct. Anal. **40** (1981), 1–19.
2. M. S. Berger, *Nonlinearity and functional analysis*, Academic Press, 1977.
3. X. Y. Chen, H. Matano and L. Veron, *Anisotropic singularities of solutions of nonlinear elliptic equations in \mathbf{R}^2* , J. Funct. Anal. (to appear).
4. M. Guedda and L. Veron, *Quasilinear elliptic equations involving critical Sobolev exponents*, Nonlinear Anal. (to appear).

5. M. Otani, *On certain second order differential equations associated with Sobolev-Poincaré-type inequalities*, Nonlinear Anal. **8** (1984), 1255–1270.
6. F. De Thelin, *Quelques resultats d'existence et de non-existence pour une E. D. P. elliptique non linéaire*, C. R. Acad. Sci. Paris **299** (1984), 911–914.
7. ———, *Sur l'espace propre associé à la première valeur propre du pseudo-laplacien*, C. R. Acad. Sci. Paris **303** (1986), 355–358.

DÉPARTEMENT DE MATHÉMATIQUES, FACULTÉ DES SCIENCES, PARC DE GRANDMONT,
F-37200 TOURS, FRANCE