

ON THE GENUS OF SMOOTH 4-MANIFOLDS

ALBERTO CAVICCHIOLI

ABSTRACT. The projective complex plane and the “twisted” S^3 bundle over S^1 are proved to be the unique closed prime connected (smooth or PL) 4-manifolds of genus two. Then the classification of the nonorientable 4-manifolds of genus 4 is given. Finally the genus of a manifold M is shown to be related with the 2nd Betti number of M and some applications are proved in the general (resp. simply-connected) case.

1. INTRODUCTION

Throughout the paper, we consider smooth or combinatorial (PL) closed 4-manifolds indifferently. In fact, a well-known result states that if M is a PL 4-manifold, then M possesses a C^∞ -differentiable structure compatible with the triangulation (see [20]).

For the topology of 4-manifolds we refer to [9, 15].

We recall the definition of (regular) genus for a closed (PL) n -manifold as introduced in [10].

An $(n+1)$ -colored graph is a pair (G, c) , where $G = (V, E)$ is a finite multigraph, regular of degree $n+1$, and $c: E \rightarrow \Delta_n = \{0, 1, \dots, n\}$ is an edge-coloration on G with $n+1$ colors (i.e., $c(e) \neq c(f)$ for any pair of adjacent edges $e, f \in E$). The graph (G, c) is said to be *contracted* if the partial subgraph $G_i = (V, c^{-1}(\Delta_n - \{i\}))$ is connected for each color $i \in \Delta_n$. An n -pseudocomplex (see [14]) $K = K(G)$ can be uniquely associated with (G, c) so that $|G|$ becomes its dual 1-skeleton (see [5]). A *crystallization* of a closed (PL) n -manifold M is a contracted $(n+1)$ -colored graph (G, c) which *represents* M , i.e. $|K(G)| \simeq_{\text{PL}} M$.

It is well known that each closed connected (PL) n -manifold admits a crystallization as proved in [17].

Given a crystallization (G, c) , the minimum genus of a closed (connected) surface into which (G, c) regularly imbeds (see [19]) is denoted by $g(G)$. The *regular genus* (or simply called the *genus*) of a closed (PL) n -manifold M is

Received by the editors October 28, 1988 and June 10, 1989 and, in revised form, February 1, 1990. The contents of the paper have been presented to a meeting of the G.N.S.A.G.A. of the C.N.R. (National Research Council) of Italy, held in Trieste, Italy, Octob 6–8, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 57N13, 57Q05, 57R60.

Key words and phrases. Smooth 4-manifold, crystallization, genus, collapsing, intersection form, homotopy sphere.

Work performed under the auspices of the G.N.S.A.G.A. of the C.N.R. and financially supported by the M.P.I. of Italy within the project “Geometria delle Varietà Differenziabili”.

defined as the nonnegative integer (see [10])

$$g(M) = \min\{g(G)/(G, c) \text{ is a crystallization of } M\}.$$

Given a closed smooth connected 4-manifold M , the following results are well known. If $g(M) = 0$, then $M \simeq_{\text{PL}} S^4$ (4-sphere) (see [6]). If $g(M) = 1$, then $M \simeq_{\text{PL}} S^1 \times S^3$ (see [2]). If $\Pi_1(M)$ is the fundamental group of M , then $g(M) \geq \text{rank } \Pi_1(M)$ (see [1]). If M is nonorientable, then $g(M)$ is even (see [10]). In [7] bounds were determined for the genus of any 4-manifold which is a product of S^1 by a closed 3-manifold or a product of two closed surfaces.

Remark. Let (G_i, c_i) be a crystallization of a closed connected n -manifold M_i ($i = 1, 2$), and let $f_i: |G_i| \rightarrow S_i$ be a regular imbedding of G_i into the closed connected surface S_i or genus $g(M_i)$. By direct construction it is very easy to obtain a regular imbedding of $G_1 \# G_2$ into the surface $S_1 \# S_2$. If M_1 and M_2 are both orientable (resp. nonorientable), then the genus is subadditive, i.e. $g(M_1 \# M_2) \leq g(M_1) + g(M_2)$, since the genus of $S_1 \# S_2$ is exactly $g(M_1) + g(M_2)$. If M_1 is orientable and M_2 is nonorientable, then we have $g(M_1 \# M_2) \leq 2g(M_1) + g(M_2)$ since the genus of $S_1 \# S_2$ is just $2g(M_1) + g(M_2)$.

Now we state the main results of the present paper:

Proposition 1. *Let M^4 be a smooth (or PL) closed connected 4-manifold of genus g .*

(a) *If $g = 2$ and M is orientable, then M is (PL) homeomorphic to either the projective complex plane CP^2 or the connected sum $\#_2(S^1 \times S^3)$.*

(b) *If $g = 2$ and M is nonorientable, then M is (PL) homeomorphic to $S^1 \times S^3$ (the “twisted” S^3 bundle over S^1).*

(c) *If $g = 4$ and M is nonorientable, then M is (PL) homeomorphic to either $\#_2(S^1 \times S^3)$ or $(S^1 \times S^3) \# (S^1 \times S^3)$.*

Corollary 1.

$$g(RP^4) = g\left(\#_2(S^1 \times S^3) \# (S^1 \times S^3)\right) = g(\pm CP^2 \# S^1 \times S^3) = 6,$$

where RP^4 denotes the real projective 4-dimensional space.

The formulae, used in the proof of Proposition 1, imply the following

Proposition 2. *Let M^4 be a closed connected smooth (or PL) orientable 4-manifold of genus g . Then $b_2(M) \leq [(5/2)g]$. If M is simply-connected, then $b_2(M) \leq [g/2]$. Here $[x]$ and $b_k(M)$ denote the integer part of x and the k th Betti number of M respectively.*

Corollary 2.

$$(1) \quad g\left(\#_k(S^1 \times S^3)\right) = k,$$

$$(2) \quad g\left(\#_k \pm CP^2\right) = 2k,$$

$$(3) \quad g\left(\#_k(S^2 \times S^2)\right) = 4k.$$

Now we can apply the Freedman classification of simply-connected smooth 4-manifolds (see [9]) to obtain the following consequence of Proposition 2:

Proposition 3. *If M^4 is a closed simply-connected smooth 4-manifold of genus $g \leq 31$, then we have either*

$$M \cong_{\text{TOP}} \#_r (\pm CP^2) \quad \text{or} \quad M \cong_{\text{TOP}} \#_r (S^2 \times S^2)$$

where $r = b_2(M)$.

Other applications of these results and some open questions related with the 4-dimensional Poincaré conjecture complete the paper.

Acknowledgement. The author wishes to thank the referee for his useful suggestions.

2. PROOF OF PROPOSITION 1: THE ORIENTABLE CASE

In order to prove Proposition 1.a, we recall some constructions and results given in [2].

Let M be a closed connected orientable smooth (or PL) 4-manifold of genus g . Let (G, c) be a crystallization of M and $\{v_i | i \in \Delta_4\}$ the vertex-set of $K = K(G)$. If $\{i, j\} = \Delta_4 - \{r, s, t\}$, then $K(i, j)$ (resp. $K(r, s, t)$) represents the subcomplex of K generated by the vertices v_i and v_j (resp. v_r, v_s and v_t). By g_{rst} (resp. g_{ij}) we denote the number of edges (resp. triangles) of $K(i, j)$ (resp. $K(r, s, t)$). It is very easy to see that g_{rst} and g_{ij} also represent the numbers of components of the subgraphs $G_{\{r,s,t\}}$ and $G_{\{i,j\}}$ respectively.

Here G_B denotes the subgraph $(V, c^{-1}(B))$ for any subset B of Δ_4 .

If $\text{Sd } K$ is the first barycentric subdivision of K , let $H(i, j)$ be the largest subcomplex of $\text{Sd } K$ disjoint from $\text{Sd } K(i, j) \cup \text{Sd } K(r, s, t)$. Then the polyhedron $|H(i, j)|$ is a closed orientable 3-manifold which splits M into two complementary 4-manifolds $N(i, j)$ and $N(r, s, t)$ with common boundary. Further $N(i, j)$ and $N(r, s, t)$ are regular neighborhoods in M of $|\text{Sd } K(i, j)|$ and $|\text{Sd } K(r, s, t)|$ respectively.

Following [2], we can always assume that (G, c) regularly imbeds into the closed orientable surface of genus g and of Euler characteristic $2 - 2g = g_{01} + g_{12} + g_{23} + g_{34} + g_{40} - 3p$, where p is the order of G divided by 2.

As proved in [2], we have

$$\begin{aligned} (1) \quad g_{013} &= 1 + g - g_2 - g_4, & (6) \quad g_{14} &= g_{014} + g - g_0, \\ (2) \quad g_{023} &= 1 + g - g_1 - g_4, & (7) \quad g_{02} &= g_{012} + g - g_1, \\ (3) \quad g_{024} &= 1 + g - g_1 - g_3, & (8) \quad g_{13} &= g_{123} + g - g_2, \\ (4) \quad g_{124} &= 1 + g - g_0 - g_3, & (9) \quad g_{24} &= g_{234} + g - g_3, \\ (5) \quad g_{134} &= 1 + g - g_0 - g_2, & (10) \quad g_{03} &= g_{034} + g - g_4, \end{aligned}$$

$$(11) \quad \chi(M) = 2 - 2g + \sum_i g_i$$

where g_i ($0 \leq g_i \leq g$) is the genus of an orientable closed surface into which the subgraph G_i ($i \in \Delta_4$) regularly imbeds and $\chi(M)$ is the Euler characteristic of M .

If $g = 2$, then the sum

$$R = g_{013} + g_{023} + g_{024} + g_{124} + g_{134} = 5 + 5g - 2 \sum_i g_i$$

belongs to the set $\{5, 7, 9, 11, 13, 15\}$ since R ($R \geq 5$) is odd.

Now we show that the cases $R \in \{9, 13, 7, 11\}$ give a contradiction, while the cases $R = 5$ and $R = 15$ imply that M is (PL) homeomorphic to the complex projective plane and the connected sum $\#_2(S^1 \times S^3)$ respectively

Case $R = 9$. If $R = 9$, then $\sum_i g_i = 3$ and the manifold M is simply-connected because at least one of the above g_{ijk} 's equals 1 (see [11] for the generators of the fundamental group of M). This implies that $\chi(M) = 2 + b_2(M) \geq 2$, i.e. a contradiction since $\chi(M) = 1$ by (11).

Case $R = 13$. If $R = 13$, then $\sum_i g_i = 1$ and $\chi(M) = -1$. The fundamental group of M is trivial or cyclic since at least one of the above g_{ijk} 's must be ≤ 2 (see [11]). The inequality $b_1(M) \leq 1$ implies that $\chi(M) = 2 - 2b_1(M) + b_2(M) \geq b_2(M) \geq 0$, i.e. a contradiction.

Case $R = 7$. If $R = 7$, then $\sum_i g_i = 4$ and $\chi(M) = 2$. Because at least one of the above g_{ijk} 's equals 1, the manifold M is simply-connected, whence $\chi(M) = 2$ implies that $b_2(M) = 0$.

Now the addendum of the sum R may assume the values listed in Table 1.

The fifteen cases to verify can be reduced to three by cyclic permutations in the color set. In fact, doing this type of change of names in the color set Δ_4 the permutation of Δ_4 giving regular imbedding of G does not change. Thus we can only examine the cases 7.1, 7.6 and 7.7.

(n. 7.1) If $g_{023} = g_{024} = g_{124} = g_{134} = 1$ and $g_{013} = 3$, then the relations (1), ..., (5) imply that $g_0 = g_1 = 2$ and $g_2 = g_3 = g_4 = 0$. By (6), ..., (10) we obtain $g_{14} = g_{014}$, $g_{02} = g_{012}$, $g_{13} = g_{123} + 2$, $g_{24} = g_{234} + 2$ and $g_{03} = g_{034} + 2$. Since $g_{134} = 1$, then $K(0, 2)$ consists of exactly one edge, hence $N(0, 2)$ is a 4-ball. Furthermore $K(1, 3)$ and $K(1, 4)$ are also formed by one edge, each one as $g_{024} = g_{023} = 1$. Thus all triangles of $K(1, 3, 4)$ have two edges of $K(1, 3)$ and $K(1, 4)$ in common. The other edge of each triangle of $K(1, 3, 4)$ is free since $g_{02} = g_{012}$. Therefore $K(1, 3, 4)$ is a cone which collapses to a point, i.e. $N(1, 3, 4)$ is a 4-ball. This implies that M is (PL) homeomorphic to the 4-sphere S^4 which is a contradiction since $g(S^4) = 0$.

(n. 7.6) If $g_{013} = g_{023} = 2$ and $g_{024} = g_{124} = g_{134} = 1$, then $g_0 = g_1 = g_2 = g_3 = 1$, $g_4 = 0$, $g_{14} = g_{014} + 1$, $g_{02} = g_{012} + 1$, $g_{13} = g_{123} + 1$, $g_{24} = g_{234} + 1$ and $g_{03} = g_{034} + 2$. The manifold $N(1, 3)$ is a 4-ball since $g_{024} = 1$. Further $K(0, 2)$ (resp. $K(2, 4)$) consists of exactly one (resp. two) edge since $g_{134} = 1$ (resp. $g_{013} = 2$). Because $g_{13} = g_{123} + 1$, the pseudocomplex $K(0, 2, 4)$ contains many triangles but one as there are edges in $K(0, 4)$. The two triangles of $K(0, 2, 4)$, which have two edges $\langle v_2, v_4 \rangle$ and $\langle v_0, v_4 \rangle$ in common, cannot have the same boundary since $FH_2(M) \simeq 0$ and $H_2(N(1, 3)) \simeq 0$ imply that $H_2(N(0, 2, 4)) \simeq 0$ (use the Mayer-Vietoris sequence of the pair $(N(1, 3), N(0, 2, 4))$). Thus $K(0, 2, 4)$ collapses to a point, i.e. $N(0, 2, 4)$ is (PL) homeomorphic to a 4-ball. This gives a contradiction since $M \simeq_{\text{PL}} S^4$ (4-sphere) and $g(M) = 2$.

(n. 7.7) If $g_{013} = g_{024} = 2$ and $g_{023} = g_{124} = g_{134} = 1$, then we have $g_0 = 2$, $g_2 = g_3 = 0$, $g_1 = g_4 = 1$, $g_{14} = g_{014}$, $g_{02} = g_{012} = 1$, $g_{03} = g_{034} = 1$,

TABLE 1

n.	g_{013}	g_{023}	g_{024}	g_{124}	g_{134}
7.1	3	1	1	1	1
7.2	1	3	1	1	1
7.3	1	1	3	1	1
7.4	1	1	1	3	1
7.5	1	1	1	1	3
7.6	2	2	1	1	1
7.7	2	1	2	1	1
7.8	2	1	1	2	1
7.9	2	1	1	1	2
7.10	1	2	2	1	1
7.11	1	2	1	2	1
7.12	1	2	1	1	2
7.13	1	1	2	2	1
7.14	1	1	2	1	2
7.15	1	1	1	2	2

$g_{13} = g_{123} + 2$ and $g_{24} = g_{234} + 2$. Because $g_{023} + 1$ and $(g_{124} = g_{134} + 1, g_{14} = g_{014})$, the contradiction follows as in the case n. 7.1 by using the pair $(N(1, 4), N(0, 2, 3))$.

Case $R = 11$. If $R = 11$, then $\sum_i g_i = 2$ and $\chi(M) = 0$. The manifold M cannot be simply-connected since $\chi(M) = 0$. Thus we have $g_{ijk} \geq 2$ for any i, j, k , whence the addendum of R may assume the values listed in Table 2.

TABLE 2

n.	g_{013}	g_{023}	g_{024}	g_{124}	g_{134}
11.1	2	2	2	2	3
11.2	2	2	2	3	2
11.3	2	2	3	2	2
11.4	2	3	2	2	2
11.5	3	2	2	2	2

All these cases are equivalent to one by the same reason that $R = 7$.

The fundamental group of M is cyclic and nontrivial since $b_1(M) \neq 0$ and at least one of the above g_{ijk} 's equals 2. This implies that $H_1(M) \simeq H_3(M) \simeq \mathbb{Z}$ and $FH_2(M) \simeq 0$ (use $\chi(M) = 0$).

(n. 11.1) If $g_{013} = g_{023} = g_{024} = g_{124} = 2$ and $g_{134} = 3$, then we have $g_0 = g_1 = g_2 = 0$, $g_3 = g_4 = 1$, $g_{03} = g_{034} + 1$, $g_{24} = g_{234} + 1$, $g_{13} = g_{123} + 2$, $g_{02} = g_{012} + 2$ and $g_{14} = g_{014} + 2$ (use (1), ..., (10)). Since $g_{013} = 2$, then $K(2, 4)$ is formed by two vertices joined by exactly two edges, whence $N(2, 4)$ is (PL) homeomorphic to $S^1 \times B^3$. Here B^3 represents a (closed) 3-ball. Furthermore $K(1, 3)$ and $K(0, 3)$ consist of exactly two edges each one as $g_{124} = g_{024} = 2$. Because $g_{24} = g_{234} + 1$, the pseudocomplex $K(0, 1, 3)$ contains many triangles but one as there are edges in $K(0, 1)$. However any two triangles of $K(0, 1, 3)$ cannot have the same boundary since $FH_2(M) \simeq 0$ and $H_2(N(2, 4)) \simeq 0$ imply that $H_2(N(0, 1, 3)) \simeq 0$ (use the Mayer-Vietoris sequence of the pair $(N(2, 4), N(0, 1, 3))$). Thus $K(0, 1, 3)$ collapses to a circle, whence $N(0, 1, 3)$ is (PL) homeomorphic to $S^1 \times B^3$. By [16, Theorem 2], we have that $M \simeq_{\text{PL}} S^1 \times S^3$ which is a contradiction since $g(S^1 \times S^3) = 1$ by [7].

Case $r = 15$: The connected sum. $\#_2(S^1 \times S^3)$. If $R = 15$, then it follows that $g_{013} = g_{023} = g_{024} = g_{124} = g_{134} = 3$ and $g_i = 0$ for each $i \in \Delta_4$. The rank of $\Pi_1(M)$ is ≤ 2 as each one of the above g_{ijk} 's equals 3. This implies that $b_1(M) = b_3(M) = 2$ and $b_2(M) = 0$ since $\chi(M) = -2$. Because $g_{024} = 3$, the pseudocomplex $K(1, 3)$ consists of exactly three edges, hence $N(1, 3)$ is (PL) homeomorphic to the connected sum $\#_2(S^1 \times B^3)$. Furthermore $K(0, 2)$ and $K(2, 4)$ are also formed by three edges each one as $g_{134} = g_{013} = 3$. Because $g_{13} = g_{123} + 2$ (use (8)), the complex $K(0, 4)$ has many edges but two as there are triangles in $K(0, 2, 4)$. Since $H_1(M) \simeq H_3(M) \simeq \mathbb{Z} \oplus \mathbb{Z}$, $FH_2(M) \simeq 0$ and $H_2(N(1, 3)) \simeq 0$, the Mayer-Vietoris sequence of the pair $(N(1, 3), N(0, 2, 4))$ gives

$$0 \rightarrow \mathbb{Z} \oplus \mathbb{Z} \rightarrow \mathbb{Z} \oplus \mathbb{Z} \rightarrow H_2(N(0, 2, 4)) \rightarrow 0,$$

whence $H_2(N(0, 2, 4)) \simeq 0$, i.e. there are no two triangles of $K(0, 2, 4)$ with common boundary. Now it is very easy to see that the pseudocomplex $K(0, 2, 4)$ collapses to a one-dimensional subcomplex formed by two vertices joined by three edges. Therefore $N(0, 2, 4)$ is (PL) homeomorphic to the connected sum $\#_2(S^1 \times B^3)$.

Thus $M \simeq_{\text{PL}} \#_2(S^1 \times S^3)$ by Theorem 2 of [16]. Now the result in Proposition 1 follows as $g(\#_2(S^1 \times S^3)) = 2$ by [7] and the subadditivity of the genus.

Case $R = 5$: The projective complex plane. If $R = 5$, then $g_{013} = g_{023} = g_{024} = g_{124} = g_{134} = 1$ and $\chi(M) = 3$. By (1), ..., (5) it follows that $g_i = 1$ for each $i \in \Delta_4$. In fact, suppose on the contrary that $g_i = 2$ (recall that $g_i \leq 2$). Then we have $g_0 = g_2 = 2$ and $g_3 = g_4 = 0$. This is a contradiction since $\sum_i g_i = 5$.

Furthermore, if $g_i = 0$, then $g_0 = g_2 = 0$ and $g_3 = g_4 = 2$ which give a contradiction as shown above.

Therefore we must have $g_i = 1$, whence $g_0 = g_2 = g_3 = g_4 = 1$. By

(6), ..., (10) the relations $g_{14} = g_{014} + 1$, $g_{02} = g_{012} + 1$, $g_{13} = g_{123} + 1$, $g_{24} = g_{234} + 1$, and $g_{03} = g_{034} + 1$ hold. Since $g_{024} = 1$, then $K(1, 3)$ consists of exactly one edge, hence $N(1, 3)$ is a 4-ball. Further $K(0, 2)$ and $K(2, 4)$ are also formed by one edge each one since $g_{134} = g_{013} = 1$. Thus all triangles of $K(0, 2, 4)$ have two edges in common. Because $g_{13} = g_{123} + 1$, the complex $K(0, 2, 4)$ has many triangles but one as there are edges in $K(0, 4)$.

Therefore $K(0, 2, 4)$ collapses to a combinatorial 2-sphere, formed by exactly two triangles T_1, T_2 of $K(0, 2, 4)$, with common boundary. Then we have that $H_0(N(0, 2, 4)) \simeq H_2(N(0, 2, 4)) \simeq \mathbb{Z}$ and $H_1(N(0, 2, 4)) \simeq H_3(N(0, 2, 4)) \simeq 0$. By isotopy we can always suppose that T_1 is the standard 2-simplex in M . Let $\hat{T}_1$ be the barycenter of T_1 and $\text{Sd}^2 K$ be the second barycentric subdivision of $K = K(G)$. Then $N(0, 2, 4)$ is the orientable bordered 4-manifold obtained by adding a 2-handle (a regular neighborhood of $\hat{T}_1$ in $\text{Sd}^2 K$) onto the boundary of a 4-ball (a small regular neighborhood of T_2 in M) along a knot N . Since the surgery is given by attaching 2-handles in dimension 4, the surgery coefficient associated to N must be an integer and by homological reasons equal to ± 1 . Therefore $\partial N(0, 2, 4) = S^3$ so by Gordon and Luecke¹, Theorem 2, N is the trivial knot and the manifold $N(0, 2, 4)$ is (PL) homeomorphic to $\pm CP^2 - (4\text{-ball})$, whence M is the projective complex plane as required. Now the proof is completed because a crystallization of CP^2 with genus 2 is really constructed in [12].

3. PROOF OF PROPOSITION 1: THE NONORIENTABLE CASE

Let M be a closed smooth (or PL) nonorientable connected 4-manifold of genus h . As proved in the orientable case, we can obtain the following relations:

$$\begin{aligned} g_{013} &= 1 + h/2 - g_2 - g_4, & g_{14} &= g_{014} + h/2 - g_0, \\ g_{023} &= 1 + h/2 - g_1 - g_4, & g_{02} &= g_{012} + h/2 - g_1, \\ g_{024} &= 1 + h/2 - g_1 - g_3, & g_{13} &= g_{123} + h/2 - g_2, \\ g_{124} &= 1 + h/2 - g_0 - g_3, & g_{24} &= g_{234} + h/2 - g_3, \\ g_{134} &= 1 + h/2 - g_0 - g_2, & g_{03} &= g_{034} + h/2 - g_4, \\ \chi(M) &= 2 - h + \sum_i g_i. \end{aligned}$$

(Statement 1.b). If $h = 2$, then the sum

$$R = g_{013} + g_{023} + g_{024} + g_{124} + g_{134} = 10 - 2 \sum_i g_i$$

is even, hence R belongs to the set $\{6, 8, 10\}$ as $R \geq 6$ and $g_i \geq 0$. Now we show that the cases $R = 6$ and $R = 8$ give a contradiction, while the case $R = 10$ implies that the manifold M is (PL) homeomorphic to the "twisted" S^3 bundle over S^1 , i.e. $M \simeq_{\text{PL}} S^1 \times S^3$. If $R = 6$ or $R = 8$, then at least one of the above g_{ijk} 's equals 1. Thus the fundamental group of M is trivial so that M is orientable, i.e. a contradiction.

If $R = 10$, then we have $\chi(M) = 0$ and $g_i = 0$ for each $i \in \Delta_4$. Furthermore the relations $g_{124} = g_{013} = g_{023} = g_{024} = g_{134} = 2$ imply that

¹C. McA. Gordon and J. Luecke, *Knots are determined by their complements*, Bull. Amer. Math. Soc. (N.S.) **20** (1989), 83–89.

$\Pi_1(M) \simeq H_1(M) \simeq Z$. Since $g_{024} = 2$, then $K(1, 3)$ consists of two vertices joined by two edges, hence $N(1, 3) \simeq_{\text{PL}} (S^1 \times B^3)$ or $S^1 \times \tilde{B}^3$ (the “twisted” B^3 bundle over S^1) and $\partial N(1, 3) \simeq_{\text{PL}} S^1 \times S^2$ or $S^1 \times \tilde{S}^2$ respectively. The Mayer-Vietoris sequence of the pair $(N(1, 3), N(0, 2, 4))$ gives

$$\begin{aligned} 0 \rightarrow FH_3(M) \rightarrow FH_2(\partial N(1, 3)) \simeq Z \rightarrow H_2(N(0, 2, 4)) \\ \rightarrow H_2(M) \rightarrow Z \rightarrow Z \oplus H_1(N(0, 2, 4)) \rightarrow Z \rightarrow 0, \end{aligned}$$

whence $H_2(N(0, 2, 4)) \simeq 0$ since $FH_3(M) \simeq Z$ and $FH_2(M) \simeq H_2(M) \simeq 0$ (note that $H_1(N(0, 2, 4)) \simeq Z$). In this case the relations $g_{14} = g_{014} + 1$, $g_{02} = g_{012} + 1$, $g_{13} = g_{123} + 1$, $g_{24} = g_{234} + 1$ and $g_{03} = g_{034} + 1$ hold too. Now the formulae $g_{013} = g_{134} = 2$ and $g_{13} = g_{123} + 1$ imply that $K(0, 2, 4)$ collapses to a circle since $H_2(K(0, 2, 4)) \simeq 0$. Therefore the manifold $N(0, 2, 4)$ is (PL) homeomorphic to either $S^1 \times B^3$ or $S^1 \times \tilde{B}^3$. Since M is nonorientable, it follows that $M \simeq_{\text{PL}} S^1 \times \tilde{S}^3$ by Theorem 2 of [16]. Now the proof is completed because a crystallization of $S^1 \times \tilde{S}^3$ with genus 2 is shown in [13, Figure 1, p. 155].

(Statement 1.c) By the “subadditivity” of the genus we have that

$$g\left(\#_2(S^1 \times \tilde{S}^3)\right) \leq 2g(S^1 \times \tilde{S}^3) = 4$$

and $g(S^1 \times S^3 \# S^1 \times \tilde{S}^3) \leq 2g(S^1 \times S^3) + g(S^1 \times \tilde{S}^3) = 4$. If $h = 4$, then the sum $R = 15 - 2 \sum_i g_i$ is odd, whence $R \in \{5, 7, 9, 11, 13, 15\}$. We show that the cases $R \in \{5, 7, 9, 11, 13\}$ give a contradiction while the case $R = 15$ implies the statement (c) in Proposition 1.

If $R \in \{5, 7, 9\}$, then at least one of the g_{ijk} ’s in R must be one, so that $\Pi_1(M) = 0$ and M is orientable, i.e. a contradiction.

If $R = 11$, then $\sum_i g_i = 2$ and $\chi(M) = 0$. Thus the addendum of R may assume the values listed in the table n. 11 (§2). Since these cases are equivalent to one, we can suppose that $g_{013} = g_{023} = g_{024} = g_{124} = 2$ and $g_{134} = 3$. As in the case n. 11.1 (§2), the formulae $g_0 = g_1 = g_2 = 0$, $g_3 = g_4 = 1$, $g_{03} = g_{034} + 1$, $g_{24} = g_{234} + 1$, $g_{13} = g_{123} + 2$, $g_{02} = g_{012} + 2$ and $g_{14} = g_{014} + 2$ hold too. Since $g_{013} = 2$, then $\text{rk } \Pi_1(M) \leq 1$, whence $b_1(M; Z_2) \leq 1$ and $\chi(M) = 0 = 2 - 2b_1(M; Z_2) + b_2(M; Z_2)$ implies that $H_2(M; Z_2) = 0$. Then by the same arguments used in the proof of n. 11.1 (§2), we have that $N(1, 4)$ and $N(0, 2, 3)$ are both (PL) homeomorphic to either $S^1 \times B^3$ or $S^1 \times \tilde{B}^3$.

Since M is nonorientable, it follows that $M \simeq_{\text{PL}} S^1 \times \tilde{S}^3$ by Theorem 2 of [16]. This is a contradiction because $g(S^1 \times \tilde{S}^3) = 2$.

If $R = 13$, then $\sum_i g_i = 1$ and $\chi(M) = -1$. Since the manifold M cannot be simply-connected, at least one of g_{ijk} ’s in R equals 2. Thus we have $\text{rk } \Pi_1(M) \leq 1$, $b_1(M; Z_2) \leq 1$ and $\chi(M) = -1 = 2 - 2b_1(M; Z_2) + b_2(M; Z_2) \geq 0$, which is a contradiction.

If $R = 15$, then $g_{013} = g_{023} = g_{024} = g_{124} = g_{134} = 3$ and $g_i = 0$ for each $i \in \Delta_4$. The rank of $\Pi_1(M)$ is ≤ 2 as each one of the above g_{ijk} ’s equals 3. This implies that $H_1(M; Z_2) = H_3(M; Z_2) = Z_2 \oplus Z_2$ and $H_2(M; Z_2) = 0$ since $\chi(M) = -2$. Then by the same arguments used in the case $R = 15$ (§2),

it follows that $N(1, 3)$ and $N(0, 2, 4)$ are both (PL) homeomorphic to either $\#_2(S^1 \times B^3)$ or $S^1 \times B^3 \# S^1 \times B^3$. Note that the case $\#_2(S^1 \times B^3)$ can be avoided since M is nonorientable. Thus we have $M \simeq_{\text{PL}} \#_2(S^1 \times S^3)$ or $M \simeq_{\text{PL}} S^1 \times S^3 \# S^1 \times S^3$ as required.

Now Corollary 1 directly follows from Proposition 1 and the “subadditivity” of the genus (see §1). Furthermore there exists a crystallization of RP^4 with genus six as shown in [11].

4. THE GENERAL CASE

Proof of Proposition 2. The sum

$$R = g_{134} + g_{124} + g_{024} + g_{023} + g_{013} = 5 + 5g - 2 \sum_i g_i$$

is odd (resp. even) whenever g is even (resp. odd). Therefore it follows that

$$0 \leq \sum_i g_i \leq [(5/2)g],$$

whence

$$\begin{aligned} 2 - 2g + b_2(M) &\leq 2 - 2b_1(M) + b_2(M) \\ &= \chi(M) = 2 - 2g + \sum_i g_i \leq 2 - 2g + [(5/2)g] \end{aligned}$$

since $b_1(M) \leq g$ (see [1]). If M is simply-connected, then the inequality

$$2 + b_2(M) = \chi(M) \leq 2 - 2g + [(5/2)g]$$

implies that $b_2(M) \leq [g/2]$ as required. $\square$

Proof of Corollary 2. (1) Use $g(S^1 \times S^3) = 1$ (see [2]), the subadditivity of the genus and

$$g\left(\#_k(S^1 \times S^3)\right) \geq \text{rank } \Pi_1\left(\#_k(S^1 \times S^3)\right) = k$$

(see [1]).

(2) Since CP^2 is simply-connected, we have

$$k = b_2\left(\#_k \pm CP^2\right) \leq [g/2],$$

whence $g \geq 2k$. Now the relation $g(CP^2) = 2$ (see [2]) and the subadditivity of the genus prove the statement.

(3) Since $\#_k(S^1 \times S^2)$ is simply-connected, we have

$$2k = b_2\left(\#_k(S^2 \times S^2)\right) \leq [g/2],$$

whence $g \geq 4k$. Then the statement follows by $g(S^2 \times S^2) \leq 4$ (use the crystallization shown in [7]) and the subadditivity of the genus. $\square$

Proof of Proposition 3. Since $r \leq [g/2] \leq [31/2] = 15$, the intersection form ω_M , induced on M by the cup product, has rank ≤ 15 . Recall that ω_M may only be either odd or indefinite even whenever M is smooth. In the last case the

signature $\sigma(M)$ of M is divisible by 16 (see [3, 4, 9] and the Rohlin theorem). Furthermore ω_M is equivalent (over the integers) to either $(\pm 1) \oplus \cdots \oplus (\pm 1)$ (r times) or a $E_8 \oplus b \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ where a is even, $b \neq 0$ and $a, b \in \mathbb{Z}$ (see [3, 4]). Since the rank of a $E_8 \oplus b \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ is $8|a| + 2|b|$, it follows that $a = 0$, (use $\text{rank } \omega_M \leq 15$ and a even). Now the Freedman classification of simply-connected smooth 4-manifolds implies the statement (see [9]). $\square$

Corollary 3. (1) *Let M^4 be a smooth homotopy 4-sphere. Then there exists an integer k such that*

$$g \left(M \# \#_k (S^2 \times S^2) \right) = 4k.$$

(2) *If V_n is an algebraic nonsingular hypersurface of degree n in CP^3 , then $n^3 - 4n^2 + 6n - 2 \leq [g/2]$.*

(3) *If V is a simply-connected complex surface, then $12p_g + 10 - c_1^2[V] \leq [g/2]$.*

If V is minimal elliptic, then $12p_g + 10 \leq [g/2]$.

Here p_g and $c_1[V]$ denote the geometric genus and the first Chern class of V respectively.

Proof. (1) If M is a homotopy 4-sphere, then there exists an integer k such that $M \# \#_k (S^2 \times S^2) \simeq_{\text{DIFF}} \#_k (S^2 \times S^2)$. Now use Corollary 2.

(2) Recall that $b_2(V_n) = n^3 - 4n^2 + 6n - 2$ (see [15]). Now use Proposition 2 as V_n is simply-connected.

(3) It is well known that $b_2(V) = 12p_g + 10 - c_1^2[V]$ (see [15]). Furthermore we have that $c_1^2[V] = 0$ for the minimal elliptic case. Now we use Proposition 2. $\square$

5. RELATIONS WITH THE POINCARÉ CONJECTURE

Let $P(4)$ be the Poincaré conjecture of 4-dimension. Now we state some conjectures which are related with $P(4)$.

Conjecture $C(1)$. If M is a closed smooth simply-connected 4-manifold, then $g(M) = 2b_2(M)$.

Conjecture $C(2)$. The genus is additive with respect to the connected sum of simply-connected smooth (or PL) 4-manifolds.

The conjecture $C(2)$ and its relation with $P(4)$ are also established in [8]. We prove that $C(1)$ and $C(2)$ are equivalent:

$C(1) \Rightarrow C(2)$. Let M_1, M_2 be closed smooth (or PL) simply-connected 4-manifolds. Then we have $\frac{1}{2}g(M_1 \# M_2) = b_2(M_1 \# M_2) = b_2(M_1) + b_2(M_2) = \frac{1}{2}g(M_1) + \frac{1}{2}g(M_2)$, whence $g(M_1 \# M_2) = g(M_1) + g(M_2)$.

$C(2) \Rightarrow C(1)$. If M is a closed smooth (or PL) simply-connected 4-manifold, then there exist two integers $p, q \in \mathbb{Z}$ such that

$$M \# \#_p CP^2 \# \#_q (-CP^2) \simeq_{\text{DIFF}} \#_a CP^2 \# \#_b (-CP^2)$$

where

$$a = p + \frac{1}{2}(b_2(M) + \sigma(M)),$$

$$b = q + \frac{1}{2}(b_2(M) - \sigma(M)),$$

$$\sigma(M) = \text{signature of } M,$$

(see [15]). Thus we have

$$\begin{aligned} g\left(M \#_p \#_q CP^2 \#_q (-CP^2)\right) &= g\left(\#_a CP^2 \#_b (-CP^2)\right) = 2(a+b) \\ &= 2(p+q) + 2b_2(M) = g(M) + g\left(\#_p CP^2 \#_q (-CP^2)\right) \\ &= g(M) + 2(p+q), \end{aligned}$$

whence $g(M) = 2b_2(M)$ as required. $\square$

Obviously $C(1)$ (or $C(2)$) implies $P(4)$ as follows. Let M be a closed smooth (or PL) homotopy 4-sphere. Since $b_2(M) = 0$, we have $g(M) = 2b_2(M) = 0$, whence $M \simeq_{\text{PL}} S^4$ (4-sphere).

We also have the following

Conjecture $C(3)$. If M is a smooth (or PL) homotopy 4-sphere, then

$$g((M) \#_k (S^2 \times S^2)) \geq g(M) + 4k$$

for any integer k .

We prove that $P(4)$ and $C(3)$ are equivalent.

$C(3) \Rightarrow P(4)$. If M is a homotopy 4-sphere, then there exists an integer k (see Corollary 3) such that $4k = g(M \#_k (S^2 \times S^2)) \geq g(M) + 4k$, whence $g(M) = 0$ and $M \simeq_{\text{PL}} S^4$.

$P(4) \Rightarrow C(3)$. If M is a smooth homotopy 4-sphere, then $M \simeq_{\text{PL}} S^4$, whence

$$g\left(M \#_k (S^2 \times S^2)\right) = g\left(\#_k (S^2 \times S^2)\right) = 4k \geq g(M) + 4k$$

since $g(M) = 0$.

We conclude the paper by noting that it is an open question whether the number of closed smooth (or PL) connected 4-manifolds of a fixed genus g is finite or not.

REFERENCES

1. J. Bracho and L. Montejano, *The combinatorics of colored triangulations of manifolds*, Geom. Dedicata **22** (1987), 303–328.
2. A. Cavicchioli, *A combinatorial characterization of $S^3 \times S^1$ among closed 4-manifolds*, Proc. Amer. Math. Soc. **105** (1989), 1008–1014.
3. S. K. Donaldson, *An application of gauge theory to four-dimensional topology*, J. Differential Geom. **18** (1983), 279–315.
4. —, *Connections, cohomology and the intersection forms of 4-manifolds*, J. Differential Geom. **24** (1986), 275–341.
5. M. Ferri, *Una rappresentazione delle n -varietà topologiche triangolabili mediante grafi $(n+1)$ -colorati*, Boll. Un. Mat. Ital. **13-B** (1976), 250–260.
6. M. Ferri and C. Gagliardi, *The only genus zero n -manifold is S^n* , Proc. Amer. Math. Soc. **85** (1982), 638–642.
7. —, *On the genus of 4-dimensional products of manifolds*, Geom. Dedicata **13** (1982), 331–345.
8. M. Ferri, C. Gagliardi, and L. Grasselli, *A graph-theoretical representation of PL-manifolds — A survey on crystallizations*, Aequationes Math. **31** (1986), 121–141.

9. M. H. Freedman, *The topology of four-dimensional manifolds*, J. Differential Geom. **17** (1982), 357–453.
10. C. Gagliardi, *Extending the concept of genus to dimension n* , Proc. Amer. Math. Soc. **81** (1981), 473–481.
11. —, *How to deduce the fundamental group of a closed n -manifold from a contracted triangulation*, J. Combin. Informat. Systems Sci. **4** (1979), 237–252.
12. —, *On the genus of CP^2* , Aequationes Math. **37** (1989), 130–140.
13. C. Gagliardi and G. Volzone, *Handles in graphs and sphere bundles over S^1* , European J. Combin. **8** (1987), 151–158.
14. P. J. Hilton and S. Wylie, *An introduction to algebraic topology—Homology theory*, Cambridge Univ. Press, Cambridge, 1960.
15. R. Mandelbaum, *Four-dimensional topology: an introduction*, Bull. Amer. Math. Soc. (N.S.) **2** (1980), 1–159.
16. J. M. Montesinos, *Heegaard diagrams for closed 4-manifolds*, Geometric Topology (J. Cantrell, Ed.), Proc. 1977 Georgia Conference, Academic Press, New York, 1979, pp. 219–237.
17. M. Pezzana, *Sulla struttura topologica delle varietà compatte*, Atti Sem. Mat. Fis. Univ. Modena **23** (1974), 269–277.
18. D. Rolfsen, *Knots and links*, Math. Lecture Ser. 7, Publish or Perish, Berkeley, Calif., 1976.
19. A. T. White, *Graphs, groups and surface*, North-Holland, Amsterdam, 1973.
20. J. H. C. Whitehead, *Manifolds with transverse fields in euclidean space*, Ann. of Math. (2) **73** (1961), 154–211.

ISTITUTO DI MATEMATICA, UNIVERSITÀ DELLA BASILICATA, VIA NAZARIO SAURO N. 85, 85100 POTENZA, ITALIA

Current address: Dipartimento di Matematica, Università di Modena, via Campi 213/B, 41100 Modena, Italia