

THE RECTIFIABLE METRIC ON THE SET OF CLOSED SUBSPACES OF HILBERT SPACE

LAWRENCE G. BROWN

ABSTRACT. Consider the set of selfadjoint projections on a fixed Hilbert space. It is well known that the connected components, under the norm topology, are the sets $\{p: \text{rank } p = \alpha, \text{rank}(1 - p) = \beta\}$, where α and β are appropriate cardinal numbers. On a given component, instead of using the metric induced by the norm, we can use the rectifiable metric d_r which is defined in terms of the lengths of rectifiable paths or, equivalently in this case, the lengths of ε -chains. If $\|p - q\| < 1$, then $d_r(p, q) = \sin^{-1}(\|p - q\|)$, but if $\|p - q\| = 1$, $d_r(p, q)$ can have any value in $[\frac{\pi}{2}, \pi]$ (assuming α and β are infinite). If $d_r(p, q) \neq \frac{\pi}{2}$, a minimizing path joining p and q exists; but if $d_r(p, q) = \frac{\pi}{2}$, a minimizing path exists if and only if $\text{rank}(p \wedge (1 - q)) = \text{rank}(q \wedge (1 - p))$.

1. INTRODUCTION

Let H be a Hilbert space, possibly nonseparable, and let \mathcal{P} be the set of (selfadjoint) projections on H . For p, q in \mathcal{P} , let $d_n(p, q) = \|p - q\|$, where “ $\| \cdot \|$ ” is operator norm. Since $\|p - q\| \leq 1$, we can also define $d_a(p, q) = \sin^{-1}(\|p - q\|)$. So d_n is the metric induced by the norm, and d_a is the “angular metric.” The fact that d_a is a metric will be proved later (possibly this is folklore).

If α and β are cardinal numbers such that $\alpha + \beta = \dim H$ (orthogonal dimension), let $C_{\alpha, \beta} = \{p \in \mathcal{P}: \text{rank } p = \alpha, \text{rank}(1 - p) = \beta\}$. Then the various $C_{\alpha, \beta}$ ’s are the connected components of \mathcal{P} . It is well known that $\|p - q\| < 1$ implies that p and q are in the same component.

Some of our study will rely on the availability of a classification, up to unitary equivalence, of pairs of projections. This was first given by Dixmier [6] and Kreĭn, Krasnosel’skiĭ, and Mil’man [9], independently. The note on p. 18 of [5] and the introduction to [16] give other historical information and references for this subject. Some other references are [4, 8, 10, §3, and 12].

If p and q are projections with ranges M and N , let $H_{11} = M \cap N$, $H_{10} = M \cap N^\perp$, $H_{01} = M^\perp \cap N$, $H_{00} = M^\perp \cap N^\perp$ and $H_0 = (H_{11} \oplus H_{10} \oplus H_{01} \oplus H_{00})^\perp$. If $H_0 = H$, M and N are said to be in *generic position* ([8], Dixmier uses the term “position p ”). We will also refer to H_0 as the generic part of H , $M \cap H_0$ as the generic part of M , etc.

The basic example of two subspaces in generic position occurs when $\dim H =$

Received by the editors September 10, 1990 and, in revised form, February 1, 1991.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 46C99, 47A99, 47C15.

Key words and phrases. Hilbert space, projection, rectifiable.

2, $\dim M = \dim N = 1$, and M and N have angle θ , where $0 < \theta < \frac{\pi}{2}$. The most general generic pair is a direct integral of such two-dimensional examples, for various values of θ . One way to describe an arbitrary pair of projections more explicitly is as follows:

It is possible to identify both $H_0 \cap M$ and $H_0 \cap M^\perp$, with $L^2(X)$, for some measure space X , in such a way that the generic parts of p and q are given by

$$p_0 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad q_0 = \begin{bmatrix} \cos^2 \varphi & \cos \varphi \sin \varphi \\ \cos \varphi \sin \varphi & \sin^2 \varphi \end{bmatrix},$$

where φ is a measurable function on X such that $0 < \varphi(x) < \frac{\pi}{2}$, $\forall x$. Here p_0 and q_0 operate on $L^2(X) \oplus L^2(X)$ and the matrices are operator matrices whose entries are multiplication operators.

In Theorem 12 below we use another method of describing generic pairs of subspaces, given by Halmos [8]. He shows that N is the graph of a suitable (in general unbounded) operator $T: M \rightarrow M^\perp$. By using the polar decomposition of T , we can identify M with M^\perp in such a way that T is positive and selfadjoint (this is part of Halmos' argument). In the description given above T is the multiplication operator on $L^2(X)$ induced by $\tan \varphi$. The idea of representing N as a graph can actually be used in some nongeneric cases also. It can be used whenever $H_{01} = 0$, provided we weaken the requirements for T sufficiently.

When we use the classification theory for pairs of projections below, we will often refer to the "canonical representation of the pair (p, q) ", though the use of the term "canonical" is not strictly correct. The pair consisting of the Hilbert space $L^2(X)$ and the multiplication operator induced by φ is uniquely determined up to unitary equivalence, but X itself is not uniquely determined.

We now discuss the basic concepts relating to rectifiable metrics. This discussion will be oversimplified because certain difficulties which could arise in general do not occur in \mathcal{P} . Further information on this is contained in Chapter 1, §B of [7] and Chapters 3, 4 of [13]. Let (X, d) be a metric space. The concepts of rectifiable path and arc length of a path are defined for paths in X in exactly the same way as for paths in \mathbb{R}^n . If two points x and y can be joined by a rectifiable path, we define $d_r(x, y)$ to be the infimum of the lengths of rectifiable paths joining x and y . Clearly d_r is a metric if it is everywhere defined, and $d_r \geq d$. The metric d could be called "rectifiable" if $d_r = d$. Any d -rectifiable path is also d_r -rectifiable and has the same arc lengths relative to d_r and d . (Thus d_r , if everywhere defined, is a rectifiable metric, not necessarily equivalent to d .) A *minimizing path* joining x and y is a rectifiable arc, parametrized by arc length, whose length is exactly $d_r(x, y)$. Clearly if a rectifiable path joining x and y of length $d_r(x, y)$ exists, then a minimizing path also exists.

If $\varepsilon > 0$, an ε -chain joining x and y is a finite sequence of points in X , $x = x_0, x_1, x_2, \dots, x_n = y$, such that $d(x_{i-1}, x_i) \leq \varepsilon$ for $1 \leq i \leq n$. The *length* of this ε -chain is $\sum_1^n d(x_{i-1}, x_i)$. It is well known that any two points in the same component can be joined by an ε -chain. For such x, y let $d^\varepsilon(x, y)$ be the infimum of the lengths of ε -chains joining x and y . Clearly d^ε is a metric on each component of X , $d^\varepsilon \geq d$ (also d^ε is equivalent to d), and $\varepsilon_1 < \varepsilon_2 \Rightarrow d^{\varepsilon_1} \geq d^{\varepsilon_2}$. Let $d'(x, y) = \lim_{\varepsilon \rightarrow 0+} d^\varepsilon(x, y)$. d' may take the value

∞ , but it clearly satisfies the triangle inequality. Thus d' is a metric (no longer necessarily equivalent to d) if finite-valued. Clearly $d' \leq d_r$, wherever d_r is defined. In our case $d' = d_r$ and they are equivalent to d_n on each component $C_{\alpha, \beta}$. (In general a different concept of rectifiable can be defined by saying d is "rectifiable" if $d = d'$, and it follows from a result in [7] that this concept is equivalent if (X, d) is complete, but presumably in general $(d')'$ may be unequal to d' . The only general theorems the author has heard of giving the existence of rectifiable paths between two points such that $d'(x, y) < \infty$ have compactness hypotheses. However, the author is not an expert.)

Finally, we point out that if H is finite dimensional, the concept of minimizing path used in this paper does not coincide with the usual geodesics on Grassmannians, as studied in differential geometry. The reason is that the differential-geometric geodesics, defined in terms of a Riemannian metric, are related to the Hilbert-Schmidt norm rather than the operator norm. Our minimizing paths do coincide with the differential-geometric geodesics if α or β is 1 (or 0). The reason is that for rank 2 selfadjoint operators of trace 0 the Hilbert-Schmidt norm is proportional to the operator norm.

We gratefully acknowledge that we learned about rectifiable metrics in a differential geometry course taught by H. Gluck, probably in 1967–1968. Our notation may be different. We are also grateful for bibliographical help from P. de la Harpe, G. Pedersen, N. C. Phillips, and N. Salinas.

After this paper was written we learned of some related work by other authors. This will be discussed in Remark 5 of §3.

2. MAIN RESULTS

The next two elementary lemmas are stated only for reference.

Lemma 1. Assume θ_1 and θ_2 are nonnegative numbers such that $\theta_1 + \theta_2 \leq \pi$ and u, v, w are unit vectors in H such that $\operatorname{Re}(u, v) = \cos \theta_1$ and $\operatorname{Re}(v, w) = \cos \theta_2$. Then $\operatorname{Re}(u, w) \geq \cos(\theta_1 + \theta_2)$.

Lemma 2. Assume u is a unit vector in H and M a closed subspace of H . Let θ denote the minimum of angles between u and unit vectors v in M (take $\theta = \frac{\pi}{2}$ if $M = \{0\}$). Then $\operatorname{dist}(u, M) = \sin \theta$.

Lemma 3. Assume θ_1 and θ_2 are nonnegative numbers such that $\theta_1 + \theta_2 < \frac{\pi}{2}$ and p, q, r are in \mathcal{P} such that $\|p - q\| = \sin \theta_1$ and $\|q - r\| = \sin \theta_2$. Then $\|p - r\| \leq \sin(\theta_1 + \theta_2)$.

Proof. Let L, M, N be the ranges of p, q, r . It is known that $\|p - r\|$ (for example) is the larger of $\sup\{\operatorname{dist}(u, N) : u \text{ is a unit vector in } L\}$ and $\sup\{\operatorname{dist}(w, L) : w \text{ is a unit vector in } N\}$. Let u be a unit vector in L . Since $\operatorname{dist}(u, M) \leq \sin \theta_1$, there is a unit vector v in M such that $\operatorname{Re}(u, v) \geq \cos \theta_1$ by Lemma 2. Since $\operatorname{dist}(v, N) \leq \sin \theta_2$, there is a unit vector w in N such that $\operatorname{Re}(v, w) \geq \cos \theta_2$ by Lemma 2. Then by Lemma 1, $\operatorname{Re}(u, w) \geq \cos(\theta_1 + \theta_2)$. Thus, again by Lemma 2, $\operatorname{dist}(u, N) \leq \sin(\theta_1 + \theta_2)$. Similarly, the second supremum is at most $\sin(\theta_1 + \theta_2)$.

Remark. Unless θ_1 or θ_2 is 0, $\sin(\theta_1 + \theta_2) < \sin \theta_1 + \sin \theta_2$. Thus the conclusion of Lemma 3 is stronger than the triangle inequality for $\|\cdot\|$.

Corollary 4. d_a satisfies the triangle inequality. Hence d_a is a metric on \mathcal{P} , equivalent to d_n .

Proof. Assume p, q, r are in \mathcal{P} and $d_a(p, q) = \theta_1$, $d_a(q, r) = \theta_2$. For the proof that $d_a(p, r) \leq \theta_1 + \theta_2$ it is clearly permissible to assume that $\theta_1 + \theta_2 < \frac{\pi}{2}$. Thus Lemma 3 applies.

Remark. It was already known how to calculate $\|p - q\|$ in terms of the canonical representation of the pair (p, q) . Using this, we see that $d_a(p, q) = \frac{\pi}{2}$ if H_{01} or H_{10} is not $\{0\}$ and otherwise $d_a(p, q) = \|\varphi\|_\infty$, in the notation of §1.

We now specialize to consideration of a fixed component $C_{\alpha, \beta}$. Because $\lim_{\varepsilon \rightarrow 0^+} \frac{\sin \varepsilon}{\varepsilon} = 1$, the same paths are rectifiable relative to d_a and d_n and the arc lengths are the same. Thus $(d_a)_r = (d_n)_r$, and we will write " d_r " for this metric. Similarly, $d'_a = d'_n$ and we write " d' " for this where no confusion would arise. Thus (assuming, as will later be proved, that d_r is everywhere defined) $d_r \geq d' \geq d_a \geq d_n$.

Proposition 5. *If p and q are in $C_{\alpha, \beta}$ and either $\|p - q\| < 1$ or $\|p - q\| = 1$ and $\text{rank}(p \wedge (1 - q)) = \text{rank}(q \wedge (1 - p))$, then there is a path joining p and q of arc length $d_a(p, q)$. Hence in these cases $d_r(p, q) = d'(p, q) = d_a(p, q)$, and a minimizing path exists.*

Proof. Of course the hypothesis is stated redundantly, since $\|p - q\| < 1$ implies $p \wedge (1 - q) = q \wedge (1 - p) = 0$. Both cases are proved simultaneously by using the canonical representation of (p, q) . If $\|p - q\| = 1$, we choose a unitary from H_{01} onto H_{10} and represent $H_{01} \oplus H_{10}$ in the same way as the generic part, with an angle of $\frac{\pi}{2}$. Then we can easily define a function $\gamma: [0, d_a(p, q)] \rightarrow C_{\alpha, \beta}$ such that $\gamma(0) = p$, $\gamma(d_a(p, q)) = q$, and $d_a(\gamma(s), \gamma(t)) = |s - t|$. In informal language we define γ by rotating p toward q .

Explicitly, write $H \cong H_{00} \oplus (L^2(X') \oplus L^2(X')) \oplus (L^2(X) \oplus L^2(X)) \oplus H_{11}$ so that

$$p = 0 \oplus \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \oplus \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \oplus 1$$

and

$$q = 0 \oplus \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \oplus \begin{bmatrix} \cos^2 \varphi & \cos \varphi \sin \varphi \\ \cos \varphi \sin \varphi & \sin^2 \varphi \end{bmatrix} \oplus 1,$$

where the second component (involving $L^2(X')$) is missing if $\|p - q\| < 1$. Then

$$\gamma(s) = 0 \oplus \begin{bmatrix} \cos^2 s & \cos s \sin s \\ \cos s \sin s & \sin^2 s \end{bmatrix} \oplus \begin{bmatrix} \cos^2 \varphi_s & \cos \varphi_s \sin \varphi_s \\ \cos \varphi_s \sin \varphi_s & \sin^2 \varphi_s \end{bmatrix} \oplus 1,$$

where $\varphi_s(x) = \min(s, \varphi(x))$. (Another choice would be to take $\varphi_s(x) = s\varphi(x)/d_a(p, q)$. This choice would produce a smooth curve.)

Remark. If α or β is finite, then $\text{rank}(p \wedge (1 - q)) = \text{rank}(q \wedge (1 - p))$ for all p, q in $C_{\alpha, \beta}$ and Proposition 5 is all-inclusive. Thus from now on the reader may as well assume that α and β are infinite.

Proposition 6. *If p and q are in $C_{\alpha, \beta}$, then there is a path of arc length at most π joining p and q . Hence $d_r(p, q) \leq \pi$.*

Proof. Again we use the canonical representation of (p, q) . We construct a path γ of length at most $\frac{\pi}{2}$ from p to r , where r is chosen so that (r, q) satisfies the hypothesis of Proposition 5. There are two cases:

- (i) $\dim H_{11} \leq \dim H_{00}$,

(ii) $\dim H_{11} > \dim H_{00}$.

In both cases $\gamma(\cdot)$ is defined similarly to the proof of Proposition 5, except that now we rotate p away from q , ending at an angle of $\frac{\pi}{2}$.

Explicitly, in case (i) we choose a unitary from H_{11} onto a subspace L of H_{00} . Then

$$\begin{aligned} H &\cong (H_{00} \ominus L) \oplus H_{10} \oplus H_{01} \oplus (L \oplus L) \oplus (L^2(X) \oplus L^2(X)), \\ p &= 0 \oplus 1 \oplus 0 \oplus \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \oplus \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \\ q &= 0 \oplus 0 \oplus 1 \oplus \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \oplus \begin{bmatrix} \cos^2 \varphi & \cos \varphi \sin \varphi \\ \cos \varphi \sin \varphi & \sin^2 \varphi \end{bmatrix}. \end{aligned}$$

We take

$$r = 0 \oplus 1 \oplus 0 \oplus \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \oplus \begin{bmatrix} \cos^2(\varphi - \frac{\pi}{2}) & \cos(\varphi - \frac{\pi}{2}) \sin(\varphi - \frac{\pi}{2}) \\ \cos(\varphi - \frac{\pi}{2}) \sin(\varphi - \frac{\pi}{2}) & \sin^2(\varphi - \frac{\pi}{2}) \end{bmatrix}.$$

In case (ii) we choose a unitary from a subspace L of H_{11} onto H_{00} . Then

$$\begin{aligned} H &\cong H_{10} \oplus H_{01} \oplus (L \oplus L) \oplus (L^2(X) \oplus L^2(X)) \oplus (H_{11} \ominus L), \\ p &= 1 \oplus 0 \oplus \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \oplus \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \oplus 1, \\ q &= 0 \oplus 1 \oplus \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \oplus \begin{bmatrix} \cos^2 \varphi & \cos \varphi \sin \varphi \\ \cos \varphi \sin \varphi & \sin^2 \varphi \end{bmatrix} \oplus 1. \end{aligned}$$

We take

$$r = 1 \oplus 0 \oplus \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \oplus \begin{bmatrix} \cos^2(\varphi - \frac{\pi}{2}) & \cos(\varphi - \frac{\pi}{2}) \sin(\varphi - \frac{\pi}{2}) \\ \cos(\varphi - \frac{\pi}{2}) \sin(\varphi - \frac{\pi}{2}) & \sin^2(\varphi - \frac{\pi}{2}) \end{bmatrix} \oplus 1.$$

In case (i) r is orthogonal to q . Since $\text{rank } r = \text{rank } q = \alpha$, Proposition 5 applies to (r, q) . In case (ii) $rq = qr$ and $r \vee q = 1$. Therefore

$$\text{rank}(r \wedge (1 - q)) = \text{rank}(1 - q) = \beta = \text{rank}(1 - r) = \text{rank}(q \wedge (1 - r)).$$

Thus again Proposition 5 applies. To complete the proof, we join r to q by a path of arc length $\frac{\pi}{2}$.

Lemma 7. Assume p and q are in $C_{\alpha, \beta}$ and $\text{rank}(p - p \wedge q) \neq \text{rank}(q - p \wedge q)$. Then $d'(p, q) \geq \pi$.

Proof. If $d'(p, q) = d'_a(p, q) < \pi$, then there is an r such that $d_a(p, r) < \frac{\pi}{2}$ and $d_a(r, q) < \frac{\pi}{2}$. Denote the ranges of p , r , and q by L , M , and N , respectively. Let $M_0 = r(L \cap N)$. Since $\|p - r\| < 1$, the restriction of r to L is a topological isomorphism from L onto M . Thus $L/L \cap N$ is topologically isomorphic to M/M_0 . It follows that $\text{rank}(p - p \wedge q) = \dim(L \ominus L \cap N) = \dim(M \ominus M_0)$. Similarly, $\text{rank}(q - p \wedge q) = \dim(M \ominus M_0)$. This contradiction completes the proof.

Remark. Lemma 7 applies in particular if p is a proper subprojection of q .

We are now ready to "calculate" d_r and show that $d_r = d'$. For the sake of clarity we describe the notation before stating a formal theorem. Assume that p and q are in $C_{\alpha, \beta}$ and $\text{rank}(p \wedge (1 - q)) \neq \text{rank}(q \wedge (1 - p))$. Let M and N be the ranges of p and q and refer to the canonical representation of (p, q) .

For $0 \leq \theta < \frac{\pi}{2}$, let $E_\theta = \{x \in X: \varphi(x) \geq \theta\}$ and let H_θ be the closed subspace of H_0 corresponding to the projection

$$\begin{bmatrix} \chi_{E_\theta} & 0 \\ 0 & \chi_{E_\theta} \end{bmatrix}, \quad \text{where } \chi_{E_\theta}(x) = \begin{cases} 1, & x \in E_\theta, \\ 0, & x \notin E_\theta, \end{cases}$$

and the matrix has the same meaning as in §1. Let $M_\theta = M \cap H_\theta$ and $N_\theta = N \cap H_\theta$. (Thus M_0 is the entire generic part of M .)

It may be that for some θ in $(0, \frac{\pi}{2})$ $\dim(H_{10} \oplus M_\theta) = \dim(H_{01} \oplus N_\theta)$. (Note that $\dim M_\theta = \dim N_\theta$.) If this occurs let θ_0 be the supremum of all $\theta < \frac{\pi}{2}$ such that $\dim(H_{10} \oplus M_\theta) = \dim(H_{01} \oplus N_\theta)$. (Thus θ_0 may be $\frac{\pi}{2}$.) If this does not occur, let $\theta_0 = 0$.

We remark that in the separable case there is a more suggestive way to describe θ_0 : We have $\dim(H_{10} \oplus M_\theta) = \dim(H_{01} \oplus N_\theta)$ if and only if M_θ (or equivalently H_θ) is infinite dimensional. Then the lowest point in the essential spectrum of the restriction of pqp to its initial space is $\cos^2 \theta_0$. (The initial space of pqp is $M_0 \oplus H_{11}$.) Furthermore if H_{10} is finite dimensional, $\cos^2 \theta_0$ is also the lowest point in the essential spectrum of $pqp|_M$. Since $\text{rank}(p \wedge (1 - q)) \neq \text{rank}(q \wedge (1 - p))$, at least one of H_{10} , H_{01} is finite dimensional. Thus θ_0 can be calculated from the images of p and q in the Calkin algebra. In particular $\theta_0 = 0$ if and only if the Calkin image of p is a subprojection (not necessarily proper) of the Calkin image of q , or vice-versa. Note that the hypothesis $\text{rank}(p \wedge (1 - q)) \neq \text{rank}(q \wedge (1 - p))$ cannot be verified in terms of the Calkin images of p and q .

Theorem 8. *If p and q are in $C_{\alpha, \beta}$ and $\text{rank}(p \wedge (1 - q)) \neq \text{rank}(q \wedge (1 - p))$, then $d_r(p, q) = d'(p, q) = \pi - \theta_0$.*

Proof. 1. $d_r(p, q) \leq \pi - \theta_0$:

In view of Proposition 6, we may assume $\theta_0 > 0$. Choose θ such that $0 < \theta < \theta_0$. Then $\dim(H_{10} \oplus M_\theta) = \dim(H_{01} \oplus N_\theta)$. We can find a path of length at most $\frac{\pi}{2} - \theta$ joining p to r , where the H_θ -component of r has range $H_\theta \ominus N_\theta$, and the other components of r are the same as those of p (cf. proof of Proposition 6). Then $\text{rank}(r \wedge (1 - q)) = \text{rank}(q \wedge (1 - r)) = \dim(H_{01} \oplus N_\theta)$. Thus Proposition 5 implies $d_r(r, q) \leq \frac{\pi}{2}$, and hence $d_r(p, q) \leq \pi - \theta$. Since θ can be taken arbitrarily close to θ_0 , $d_r(p, q) \leq \pi - \theta_0$.

2. $d'(p, q) \geq \pi - \theta_0$:

We already know $d'(p, q) \geq d_a(p, q) = \frac{\pi}{2}$. Thus we may assume $\theta_0 < \frac{\pi}{2}$. Choose θ such that $\theta_0 < \theta < \frac{\pi}{2}$. Then $\dim(H_{10} \oplus M_\theta) \neq \dim(H_{01} \oplus N_\theta)$. We can find a path of length at most θ joining p to r , where the $(H_0 \ominus H_\theta)$ -component of r is the same as that of q and the other components of r are the same as those of p . Then $\text{rank}(r - r \wedge q) = \dim(H_{10} \oplus M_\theta) \neq \dim(H_{01} \oplus N_\theta) = \text{rank}(q - r \wedge q)$. Thus Lemma 7 implies $d'(r, q) \geq \pi$. Then $d'(p, q) \geq d'(r, q) - d'(p, r) \geq d'(r, q) - d_r(p, r) \geq \pi - \theta$. Since θ can be taken arbitrarily close to θ_0 , $d'(p, q) \geq \pi - \theta_0$.

Remark. On the basis of what has already been said, it is obvious that when α and β are infinite, there exist examples of (p, q) satisfying the hypotheses of Theorem 8 for any θ_0 in $[0, \frac{\pi}{2}]$ (even if H is not separable).

All that remains is to settle the question of existence of minimizing paths. By Propositions 5 and 6, minimizing paths always exist if $d_r(p, q) < \frac{\pi}{2}$ or if $d_r(p, q) = \pi$.

Theorem 9. If $\text{rank}(p \wedge (1 - q)) \neq \text{rank}(q \wedge (1 - p))$ and if $d_r(p, q) = \frac{\pi}{2}$, then p and q are not joined by a minimizing path.

Proof. If a minimizing path exists, then there is an r such that $d_a(p, r) = d_a(r, q) = \frac{\pi}{4}$. Let L, M, N be the ranges of p, r, q , respectively. Referring to the canonical representation of (p, r) , we can find a unitary U from L onto M with the following properties:

1. $(Ux, x) \geq \cos \frac{\pi}{4} = 1/\sqrt{2}$, for every unit vector x in L . (In particular $(Ux, x) \geq 0$.)
2. If $(Ux, x) = \cos \frac{\pi}{4}$ then $rx = \cos \frac{\pi}{4}Ux$ and $pUx = \cos \frac{\pi}{4}x$.

U is closely related to Dixmier's " S_{V_0} " in [6]. In fact the H_0 -component of U is just the restriction of S_{V_0} to L_0 . (The H_{11} -component of U is the identity.) We also find a unitary V from M onto N with similar properties.

Then VU is a unitary from L onto N . Without loss of generality, we assume

$$\dim(L \cap N^\perp) = \text{rank}(p \wedge (1 - q)) > \text{rank}(q \wedge (1 - p)) = \dim(N \cap L^\perp).$$

Thus there is a unit vector x in $L \cap N^\perp$ such that VUx is orthogonal to $N \cap L^\perp$. Let $Ux = y$ and $VUx = z$. Since x is in N^\perp and z is in N , $x \perp z$. By property 1 for U and V , $(y, x) \geq 1/\sqrt{2}$ and $(y, z) \geq 1/\sqrt{2}$. Since $\|y\| = 1$, this implies $y = (1/\sqrt{2})x + (1/\sqrt{2})z$. Then property 2 for U applies, and we conclude that $py = (1/\sqrt{2})x$. Since $py = (1/\sqrt{2})x + (1/\sqrt{2})pz$, this implies $pz = 0$. In other words, z is in $N \cap L^\perp$, a contradiction.

The remainder of this section could be simplified significantly in the case where H is separable. The reader who wishes to assume H separable may substitute his own simpler arguments in appropriate places. We have a reason for presenting the material in this way, which is explained by Remark 3 in the next section.

Lemma 10. Let h be a positive selfadjoint operator on a Hilbert space M_0 and λ a cardinal number such that $\lambda \leq \text{rank } E_{(0, \varepsilon)}(h)$, $\forall \varepsilon > 0$, where $E_{(0, \varepsilon)}(h)$ is the spectral projection of h for the interval $(0, \varepsilon)$. Then there is a closed subspace M_1 such that $\dim M_1 = \lambda$ and M_1 is disjoint from the range of h .

Sublemma. There are a decreasing sequence (ε_n) of positive numbers and a non-decreasing sequence (λ_n) of cardinal numbers such that $\varepsilon_n \rightarrow 0$, $\sup_n \lambda_n = \lambda$, and $\lambda_n \leq \text{rank } E_{(\varepsilon_{n+1}, \varepsilon_n)}(h)$, $\forall n$.

Proof. We consider two cases:

- (i) λ is the supremum of countably many cardinal numbers, each strictly less than λ .
- (ii) λ is not such a supremum.

In case (i) we choose an increasing sequence (λ_n) such that $\sup_n \lambda_n = \lambda$ and $\lambda_n < \lambda$, $\forall n$. In case (ii) we take $\lambda_n = \lambda$, $\forall n$.

Now we can construct the ε_n 's recursively, starting with $\varepsilon_1 = 1$. If ε_n has already been chosen, we will choose ε_{n+1} such that $0 < \varepsilon_{n+1} \leq \min(\frac{1}{n+1}, \varepsilon_n)$ and $\text{rank } E_{(\varepsilon_{n+1}, \varepsilon_n)}(h) \geq \lambda_n$. This is possible because $\sup_m (\text{rank } E_{(\varepsilon_n/m, \varepsilon_n)}(h)) \geq \lambda$. In case (i), it is clearly impossible that $\text{rank } E_{(\varepsilon_n/m, \varepsilon_n)}(h) \leq \lambda_n$, $\forall m$. In case (ii), clearly $\text{rank } E_{(\varepsilon_n/m, \varepsilon_n)}(h) \geq \lambda$ for some m .

Proof of Lemma 10. Choose a sequence (c_n) of positive numbers such that $\sum_1^\infty c_n^2 = 1$ and $\sum_1^\infty (c_n^2/\varepsilon_n^2) = \infty$. Choose a Hilbert space L and a nondecreasing sequence of closed subspaces L_n such that $\sup_n L_n = L$, $\dim L = \lambda$, and $\dim L_n = \lambda_n$. Let P_n be the projection operator on L with range L_n . For each n choose a partial isometry U_n such that $U_n^* U_n = P_n$ and $U_n U_n^* \leq E_{(\varepsilon_{n+1}, \varepsilon_n)}(h)$. Let $V_n = U_n \sum_1^n c_{n-k+1} (P_k - P_{k-1})$, where $P_0 = 0$. We can define an isometry $V: L \rightarrow M_0$ by $V = \sum_1^\infty V_n$, where the sum converges strongly. Since the V_n 's have mutually orthogonal ranges, $V^* V = \sum_1^\infty V_n^* V_n$, and it is easy to check that V really is an isometry and that the sum really converges. Now we let M_1 be the range of V , and we need only show that Vx is not in the range of h , for a nonzero x in L . But clearly Vx in range h implies $\sum_1^\infty (\|V_n x\|^2/\varepsilon_n^2) < \infty$.

$$\begin{aligned} \frac{\|V_n x\|^2}{\varepsilon_n^2} &= \frac{1}{\varepsilon_n^2} \sum_1^n c_{n-k+1}^2 ((P_k - P_{k-1})x, x) \\ &\geq \sum_1^n \frac{c_{n-k+1}^2}{\varepsilon_{n-k+1}^2} ((P_k - P_{k-1})x, x). \end{aligned}$$

Thus

$$\sum_1^\infty \frac{\|V_n x\|^2}{\varepsilon_n^2} \geq \sum_{k=1}^\infty \left(\sum_{n \geq k} \frac{c_{n-k+1}^2}{\varepsilon_{n-k+1}^2} \right) ((P_k - P_{k-1})x, x).$$

The last sum is ∞ , since $((P_k - P_{k-1})x, x) > 0$ for some k and

$$\sum_{n \geq k} \frac{c_{n-k+1}^2}{\varepsilon_{n-k+1}^2} = \sum_{n=1}^\infty \frac{c_n^2}{\varepsilon_n^2} = \infty.$$

Lemma 11. Let L and M_0 be Hilbert spaces, t a positive number, and S a positive invertible selfadjoint operator on M_0 . Assume that $\|S\| = t$, that S does not achieve its norm, and that $\dim L \leq \text{rank } E_{(t-\varepsilon, t)}(S)$, $\forall \varepsilon > 0$. Then there is an operator $T: L \oplus M_0 \rightarrow M_0$ such that $\|T\| \leq \frac{1}{t}$ and $q: \text{graph } T \rightarrow \text{graph } S$ is one-to-one with dense range, where q is the projection operator on $L \oplus M_0 \oplus M_0$ whose range is $\text{graph } S$. (Thus $q|_L = 0$.)

Remark. The hypothesis that S be positive is given only to simplify the notation. We could equally well work with the spectral projections of $|S|$ instead of S .

Proof. $q|_{M_0 \oplus M_0}$ is given by the operator matrix

$$\begin{pmatrix} (1 + S^2)^{-1} & (1 + S^2)^{-1} S \\ S(1 + S^2)^{-1} & S(1 + S^2)^{-1} S \end{pmatrix}.$$

T will be given by a pair (T', T_0) , where $T': L \rightarrow M_0$ and $T_0: M_0 \rightarrow M_0$. Then $\|T\| \leq \frac{1}{t}$ is equivalent to $T_0 T_0^* + T' T'^* \leq 1/t^2$. Since $\text{graph } T$ and $\text{graph } S$ are topologically isomorphic to $L \oplus M_0$ and M_0 , respectively, instead of proving that q is one-to-one with dense range between the graphs, we may instead prove that the appropriate operator $Q: L \oplus M_0 \rightarrow M_0$ is one-to-one with dense range. So Q is given by the pair (Q', Q_0) , where $Q' = (1 + S^2)^{-1} S T'$ and $Q_0 = (1 + S^2)^{-1} (1 + S T_0)$. We may further simplify the problem by replacing Q with $S^{-1}(1 + S^2)Q$, which is given by the pair $(T', S^{-1} + T_0)$.

Now we apply Lemma 10 with $h = S^{-1} - \frac{1}{t}$ and $\lambda = \dim L$. Let P be the projection operator on M_0 whose range is the space M_1 of Lemma 10, let T' be a one-to-one operator such that $T'T'^* = P/t^2$, and let $T_0 = \frac{1}{t}(-1 + P)$. Then $T_0T_0^* + T'T'^* = 1/t^2$ so that $\|T\| \leq \frac{1}{t}$. Since $S^{-1} + T_0 = h + \frac{1}{t}P \geq h$, and since h is one-to-one (because S does not achieve its norm), $S^{-1} + T_0$ is one-to-one with dense range. Thus we need only show that the ranges of T' and $S^{-1} + T_0$ are disjoint. But if $T'x = (S^{-1} + T_0)y = hy + \frac{1}{t}Py$, then $hy = T'x - \frac{1}{t}Py$, an element of M_1 . Since M_1 is disjoint from the range of h , this implies $y = 0$. Q.E.D.

Theorem 12. *If p and q are in $C_{\alpha, \beta}$ and $\frac{\pi}{2} < d_r(p, q) < \pi$, then there is a minimizing path joining p and q . Thus to summarize: A minimizing path joining p and q always exists if $d_r(p, q) \neq \frac{\pi}{2}$; and if $d_r(p, q) = \frac{\pi}{2}$, a minimizing path exists if and only if $\text{rank}(p \wedge (1 - q)) = \text{rank}(q \wedge (1 - p))$.*

Proof. We may assume $\text{rank}(p \wedge (1 - q)) > \text{rank}(q \wedge (1 - p))$. Choose a unitary from H_{01} onto a subspace L_1 of H_{10} , and let $L = H_{10} \ominus L_1$. Let θ_0 , M , N be as in Theorem 8, so that $d_r(p, q) = \pi - \theta_0$. It may be that $\dim(H_{10} \oplus M_{\theta_0}) = \dim(H_{01} \oplus N_{\theta_0})$. If so, the method of point 1 of the proof of Theorem 8 produces the desired minimizing path. Thus we assume $\dim(H_{10} \oplus M_{\theta_0}) > \dim(H_{01} \oplus N_{\theta_0})$. Then for any θ such that $0 < \theta < \theta_0$ we have

$$\dim(L \oplus (M_{\theta} \ominus M_{\theta_0})) = \dim(N_{\theta} \ominus N_{\theta_0}).$$

(This is an easy exercise in cardinal number theory.) Now choose θ_1 such that $0 < \theta_1 < \theta_0$, and let $H_1 = L \oplus (H_{\theta_1} \ominus H_{\theta_0})$. Then H_1 is a reducing subspace for both p and q . It is another easy exercise to show that both the H_1 - and H_1^\perp -components of p and q still agree in rank and co-rank. Since the H_1^\perp -components clearly satisfy the hypothesis of Proposition 5, we may change notation and assume $H = H_1$. (In the new notation what was $M_{\theta_1} \ominus M_{\theta_0}$ becomes, simply, M_0 .)

Now we can identify $H_0 \ominus M_0$ with M_0 so that N is the graph of a positive operator $S: M_0 \rightarrow M_0$. In terms of the notation for the canonical representation given in §1, S is just the multiplication operator for $\tan \varphi$. Then $\|S\| = \tan \theta_0$, S does not achieve its norm, and S is bounded below by $\tan \theta_1$. Since $\dim(L \oplus M_{\theta}) = \dim N_{\theta}$, $\forall \theta \in (\theta_1, \theta_0)$, we have $\dim L \leq \text{rank } E_{(\tan \theta_0 - \varepsilon, \tan \theta_0)}(S)$, $\forall \varepsilon > 0$. Thus Lemma 11 applies with $t = \tan \theta_0$.

Let r be the projection whose range is $\text{graph } T$. Since $\|T\| \leq \frac{1}{t}$, $d_a(p, r) \leq \tan^{-1}(\frac{1}{t}) = \frac{\pi}{2} - \theta_0 < \frac{\pi}{2}$. Therefore there is a path of length at most $\frac{\pi}{2} - \theta_0$ joining p to r . Because the projection from $\text{graph } T$ to $\text{graph } S$ is one-to-one with dense range, $r \wedge (1 - q) = q \wedge (1 - r) = 0$. Therefore, by Proposition 5, there is a path of length at most $\frac{\pi}{2}$ joining r to q . Combining the two paths gives the desired minimizing path.

3. CONCLUDING REMARKS

1. The metric d_r is not defined on all of \mathcal{P} but only on each $C_{\alpha, \beta}$. We can easily extend d_r to a metric on \mathcal{P} . Just choose any number $\delta > \pi$ and define $d_r(p, q) = \delta$ whenever p and q are in different components of \mathcal{P} . The most natural choice would be $\delta = \infty$, but this is not allowed. The extended d_r is

rectifiable if we redefine “rectifiable” to mean “the restriction to each component is rectifiable in the original sense”.

2. It is obvious that minimizing paths in $C_{\alpha, \beta}$ are usually not unique. For completeness we state without proof the facts on uniqueness. For $p \neq q$, p and q are joined by a unique minimizing path if and only if $d_a(p, q) < \frac{\pi}{2}$, the function φ appearing in the canonical representation is essentially constant, and either $H_{11} = 0$ or $H_{00} = 0$.

3. We can generalize this paper by considering the set of projections in a von Neumann algebra M , instead of the set of projections on H . There are some obvious changes which have to be made. For example, α and β now have to be Murray-von Neumann equivalence classes of projections in M instead of cardinal numbers. The often-stated property “ $\text{rank}(p \wedge (1 - q)) = \text{rank}(q \wedge (1 - p))$ ” becomes “ $p \wedge (1 - q) \sim q \wedge (1 - p)$ ”. The hardest result to generalize to this context is Lemma 10. The new version of this replaces λ with a projection r such that r is equivalent to some subprojection of $E_{(0, \varepsilon)}(h)$ for each $\varepsilon > 0$. The conclusion is that there is a projection r_1 such that $r_1 \sim r$, $r_1 \leq E_{(0, \infty)}(h)$, and the range of r_1 is disjoint from the range of h . It is just a somewhat tedious exercise to prove this generalization, using the given proof of Lemma 10 as a model, though the generalized version does present some additional complications.

4. The basic results of this paper, including especially Theorem 9, can be used to prove a sort of metric index zero theorem: If p , q , and r are distinct projections such that

$$\sin^{-1}(\|p - r\|) + \sin^{-1}(\|r - q\|) \leq \frac{\pi}{2},$$

then $\text{rank}(p \wedge (1 - q)) = \text{rank}(q \wedge (1 - p))$.

5. The often-stated property “ $\text{rank}(p \wedge (1 - q)) = \text{rank}(q \wedge (1 - p))$ ” was used by Davis in [4], where p and q are said to be “equivalently positioned”. The concept of direct rotation from [4] and Propositions 3.1 and 3.2 of Davis and Kahane [5] probably have some relation to the underlying ideas of some of our proofs, including our proof of Proposition 5. Sano and Watatani [15] have applied angles between projections in the context of subfactors. Several authors have previously considered rectifiable metrics (also called Finsler metrics) for various spaces of operators from a differential geometric point of view. Atkin [1, 2] considered various groups of invertible operators instead of \mathcal{P} and Wilkins [17] considered Grassmannians of C^* -algebras. Various works of Salinas, Corach, Porta, and Recht have some overlap with Proposition 5 above and implicitly also Corollary 4, and we proceeded to give more detail. In the works of these authors a different metric is used, so that all of their arc lengths are exactly the doubles of ours. Also these authors consider the set of projections in an arbitrary C^* -algebra, not necessarily a von Neumann algebra as in this paper. Thus their results similar to Proposition 5 above are more general, but they do not have results analogous to Theorems 8, 9, and 12 above, which are not valid in the more general context. Porta and Recht [11] and Salinas [14] have results which include the case of Proposition 5 where $\|p - q\| < 1$. This could be used to prove that d_a is a metric. Corach, Porta, and Recht [3] consider the space of not necessarily selfadjoint idempotents in a C^* -algebra. We also note that the answer to the question in Remark 2.22(b) of [14] is “no”. (See the proof of Proposition 5 above and especially the last sentence.)

REFERENCES

1. C. J. Atkin, *The Finsler geometry of groups of isometries of Hilbert space*, J. Austral. Math. Soc. **42** (1987), 196–222.
2. —, *The Finsler geometry of certain covering groups of operator groups*, Hokkaido Math. J. **18** (1989), 45–77.
3. G. Corach, H. Porta and L. Recht, *The geometry of spaces of projections in C^* -algebras*, Adv. in Math. (to appear).
4. C. Davis, *Separation of two linear subspaces*, Acta Sci. Math. (Szeged) **19** (1958), 172–187.
5. C. Davis and W. M. Kahane, *The rotation of eigenvalues by a perturbation. III*, SIAM J. Numer. Anal. **7** (1970), 1–46.
6. J. Dixmier, *Position relative de deux variétés linéaires fermées dans un espace de Hilbert*, Rev. Sci. **86** (1948), 387–399.
7. M. Gromov, *Structure métriques pour les variétés riemanniennes*, CEDIC, Paris, 1981.
8. P. Halmos, *Two subspaces*, Trans. Amer. Math. Soc. **144** (1969), 381–389.
9. M. G. Krein, M. A. Krasnosel'skii, and D. P. Mil'man, *Defect numbers of linear operators in Banach space and some geometrical problems*, Sbor. Trudov Inst. Mat. Akad. Nauk SSSR, no. 11, 1948, pp. 97–112. (Russian)
10. G. Pedersen, *Measure theory for C^* -algebras. II*, Math. Scand. **22** (1968), 63–74.
11. H. Porta and L. Recht, *Minimality of geodesics in Grassmann manifolds*, Proc. Amer. Math. Soc. **100** (1987), 464–466.
12. I. Raeburn and A. Sinclair, *The C^* -algebra generated by two projections*, Math. Scand. **65** (1989), 278–290.
13. W. Rinow, *Die innere geometrie der metrischen räume*, Springer-Verlag, Berlin, Göttingen, and Heidelberg, 1961.
14. N. Salinas, *The Grassmann manifold of a C^* -algebra and Hermitian holomorphic bundles*, Operator Theory: Adv. Appl. **28** (1988), 267–289.
15. T. Sano and Y. Watatani, *Angles between two subfactors*, preprint.
16. V. S. Sunder, *N subspaces*, Canad. J. Math. **40** (1988), 38–54.
17. D. R. Wilkins, *The Grassmann manifold of a C^* -algebra*, Proc. Roy. Irish Acad. Sect A **90** (1990), 99–116.

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, WEST LAFAYETTE, INDIANA 47907
 E-mail address: lgb@math.purdue.edu