

EIGENFUNCTIONS OF THE LAPLACIAN ON ROTATIONALLY SYMMETRIC MANIFOLDS

MICHEL MARIAS

ABSTRACT. Eigenfunctions of the Laplacian on a negatively curved, rotationally symmetric manifold $M = (\mathbf{R}^n, ds^2)$, $ds^2 = dr^2 + f(r)^2 d\theta^2$, are constructed explicitly under the assumption that an integral of $f(r)$ converges. This integral is the same one which gives the existence of nonconstant harmonic functions on M .

1. INTRODUCTION

Let us equip \mathbf{R}^n , $n \geq 2$, with a Riemannian metric which can be written in polar coordinates as

$$(1.1) \quad \begin{aligned} ds^2 &= dr^2 + f(r)^2 d\theta^2, \\ f(0) &= 0, \quad f'(0) = 1, \quad \text{and } f(r) > 0, \quad \forall r > 0. \end{aligned}$$

\mathbf{R}^n with the metric (1.1) becomes a rotationally symmetric manifold M . M is called a weak model in [9] and a Ricci model in [1]. The Laplace operator in these coordinates is

$$\Delta = f(r)^2 \Delta_\theta + \partial_r^2 + (n-1)f(r)^{-1}f'(r)\partial_r,$$

where Δ_θ is the Laplace operator of the unit sphere $S_{n-1}(0, 1) = S$. In this paper we assume that the radial curvature $k(r) = -f''(r)f(r)^{-1}$ is negative.

In this paper we give a method of constructing eigenfunctions of the Laplacian Δ with eigenvalue $\lambda > 0$, provided that

$$(1.2) \quad J(f) := \int_1^\infty f(r)^{n-3} dr \int_r^\infty f(\rho)^{1-n} d\rho < +\infty.$$

The integral $J(f)$ is the same which gives the existence of nonconstant bounded harmonic functions on M . In [12], March, using a probabilistic approach, proved the following alternative: If $J(f) < \infty$, then there exist nonconstant bounded harmonic functions on M , while if $J(f) = \infty$, there are none such.

The integral $J(f)$ is closely related to the radial curvature. Indeed, if $c_2 = 1$, $c_n = 1/2$, $n \geq 3$, then, [12], under the assumption that $k(r)$ is nonpositive, it follows that

$$J(f) < +\infty, \quad \text{if } k(r) \leq \frac{-c}{r^2 \log r}, \quad \text{for } c > c_n \text{ and large } r,$$

Received by the editors July 16, 1995 and, in revised form, January 18, 1996.

1991 *Mathematics Subject Classification*. Primary 58G25, 60J45.

Key words and phrases. Rotationally symmetric manifolds, radial curvature, eigenfunctions, harmonic functions, heat kernels, diffusion processes, subordination measure.

while

$$J(f) = +\infty, \text{ if } k(r) \geq \frac{-c}{r^2 \log r}, \text{ for } c < c_n \text{ and large } r.$$

This alternative had been observed before March by Milnor, [14], in dimension 2 and in part by Choi, [3], in dimension 3. In [13], using generalised Poisson kernels, it is proved that $J(f) < +\infty$ is a sufficient condition for the existence of nonconstant bounded harmonic functions on M .

Under the assumption (1.2), eigenfunctions W_λ , $\lambda > 0$, of Δ with eigenvalue λ , are constructed explicitly in Section 3 by means of the heat kernel K_S of the unit sphere S and a subordination measure $\mu(\lambda, r, t)dt$, $t > 0$. More precisely, in Section 2, we show that there exists a positive and nondecreasing θ -independent eigenfunction $g(\lambda, r)$, $r > R$, with eigenvalue $\lambda > 0$, such that $g(\lambda, R) = 1$. The measure $\mu(\lambda, r, t)$ is a solution of the parabolic equation $\partial_t \mu = B_{r,\lambda} \mu$, where

$$B_{r,\lambda} = f(r)^2 \partial_r^2 + \{(n-1)f(r)f'(r) + 2f(r)^2 g'(\lambda, r)g(\lambda, r)^{-1}\} \partial_r.$$

The measure $\mu(\lambda, r, t)$ is in fact the law of the first hitting time of the boundaries of (R, ∞) by the diffusion associated with the operator $B_{r,\lambda}$. This gives the splitting $\mu = \mu_1 + \mu_2$, where μ_1 (resp. μ_2) is related to the first hitting time of ∞ (resp. R).

In Section 3, Theorem 9, we show that if $J(f) < \infty$, then for any $h_1, h_2 \in C^\infty(S)$ and $\lambda > 0$,

$$(1.3) \quad W_\lambda(r, \theta) := g(\lambda, r) \sum_{j=1,2} \int_0^\infty \int_S K_S(t, \theta, \theta') h_j(\theta') \mu_j(\lambda, r, t) d\theta' dt,$$

is an eigenfunction of Δ with eigenvalue λ , outside the ball $B(0, R)$, such that

$$(1.4) \quad \lim_{r \rightarrow \infty} g(\lambda, r)^{-1} W_\lambda(r, \theta) = h_1(\theta) \quad \text{and} \quad \lim_{r \rightarrow R} g(\lambda, r)^{-1} W_\lambda(r, \theta) = h_2(\theta).$$

In a second step, using the probabilistic interpretation of harmonic functions, it is proved, Theorem 11, that an eigenfunction on $B(0, R)^c$ which satisfies (1.4) is given by (1.3). This allows us to extend uniquely to the whole of M the eigenfunction $W_\lambda(r, \theta)$ given by (1.3), Theorem 12.

Finally, in Section 4, we treat the case of the Euclidean upper-half space \mathbf{R}_+^{n+1} . The Poisson kernels which give rise to the eigenfunctions with eigenvalue $\lambda > 0$ are computed explicitly and reduce to the classical one as $\lambda \rightarrow 0$.

2. THE SUBORDINATION MEASURES

In this section we construct a pair of measures $\mu_j(\lambda, r, t)dt$, $j = 1, 2$, which enable one to write down explicitly eigenfunctions of the Laplacian with eigenvalue $\lambda > 0$.

We start with the existence of a θ -independent eigenfunction $g(\lambda, r)$.

Lemma 1. *For any $\lambda > 0$, there exists a positive and nondecreasing θ -independent eigenfunction $g(\lambda, r)$, $r > 0$, with eigenvalue $\lambda > 0$, such that $g(\lambda, 0) = 1$.*

Proof. The radial part of Δ is given by $\partial_r^2 + (n-1)f(r)^{-1}f'(r)\partial_r$ and $f(r) \sim r$, for r small enough. So, if

$$B(r) = \int_1^r (n-1)f'(s)f(s)^{-1} ds = (n-1) \log f(r)f(1)^{-1},$$

then

$$e^{B(r)} = cf(r)^{n-1} \sim r^{n-1}, \quad \text{for } r \text{ small enough.}$$

The integral tests as in [5], p. 408, give that 0 is an entrance boundary and the result follows. \square

Let $W_\lambda(r, \theta)$ be an eigenfunction of Δ with eigenvalue $\lambda > 0$. If we set

$$w_\lambda(r, \theta) = g(\lambda, r)^{-1} W_\lambda(r, \theta),$$

then it is straightforward to check that $w_\lambda(r, \theta)$ is a harmonic function for the operator $\Delta_\theta + B_{r,\lambda}$, where

$$\begin{aligned} B_{r,\lambda} &= f(r)^2 \partial_r^2 + \{(n-1)f(r)f'(r) + 2f(r)^2 g'(\lambda, r)g(\lambda, r)^{-1}\} \partial_r \\ &= a(r) \partial_r^2 + b(r) \partial_r. \end{aligned}$$

Lemma 2. *If $J(f) < \infty$, then for any $\lambda > 0$, the origin is an entrance boundary for the operator $B_{r,\lambda}$, while $+\infty$ is an exit boundary.*

Proof. As in Lemma 1 we set

$$W(r) = \int_1^r b(s)a(s)^{-1} ds = (n-1) \log f(r)f(1)^{-1} + 2 \log g(\lambda, r)g(\lambda, 1)^{-1}.$$

So,

$$e^{W(r)} = cf(r)^{n-1}g(\lambda, r)^2 \sim r^{n-1},$$

for r small enough, since $g(\lambda, r) \sim 1$, as $r \rightarrow 0$, by Lemma 1. It follows ([7], p.515), that 0 is an entrance boundary.

On the other hand, since the radial curvature is negative, we get that $f(r) \geq cr$ ([9], p.36). This yields

$$\int_1^\infty e^{W(r)} dr = c \int_1^\infty f(r)^{n-1} g(\lambda, r)^2 dr \geq cg(\lambda, 1)^2 \int_1^\infty r^{n-1} dr = +\infty,$$

since $g(\lambda, r)$ is nondecreasing by Lemma 1. Therefore, $+\infty$ is not a regular boundary. But

$$\begin{aligned} & \int_1^\infty e^{-W(r)} dr \int_1^r e^{W(s)} a(s)^{-1} ds \\ &= c \int_1^\infty f(r)^{1-n} g(\lambda, r)^{-2} dr \int_1^r f(s)^{n-3} g(\lambda, s)^2 ds \\ &\leq c \int_1^\infty f(r)^{1-n} dr \int_1^r f(s)^{n-3} ds := I(f), \end{aligned}$$

since $g(\lambda, r)$ is nondecreasing. From Fubini's theorem we get that $I(f) = J(f) < +\infty$, so $+\infty$ is an exit boundary. \square

From now on we fix a positive R . As in Lemma 2, one can show that R is a regular boundary for $B_{r,\lambda}$. Therefore the situation is the same as in [11], Section 3. Following the method developed in [11], one can prove the statements below, which we give without proof.

Lemma 3. *If $J(f) < \infty$, then for any $\lambda > 0$ and $k > 0$, there exists a positive and nondecreasing solution $\varphi(\lambda, k, r)$ of the equation*

$$(2.1) \quad B_{r,\lambda} \varphi(\lambda, k, r) = k \varphi(\lambda, k, r), \quad r \in (R, \infty),$$

such that $\varphi(\lambda, k, R) = 0$ and $\varphi(\lambda, k, r) \rightarrow 1$, as $r \rightarrow \infty$.

The function $\varphi(\lambda, k, r)$ appearing in Lemma 3 has the following probabilistic interpretation. Let us denote by r_t^λ the diffusion on (R, ∞) associated with the operator $B_{r,\lambda}$, and let σ (resp. τ) be the first hitting time of ∞ (resp. R) by r_t^λ starting at $r > R$. The explosion time of r_t^λ is then given by $\zeta = \sigma \wedge \tau$. Let us also denote by $P_{\lambda,r}$ the probability attached to the motion r_t^λ and by $E_{\lambda,r}$ the corresponding expectation.

Lemma 4. *If $J(f) < \infty$, then for any $\lambda > 0$,*

$$P_{\lambda,r} \{ \zeta < +\infty \} = 1, \quad \forall r \in (R, \infty),$$

and

$$(2.2) \quad \varphi(\lambda, k, r) = E'_{\lambda,r} (e^{-k\sigma}), \quad \forall r \in (R, \infty),$$

where $P'_{\lambda,r}$ is the restriction of $P_{\lambda,r}$ to the set Ω_1 of the paths r_t^λ such that $r_\sigma^\lambda = \infty$.

As in [11], Section 3, using (2.2), one can show the following

Proposition 5. *If $J(f) < \infty$, then for any $\lambda > 0$ and $r \in (R, \infty)$, there exists a positive, bounded and C^∞ function $t \rightarrow \mu_1(\lambda, r, t)$, $t > 0$, such that*

$$(2.3) \quad \varphi(\lambda, k, r) = \int_0^\infty e^{-kt} \mu_1(\lambda, r, t) dt,$$

$$(2.4) \quad \partial_t \mu_1 = B_{r,\lambda} \mu_1,$$

$$(2.5) \quad \mu_1(\lambda, r, t) = o(t^m), \quad \forall m > 0, \text{ as } t \rightarrow 0, \text{ and } \lim_{t \rightarrow \infty} \mu_1(\lambda, r, t) = 0,$$

$$(2.6) \quad \int_0^\infty \mu_1(\lambda, r, t) dt < 1, \text{ for any } \lambda > 0, \text{ and } r \in (R, \infty).$$

Remark 6. *From (2.3) of Proposition 5 and the boundary values of $\varphi(\lambda, k, r)$, we deduce that*

$$(2.7) \quad \mu_1(\lambda, r, t) \longrightarrow \delta_0(t), \text{ as } r \rightarrow \infty,$$

in the sense of distributions, and

$$(2.8) \quad \mu_1(\lambda, r, t) \longrightarrow 0, \text{ as } r \rightarrow R.$$

Remark 7. *The subordination measure μ_1 admits a companion μ_2 which, like μ_1 , is related to the first hitting time τ of R . Indeed, since ∞ is an exit boundary and R is a regular one, there exists a positive and nonincreasing solution $\psi(\lambda, k, r)$ of (2.1) such that $\psi(\lambda, k, R) = 1$ and $\psi(\lambda, k, r) \longrightarrow 0$, as $r \rightarrow \infty$. As in the case of φ , the subordination measure μ_2 associated with ψ satisfies the properties stated for μ_1 in Proposition 5. From the boundary values of ψ we get that*

$$(2.9) \quad \mu_1(\lambda, r, t) \longrightarrow \delta_0(t), \text{ as } r \rightarrow R,$$

and

$$(2.10) \quad \mu_1(\lambda, r, t) \longrightarrow 0, \text{ as } r \rightarrow \infty.$$

Remark 8. *For any $\lambda > 0$ and $r \in (R, \infty)$, we set $\mu = \mu_1 + \mu_2$. The measure μ is the law of the explosion time $\zeta = \sigma \wedge \tau$ of the diffusion r_t^λ . Indeed, as in Lemma 4, we have*

$$\begin{aligned}
E_{\lambda,r}(e^{-k\zeta}) &= E_{\lambda,r}(e^{-k\zeta}\chi_{\Omega_1}) + E_{\lambda,r}(e^{-k\zeta}\chi_{\Omega_1^c}) \\
&= E_{\lambda,r}(e^{-k\sigma}\chi_{\Omega_1}) + E_{\lambda,r}(e^{-k\tau}\chi_{\Omega_1^c}) = \varphi(\lambda, k, r) + \psi(\lambda, k, r) \\
&= \int_0^\infty e^{-kt} \mu(\lambda, r, t) dt.
\end{aligned}$$

3. A CLASS OF EIGENFUNCTIONS

In this section, using the subordination measures, we construct, in a first step, eigenfunctions of Δ outside the ball $B(0, R)$, with eigenvalue $\lambda > 0$. In a second step, we show that they can be extended uniquely to the whole of M .

Theorem 9. *If $J(f) < \infty$, then for any $\lambda > 0$ and $h_1, h_2 \in C^\infty(S)$,*

$$(3.1) \quad W_\lambda(r, \theta) := g(\lambda, r) \sum_{j=1,2} \int_0^\infty \int_S K_S(t, \theta, \theta') h_j(\theta') \mu_j(\lambda, r, t) d\theta' dt,$$

is an eigenfunction of Δ outside the ball $B(0, R)$, with eigenvalue $\lambda > 0$, such that

$$(3.2) \quad \lim_{r \rightarrow \infty} g(\lambda, r)^{-1} W_\lambda(r, \theta) = h_1(\theta) \quad \text{and} \quad \lim_{r \rightarrow R} g(\lambda, r)^{-1} W_\lambda(r, \theta) = h_2(\theta).$$

Let us set

$$(3.3) \quad W_\lambda^j(r, \theta) := g(\lambda, r) \int_0^\infty \int_S K_S(t, \theta, \theta') h_j(\theta') \mu_j(\lambda, r, t) dt, \quad j = 1, 2.$$

According to the boundary behaviour of the measures μ_1, μ_2 , given in Remark 6 and Remark 7 of Section 2, for the proof of the Theorem 9, it suffices to show that $W_\lambda^j(r, \theta)$ are eigenfunctions of Δ outside the ball $B(0, R)$, with eigenvalue $\lambda > 0$, such that

$$(3.4) \quad \lim_{r \rightarrow \infty} g(\lambda, r)^{-1} W_\lambda^1(r, \theta) = h_1(\theta) \quad \text{and} \quad \lim_{r \rightarrow R} g(\lambda, r)^{-1} W_\lambda^1(r, \theta) = 0,$$

and

$$(3.5) \quad \lim_{r \rightarrow \infty} g(\lambda, r)^{-1} W_\lambda^2(r, \theta) = 0 \quad \text{and} \quad \lim_{r \rightarrow R} g(\lambda, r)^{-1} W_\lambda^2(r, \theta) = h_2(\theta).$$

Below, we treat the case of $W_\lambda^1(r, \theta)$. The treatment of $W_\lambda^2(r, \theta)$ is similar.

Let us set $w_\lambda^1(r, \theta) = g(\lambda, r)^{-1} W_\lambda^1(r, \theta)$. For the proof of Theorem 9, it suffices to show that w_λ^1 satisfies the following Dirichlet problem:

$$(3.6) \quad \begin{aligned} &(\Delta_\theta + B_{r,\lambda}) w_\lambda^1(r, \theta) = 0, \quad r \in (R, \infty), \quad \theta \in S, \\ &\lim_{r \rightarrow \infty} w_\lambda^1(r, \theta) = h_1(\theta), \quad \text{and} \quad \lim_{r \rightarrow R} w_\lambda^1(r, \theta) = 0. \end{aligned}$$

In general, the proof of (3.6) is probabilistic, [12], [11]. Here, we give an analytic proof since estimates of $\partial_t K_S$ are available. Indeed, since S is a complete manifold with strictly positive Ricci curvature, then

$$(3.7) \quad |\partial_t^l K_S(t, \theta, \theta')| \leq cV(\theta, \sqrt{t})^{-1} t^{-l} (1 + d^2/t)^{l+\frac{n}{2}} e^{-d^2/4t},$$

for all $t > 0$, $\forall \theta, \theta' \in S$, where $V(\theta, r)$ is the volume of the ball $B(\theta, r)$ and $d = d(\theta, \theta')$ is the distance of θ, θ' . (See for instance the survey article [15], Theorem 4.2 combined with Example 1.)

Using the volume estimates of [2]:

$$ct^n V(\theta, 1) \leq V(\theta, t) \leq V(\theta, 1), \quad \forall \theta \in S, \quad \forall t \leq 1,$$

we get from (3.7) that for $t \leq 1$

$$(3.8) \quad |\partial_t^l K_S(t, \theta, \theta')| \leq cV(\theta, 1)^{-1} t^{-l-\frac{n}{2}} (1 + d^2/t)^{l+\frac{n}{2}} e^{-d^2/4t}.$$

On the other hand, from (3.7) it follows that for $t > 1$,

$$(3.9) \quad |\partial_t^l K_S(t, \theta, \theta')| \leq cVol(S)^{-1} t^{-l} (1 + d^2/t)^{l+\frac{n}{2}} e^{-d^2/4t}.$$

Lemma 10. *If $J(f) < \infty$, then for any $\lambda > 0$ and $h_1 \in C^\infty(S)$:*

(i) *The integral*

$$w_\lambda^1(r, \theta) := \int_0^\infty \int_S K_S(t, \theta, \theta') h_1(\theta') \mu_1(\lambda, r, t) d\theta' dt$$

is absolutely convergent and satisfies

$$\lim_{r \rightarrow \infty} w_\lambda^1(r, \theta) = h_1(\theta), \quad \text{and} \quad \lim_{r \rightarrow R} w_\lambda^1(r, \theta) = 0.$$

(ii) *The integral*

$$\int_0^\infty \int_S \partial_t K_S(t, \theta, \theta') h_1(\theta') \mu_1(\lambda, r, t) d\theta' dt,$$

is absolutely convergent.

Proof. (i) For any $h_1 \in C^\infty(S)$,

$$\begin{aligned} & \int_0^\infty \int_S K_S(t, \theta, \theta') |h_1(\theta')| \mu_1(\lambda, r, t) d\theta' dt \\ & \leq \|h_1\|_\infty \int_0^\infty \mu_1(\lambda, r, t) dt \leq \|h_1\|_\infty, \end{aligned}$$

since $\int_S K_S(t, \theta, \theta') d\theta' = 1$, by [16], and $\int_0^\infty \mu_1(\lambda, r, t) dt < 1$, by (2.6).

So, $t \rightarrow \int_S K_S(t, \theta, \theta') h_1(\theta') d\theta'$ belongs in $L^1(\mu_1)$. Thus $\lim_{r \rightarrow \infty} w_\lambda^1(r, \theta) = h_1(\theta)$ by (2.7) since C_0^∞ is dense in $L^1(\mu_1)$. From (2.8) we get that $\lim_{r \rightarrow R} w_\lambda^1(r, \theta) = 0$.

(ii) Suppose for instance that $d \leq 1$. Bearing in mind that $V(\theta, T) > c$, $\forall \theta \in S$, $\forall T \geq 1$, the estimates (3.7), (3.8) and (3.9) of $\partial_t K_S$ give

$$\begin{aligned} & \int_0^\infty |\partial_t K_S(t, \theta, \theta')| \mu_1(\lambda, r, t) dt \\ & \leq c \int_0^{d^2} t^{-1-\frac{n}{2}} (1 + d^2/t)^{1+\frac{n}{2}} e^{-d^2/4t} \mu_1(\lambda, r, t) dt \\ & + c \int_{d^2}^1 t^{-1-\frac{n}{2}} e^{-d^2/4t} \mu_1(\lambda, r, t) dt + c vol(S)^{-1} \int_1^\infty t^{-1} \mu_1(\lambda, r, t) dt \\ & \leq cd^{n+2} \int_0^1 t^{-n-2} \mu_1(\lambda, r, t) dt \\ & + c \int_0^1 t^{-\frac{n}{2}-1} \mu_1(\lambda, r, t) dt + c \int_1^\infty t^{-1} \mu_1(\lambda, r, t) dt \leq c, \end{aligned}$$

since $t \rightarrow \mu_1(\lambda, r, t)$ is infinitely flat near $t = 0$ by (2.5), and

$$\begin{aligned} \int_1^\infty t^{-1} \mu_1(\lambda, r, t) dt &\leq \left(\int_1^\infty t^{-2} dt \right)^{\frac{1}{2}} \left(\int_1^\infty \mu_1(\lambda, r, t)^2 dt \right)^{\frac{1}{2}} \\ &\leq c \|\mu\|_\infty^{\frac{1}{2}} \left(\int_1^\infty \mu_1(\lambda, r, t) dt \right)^{\frac{1}{2}} \leq c, \end{aligned}$$

by (2.6). This gives that

$$\int_0^\infty \int_S |\partial_t K_S(t, \theta, \theta') h_1(\theta')| \mu_1(\lambda, r, t) d\theta' dt \leq c \|h_1\|_1,$$

if $d \leq 1$. The case $d > 1$ is similar. \square

End of the proof of Theorem 9. It remains to show that

$$(\Delta_\theta + B_{r,\lambda}) w_\lambda^1(r, \theta) = 0, \quad r \in (R, \infty), \quad \theta \in S.$$

From Lemma 10 we get

$$\begin{aligned} \Delta_\theta w_\lambda^1(r, \theta) &= \int_0^\infty \int_S \Delta_\theta K_S(t, \theta, \theta') h_1(\theta') \mu_1(\lambda, r, t) d\theta' dt \\ &= - \int_0^\infty \int_S K_S(t, \theta, \theta') h_1(\theta') \partial_t \mu_1(\lambda, r, t) d\theta' dt = -B_{r,\lambda} w_\lambda^1(r, \theta), \end{aligned}$$

since $\partial_t \mu_1 = B_{r,\lambda} \mu_1$, by (2.4). \square

Theorem 11. *If $J(f) < \infty$, then for any $\lambda > 0$ the eigenfunction $W_\lambda(r, \theta)$ defined by (3.1) is the unique eigenfunction satisfying (3.2).*

Proof. The proof of the theorem is probabilistic and is based on the integral representation of bounded harmonic functions.

Let $U_\lambda(r, \theta)$ be an eigenfunction of Δ outside the ball $B(0, R)$, with eigenvalue $\lambda > 0$, satisfying (3.2). Then $u_\lambda(r, \theta) = g(\lambda, r)^{-1} U_\lambda(r, \theta)$ satisfies the Dirichlet problem (3.6).

Let us denote by B_t the Brownian motion on S . The diffusion on $B(0, R)^c$ associated with the operator $\Delta_\theta + B_{r,\lambda}$ is (B_t, r_t^λ) . We choose r_t^λ independent of B_t . Let us also denote by P_θ the probability attached to the Brownian motion B_t . From the independence of the motions B_t, r_t^λ , it follows that the probability $P_{(r,\theta)}^\lambda$ attached to the motion (B_t, r_t^λ) splits to the product $P_\theta P_r^\lambda$. Finally, from the fact that B_t does not explode, we get that the explosion time of the product motion (B_t, r_t^λ) is in fact the explosion time $\zeta = \sigma \wedge \tau$ of r_t^λ , which satisfies

$$(3.10) \quad P_{(r,\theta)}^\lambda \{\zeta < +\infty\} = P_r^\lambda \{\zeta < +\infty\} = 1,$$

by Lemma 4 of Section 2.

From (3.10) and [8], Theorem 2.1, p.127, and Remark 2, p.130, it follows that every bounded harmonic function $u_\lambda(r, \theta)$ for the operator $\Delta_\theta + B_{r,\lambda}$ on $B(0, R)^c$ has the following integral representation:

$$(3.11) \quad u_\lambda(r, \theta) = E_\theta E_r^\lambda \{\Psi(B_\zeta, r_\zeta^\lambda)\},$$

where Ψ is the boundary value of $u_\lambda(r, \theta)$. But, the boundary of $B(0, R)^c$ is $S(0, R) \cup S_\infty$, where the "sphere at infinity" S_∞ is isomorphic to the unit sphere S , [4].

Therefore

$$(3.12) \quad \Psi(r, \theta) = \begin{cases} h_1(\theta), & \text{if } r = +\infty, \\ h_2(\theta), & \text{if } r = R. \end{cases}$$

From (3.11), (3.12) and the fact that $\mu(\lambda, r, t)$ is the law of the explosion time ζ , we obtain that

$$u_\lambda(r, \theta) = \sum_{j=1,2} \int_0^\infty \int_S K_S(t, \theta, \theta') h_j(\theta') \mu_j(\lambda, r, t) d\theta' dt,$$

and the result follows. \square

Theorem 12. *The eigenfunction $W_\lambda(r, \theta)$ defined by (3.1) extends uniquely to the whole of M .*

Proof. By (1.1) and Lemma 1, the operator $\Delta_\theta + B_{r,\lambda}$ has bounded coefficients on the bounded domain $B(0, R)$. Further, its boundary $S(0, R)$, is regular. So, by [8], Theorem 3.1, p.142, the Dirichlet problem

$$\begin{aligned} \{\Delta_\theta + B_{r,\lambda}\} u &= 0, \quad \forall (r, \theta) \in B(0, R), \\ u(r, \theta) &\longrightarrow h_2(\theta), \quad \text{as } r \rightarrow R, \end{aligned}$$

admits a unique solution which is given by

$$u(r, \theta) = E_\theta E_r^\lambda \{h_2(r_T^\lambda)\},$$

where $T = \inf \{t > 0 \mid r_t^\lambda = R\}$ is the first hitting time of $S(0, R)$. \square

4. THE EUCLIDEAN UPPER HALF-SPACE

In this section we treat the case of the Euclidean upper half-space \mathbf{R}_+^{n+1} . The Poisson kernels which give rise to the eigenfunctions with eigenvalue $\lambda > 0$ are computed explicitly and reduce to the classical one as $\lambda \rightarrow 0$.

For any $\lambda > 0$, $g(\lambda, y) = e^{-y\sqrt{\lambda}}$, $y > 0$, is an x -independent eigenfunction of the Laplacian $\Delta = \Delta_x + \partial_y^2$ with eigenvalue λ . So, if $U_\lambda(x, y)$ is an eigenfunction of Δ with eigenvalue λ , then $u_\lambda(x, y) = g(\lambda, y)^{-1} U_\lambda(x, y)$ is a harmonic function for the operator $\Delta_x + \partial_y^2 - 2\sqrt{\lambda}\partial_y$. Therefore, the subordination measure $\nu(\lambda, y, t)dt$ is a solution of the parabolic equation

$$\partial_t \nu = \partial_y^2 \nu - 2\sqrt{\lambda}\partial_y \nu, \quad y > 0, \quad t > 0,$$

which satisfies $\nu(\lambda, y, t) \rightarrow \delta_0(t)$, as $y \rightarrow 0$, and $\nu(\lambda, y, t) \rightarrow 0$, as $y \rightarrow \infty$.

Passing to the Laplace transform variables, we get that $\nu(\lambda, y, t)$ is the Laplace transform of the function

$$\varphi(\lambda, y, k) = e^{-y\sqrt{\lambda}} e^{-y\sqrt{\lambda+k}}.$$

But, $e^{-y\sqrt{k}}$ is the Laplace transform of

$$(4\pi)^{-1/2} y t^{-3/2} e^{-y^2/4t},$$

so $k \rightarrow \varphi(\lambda, y, k)$ is the Laplace transform of

$$\nu(\lambda, y, t) = \frac{y e^{-y\sqrt{\lambda}}}{2\sqrt{\pi}} \frac{e^{-\lambda t} e^{-y^2/4t}}{t^{3/2}}.$$

The Poisson kernels $Q_\lambda(x, y, x')$ which give rise to the eigenfunctions with eigenvalue $\lambda > 0$ are obtained by

$$\begin{aligned} Q_\lambda(x, y, x') &= g(\lambda, y) \int_0^\infty K_{R^n}(t, x, x') \nu(\lambda, y, t) dt \\ &= \frac{ye^{-2y\sqrt{\lambda}}}{2\sqrt{\pi}} \int_0^\infty \frac{e^{-\|x-x'\|^2/4t}}{(4\pi t)^{n/2}} \frac{e^{-\lambda t} e^{-y^2/4t}}{t^{3/2}} dt \\ &= 2 \frac{y\lambda^{\frac{n+1}{4}} e^{-2y\sqrt{\lambda}}}{\pi^{\frac{n+1}{2}}} \left(\|x-x'\|^2 + y^2 \right)^{-\frac{n+1}{4}} K_{\frac{n+1}{2}} \left(\sqrt{\lambda} \sqrt{\|x-x'\|^2 + y^2} \right), \end{aligned}$$

[6], p.146, formula (29), where K_m is the modified Bessel function.

We note that, since $K_m(x) \sim \frac{\Gamma(m)}{2} \left(\frac{2}{x}\right)^m$, $m > 0$, as $x \rightarrow 0$, we get

$$Q_\lambda(x, y, x') \longrightarrow \left(\frac{2}{\pi}\right)^{\frac{n+1}{2}} \Gamma\left(\frac{n+1}{2}\right) \frac{y}{\left(\|x-x'\|^2 + y^2\right)^{\frac{n+1}{2}}}, \text{ as } \lambda \rightarrow 0,$$

i.e. the classical Poisson kernel of the Euclidean upper-half space.

REFERENCES

- [1] J. Cheeger, S.T. Yau, A lower bound for the heat kernel, *Comm. Pure Appl. Math.*, 34, 465-480, (1981). MR **82i**:58065
- [2] J. Cheeger, M. Gromov, M. Taylor, Finite propagation speed, kernel estimates for functions of the Laplace operator, and the geometry of a complete Riemannian manifold, *J. Diff. Geom.*, 17, 15-53, (1982). MR **84b**:58109
- [3] H.I. Choi, Asymptotic Dirichlet problems for harmonic functions on Riemannian manifolds, *Trans. Amer. Math. Soc.*, 281, 691-716, (1984). MR **85b**:53040
- [4] P. Eberlein, B. O'Neill, Visibility manifolds, *Pac. J. Math.*, 46, 45-109, (1973). MR **49**:1421
- [5] J. Elliott, Eigenfunction expansions associated with singular differential operators, *Trans. Amer. Math. Soc.*, 78, 406-425, (1955). MR **16**:927a
- [6] A. Erdelyi, *Tables of Integral Transforms*, Vol. 1, McGraw-Hill, 1954. MR **15**:868a
- [7] W. Feller, The parabolic differential equation and the associated semi-groups of transformations, *Ann. of Math.*, 55, 468-519, (1952). MR **13**:948a
- [8] M. Freidlin, *Functional Integration and Partial Differential Equations*, *Annals of Math. Studies*, Princeton University Press, Princeton, 1985. MR **87g**:60066
- [9] R.E. Greene, H. Wu, *Function Theory on Manifolds Which Possess a Pole*, L.N.M. 699, Springer, Berlin, 1979. MR **81a**:53002
- [10] N. Ikeda, S. Watanabe, *Stochastic Differential Equations and Diffusion Processes*, North-Holland, Amsterdam, and Kodansha, Tokyo, 1981. MR **84b**:60080
- [11] N. Mandouvalos, M. Marias, Eigenfunctions of the Laplacian and boundary behaviour on manifolds of hyperbolic type, *Math. Proc. Camb. Phil. Soc.*, 122, 551-564, (1997). CMP 97:17
- [12] P. March, Brownian motion and harmonic functions on rotationally symmetric manifolds, *Annals of Prob.*, 14, 793-801, (1986). MR **87m**:60181
- [13] M. Marias, Generalised Poisson Kernels and Applications, *Rev. Roumaine Math. Pures Appl.*, 35, 417-429, (1990). CMP 91:04
- [14] J. Milnor, On deciding whether a surface is parabolic or hyperbolic, *Amer. Math. Monthly*, 84, 43-46, (1977). MR **55**:1257
- [15] L. Saloff-Coste, A note on Poincaré, Sobolev and Harnack inequalities, *Inter. Math. Research Notices* 1992, No. 2, 27-38. MR **93d**:58158
- [16] S.T. Yau, On the heat kernel of a complete Riemannian manifold, *J. Math. Pures et Appl.*, 57, 191- 201, (1978). MR **81b**:58041

DEPARTMENT OF MATHEMATICS, ARISTOTLE UNIVERSITY OF THESSALONIKI, THESSALONIKI 54.006, GREECE

E-mail address: marias@ccf.auth.gr