

ON THE DIOPHANTINE EQUATION

$$(x^3 - 1)/(x - 1) = (y^n - 1)/(y - 1)$$

MAOHUA LE

ABSTRACT. In this paper we prove that the equation $(x^3 - 1)/(x - 1) = (y^n - 1)/(y - 1)$, $x, y, n \in \mathbb{N}$, $x > 1$, $y > 1$, $n > 3$, has only the solutions $(x, y, n) = (5, 2, 5)$ and $(90, 2, 13)$ with y is a prime power. The proof depends on some new results concerning the upper bounds for the number of solutions of the generalized Ramanujan-Nagell equations.

1. INTRODUCTION

Let $\mathbb{Z}, \mathbb{N}, \mathbb{Q}$ be the sets of integers, positive integers and rational numbers respectively. For any positive integer N with $N > 2$, let $s(N)$ denote the number of solutions (x, m) of the equation

$$(1) \quad N = \frac{x^m - 1}{x - 1}, \quad x, m \in \mathbb{N}, \quad x > 1, \quad m > 2.$$

Eighty years ago, Rataj [13] and Goormaghtigh [4] observed that $s(31) = 2$ and $s(8191) = 2$, respectively. Simultaneously, they conjectured that if $N \notin \{31, 8191\}$, then $s(N) \leq 1$. The problem can be written in the following form

Conjecture A. The equation

$$(2) \quad \frac{x^m - 1}{x - 1} = \frac{y^n - 1}{y - 1}, \quad x, y, m, n \in \mathbb{N}, \quad x > 1, \quad y > 1, \quad n > m > 2,$$

has only the solutions $(x, y, m, n) = (5, 2, 3, 5)$ and $(90, 2, 3, 13)$.

In addition, Bateman (see [5, Problem B25]) asked the following problem.

Problem B. Is $(x, y, m, n) = (5, 2, 3, 5)$ the only solution of (2) for which x, y and $(x^m - 1)/(x - 1)$ are all prime powers?

These are two rather difficult questions. By the results of Baker, Davenport, Lewis, Schinzel, Shorey and Tijdeman (see the references of [14]), we see that (2) has only finitely many solutions (x, y, m, n) if any two out of the four variables x, y, m and n are fixed. However, to this day we do not know whether (2) has only finitely many solutions (x, y, m, n) if any one out of x, y, m and n is fixed.

In this paper we discuss the finiteness of solutions of (2) by the means of the generalized Ramanujan-Nagell equations. Let D_1, D_2 be coprime positive integers,

Received by the editors February 28, 1996 and, in revised form, September 18, 1996 and January 22, 1997.

1991 *Mathematics Subject Classification.* Primary 11D61, 11J86.

Supported by the National Natural Science Foundation of China and the Guangdong Provincial Natural Science Foundation.

and let p be a prime with $p \nmid D_1 D_2$. Further let $N(D_1, D_2, p)$ and $N'(D_1, D_2, p)$ denote the numbers of the solutions (r, s) and (r', s') of equations

$$(3) \quad D_1 r^2 + D_2 = p^s, \quad r, s \in \mathbb{N},$$

and

$$(4) \quad D_1 r'^2 + D_2 = 2p^{s'}, \quad r', s' \in \mathbb{N},$$

respectively. In this paper we prove the following two general results.

Theorem 1. $N(D_1, D_2, p) \leq 2$ except for $N(1, 7, 2) = 5$ and $N(3, 5, 2) = 3$.

Theorem 2. If $2 \nmid D_1 D_2$ and p is an odd prime, then $N'(D_1, D_2, p) \leq 2$.

On applying the above-mentioned theorems to the equation (2), we shall deduce the following result.

Corollary. The equation (2) has only the solutions $(x, y, m, n) = (5, 2, 3, 5)$ and $(90, 2, 3, 13)$ such that $m = 3$ and y is a prime power.

2. PROOF OF THEOREM 1

First we may assume that D_1 is squarefree. In [1] and [2], Apéry proved that $N(1, D_2, p) \leq 2$ except for $N(1, 7, 2) = 5$. In [7] and [8], the author proved that $N(D_1, 1, p) \leq 2$ and $N(D_1, D_2, 2) \leq 2$ except for $N(1, 7, 2) = 5$ and $N(3, 5, 2) = 3$, respectively. Therefore, we may assume that $D_1 > 1$, $D_2 > 1$ and p is an odd prime.

Lemma 1 ([10, Formula 3.76]). For any positive integer t and any complex numbers α and β , we have

$$\alpha^t + \beta^t = \sum_{i=0}^{\lfloor t/2 \rfloor} (-1)^i \binom{t}{i} (\alpha + \beta)^{t-2i} (\alpha\beta)^i,$$

where $\lfloor t/2 \rfloor$ is the largest integer which does not exceed $t/2$,

$$\binom{t}{i} = \frac{(t-i-1)!t}{(t-2i)!i!}, \quad i = 0, 1, \dots, \left\lfloor \frac{t}{2} \right\rfloor,$$

are positive integers.

Lemma 2 ([9, Lemma 4]). If the equation

$$(5) \quad D_1 X^2 + D_2 Y^2 = p^Z, \quad X, Y, Z \in \mathbb{Z}, \quad \gcd(X, Y) = 1, \quad Z > 0,$$

has solutions (X, Y, Z) , then it has a unique solution (X_1, Y_1, Z_1) such that $X_1 > 0$, $Y_1 > 0$ and $Z_1 \leq Z$, where Z runs through all solutions (X, Y, Z) of (5). (X_1, Y_1, Z_1) is called the least solution of (5). Further, every solution (X, Y, Z) of (5) can be expressed as

$$Z = Z_1 t, \quad X\sqrt{D_1} + Y\sqrt{-D_2} = \lambda_1(X_1\sqrt{D_1} + \lambda_2 Y_1\sqrt{-D_2})^t, \\ t \in \mathbb{N}, 2 \nmid t, \lambda_1, \lambda_2 \in \{-1, 1\}.$$

Lemma 3. If $N(D_1, D_2, p) > 2$, then (3) has two solutions (r_1, s_1) and (r_2, s_2) such that

$$(6) \quad s_1 = Z_1 t_1, \quad s_2 = Z_1 t_2, \quad 1 < t_1 < t_2,$$

where (X_1, Y_1, Z_1) is the least solution of (5), t_1 and t_2 are odd primes satisfying

$$(7) \quad \sum_{i=0}^{(t_j-1)/2} \begin{bmatrix} t_j \\ i \end{bmatrix} (-4D_2)^{(t_j-1)/2-i} p^{Z_1 i} = (-1)^{(t_j-1)(p^{Z_1}-1)/4}, \quad j = 1, 2,$$

$$(8) \quad \frac{\pi}{2 \arcsin(D_2/p^{s_1})^{1/2}} < t_2$$

and

$$(9) \quad \left| \arcsin \left(\frac{D_2}{p^{Z_1}} \right)^{1/2} - \frac{k\pi}{t_j} \right| < \frac{\pi}{2t_j p^{Z_1(t_j-1)/2}}, \quad j = 1, 2,$$

where k is a positive integer satisfying $k \leq (t_j - 1)/2$.

Proof. We see that if (3) has a solution (r, s) , then (5) has a solution $(X, Y, Z) = (r, 1, s)$. It follows from Lemma 2 that the least solution (X_1, Y_1, Z_1) of (5) satisfies $Y_1 = 1$. Then $(r, s) = (X_1, Z_1)$ is a solution of (3). Further, let

$$(10) \quad \varepsilon = X_1 \sqrt{D_1} + \sqrt{-D_2}, \quad \bar{\varepsilon} = X_1 \sqrt{D_1} - \sqrt{-D_2}.$$

By the proof of [9, Theorem 1], if (r, s) is a solution of (3) with $(r, s) \neq (X_1, Z_1)$, then there exists a positive integer t satisfying $t > 1$, $2 \nmid t$,

$$(11) \quad s = Z_1 t$$

and

$$(12) \quad \left| \frac{\varepsilon^t - \bar{\varepsilon}^t}{\varepsilon - \bar{\varepsilon}} \right| = 1.$$

Therefore, by (11), if $N(D_1, D_2, p) > 2$, then (3) has two solutions (r_1, s_1) and (r_2, s_2) satisfying (6). Moreover, we may assume that (3) has no solutions (r, s) satisfying

$$(13) \quad Z_1 < s < s_1, \quad s_1 < s < s_2.$$

Since $\varepsilon - \bar{\varepsilon} = 2\sqrt{-D_2}$ and $\varepsilon\bar{\varepsilon} = p^{Z_1}$, by Lemma 1, we get from (12) that

$$(14) \quad \begin{aligned} \frac{\varepsilon^t - \bar{\varepsilon}^t}{\varepsilon - \bar{\varepsilon}} &= \sum_{i=0}^{(t-1)/2} (-1)^i \begin{bmatrix} t \\ i \end{bmatrix} (\varepsilon - \bar{\varepsilon})^{t-2i-1} (-\varepsilon\bar{\varepsilon})^i = \sum_{i=0}^{(t-1)/2} \begin{bmatrix} t \\ i \end{bmatrix} (-4D_2)^{(t-1)/2-i} p^{Z_1 i} \\ &= \pm 1. \end{aligned}$$

Since $t = (-1)^{(t-1)/2} \pmod{4}$ and $p^{Z_1} \equiv (-1)^{(p^{Z_1}-1)/2} \pmod{4}$, we obtain (7) from (14).

For any positive integer t with $2 \nmid t$, let

$$(15) \quad X_t = \left\lfloor \frac{\varepsilon^t + \bar{\varepsilon}^t}{2\sqrt{D_1}} \right\rfloor, \quad Y_t = \left\lfloor \frac{\varepsilon^t - \bar{\varepsilon}^t}{\varepsilon - \bar{\varepsilon}} \right\rfloor.$$

By Lemma 1, X_t and Y_t are positive integers satisfying

$$(16) \quad D_1 X_t^2 + D_2 Y_t^2 = p^{Z_1 t}.$$

Further, by Lemma 2, we see from (16) that (r, s) is a solution of (3) satisfying (11) if and only if $Y_t = 1$. From (7), we get

$$(17) \quad Y_{t_1} = Y_{t_2} = 1.$$

If t_1 is not an odd prime, then $t_1 = k_1 k_2$, where k_1 and k_2 are positive integers satisfying $k_1 > 1$, $k_2 > 1$ and $2 \nmid k_1 k_2$. On applying Lemma 1, we find from (17) that

$$(18) \quad 1 = Y_{t_1} = Y_{k_1} \left| \frac{(\varepsilon^{k_1})^{k_2} - (\bar{\varepsilon}^{k_1})^{k_2}}{\varepsilon^{k_1} - \bar{\varepsilon}^{k_1}} \right| = Y_{k_1} \left| \sum_{i=0}^{(k_2-1)/2} \begin{bmatrix} k_2 \\ i \end{bmatrix} (-4D_2 Y_{k_1}^2)^{(k_2-1)/2-i} p^{Z_1 k_1 i} \right|.$$

Hence, we get from (18) that $Y_{k_1} = Y_{t_1}/Y_{k_1} = 1$. It implies that (3) has a solution $(r, s) = (X_{k_1}, Z_1 k_1)$ satisfying $Z_1 < s < s_1$, which contradicts (13). Thus t_1 must be an odd prime.

By the same argument, if t_2 is not an odd prime, then from (13) we get $t_2 = t_1^2$ and $Y_{t_2}/Y_{t_1} = 1$. In this case, by Lemma 1, we have

$$(19) \quad \frac{Y_{t_2}}{Y_{t_1}} = \left| \frac{\varepsilon^{t_1^2} - \bar{\varepsilon}^{t_1^2}}{\varepsilon^{t_1} - \bar{\varepsilon}^{t_1}} \right| = \left| \sum_{i=0}^{(t_1-1)/2} \begin{bmatrix} t_1 \\ i \end{bmatrix} (-4D_2)^{(t_1-1)/2-i} p^{Z_1 t_1 i} \right| = 1.$$

Since $(-1)^{(t_1-1)(p^{Z_1}+1)/4} = (-1)^{(t_1-1)(p^{Z_1 t_1}+1)/4}$, we get from (19) that

$$(20) \quad \sum_{i=0}^{(t_1-1)/2} \begin{bmatrix} t_1 \\ i \end{bmatrix} (-4D_2)^{(t_1-1)/2-i} p^{Z_1 t_1 i} = (-1)^{(t_1-1)(p^{Z_1}+1)/4}.$$

Further, since $t_1 \geq 3$, we see from (20) that

$$(21) \quad (-4D_2)^{(t_1-1)/2} - (-1)^{(t_1-1)(p^{Z_1}+1)/4} \equiv 0 \pmod{p^{3Z_1}}.$$

On the other hand, by (7), we have

$$(22) \quad \begin{aligned} & \left((-4D_2)^{(t_1-1)/2} - (-1)^{(t_1-1)(p^{Z_1}+1)/4} \right) + \begin{bmatrix} t_1 \\ 1 \end{bmatrix} (-4D_2)^{(t_1-3)/2} p^{Z_1} \\ & + \begin{bmatrix} t_1 \\ 2 \end{bmatrix} (-4D_2)^{(t_1-5)/2} p^{2Z_1} \equiv 0 \pmod{p^{3Z_1}}. \end{aligned}$$

The combination of (21) and (22) yields

$$(23) \quad t_1 \left(4D_2 - \left(\frac{t_1-3}{2} \right) p^{Z_1} \right) \equiv 0 \pmod{p^{2Z_1}}.$$

Since $p \nmid D_2$ and t_1 is an odd prime, (23) is impossible. Thus t_2 is an odd prime too.

By (10), we have

$$(24) \quad \varepsilon = p^{Z_1/2} e^{\theta\sqrt{-1}}, \quad \bar{\varepsilon} = p^{Z_1/2} e^{-\theta/\sqrt{-1}},$$

where θ is a real number satisfying

$$(25) \quad \sin \theta = \frac{\varepsilon - \bar{\varepsilon}}{2p^{Z_1/2}\sqrt{-1}} = \left(\frac{D_2}{p^{Z_1}} \right)^{1/2}.$$

Since $0 < D_2/p^{Z_1} < 1$, we may assume that θ satisfies

$$(26) \quad 0 < \theta < \frac{\pi}{2}.$$

Further, since $Y_{t_1} = Y_{t_2} = 1$ by (17), we get from (24) and (25) that

$$(27) \quad \sin t_j \theta = (-1)^{(t_j-1)(p^{Z_1}+1)/4} \frac{\sin \theta}{p^{Z_1(t_j-1)/2}}, \quad j = 1, 2.$$

Hence

$$(28) \quad t_j \theta = k\pi + (-1)^{k+(t_j-1)(p^{Z_1}+1)/4} \phi_j, \quad j = 1, 2,$$

where k is an integer, $\phi_j (j = 1, 2)$ are positive numbers satisfying

$$(29) \quad \sin \phi_j = \frac{\sin \theta}{p^{Z_1(t_j-1)/2}}, \quad 0 < \phi_j < \frac{\pi}{2}, \quad j = 1, 2.$$

Notice that $0 < \theta < \pi/2$ by (26). Since $0 < \phi_j < \theta < \pi/2$ ($j = 1, 2$) by (26) and (29), we see from (28) that k satisfies $1 \leq k \leq (t_j - 1)/2$. Thus, by (25), (28) and (29), we get

$$\begin{aligned} \left| \arcsin \left(\frac{D_2}{p^{Z_1}} \right)^{1/2} - \frac{k\pi}{t_j} \right| &= \left| \theta - \frac{k\pi}{t_j} \right| = \frac{\phi_j}{t_j} = \frac{\arcsin((\sin \theta)/p^{Z_1(t_j-1)/2})}{t_j} \\ &< \frac{\arcsin(p^{-Z_1(t_j-1)/2})}{t_j} < \frac{\pi}{2t_j p^{Z_1(t_j-1)/2}}, \quad j = 1, 2. \end{aligned}$$

The inequality (9) is proved.

From (25) and (27), we get

$$(30) \quad |\sin t_1 \theta| = \left(\frac{D_2}{p^{s_1}} \right)^{1/2}, \quad |\sin t_2 \theta| = \left(\frac{D_2}{p^{s_2}} \right)^{1/2}.$$

By (30), there exist suitable nonnegative integers k_1 and k_2 such that

$$(31) \quad |k_1 \pi - t_1 \theta| = \arcsin \left(\frac{D_2}{p^{s_1}} \right)^{1/2}, \quad |k_2 \pi - t_2 \theta| = \arcsin \left(\frac{D_2}{p^{s_2}} \right)^{1/2}.$$

Since $s_1 < s_2$ and $\arcsin(D_2/p^{s_1})^{1/2} > \arcsin(D_2/p^{s_2})^{1/2}$, we get from (31) that

$$(32) \quad 0 < \left| \frac{k_1}{t_1} - \frac{k_2}{t_2} \right| \pi \leq \frac{1}{t_1} \arcsin \left(\frac{D_2}{p^{s_1}} \right)^{1/2} + \frac{1}{t_2} \arcsin \left(\frac{D_2}{p^{s_2}} \right)^{1/2} < \frac{2}{t_1} \arcsin \left(\frac{D_2}{p^{s_1}} \right)^{1/2}.$$

Notice that $|k_1/t_1 - k_2/t_2| \geq 1/t_1 t_2$ if $k_1/t_1 \neq k_2/t_2$. We obtain (8) from (32). The lemma is proved.

Lemma 4. *If (3) has two solutions (r_1, s_1) and (r_2, s_2) satisfying (6) with $t_1 = 3$, then we have*

$$(33) \quad 4D_2 = 3p^{Z_1} + (-1)^{(p^{Z_1}-1)/2}$$

and

$$(34) \quad t_2 \geq 2p^{2Z_1} + 3.$$

Proof. Since $t_1 = 3$, we get (33) from (7). Let $\delta = (-1)^{(p^{Z_1}+1)/2}$. We get from (17) and (33) that

$$(-3p^{Z_1} + \delta)^{(t_2-1)/2} + t_2(-3p^{Z_1} + \delta)^{(t_2-3)/2}(3p^{Z_1}) \equiv \delta^{(t_2-1)/2} \pmod{p^{2Z_1}},$$

whence we obtain $(t_2 - 3)/2 \equiv 0 \pmod{p^{Z_1}}$. By (7) again, we get

$$(35) \quad -\left(\frac{t_2-3}{2}\right) + \left(9 \binom{(t_2-1)/2}{2} - 3 \begin{bmatrix} t_2 \\ 1 \end{bmatrix} \binom{(t_2-3)/2}{1} + \begin{bmatrix} t_2 \\ 2 \end{bmatrix}\right) \delta p^{Z_1} \equiv 0 \pmod{p^{2Z_1}}.$$

Notice that

$$\binom{(t_2-1)/2}{2} \equiv \binom{(t_2-3)/2}{1} \equiv \begin{bmatrix} t_2 \\ 2 \end{bmatrix} \equiv 0 \pmod{\frac{t_2-3}{2\lambda}},$$

where $\lambda = 1$ or 2 according to whether $t_2 \equiv 1 \pmod{4}$ or not. We find from (35) that $(t_2 - 3)/2 \equiv 0 \pmod{p^{2Z_1}}$. Since $t_2 > 3$, it implies (34). The lemma is proved.

Let α be an algebraic number of degree d with conjugates $\sigma_1\alpha, \sigma_2\alpha, \dots, \alpha_d\alpha$ and the minimal polynomial

$$a_0z^d + a_1z^{d-1} + \dots + a_d = a_0 \prod_{i=1}^d (z - \sigma_i\alpha) \in \mathbb{Z}[z], \quad a_0 > 0.$$

Then

$$h(\alpha) = \frac{1}{d} \left(\log a_0 + \sum_{i=1}^d \log \max(1, |\sigma_i\alpha|) \right)$$

is called the logarithmic absolute height of α .

Lemma 5 ([6, Théorème 3]). *Let α be an algebraic number with $|\alpha| = 1$. Let b_1, b_2 be positive integers, and let $\Lambda = b_1 \log \alpha - b_2 \pi \sqrt{-1}$. If α is not a root of unity, then we have*

$$\log |\Lambda| \geq -8.87AB^2,$$

where $A = \max(20, 10.98|\log \alpha| + rh(\alpha))$, $B = \max(17, r^{1/2/10}, 5.03 + 2.35r + r \log(b_1/68.9 + b_2/2A))$, $r = [\mathbb{Q}(\alpha) : \mathbb{Q}]/2$.

Lemma 6. *If (3) has two solutions (r_1, s_1) and (r_2, s_2) satisfying (6), then we have*

$$(36) \quad t_2 < 2 + 2563.43 \left(1 + \frac{10.98\pi}{\log p^{Z_1/2}} \right).$$

Proof. Under the assumption, we get from (15) and (17) that

$$(37) \quad 2p^{Z_1/2} > 2D_2^{1/2} |\varepsilon - \bar{\varepsilon}| = |\varepsilon^{t_2} - \bar{\varepsilon}^{t_2}|.$$

Let $\alpha = \varepsilon/\bar{\varepsilon}$. Then from (37) we get

$$(38) \quad \log 2p^{Z_1/2} > \log |\bar{\varepsilon}^{t_2}| + \log |\alpha^{t_2} - 1| = t_2 \log p^{Z_1/2} + \log |\alpha^{t_2} - 1|.$$

Since $t_2 \geq 5$, if $|\alpha^{t_2} - 1| \geq 1$, then (38) is impossible. Therefore, we have

$$(39) \quad |\alpha^{t_2} - 1| \geq |t_2 \log \alpha - k\pi\sqrt{-1}|$$

for some positive integers k with $k \leq t_2$. Let $\Lambda = t_2 \log \alpha - k\pi\sqrt{-1}$. From (38) and (39), we get

$$(40) \quad \log 2p^{Z_1/2} - \log |\Lambda| > t_2 \log p^{Z_1/2}.$$

We see from (10) that α satisfies

$$p^{Z_1}\alpha^2 - 2(D_1X_1^2 - D_2)\alpha + p^{Z_1} = 0.$$

It implies that $[\mathbb{Q}(\alpha) : \mathbb{Q}] = 2$, $h(\alpha) = \log p^{Z_1/2}$ and α is not a root of unity. Further, by (24) and (26), we have $|\alpha| = 1$ and $|\log \alpha| = 2\theta < \pi$. Since $k \leq t_2$, by Lemma 5, we get

$$(41) \quad \log |\Lambda| \geq -8.87(10.98\pi + \log p^{Z_1/2}) \\ \times \left(\max \left(17, 7.38 + \log \left(\frac{t_2}{68.9} + \frac{t_2}{2(10.98\pi + \log p^{Z_1/2})} \right) \right) \right)^2.$$

If

$$(42) \quad 7.38 + \log \left(\frac{t_2}{68.9} + \frac{t_2}{2(10.98\pi + \log p^{Z_1/2})} \right) \geq 17,$$

then we have

$$\log 0.0288015t_2 \geq 9.62,$$

whence we conclude that

$$(43) \quad t_2 > 523063.$$

On the other hand, by (40), (41) and (42), we get

$$(44) \quad 3 + 565.8842(7.38 + \log 0.0288015t_2)^2 > 1 + \frac{\log 2}{\log p^{Z_1/2}} + 8.87 \left(1 + \frac{10.98\pi}{\log p^{Z_1/2}} \right) \\ \left(7.38 + \log \left(\frac{t_2}{68.9} + \frac{t_2}{2(10.98\pi + \log p^{Z_1/2})} \right) \right)^2 > t_2.$$

We calculate from (44) that $t_2 < 150000$. It contradicts with (43). So we have

$$(45) \quad 7.38 + \log \left(\frac{t_2}{68.9} + \frac{t_2}{2(10.98\pi + \log p^{Z_1/2})} \right) < 17.$$

Thus, by (40), (41) and (45), we obtain (36). The lemma is proved.

Lemma 7 ([3]). *The equation*

$$X^3 + X^2Y - 2XY^2 - Y^3 = 1, \quad X, Y \in \mathbb{Z}$$

has only the solutions $(X, Y) = (1, 0), (0, -1), (-1, 1), (2, -1), (-1, 2), (5, 4), (4, -9)$ *and* $(-9, 5)$.

Proof of Theorem 1. We now suppose that $N(D_1, D_2, p) > 2$. By Lemma 3, then (3) has two solutions (r_1, s_1) and (r_2, s_2) satisfying (6).

By Lemmas 4 and 6, if $t_1 = 3$, then we have

$$(46) \quad 2p^{2Z_1} + 3 \leq t_2 < 2 + 2563.43 \left(1 + \frac{10.98\pi}{\log p^{Z_1/2}} \right),$$

whence we conclude that $p^{Z_1} \leq 137$. Recall that $D_1 > 1$, $D_2 > 1$ and D_1 is square free. By (33), we only need to consider the cases $(D_1, D_2, p^{Z_1}) = (2, 5, 7), (2, 7, 9), (3, 8, 11), (3, 10, 13), (5, 14, 19), (6, 17, 23), (6, 19, 25), (7, 20, 27), (7, 22, 29), (2, 23, 31), (10, 31, 41), (11, 32, 43), (3, 35, 47), (3, 37, 49), (13, 40, 53), (15, 44, 59), (15, 46, 61), (17, 50, 67), (2, 53, 71), (2, 55, 73), (5, 59, 79), (5, 61, 81), (21, 62, 83), (22, 67, 89), (6, 73, 97),$

(26,77,103), (3,80,107), (3,82,109), (7,85,113), (30,91,121), (31,94,125), (2,95,127), (33,98,131) and (34,103,137). On the other hand, by (9) and (33), we get

$$(47) \quad \left| \arcsin \left(\frac{3}{4} + \frac{(-1)^{(p^{Z_1}-1)/2}}{4p^{Z_1}} \right)^{1/2} - \frac{k\pi}{t_2} \right| < \frac{\pi}{2t_2 p^{Z_1(t_2-1)/2}}$$

for some positive integers k with $k \leq (t_2 - 1)/2$. However, by (46), (47) is false for the above-mentioned cases.

If $t_1 = 5$, then from (7) we get

$$(48) \quad 16D_2^2 - 20D_2p^{Z_1} + 5p^{2Z_1} = 1.$$

Since p^{Z_1} is an odd prime power, we see from (48) that $4D_2 \equiv \lambda \pmod{p^{Z_1}}$, where $\lambda \in \{-1, 1\}$. So we have $4D_2 = kp^{Z_1} + \lambda$, where $k = 1$ or 3 . Hence, by (48), we get

$$p^{Z_1}(k^2 - 5k + 5) = (5 - 2k)\lambda.$$

This implies that $p^{Z_1} = 3$. Since D_2 is an integer with $D_2 > 1$, this is impossible.

If $t_1 = 7$, then we have

$$(49) \quad (2p^{Z_1} - 4D_2)^3 + (2p^{Z_1} - 4D_2)^2 p^{Z_1} - 2(2p^{Z_1} - 4D_2)p^{2Z_1} - p^{3Z_1} = \pm 1$$

by (7). However, by Lemma 7, (49) is impossible.

From (8) and (36), if $t_1 > 7$, then we have $t_1 \geq 11$ and

$$\begin{aligned} p^{5Z_1} &\leq p^{Z_1(t_1-1)/2} < \left(\frac{p^{Z_1 t_1}}{D_2} \right)^{1/2} < \frac{\pi}{2 \arcsin(D_2/p^{Z_1 t_1})^{1/2}} < t_2 \\ &< 2 + 2563.42 \left(1 + \frac{10.98\pi}{\log p^{Z_1/2}} \right), \end{aligned}$$

whence we calculate that $p^{Z_1} \leq 9$. Since, if $(D_1, D_2, p^{Z_1}) = (2, 5, 7)$ or $(2, 7, 9)$, then $t_1 = 3$. So we only need to consider the cases $(D_1, D_2, p^{Z_1}) = (2, 3, 5)$, $(3, 2, 5)$, $(3, 4, 7)$, $(5, 2, 7)$, $(5, 4, 9)$ and $(7, 2, 9)$. For the above-mentioned cases, (7) is false if $t_1 = 11$ or 13 . So we have $t_1 \geq 17$. Then, by (8) and (36) again, we get

$$390625 \leq p^{8Z_1} \leq p^{Z_1(t_1-1)/2} < t_2 < 2 + 2563.42 \left(1 + \frac{10.98\pi}{\log 5^{1/2}} \right) < 112451,$$

a contradiction. All cases have been considered, the proof is complete.

3. PROOF OF THEOREM 2

By the same method as in the proofs of Lemmas 2, 3, 4 and 6, we can prove the corresponding lemmas about the equation (4) without any difficulty.

Lemma 8. *If the equation*

$$(50) \quad D_1 X'^2 + D_2 Y'^2 = 2p^{Z'}, \quad X', Y', Z' \in \mathbb{Z}, \quad \gcd(X', Y') = 1, \quad Z' > 0,$$

has solutions (X', Y', Z') , then it has a unique solution (X'_1, Y'_1, Z'_1) such that $X'_1 > 0$, $Y'_1 > 0$ and $Z'_1 \leq Z'$, where Z' runs through all solutions (X', Y', Z') of (50). (X'_1, Y'_1, Z'_1) is called the least solution of (50). Moreover, every solution (X', Y', Z') of (50) can be expressed as

$$\begin{aligned} Z' &= Z'_1 t', \quad \frac{X' \sqrt{D_1} + Y' \sqrt{-D_2}}{\sqrt{2}} = \lambda_1 \left(\frac{X'_1 \sqrt{D_1} + \lambda_2 Y'_1 \sqrt{-D_2}}{\sqrt{2}} \right)^{t^0}, \\ &\quad t' \in \mathbb{N}, \quad 2 \nmid t', \quad \lambda_1, \lambda_2 \in \{-1, 1\}. \end{aligned}$$

Lemma 9. *The equation (4) has solutions (r', s') if and only if (50) has solutions (X', Y', Z') and its least solution (X'_1, Y'_1, Z'_1) satisfies $Y'_1 = 1$. Moreover, if $N'(D_1, D_2, p) > 2$, then (4) has two solutions (r'_1, s'_1) and (r'_2, s'_2) such that*

$$(51) \quad s'_1 = Z'_1 t'_1, \quad s'_2 = Z'_1 t'_2, \quad 1 < t'_1 < t'_2,$$

where t'_1 and t'_2 are odd primes satisfying

$$(52) \quad \sum_{i=0}^{(t'_j-1)/2} \begin{bmatrix} t'_j \\ i \end{bmatrix} (-2D_2)^{(t'_j-1)/2-i} p^{Z'_1 i} = -(-1)^{(t'_j-1)(p^{Z'_1}+1)/4}, \quad j = 1, 2,$$

$$(53) \quad \frac{\pi}{2 \arcsin(D_2/2p^{s'_1})^{1/2}} < t'_2$$

and

$$(54) \quad \left| \arcsin \left(\frac{D_2}{2p^{Z'_1}} \right)^{1/2} - \frac{k\pi}{t'_j} \right| < \frac{\pi}{2t'_j p^{Z'_1(t'_j-1)/2}}, \quad j = 1, 2,$$

where k is a positive integer satisfying $k \leq (t'_j - 1)/2$.

Lemma 10. *If (4) has two solutions (r'_1, s'_1) and (r'_2, s'_2) satisfying (51) with $t'_1 = 3$, then we have*

$$(55) \quad 2D_2 = 3p^{Z'_1} + (-1)^{(p^{Z'_1}+1)/2}$$

and

$$(56) \quad t'_2 \geq 2p^{2Z'_1} + 3.$$

Lemma 11. *If (4) has two solutions (r'_1, s'_1) and (r'_2, s'_2) satisfying (51), then we have*

$$(57) \quad t'_2 < 3 + 2563.43 \left(1 + \frac{10.98\pi}{\log p^{Z'_1/2}} \right).$$

Proof of Theorem 2. We now suppose that $N'(D_1, D_2, p) > 2$. By Lemma 9, then (4) has two solutions (r'_1, s'_1) and (r'_2, s'_2) satisfying (51).

By Lemmas 10 and 11, if $t'_1 = 3$, then we have

$$(58) \quad 2p^{2Z'_1} + 3 \leq t'_2 < 3 + 2563.43 \left(1 + \frac{10.98\pi}{\log p^{Z'_1/2}} \right),$$

whence we calculate that $p^{Z'_1} \leq 137$. Hence, by (55), we only need to consider the cases $(D_1, D_2, p^{Z'_1})$ such that $p^{Z'_1} \leq 137$ and $2D_2 = 3p^{Z'_1} + (-1)^{(p^{Z'_1}+1)/2}$. On the other hand, by (54) and (55), we get

$$(59) \quad \left| \arcsin \left(\frac{3}{4} + \frac{(-1)^{(p^{Z'_1}+1)/2}}{4p^{Z'_1}} \right)^{1/2} - \frac{k\pi}{t'_2} \right| < \frac{\pi}{2t'_2 p^{Z'_1(t'_2-1)/2}}$$

for some positive integers k with $k \leq (t'_2 - 1)/2$. However, for the above-mentioned cases, if t'_2 satisfies (58), then (59) is impossible.

If $t'_1 = 5$, then from (52) we get

$$4D_2^2 - 10D_2 p^{Z'_1} + 5p^{2Z'_1} = -1.$$

It implies that

$$(60) \quad (4D_2 - 5p^{Z'_1})^2 - 5p^{2Z'_1} = -4.$$

For any nonnegative integer m , let F_m and L_m denote the m th Fibonacci number and the m th Lucas number, respectively. Notice that all solutions (X, Y) of the equation

$$X^2 - 5Y^2 = -4, \quad X, Y \in \mathbb{N}, \quad \gcd(X, Y) = 1$$

are given by $(X, Y) = (L_{6l}, F_{6l+1})$ and (L_{6l+5}, F_{6l+5}) , where l runs through all nonnegative integers. We see from (60) that

$$(61) \quad (D_2, p^{Z'_1}) = (\tfrac{1}{2}L_{6l}, F_{6l+1}) \quad \text{or} \quad (\tfrac{1}{2}L_{6l+5}, F_{6l+5}).$$

On the other hand, by (53) and (57), we have

$$(62) \quad p^{2Z'_1} < \left(\frac{2p^{5Z'_1}}{D_2} \right)^{1/2} < \frac{\pi}{2 \arcsin(D_2/2p^{5Z'_1})^{1/2}} < t'_2 < 3 + 2563.43 \left(1 + \frac{10.98\pi}{\log p^{Z'_1/2}} \right),$$

whence we calculate that $p^{Z'_1} < 200$. Therefore, we find from (61) that $(D_1, D_2, p^{Z'_1}) = (1, 9, 5)$, $(17, 9, 13)$ and $(17, 161, 89)$. Further, by (54), we get

$$(63) \quad \left| \arcsin \left(\frac{D_2}{2p^{Z'_1}} \right)^{1/2} - \frac{k\pi}{t'_2} \right| < \frac{\pi}{2t'_2 p^{Z'_1(t'_2-1)/2}}$$

for some positive integers k with $k \leq (t'_2 - 1)/2$. However, if t'_2 satisfies (62), then (63) is impossible.

If $t'_2 = 7$, then from (52) we get

$$(64) \quad (2p^{Z'_1} - 2D_2)^3 + (2p^{Z'_1} - 2D_2)^2 p^{Z'_1} - 2(2p^{Z'_1} - 2D_2)p^{2Z'_1} - p^{3Z'_1} = (-1)^{(p^{Z'_1}+1)/2}.$$

By Lemma 7, we obtain from (64) that $(D_1, D_2, p^{Z'_1}) = (7, 11, 9)$. For this case, by (53), (54) and (57), we get

$$(65) \quad 729 < t'_2 < 83099$$

and

$$(66) \quad \left| \arcsin \left(\frac{11}{18} \right)^{1/2} - \frac{k\pi}{t'_2} \right| < \frac{\pi}{2 \cdot 3^{t'_2-1} t'_2}$$

for some positive integers k with $k \leq (t'_2 - 1)/2$. However, (66) is impossible if t'_2 satisfies (65).

By (53) and (57), if $t'_1 > 7$, then $t'_1 \geq 11$ and $p^{Z'_1} \leq 9$. Except for the already considered cases, we have $(D_1, D_2, p^{Z'_1}) = (5, 1, 3)$, $(1, 9, 5)$, $(9, 1, 5)$, $(1, 13, 7)$, $(5, 9, 7)$, $(9, 5, 7)$, $(11, 3, 7)$, $(13, 1, 7)$, $(1, 17, 9)$, $(11, 7, 9)$, $(13, 5, 9)$ or $(17, 1, 9)$. For these cases, (52) is false when $t'_1 = 11, 13, 17$ and 19 . So we have $t'_1 \geq 23$. Then, by (53) and (57) again, we get

$$\begin{aligned} 177147 \leq p^{11Z'_1} \leq p^{Z'_1(t'_1-1)/2} &< \frac{\pi}{2 \arcsin(D_2/2p^{s'_1})^{1/2}} < t'_2 < 3 + 2563.43 \\ &\times \left(1 + \frac{10.98\pi}{\log 3^{1/2}} \right) < 153544, \end{aligned}$$

a contradiction. The theorem is proved.

4. PROOF OF THE COROLLARY

Lemma 12 ([11]). *The equation*

$$z^2 + 4 = y^n, \quad x, y, n \in \mathbb{N}, \quad 2 \nmid y, \quad n > 3,$$

has no solution (z, y, n) .

Proof of Corollary. Let (x, y, m, n) be a solution of (2) such that $m = 3$ and y is a prime power. Then we have

$$(67) \quad (y - 1)(2x + 1)^2 + (3y + 1) = 4y^n, \quad n > 3.$$

If $y = 2$, then from (67) we get

$$(68) \quad (2x + 1)^2 + 7 = 2^{n+2}, \quad n > 3.$$

By [12], we find from (68) that $(x, y, m, n) = (5, 2, 3, 5)$ and $(90, 2, 3, 13)$.

If $y = 2^k$, where k is a positive integer with $k > 1$, then

$$(69) \quad (2^k - 1)(2x + 1)^2 + (3 \cdot 2^k + 1) = 2^{kn+2}, \quad n > 3.$$

Let $D_1 = 2^k - 1$, $D_2 = 3 \cdot 2^k + 1$ and $p = 2$. We see from (69) that $(r, s) = (2x + 1, kn + 2)$ is a solution of (3) with $s > 3k + 2$. However, by [8], then (3) has exactly two solutions $(r, s) = (1, k + 2)$ and $(2^{k+1} + 1, 3k + 2)$. Therefore, (69) is impossible.

If $2 \nmid y$, then $y = p^k$, where p is an odd prime and k is a positive integer.

If $y = 5$, then from (67) we get

$$(70) \quad (2x + 1)^2 + 4 = 5^n, \quad n > 3.$$

By Lemma 12, (70) is impossible.

If $y \equiv 1 \pmod{4}$ and $y > 5$, then we have

$$(71) \quad \left(\frac{p^k - 1}{4} \right) (2x + 1)^2 + \left(\frac{3p^k + 1}{4} \right) = p^{kn}, \quad n > 3.$$

Let $D_1 = (p^k - 1)/4$ and $D_2 = (3p^k + 1)/4$. We see from (71) that $(r, s) = (2x + 1, kn)$ is a solution of (3) with $s > 3k$. Notice that (3) has two solutions $(r, s) = (1, k)$ and $(2p^k + 1, 3k)$ in this case. Therefore, by Theorem 1, (71) is impossible.

Similarly, if $y \equiv 3 \pmod{4}$, then we have

$$(72) \quad \left(\frac{p^k - 1}{2} \right) (2x + 1)^2 + \left(\frac{3p^k + 1}{2} \right) = 2p^{kn}, \quad n > 3.$$

Let $D_1 = (p^k - 1)/2$ and $D_2 = (3p^k + 1)/2$. Since (4) has two solutions $(r', s') = (1, k)$ and $(2p^k + 1, 3k)$ in this case, by Theorem 2, (72) is impossible. Now, the corollary is proved.

ACKNOWLEDGEMENTS

The author is grateful to the referees for their valuable suggestions.

REFERENCES

- [1] R. Apéry, *Sur une équation diophantienne*, C. R. Acad. Sci. Paris Sér. A **251** (1960), 1263–1264. MR **22**:10951
- [2] R. Apéry, *Sur une équation diophantienne*, C. R. Acad. Sci. Paris Sér. A **251** (1960), 1451–1452. MR **22**:10950
- [3] V. I. Baulin, *On an indeterminate equation of the third degree with least positive discriminant*, Tul'sk Gos. Ped. Inst. Učen. Zap. Fiz. Mat. Nauk Vyp. **7** (1960), 138–170. (in Russian) MR **33**:7298
- [4] R. Goormaghtigh, *L'intermédiaire des Mathématiciens*, **24** (1917), 88.
- [5] R. K. Guy, *Unsolved Problems in Number Theory*, Springer-Verlag, New York, 1981. MR **83k**:10002
- [6] M. Laurent, M. Mignotte and Y. Nesterenko, *Formes linéaires en deux logarithmes et déterminants d'interpolation*, J. Number Theory, **55** (1995), 285–321. MR **96h**:11073
- [7] M.-H. Le, *The divisibility of the class number for a class of imaginary quadratic fields*, Kexue Tongbao, **32** (1987), 724–727. (in Chinese) CMP 20:01
- [8] M.-H. Le, *The diophantine equation $D_1x^2 + D_2 = 2^{n+2}$* , Acta Arith., **64** (1993), 29–41. MR **94**:11030
- [9] M.-H. Le, *A note on the generalized Ramanujan-Nagell equation*, J. Number Theory, **50** (1995), 193–201. MR **96f**:11051
- [10] R. Lidl and H. Niederreiter, *Finite Fields*, Addison-Wesley, Reading, MA, 1983. MR **86c**:11106
- [11] T. Nagell, *Contributions to the theory of a category of diophantine equations of the second degree with two unknowns*, Nova Acta Soc. Sci. Upsal., (4) **16**, no. 2, 38pp. (1955). MR **17**:13b
- [12] T. Nagell, *The diophantine equation $x^2 + 7 = 2^n$* , Arkiv. Mat., **4** (1960), 185–187.
- [13] R. Ratat, *L'Intermédiaire des Mathématiciens*, **23** (1916), 150.
- [14] T. N. Shorey, *Some exponential diophantine equations*, in *Number Theory and Related Topics*, pp. 217–229, Bombay, 1989. MR **98d**:11038

DEPARTMENT OF MATHEMATICS, ZHANJIANG TEACHERS COLLEGE, POSTAL CODE 524048, ZHAN-
JIANG, GUANGDONG, P. R. CHINA