

# THE MONOPOLE EQUATIONS AND $J$ -HOLOMORPHIC CURVES ON WEAKLY CONVEX ALMOST KÄHLER 4-MANIFOLDS

YUTAKA KANDA

ABSTRACT. We prove that a weakly convex almost Kähler 4-manifold contains a compact, non-constant  $J$ -holomorphic curve if the corresponding monopole invariant is not zero and if the corresponding line bundle is non-trivial.

## 0. INTRODUCTION

The theory of pseudo holomorphic curves has been bringing remarkable progress to both symplectic topology and contact topology since it was initiated by Gromov in [Gr].

On the other hand, Witten introduced the monopole equations and defined a new invariant of closed orientable smooth 4-manifolds in [W]. Further, he showed that if the 4-manifold  $X$  is Kähler, the computation of its invariant can be easily done by using algebraic geometry. The key is the fact that there is a some kind of correspondence between the solutions of the monopole equations on  $X$  and the divisors of  $X$ .

After that, Taubes showed in [T1], [T2], [T3] that the monopole invariant of a closed symplectic 4-manifold  $(X, \omega)$  with  $b_2^+ > 1$  is equivalent to its Gromov-Witten invariant that counts the “number” of codimension-1 symplectic submanifolds contained in it.

After that, Kronheimer and Mrowka [K-M2] introduced a suitable analytic setting for the monopole equations on a certain class of non-compact almost Kähler 4-manifolds called A.F.A.K. and extended the definition of monopole invariants to them. Further, as an application, they obtained a striking result on symplectically fillable contact 3-manifolds.

Our main aim is to extend the main result in [T1] to *weakly convex* almost Kähler manifolds, which are non-compact in general by the definition. Namely, such a manifold contains a compact, non-constant  $J$ -holomorphic curve if the corresponding monopole invariant is non-zero and if the corresponding line bundle is non-trivial. See Theorem 4.1 in Section 4 for the precise statement. The notion of weak convexity is a slightly stronger condition than that of A.F.A.K. See Definition 1.1.

Further, in Section 10 we give an application of the main result to contact topology. See Theorem 10.1.

---

Received by the editors March 8, 1999 and, in revised form, February 28, 2000.

2000 *Mathematics Subject Classification*. Primary 57R57.

*Key words and phrases*. Symplectic structure, monopole equation,  $J$ -holomorphic curve.

The author would like to thank K.Ono and H.Ohta for several helpful discussions and encouragement.

# 1. THE MONOPOLE INVARIANTS OF WEAKLY CONVEX ALMOST KÄHLER 4-MANIFOLDS

Let  $(X, \omega)$  be a symplectic manifold. An almost complex structure  $J$  is said to be *compatible* and the triple  $(X, \omega, J)$  is called *almost Kähler* if the bilinear form  $g(*, *) := \omega(*, J*)$  is a  $J$ -invariant Riemannian metric. It is well known that the space of smooth almost complex structures is contractible under a suitable choice of topology, such as the Whitney topology.

In this paper, we will mainly work on *weakly convex* almost Kähler 4-manifolds which are defined as follows:

**Definition 1.1.** An almost Kähler manifold  $(X, \omega, J)$  is *weakly convex* if there exists a proper function  $\sigma : X \mapsto [h, \infty)$  with  $h > 0$  which has the following properties:

**Property (A).** Any  $x \in X$  obeys the conditions below.

1. The injective radius at  $x$  is no less than  $\sigma(x)$ .
2. Let  $e_x$  be the map  $e_x : TX_x \mapsto X$  defined by  $e_x(v) := \exp_x(\sigma(x)v)$  and let  $\gamma_x$  be the Riemannian metric on the unit ball in  $TX_x$  defined by  $\gamma_x := \frac{e_x^*(g)}{\sigma(x)^2}$ . There exists a sequence of non-negative constants  $\{c_k\}_{k \in \mathbb{N}}$  which is independent of  $x$  such that the  $C^0$  norm of the covariant derivatives of order  $k$  of  $\gamma_x$  is bounded by  $c_k$  for each  $k \in \mathbb{N}$ .
3. Let  $o_x$  be the 2-form on the unit ball defined by  $o_x := \frac{e_x^*(\omega)}{\sigma(x)^2}$ . There exists a sequence of non-negative constants  $\{c'_k\}_{k \in \mathbb{Z}^{\geq 0}}$  which is independent of  $x$  such that the  $C^0$  norm of the covariant derivatives of order  $k$  of  $o_x$  is bounded by  $c'_k$  for each  $k \in \mathbb{Z}^{\geq 0}$ .
4. Let  $\hat{\sigma}_x$  be the function on the unit ball defined by  $\hat{\sigma}_x := \frac{e_x^*(\sigma)}{\sigma(x)}$ . There exists a positive constant  $\acute{c}$  which is independent of  $x$  such that  $\hat{\sigma}_x \geq \acute{c}$ .

**Property (B).** There exists a non-negative, integrable function  $g_\sigma$  of  $\mathbb{R}^{\geq 0}$  such that

$$\int_{\mathbb{R}^{\geq 0}} f g_\sigma dy = \int_X f \circ \sigma d \text{vol}_X$$

for an arbitrary function  $f \in C_0^\infty(\mathbb{R}^{\geq 0})$ . Moreover, there exist constants  $C > 0, \epsilon_0 > 0$  such that  $g_\sigma \leq C y^{\epsilon_0}$ . Notice that  $g_\sigma \equiv 0$  on  $[0, h)$ .

*Remark 1.2.* The condition of weak convexity is stronger than that of A.F.A.K. manifolds dealt with in [K-M2].

A typical example of weakly convex almost Kähler 4-manifolds is described below:

Let  $X$  be an orientable 4-manifold endowed with conical end, namely,  $X$  is diffeomorphic to  $X_0 \cup_\phi \partial X_0 \times [1, \infty)$  where  $X_0$  is compact with smooth boundary and  $\phi$  means the natural identification. Denote  $\partial X_0 \times [1, \infty) \subset X$  by  $X^+$  and  $\partial X_0$  by  $M$ . Let  $\omega$  be a symplectic form of  $X$  which restricts to  $X^+$  as the symplectization of some contact form  $\alpha$  of  $M$ , that is,  $\omega|_{X^+} = d(t^2\alpha)$ . Denote by  $e_0$  the Reeb vector field of  $\alpha$ . If one chooses a Cauchy-Riemann structure  $J'$  of the contact plane field  $\zeta := \text{Ker}(\alpha)$  so that  $J'$  is compatible with  $d\alpha|_\zeta$ , this would induce an

almost complex structure  $J$  compatible with  $\omega$  over  $X^+$ . In fact, if we identify  $TX^+$  with  $TM \oplus T\mathbb{R}^{\geq 1}$ ,  $J$  is determined by the rules that  $J(\partial_t) = e_0$ ,  $J(e_0) = -\partial_t$  and that  $J(Y) = J'(Y)$  if  $Y$  is tangent to  $\zeta$ . Then  $J$  extends to the interior of  $X_0$  so that it is compatible with  $\omega$ . We can easily check that the pair  $(\omega, J)$  satisfies all the conditions of weak convexity. In fact, let  $f$  be a smooth Morse function of  $X$  such that  $f|_{X^+} = t$  and  $f \geq \frac{1}{2}$  on  $X$ . We may define the function  $\sigma$  by  $\sigma := \kappa f$  where  $\kappa$  is a sufficiently small positive constant.

This example brings us to the following definition.

**Definition 1.2** ([E]). A symplectic 4-manifold  $(X_0, \omega)$  is a *symplectic filling* of a contact 3-manifold  $(M, \zeta)$  if  $(X_0, \omega)$  satisfies the following:

1.  $X_0$  is compact.
2.  $\partial X_0 = M$  as oriented manifolds.
3.  $\omega|_\zeta$  is non-degenerate.

Recall that in general a contact structure induces a canonical orientation to the base manifold if its dimension is  $4n+3$ . (But in this case, there is no canonical orientation for the contact plane field.)

**Proposition 1.3** ([K-M2]). *Let  $(X_0, \omega)$  be a symplectic filling of a contact manifold  $(M, \zeta)$ . We can construct a weakly convex almost Kähler 4-manifold  $(X, \tilde{\omega}, J)$  which admits an embedding  $\iota : (X_0, \omega) \hookrightarrow (X, \tilde{\omega})$  such that  $\zeta$  is invariant under the action of  $\iota^*(J)$  and such that  $X - \iota(\text{Int}X_0)$  is diffeomorphic to  $\partial X_0 \times [1, \infty)$ . Moreover, this construction is unique in the sense that two such weakly convex almost Kähler structures can be connected by a smooth 1-parameter family. In particular, if  $\omega|_M$  is exact,  $(X, \tilde{\omega}, J)$  can be made so that it is as described above as a typical example.*

Let  $(X, \omega, J)$  be an almost Kähler 4-manifold.  $X$  has a  $\text{Spin}^c$  structure  $s_\omega$  that is determined canonically by  $\omega$ . With this understood, define the set  $S(X, \omega)$  as follows:

**Definition 1.4.**  $S(X, \omega)$  consists of the isomorphism classes of the pairs  $(s, \varrho)$  where  $s$  is a  $\text{Spin}^c$  structure of  $X$  being identified with  $s_\omega$  outside some compact set through the isomorphism  $\varrho$ .

As we will see later,  $S(X, \omega)$  can be identified with the set of isomorphism classes of complex line bundles that have trivializations outside some compact sets. See Section 3.

If  $(X, \omega, J)$  is weakly convex, it is A.F.A.K. by the very definition. Therefore, following [K-M2], we can define its *monopole invariant*. In our terminology, this invariant is a map  $SW : \{(X, \omega, J, s, \varrho)\} \mapsto \mathbb{Z}$  obeying the properties below.

**Property (1).**  $SW(X, \omega_0, J_0, s, \varrho_0) = \pm SW(X, \omega_1, J_1, s, \varrho_1)$  if there exists a smooth 1-parameter family  $\{(\omega_t, J_t, \varrho_t)\}_{0 \leq t \leq 1}$  outside some compact set  $K$  such that  $\varrho_t : s|_K \mapsto s_{\omega_t}|_K$  are isomorphisms and such that  $(\omega_t, J_t)$  are almost Kähler structures being weakly convex in the following sense: For some compact  $K'$  with  $K \subset \text{Int}K'$ , there exists a family of proper functions  $\sigma_t : X \setminus \text{Int}K' \mapsto [h, \infty)$  with  $h > 0$  such that  $(\omega_t, J_t, \sigma_t)$  satisfies Property (A) for any  $x \in X \setminus \text{Int}K'$  and Property (B) with  $X$  replaced by  $X \setminus \text{Int}K'$ .

In a word, the invariant up to sign depends only on the choice of a  $\text{Spin}^c$  structure and on the “boundary condition”.

**Property (2).**  $SW(X, \omega, J, s_\omega, \text{id}) = 1$ .

**Property (3).** Suppose that  $SW(X, \omega, J, s, \varrho) \neq 0$ . Then  $\langle c_1^2(L_s, \tilde{\varrho}) - c_1(L_s, \tilde{\varrho}) \cup c_1(K), [X] \rangle = 0$ . Furthermore,  $\langle c_1(L_s, \tilde{\varrho}) \cup [\omega], [X] \rangle \geq 0$  with equality only if  $(s, \varrho) \cong (s_\omega, \text{id})$ .

Here  $L_s$  stands for the corresponding line bundle to  $s$  and  $\tilde{\varrho}$  is the trivialization of  $L_s$  induced by  $\varrho$  outside a compact set. The first Chern class of  $L_s$  is regarded as an element of the compactly support cohomology group of  $X$  through  $\tilde{\varrho}$ . Similarly,  $[X]$  denotes the generator of the fourth homology group of a locally finite singular chain over  $\mathbb{Z}$  whose orientation is compatible with  $\omega$ .  $K$  in (3) denotes the canonical line bundle of  $(X, J)$ .

*Remark 1.5.* Proposition 1.3 means that we can well define  $SW$  for the pair of contact 3-manifolds and its symplectic filling.

## 2. MONOPOLE EQUATIONS ON SYMPLECTIC 4-MANIFOLDS

We will review some basic facts about monopole equations, especially those on symplectic manifolds.

1. Let  $(X, g)$  be a Riemannian 4-manifold. A monopole equation on  $(X, g)$  is a non-linear P.D.E. depending on the choice of a  $\text{Spin}^c$  structure  $s$  of  $X$ . So we will review the definitions of  $\text{Spin}^c$  structures and Dirac operators first.

a)  $\text{Spin}^c$  structure  $s$  is a  $\text{Spin}^c(4) = \frac{\text{Spin}(4) \times \text{U}(1)}{\pm 1}$  lift of the oriented orthonormal frame bundle  $\text{Fr}(TX)$ . Through the standard representations of  $\text{Spin}^c(4)$ ,  $s$  associates the positive (resp. negative) spinor bundle  $W_s^+$  (resp.  $W_s^-$ ).  $W_s^\pm$  is a complex, Hermitian, rank-2 vector bundle endowed with the linear map  $\rho : TX \mapsto \text{Hom}(W_s^+, W_s^-)$  called Clifford multiplication that obeys the relation  $\rho(v)^* \circ \rho(v) = -g(v)\text{id}$ . The signs of  $W_s^\pm$  are canonically determined by the orientation of  $X$ .

The  $\text{Spin}^c(4)$  group appears as the structure group of a 4-tuple  $(TX, W_s^+, W_s^-, \rho)$ , which is just the central extension by  $\text{U}(1)$  of the structure group  $\text{SO}(4)$  of  $TX$ . Thus we can recover the principle bundle  $s$  from the 4-tuple according to the standard argument. Therefore,  $\text{Spin}^c$  structures are in one-to-one correspondence with the isomorphism classes of spinor bundles.

b) A  $\text{Spin}^c(4)$  connection of  $s$  is said to be compatible if the associated connection of  $\text{Fr}(TX)$  agrees with the Levi-Civita connection. Let  $\nabla^W$  be a  $\text{U}(2) \times \text{U}(2)$  connection on the spinor bundle  $W_s := W_s^+ \oplus W_s^-$ . Then  $\nabla^W$  is a  $\text{Spin}^c(4)$  connection if and only if the subbundle  $\rho(TX) \subset \text{Hom}(W_s^+, W_s^-)$  is preserved by the induced connection  $\nabla^{\text{Hom}}$  and is compatible if and only if  $\nabla^{\text{Hom}}|_{\rho(TX)}$  agrees with (the push-forward of) the Levi-Civita connection.

The splitting  $\mathfrak{spin}^c(4) = \mathfrak{so}(4) \oplus \mathfrak{u}(1)$  implies that a  $\text{Spin}^c(4)$  connection is determined by choosing a  $\text{U}(1)$  connection of the determinant line bundle  $L_s := \det(W_s^+) (= \det(W_s^-))$ . Therefore, the space of a compatible  $\text{Spin}^c$  connection is an affine space modelled by the space of pure imaginary 1-forms.

c) With a compatible  $\text{Spin}^c(4)$  connection  $\nabla_B$  given, where  $B$  stands for the corresponding  $\text{U}(1)$  connection of the determinant line bundle, the Dirac operator  $\mathcal{D}_B$  is defined to be the composition of the sequence

$$(2.1) \quad \Gamma(W_s^+) \xrightarrow{\nabla_B} \Gamma(TX^* \otimes W_s^+) \xrightarrow{\text{Cont}(\rho \otimes \text{id})} \Gamma(W_s^-),$$

where we identify  $TX$  and  $T^*X$  with each other and  $\text{Cont}$  stands for the contraction.

**d)** The monopole equation for a chosen  $\text{Spin}^c$  structure  $s$  is the one with variables  $(\Phi, B) \in \Gamma(W_s^+) \times \mathcal{A}(L_s)$  written as follows:

$$(2.2.1) \quad \mathcal{D}_B \Phi = 0,$$

$$(2.2.2) \quad \rho(F_B^+) = (\Phi \Phi^*)_0.$$

Here  $\mathcal{A}(L_s)$  denotes the space of  $U(1)$  connections of the determinant line bundle,  $F_B^+$  is the self-dual part of the curvature 2-form of  $B$  and  $\rho: \Lambda^2 \mapsto \text{End}(W_s)$  is the natural extension of the Clifford multiplication. The subscript ‘0’ means the traceless part of the said endomorphism.

This equation is equivariant under the action of the gauge group

$$\mathcal{G} := \text{Map}(X, U(1))$$

which acts on  $\Gamma(W_s^+)$  by the multiplication of a complex number and on  $\mathcal{A}(L_s)$  by the pulling-back of connections. We can regard  $\mathcal{G}$  as the subgroup of the bundle automorphism of  $W_s$  that respects the Clifford multiplication.

Notice that (2.2.2) consists of gauge invariant terms. If we add an arbitrary pure imaginary self-dual 2-form to the right-hand side of (2.2.2) to perturb the equation, it remains gauge equivariant.

**2.** Let  $(X, \omega, J)$  be an almost Kähler 4-manifold. Denote by  $g_J$  the corresponding Riemannian metric. We will see that the monopole equations (2.2) on the Riemannian manifold  $(X, g_J)$  can be written in terms of differential forms and Dolbeaut operators.

**a)** There is a  $\text{Spin}^c$  structure  $s_\omega$  canonically determined by  $\omega$ . This derives from the fact that the natural projection homomorphism  $\text{pr}: \text{Spin}^c(4) \mapsto \text{SO}(4)$  has a canonical inverse homomorphism over the subgroup  $U(2) \subset \text{SO}(4)$ . The spinor bundle and the Clifford multiplication for  $s_\omega$  can be explicitly written in terms of differential forms. In fact, define  $W_{s_\omega}^+ := \Lambda^{0,0} \oplus \Lambda^{0,2}$  and  $W_{s_\omega}^- := \Lambda^{0,1}$ . The metrics on them are the ones induced by  $g_J$ . The Clifford multiplication is given for  $v \in TX_x$  by  $\rho(v) := \sqrt{2}((v^{0,1} \wedge) + (v^{0,1} \wedge)^*)$ . From the more intrinsic viewpoint, the decomposition  $W_{s_\omega}^+ = \Lambda^{0,0} \oplus \Lambda^{0,2}$  is just the eigenspace decomposition of  $\rho(\omega) \in \text{End}(W^+)$ . The corresponding eigenvalues are  $-2\sqrt{-1}$  and  $2\sqrt{-1}$ , respectively. Notice that  $\omega$  is a self-dual 2-form with length  $\sqrt{2}$ .

**b)** We will see that there are two natural  $\text{Spin}^c$  connections for  $W_s$ .

Let  $\nabla_J^1$  be the compatible  $\text{Spin}^c$  connection that projects to  $\Lambda^{0,0}$  as the trivial connection  $d$ .  $\nabla_J^1$  preserves the decomposition above if and only if the pair  $(\omega, J)$  is Kähler. In fact, if it preserves the decomposition,  $\rho(\omega)$  is parallel with respect to the induced connection, which implies that  $\nabla^{L.C} \omega \equiv 0$ . Then it follows that  $\nabla^{L.C} J \equiv 0$ , that is,  $J$  is integrable.

Let  $\nabla_J^2$  be the  $U(2) \times U(2)$  connection on  $W_{s_\omega}$  defined by the composition of the sequence

$$(2.3) \quad \bigoplus_p \Lambda^{0,p} \xrightarrow{\nabla^{L.C.}} T^*X \otimes \left( \bigoplus_p^p T^*X \otimes \mathbb{C} \right) \xrightarrow{id \otimes \text{pr}} T^*X \otimes \left( \bigoplus_p \Lambda^{0,p} \right).$$

We can check after a short calculation that  $\nabla_J^2$  is indeed a  $\text{Spin}^c(4)$  connection. This preserves the decomposition  $W_{s_\omega}^+ = \Lambda^{0,0} \oplus \Lambda^{0,2}$  and restricts it to  $\Lambda^{0,0}$  as the trivial connection  $d$ , but it is not compatible unless  $(\omega, J)$  is Kähler as we have seen before.

**c)** Define the operators  $\mathcal{D}^1$  and  $\mathcal{D}^2$  to be the compositions of the sequence (2.1) with  $\nabla^B$  being replaced by  $\nabla_J^1$  and  $\nabla_J^2$ , respectively. The former is one of the usual

Dirac operators for  $\text{Spin}^c$  structures and the latter is written in the following form

$$(2.4) \quad \mathcal{D}^2 = \sqrt{2}(\bar{\partial} + \bar{\partial}^*),$$

where  $\bar{\partial}$  is the Dolbeaut operator. Then we can easily check after a short calculation that  $\mathcal{D}^1 = \mathcal{D}^2$  if and only if  $d\omega = 0$ .

**d)** Fix a  $\text{Spin}^c$  structure  $s$ . Then its spinor bundle is given as the tensor product over  $\mathbb{C}$  of  $W_{s_\omega}$  with a suitable complex line bundle  $L$ . The Dirac operator for  $s$  is given by choosing a  $U(1)$  connection  $a$  for  $L_s$  and is written as

$$(2.5) \quad \sqrt{2}(\bar{\partial}_a + \bar{\partial}_a^*),$$

where  $\bar{\partial}_a$  means the usual coupled Dolbeaut operator. Note that

$$(2.6.1) \quad \Lambda^+ \otimes \mathbb{C} = \Lambda^{2,0} \oplus \Lambda^{0,2} \oplus \mathbb{C}\langle\omega\rangle,$$

$$(2.6.2) \quad \Lambda^- \otimes \mathbb{C} = \Lambda^{1,1} \cap (\mathbb{C}\langle\omega\rangle)^\perp.$$

With this understood, the monopole equation corresponding to  $s$  is written as

$$(2.7.1) \quad \bar{\partial}_a \alpha + \bar{\partial}_a^* \beta = 0,$$

$$(2.7.2) \quad 2F_a^{0,2} + F_{\bar{K}}^{0,2} = \frac{1}{2}\alpha^* \beta,$$

$$(2.7.3) \quad \Lambda(2F_a + F_{\bar{K}}) = \frac{\sqrt{-1}}{2}(|\alpha|^2 - |\beta|^2),$$

where

$$(\alpha, \beta, a) \in \Gamma(\Lambda^{0,0} \otimes L) \times \Gamma(\Lambda^{0,2} \otimes L) \times \mathcal{A}(L).$$

Here  $F_{\bar{K}}$  is the curvature of the connection  $\nabla_J^2|_{\Lambda^{0,2}}$  and  $\Lambda : \Lambda^{p,q} \mapsto \Lambda^{p-1,q-1}$  denotes the adjoint of  $\omega \wedge : \Lambda^{p-1,q-1} \mapsto \Lambda^{p,q}$ . Notice that  $\rho(\gamma_2)\gamma_1 = 2\gamma_1\gamma_2$  for  $\gamma_1 \in \Lambda^{0,0}$  and  $\gamma_2 \in \Lambda^{0,2}$ .

### 3. THE MODULI SPACES OF MONOPOLE EQUATIONS ON WEAKLY CONVEX ALMOST KÄHLER MANIFOLDS

Let  $(X, \omega, J)$  be a weakly convex almost Kähler 4-manifold.

Our main object is the following equation, which was introduced for the first time by Taubes:

$$(3.1.1) \quad \bar{\partial}_a \alpha + \bar{\partial}_a^* \beta = 0,$$

$$(3.1.2) \quad F_a^{0,2} = \frac{r}{4}\alpha^* \beta + \eta^{0,2},$$

$$(3.1.3) \quad \Lambda F_a = \frac{\sqrt{-1}r}{4}(-1 + |\alpha|^2 - |\beta|^2) + \Lambda\eta.$$

Here  $\eta$  is a pure imaginary self-dual 2-form introduced for the equation to be transverse.  $r$  is a positive constant which we will call the *rescaling parameter*.

*Remark 3.0.* The equation above is obtained from (2.7) by dropping the terms derived from the curvature of the anti-canonical line bundle, adding  $-\frac{\sqrt{-1}r}{2}$  to the right-hand side of (2.7.3) and rescaling  $(\alpha, \beta)$  by the factor  $\sqrt{r}$ . From the more

intrinsic viewpoint, it is equivalent to the following equation:

$$(3.2.1) \quad \mathcal{D}_a \Phi = 0,$$

$$(3.2.2) \quad 2\rho(F_a^+) = r\left\{(\Phi\Phi^*)_0 - \frac{\sqrt{-1}}{2}\rho(\omega)\right\}.$$

Fix an element  $(s, \varrho) \in S(X, \omega)$  and suppose that  $W_s = W_{s_\omega} \otimes L$  for a complex line bundle  $L$ .  $\varrho: W_s|_{X \setminus K} \xrightarrow{\cong} W_{s_\omega}|_{X \setminus K}$  induces  $\tilde{\varrho}: L|_{X \setminus K} \xrightarrow{\cong} (X \setminus K) \times \mathbb{C}$  where  $K$  is some compact set. Denote by  $L_\varrho$  the line bundle  $L$  endowed with the trivialization  $\tilde{\varrho}$  outside some compact set.

With this understood, we will introduce a suitable analytic setting for the equation (3.1) following [K-M2].

The equation (3.1) for  $s_\omega$  has the element  $(\mathbb{I}, 0, d) \in \Gamma(\Lambda^{0,0}) \times \Gamma(\Lambda^{0,2}) \times \mathcal{A}(X \times \mathbb{C})$  as a special solution for any choice of  $r$ . (Here  $X \times \mathbb{C}$  means the trivial line bundle over  $X$ .) We will adopt it as an asymptotic solution and define the function spaces  $\Gamma_0$ ,  $\mathcal{A}_0$  and  $\mathcal{G}_0$  as follows:

$$(3.3.1) \quad \Gamma_0 := \{(\alpha, \beta) \in \Gamma^\infty(\Lambda^{0,0}(L_\varrho) \oplus \Lambda^{0,2}(L_\varrho)) \mid (\alpha, \beta) - (\mathbb{I}, 0) \text{ has a compact support}\},$$

$$(3.3.2) \quad \mathcal{A}_0 := \{a \in \mathcal{A}^\infty(L_\varrho) \mid a - d \text{ has a compact support}\},$$

$$(3.3.3) \quad \mathcal{G}_0 := \{u \in C^\infty(X; \mathbb{C}) \mid |u| = 1, u - 1 \text{ has a compact support}\}.$$

$\tilde{\varrho}$  allows us to identify  $\Lambda^{0,0}(L_\varrho) \oplus \Lambda^{0,2}(L_\varrho)$  with  $\Lambda^{0,0} \oplus \Lambda^{0,2}$  outside some compact set  $K$ . Similarly,  $a|_{X \setminus K}$  can be regarded as a  $U(1)$  connection of the trivial complex line bundle over  $X \setminus K$ . These identifications are implicit in (3.3). We will adopt some suitable completions of the spaces  $\Gamma$ ,  $\mathcal{A}$  and  $\mathcal{G}$  as our function spaces.

$\mathcal{G}$  is the completion of  $\mathcal{G}_0$  with respect to the Sobolev  $W^{k+1,2}$  norm defined by the Riemannian metric  $g_J$  and the covariant derivatives.

$\mathcal{A}_0$  can be identified with the space of compact support, smooth and pure imaginary self-dual 2-forms by choosing a base point  $a_0$ .  $\mathcal{A}$  is the completion of  $\mathcal{A}_0$  with respect to the usual  $W^{k,2}$  norm for differential forms.

Define the Sobolev  $W^{k,2}$ -norm for  $\Gamma_0$  by making use of the Riemannian metric  $g_J$ , the Hermitian metric of  $W_s \equiv \Lambda^{0,0}(L_\varrho) \oplus \Lambda^{0,2}(L_\varrho)$  and the covariant derivative  $\nabla_J^1 \otimes \text{id} + \text{id} \otimes \nabla_{a_0}$ .  $\Gamma$  is the completion of  $\Gamma_0$  with respect to this norm.

Let us fix  $k$  sufficiently large so that the Sobolev embedding theorem implies that these function spaces belong to  $C^1$ . Then the Sobolev multiplication theorem implies that the gauge group  $\mathcal{G}$  acts naturally on  $\Gamma \times \mathcal{A}$ . In fact, these spaces are smooth Hilbert manifolds with the former acting as a Hilbert Lie group. Furthermore, the action is free. Thus, the quotient space  $\mathcal{B}$  is also a Hilbert manifold. The standard argument in gauge theory shows that  $\mathcal{B}$  is Hausdorff. See [K-M2].

To have the equation transverse, we introduce a Banach space  $\mathcal{N}$  as a completion of the space of compact support, smooth, pure imaginary self-dual 2-forms. The norm  $\|\cdot\|_{\mathcal{N}}$  is given by  $\|\eta\|_{\mathcal{N}} := \|\exp(\epsilon_1 \sigma) \cdot \eta\|_{C^l(X)}$  where  $\epsilon_1 > 0$  and  $l \geq k+1$  are fixed. We always assume that the  $\eta$  in the equation (3.1) is chosen from this Banach space.

With this understood, we will give some results needed later and the definition of the monopole invariant in [K-M2] in the form suitable for our terminology.

**Proposition 3.1** ([K-M2]). *Let  $(\alpha, \beta, a) \in \Gamma \times \mathcal{A}$  be a solution of the equation (3.1). There exist positive constants  $C_0$  and  $C'_0$  which depend only on  $(\omega, J, c_1(L_\varrho))$*

and have the following significance: If  $r \geq C_0$ , then

$$\begin{aligned} \int_X |\nabla_a \alpha|^2 + 2|\tilde{\nabla}_a \beta|^2 + \frac{r}{2}(1 - |\alpha|^2)^2 + \frac{r}{2}(1 + 2|\alpha|^2 + |\beta|^2)|\beta|^2 \\ \leq C'_0 \|\eta\|_{L^1(X)} + 2\pi \langle c_1(L_\varrho) \cup [\omega], [X] \rangle. \end{aligned}$$

**Proposition 3.2** ([K-M2]). *Let  $(\alpha, \beta, a) \in \Gamma \times \mathcal{A}$  be a solution of equations (3.1) with  $r \geq 1$  and  $\|\eta\|_{\mathcal{N}} \leq 1$ . There exist positive constants  $\nu_r$  and  $\chi_r$  which depend only on  $r$  and on  $(\omega, J, c_1(L_\varrho))$  and have the following significance:*

$$\left\{ \left| |\alpha|^2 - 1 \right| + |\nabla_a \alpha|^2 + |\beta|^2 + |\tilde{\nabla}_a \beta|^2 + |P^+ F_a| + |P^- F_a| \right\}_x \leq \chi_r e^{-\nu_r \sigma(x)}$$

for any  $x \in X$ .

**Theorem 3.3** ([K-M2]). *Define  $\mathcal{M}(\omega, J, s, \varrho)$  to be the set*

$$\{([\alpha, \beta, a], \eta, r) \in \mathcal{B} \times \mathcal{N} \times \mathbb{R}^{>0} \mid (\alpha, \beta, a) \text{ obeys (3.1) with these } \eta \text{ and } r\},$$

where  $[(*)]$  means the gauge equivalence class of  $(*)$ . Then  $\mathcal{M}(\omega, J, s, \rho)$  is a Banach submanifold of  $\mathcal{B} \times \mathcal{N} \times \mathbb{R}^{>0}$ . The projection  $\text{Pr}: \mathcal{M}(\omega, J, s, \rho) \mapsto \mathcal{N} \times \mathbb{R}^{>0}$  is a proper Fredholm map of index  $\langle (c_1^2(L_\varrho) - c_1(L_\varrho) \cup c_1(K), [X]) \rangle$  and the index line bundle has a canonical orientation determined by  $(\omega, J)$ .

Here  $c_1(L_\varrho)$  means  $c_1(L, \tilde{\varrho})$ .  $c_1(L, \tilde{\varrho})$  and  $[X]$  are as explained in Section 1.  $K$  denotes the canonical line bundle of  $(X, J)$ . Theorem 3.3 implies that if  $(\eta, r)$  is generic, namely, if it is chosen from a suitable Baire subset of  $\mathcal{N} \times \mathbb{R}^{>0}$ , then  $\text{Pr}^{-1}(\eta, r)$  is a compact oriented manifold of dimension  $\langle (c_1^2(L_\varrho) - c_1(L_\varrho) \cup c_1(K), [X]) \rangle$ . We will refer to  $\text{Pr}^{-1}(\eta, r)$  as the moduli space.

**Definition 3.4.** The monopole invariant  $SW: S(X, \omega) \mapsto \mathbb{Z}$  is defined as follows:

- (1) If  $\langle (c_1^2(L_\varrho) - c_1(L_\varrho) \cup c_1(K), [X]) \rangle \neq 0$ , then  $SW(s, \varrho) = 0$ .
- (2) If  $\langle c_1^2(L_\varrho) - c_1(L_\varrho) \cup c_1(K), [X] \rangle = 0$ , then  $SW(s, \varrho)$  is the sum of the suitable signs that are imposed to each connected component of the 0-dimensional manifold  $\text{Pr}^{-1}(\eta, r)$  for generic  $(\eta, r)$ . It does not depend on the choice of the pair  $(\eta, r)$ .

#### 4. THE STATEMENT OF THE MAIN RESULT

Let  $(X, \omega, J)$  be a weakly convex almost Kähler 4-manifold. Let  $\mathcal{M}(\omega, J, s, \rho)$  be the space as given in Section 3. Our main result follows:

**Theorem 4.1.** *Let  $\{r_n\}_{n \in \mathbb{N}}$  be a sequence of positive numbers which tends to infinity when  $n$  tends to infinity. Suppose there exists a sequence  $\{(\alpha_n, \beta_n, a_n, \eta_n)\}_{n \in \mathbb{N}}$  such that  $([\alpha_n, \beta_n, a_n], \eta_n, r_n) \in \mathcal{M}(\omega, J, s, \rho)$  obeying  $\|\eta_n\|_{\mathcal{N}} \leq e^{-r_n}$ . Then, after passing to a suitable subsequence,  $\{\alpha_n^{-1}(0)\}_{n \in \mathbb{N}}$  converges in the Hausdorff topology to a compact  $J$ -holomorphic curve  $D$  (which may have multiple irreducible components) whose homology class  $[D] \in H_2(X, \mathbb{Z})$  is the Poincaré dual of  $c_1(L_\varrho)$ .*

This theorem is an extension of the main result in [T1] where  $X$  is supposed to be closed. When  $X$  is non-compact, we must overcome the following problems:

The first one is that the sets  $\alpha_n^{-1}(0)$  may possibly escape to the infinity of the end when  $n$  tends to infinity. The monotonicity formula for local energy integral can settle this problem as long as we have an a priori bound for the total energy integral  $\frac{1}{4} \int_X r |1 - |\alpha|^2|$ , the bound which is independent of  $r$ .



The second one is that it is not obvious at first whether the a priori bound for the total energy integral does exist.

The third one, which is related to the second one, is that the argument in [T1] to find the a priori  $C^0$  bound of the anti-self-dual part of the curvature does not work directly in our case.

Our strategy is divided into 3 steps:

**Step 1.** We will show in Sections 5 and 6 that the  $C^0$  estimates of the terms  $|1 - |\alpha|^2|$ ,  $|\beta|^2$ ,  $|\nabla_a \alpha|^2$ ,  $|\tilde{\nabla}_a \beta|^2$  and  $|F_a^\pm|^2$  given in [T1] are also valid in our case. The major difference from [T1] is in the proof of the  $C^0$  estimate for  $F_a^-$ , which is given in Section 6.

**Step 2.** We will derive in Section 7 an a priori estimate of the total energy integral.

**Step 3.** We will derive in Section 8 a slightly refined monotonicity formula for local energy integral.

With these achieved, we can easily show that  $\alpha_n^{-1}(0)$  does remain in some compact set when  $n$  tends to infinity. This will be done in Section 9 and allows us to handle the issue as if our manifold  $X$  were compact. Thus applying the arguments in [T1] almost directly, we can prove Theorem 4.1.

Before going on to the proof, let us agree that we are subject to Assumption 1 and Conventions 1 and 2 below in Section 5, 6, 7, 8 and 9 unless otherwise specified:

**Assumption 1.** We suppose that  $r \geq 1$  and that

$$(4.1) \quad \|\eta\|_{\mathcal{N}} \leq e^{-r}.$$

**Convention 1.** We adopt the following convention for constants:

**a.** The symbol  $C$  with no subscript stands for a positive constant which depends only on the data  $(\omega, J, c_1(L_\varrho))$  and that the value which  $C$  is supposed to be may vary from line to line even in a single formula.

**b.** The symbol  $C$  with some subscript such as  $C_1$  stands for a positive constant which depends only on  $(\omega, J, c_1(L_\varrho))$  and the value which it is supposed to be is consistent in later arguments.

**Convention 2.** If we say that a constant, such as  $r$ ,  $\kappa$  and so on, is *sufficiently large*, it means that it is larger than a suitable positive constant that depends only on  $(\omega, J, c_1(L_\varrho))$ .

## 5. PRELIMINARY ESTIMATES

We will devote this section to derive preliminary estimates.

It is known that the Dolbeaut operators on an almost Kähler manifold satisfy the Kähler identities. See [Ma]. Our starting point is the following identities which derive from the Kähler identities after a short calculation (see [Ko]):

$$(5.1.1) \quad \bar{\partial}_a^* \bar{\partial}_a \alpha = \frac{1}{2} \nabla_a^* \nabla_a \alpha - \frac{1}{2} \sqrt{-1} (\Lambda F_a) \alpha,$$

$$(5.1.2) \quad \bar{\partial}_a \bar{\partial}_a \alpha = N \circ \partial_a \alpha + F_a^{0,2} \alpha,$$

$$(5.1.3) \quad \bar{\partial}_a^* \bar{\partial}_a^* \beta = \partial_a^* \circ N^* \beta + (F_a^{0,2})^* \beta,$$

$$(5.1.4) \quad \bar{\partial}_a \bar{\partial}_a^* \beta = \frac{1}{2} \tilde{\nabla}_a^* \tilde{\nabla}_a \beta + \frac{1}{2} \sqrt{-1} (\Lambda (F_a + F_{\bar{K}})) \beta,$$

where  $N \in \text{Hom}(\Lambda^{1,0}, \Lambda^{0,2})$  is the Nijenhuis tensor of  $J$ ,  $\tilde{\nabla}_a$  is the unitary connection of  $\Lambda^{0,2} \otimes L$  whose  $(1, 0)$  part agrees with the Dolbeaut operator  $\partial_a: \Omega^{0,2}(L) \mapsto \Omega^{1,2}(L)$  and  $F_a + F_{\overline{K}}$  is the curvature of  $\tilde{\nabla}_a$ .

*Remark.* In the case where  $(\omega, J)$  is Kähler, the identities (5.1) are exactly the Weitzenböck formula of a Dirac operator.

It follows from (3.1) and (5.1) after a short calculation that

$$(5.2.1) \quad \begin{aligned} \frac{1}{2} \nabla_a^* \nabla_a \alpha &= -\frac{r}{8}(-1 + |\alpha|^2 + |\beta|^2)\alpha - \partial_a^* \circ N^* \beta \\ &\quad - (\eta^{0,2})^* \beta + \frac{\sqrt{-1}}{2}(\Lambda \eta)\alpha, \end{aligned}$$

$$(5.2.2) \quad \begin{aligned} \frac{1}{2} \tilde{\nabla}_a^* \tilde{\nabla}_a \beta &= -\frac{r}{8}(+1 + |\alpha|^2 + |\beta|^2)\beta + N \circ \partial_a \alpha \\ &\quad - \frac{\sqrt{-1}}{2}(\Lambda F_{\overline{K}})\beta - \alpha \eta^{0,2} - \frac{\sqrt{-1}}{2}(\Lambda \eta)\beta. \end{aligned}$$

Taking the inner product of (5.2.1) with  $\alpha$  and making use of the identity  $\frac{1}{2}\Delta(|\alpha|^2) = \langle \nabla_a^* \nabla_a \alpha, \alpha \rangle - |\nabla_a \alpha|^2$ , it follows that

$$(5.3.1) \quad \begin{aligned} \frac{1}{4}\Delta(|\alpha|^2) &= -\frac{1}{2}|\nabla_a \alpha|^2 - \frac{r}{8}(-1 + |\alpha|^2 + |\beta|^2)|\alpha|^2 - \langle \partial_a^* \circ N^* \beta, \alpha \rangle \\ &\quad - \langle (\eta^{0,2})^* \beta, \alpha \rangle + \frac{\sqrt{-1}}{2}\Lambda \eta |\alpha|^2. \end{aligned}$$

Similarly, it follows that

$$(5.3.2) \quad \begin{aligned} \frac{1}{4}\Delta(|\beta|^2) &= -\frac{1}{2}|\tilde{\nabla}_a \beta|^2 - \frac{r}{8}(1 + |\alpha|^2 + |\beta|^2)|\beta|^2 \\ &\quad + \langle \beta, N \circ \partial_a \alpha \rangle - \frac{\sqrt{-1}}{2}(\Lambda F_{\overline{K}})|\beta|^2 \\ &\quad - \langle \eta^{0,2} \alpha, \beta \rangle - \frac{\sqrt{-1}}{2}(\Lambda \eta)|\beta|^2. \end{aligned}$$

Since  $(X, \omega, J)$  is weakly convex,  $N$ ,  $F_{\overline{K}}$  and their higher covariant derivatives are all bounded. Thus by dropping some non-positive terms and applying Schwarz' inequality, we obtain

**Lemma 5.0.** *Let  $(\alpha, \beta, a)$  be a solution of equations (3.1) with  $r \geq 1$ . It holds that*

$$(5.4.1) \quad \begin{aligned} \left(\frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2\right)(|\alpha|^2 - 1) \\ \leq -|\nabla_a \alpha|^2 + C|\tilde{\nabla}_a \beta| \cdot |\alpha| + (C + |\eta|)|\alpha| \cdot |\beta| + |\eta| \cdot |\alpha|^2, \end{aligned}$$

$$(5.4.2) \quad \begin{aligned} \left(\frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2\right)|\beta|^2 &\leq -|\tilde{\nabla}_a \beta|^2 - \frac{r}{4}\left(1 - \frac{C}{r}\right)|\beta|^2 \\ &\quad + C|\nabla_a \alpha| \cdot |\beta| + |\eta| \cdot |\alpha| \cdot |\beta| + |\eta| \cdot |\beta|^2. \end{aligned}$$

By making use of it, we can show

**Proposition 5.1.** *Let  $(\alpha, \beta, a)$  be a solution of equations (3.1) with  $r \geq 1$ . There exist non-negative constants  $\kappa_1, \kappa_2$  which depend only on  $(\omega, J, c_1(L_\varrho))$  and have*

the following significance: If  $p$  and  $\zeta$  obey that  $1 \geq p \geq 0$ ,  $\zeta > 0$ , then

$$\begin{aligned} & \left(\frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2\right) \{|\alpha|^2 - 1 + \zeta r^p |\beta|^2\} \\ & \leq -\left(1 - \frac{\zeta \kappa_1}{r^{1-p}}\right) |\nabla_a \alpha|^2 - \frac{\zeta}{2} r^p |\tilde{\nabla}_a \beta|^2 - \frac{\zeta}{24} r^{1+p} |\beta|^2 + \kappa_2 \left(\frac{1}{\zeta} + \zeta\right) \frac{1}{r^p} |\alpha|^2. \end{aligned}$$

*Proof.* It follows from (4.1), (5.4.1), (5.4.2) and Hölder's inequality that

$$\begin{aligned} \text{R.H.S.} & \leq -|\nabla_a \alpha|^2 - \zeta r^p |\tilde{\nabla}_a \beta|^2 - \frac{\zeta r^{1+p}}{8} |\beta|^2 \\ & \quad + C_1 \zeta r^p |\nabla_a \alpha| |\beta| + C_2 |\tilde{\nabla}_a \beta| |\alpha| + C_3 (1 + \zeta) |\alpha| |\beta| + |\eta| |\alpha|^2 \\ & \leq -\left(1 - \frac{C_1 \zeta \epsilon_1}{2r^{1-p}}\right) |\nabla_a \alpha|^2 - r^p \left(\zeta - \frac{C_2 \epsilon_2}{2}\right) |\tilde{\nabla}_a \beta|^2 \\ & \quad - \frac{r^{1+p}}{8} \left(\zeta - \frac{4C_1 \zeta}{\epsilon_1} - \frac{4C_3}{\epsilon_3} (1 + \zeta)\right) |\beta|^2 + \frac{1}{r^p} \left(\frac{C_2}{2\epsilon_2} + \frac{C_3 \epsilon_3}{2r} (1 + \zeta) + r^p |\eta|\right) |\alpha|^2. \end{aligned}$$

By putting  $\epsilon_1 = 12C_1$ ,  $\epsilon_2 = \frac{\zeta}{C_2}$  and  $\epsilon_3 = \frac{12C_3(1+\zeta)}{\zeta}$ , we obtain the result.  $\square$

**Proposition 5.2.** *Let  $(\alpha, \beta, a)$  be a solution of equations (3.1) with  $r \geq 1$ . There exists a constant  $C$  which depends only on  $(\omega, J, c_1(L_\varrho))$  and has the following significance: It holds that*

$$(5.5) \quad |\alpha|^2 + |\beta|^2 \leq 1 + \frac{C}{r}.$$

*Proof.* Define  $f$  to be  $|\alpha|^2 - 1 + |\beta|^2 - \frac{\kappa}{r}$  where  $\kappa$  is a positive constant determined later. Proposition 5.1 implies that  $f$  obeys

$$\left(\frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2\right) f \leq (C - \frac{\kappa}{4}) |\alpha|^2.$$

By taking  $\kappa$  sufficiently large, the right-hand side is nonpositive. On the other hand,  $f$  is negative outside some compact set (that may depend on  $r$  and  $\kappa$ ). Thus a maximum principle implies  $f \leq 0$ .  $\square$

**Proposition 5.3.** *Let  $(\alpha, \beta, a)$  be a solution of equations (3.1) with  $r \geq 1$ . There exists a constant  $C_M$  which depends only on  $(\omega, J, c_1(L_\varrho))$  and has the following significance: It holds that*

$$(5.6) \quad |\beta|^2 \leq \frac{1}{r} (1 - |\alpha|^2) + \frac{C_M}{r^3}.$$

*Proof.* Let  $\zeta_1$  be a fixed positive constant such that  $\zeta_1 \kappa_1 \leq 1$ . Define  $f$  to be  $|\alpha|^2 - 1 + \zeta_1 r |\beta|^2 - \frac{\kappa}{r^2}$  where  $\kappa$  is a positive constant determined later. Proposition 5.1 implies that  $\left(\frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2\right) f \leq (C - \frac{\kappa}{4}) \frac{1}{r} |\alpha|^2$ . Thus if  $\kappa$  is sufficiently large, the same argument as in the proof of Proposition 5.2 implies  $f \leq 0$ .  $\square$

**Proposition 5.4.** *Let  $(\alpha, \beta, a)$  be a solution of equations (3.1) with  $r \geq 1$ . There exist positive constants  $\mu_1, \mu_2$  which depend only on  $(\omega, J, c_1(L_\varrho))$  and have the following significance: It holds that*

$$(5.7) \quad |F_a^+| \leq \frac{r}{4\sqrt{2}} \left(1 + \frac{\mu_1}{r}\right) (1 - |\alpha|^2) + \frac{\mu_2}{r}.$$

*Proof.* This follows directly from (3.1.2), (3.1.3) and Proposition 5.3.  $\square$

6. THE  $C^0$  ESTIMATE OF THE ANTI-SELF-DUAL PART OF THE CURVATURE

We will devote the whole of this section to prove

**Proposition 6.1.** *Let  $(\alpha, \beta, a)$  be a solution of equations (3.1) with  $r \geq 1$ . There exist non-negative constants  $\mu_3, \mu_4$  which depend only on  $(\omega, J, c_1(L_\theta))$  and have the following significance: It holds that*

$$(6.1) \quad |F_a^-| \leq \frac{r}{4\sqrt{2}}(1 + \frac{\mu_3}{r^{\frac{1}{4}}})(1 - |\alpha|^2) + \frac{\mu_4}{r^{\frac{1}{4}}}.$$

This estimate will be needed in the proofs of Proposition 7.1 and Proposition 8.1.

The proof is divided into 8 steps.

**Step (0).** Denote  $|F_a^-|$  by  $t$ . We will derive a differential inequality that  $t$  obeys.

**Lemma 6.2.**  *$t$  obeys the following inequality on  $X \setminus t^{-1}(0)$ :*

$$(6.2) \quad (\frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2)t \leq Rt + \frac{r}{4\sqrt{2}}(|\nabla_a \alpha|^2 + |\tilde{\nabla}_a \beta|^2) + Cr|\beta|^2 + |h_\eta|.$$

Here  $R$  is a non-negative function derived from the Riemannian metric and  $h_\eta$  denotes  $\frac{1}{4}P^-(d^*\eta)$  where  $P^-$  stands for the orthogonal projection  $P^-: \overset{2}{\Lambda} \mapsto \Lambda^+$ .

*Proof.* The Bianchi identity implies that

$$(6.3) \quad dF_a^+ + dF_a^- = 0.$$

Then a Bochner-Weitzenböck formula implies that

$$(6.4) \quad \frac{1}{2}\nabla^*\nabla F_a^- + \mathcal{R}F_a^- = -P^-d^*dF_a^+,$$

where  $\mathcal{R} \in \text{Hom}(\Lambda^-, \Lambda^-)$  derives from the anti-self-dual part of the curvature of the Riemannian metric and the scalar curvature. (3.1.2) and (3.1.3) imply that

$$(6.5) \quad \begin{aligned} \text{R.H.S. of (6.4)} &= P^-d^*d\left\{-\frac{\sqrt{-1}r}{8}(-1 + |\alpha|^2 - |\beta|^2)\right. \\ &\quad \left.\omega - \frac{r}{4}\alpha^*\beta + \frac{r}{4}\alpha\beta^* - \eta\right\}. \end{aligned}$$

By making use of the Kähler identities,

$$\begin{aligned} &P^-d^*d\left\{-\frac{\sqrt{-1}r}{8}(-1 + |\alpha|^2 - |\beta|^2)\omega\right\} \\ &= -\frac{\sqrt{-1}r}{8}P^-(\partial^*\partial + \bar{\partial}^*\bar{\partial})(|\alpha|^2 - |\beta|^2)\omega \\ (6.6) \quad &= -\frac{\sqrt{-1}r}{8}P^-(-\sqrt{-1}\partial\Lambda\partial + \sqrt{-1}\partial\Lambda\bar{\partial})(|\alpha|^2 - |\beta|^2)\omega \\ &\quad (\text{since Image}(\Lambda: \Lambda^{2,2} \mapsto \Lambda^{1,1}) \subset \Lambda^+) \\ &= -\frac{r}{4}P^-(\bar{\partial}\partial|\alpha|^2 + \partial\bar{\partial}|\beta|^2) \\ &\quad (\text{since } \Lambda \circ (\omega \wedge) = id \text{ on } \overset{1}{\Lambda} \text{ and since } \bar{\partial}\partial + \partial\bar{\partial} = 0 \text{ on } \Lambda^{0,0}). \end{aligned}$$

Thus we obtain the equality (6.7) below:

$$\begin{aligned} \text{R.H.S. of (6.6)} &= -\frac{r}{4}P^-\{\langle \bar{\partial}_a \partial_a \alpha, \alpha \rangle_L + \langle \alpha, \partial_a \bar{\partial}_a \alpha \rangle_L - \langle \partial_a \alpha, \partial_a \alpha \rangle_L + \langle \bar{\partial}_a \alpha, \bar{\partial}_a \alpha \rangle_L \\ &\quad + \langle \partial_{\tilde{\nabla}_a} \bar{\partial}_{\tilde{\nabla}_a} \beta, \beta \rangle_{L \otimes \bar{K}} + \langle \beta, \bar{\partial}_{\tilde{\nabla}_a} \partial_{\tilde{\nabla}_a} \beta \rangle_{L \otimes K} + \langle \partial_{\tilde{\nabla}_a} \beta, \partial_{\tilde{\nabla}_a} \beta \rangle_{L \otimes \bar{K}} - \langle \bar{\partial}_{\tilde{\nabla}_a} \beta, \bar{\partial}_{\tilde{\nabla}_a} \beta \rangle_{L \otimes K}\} \end{aligned}$$

where  $\langle \cdot, \cdot \rangle_L$  and  $\langle \cdot, \cdot \rangle_{L \otimes \overline{K}}$  are the Hermitian inner products of the line bundle  $L$  and  $L \otimes \overline{K}$ , respectively. Further, the exterior products of the forms are implicit in these expressions.  $\partial_{\tilde{\nabla}_a}$  stands for the coupled Dolbeaut operator  $\partial_{\tilde{\nabla}_a} : \Omega^{p,q}(L \otimes \overline{K}) \mapsto \Omega^{p+1,q}(L \otimes \overline{K})$ .

On the other hand, since  $\alpha^* \beta$  is a  $(0, 2)$  form, it follows that

$$\begin{aligned}
 P^- d^* d(-\frac{1}{4} \alpha^* \beta) &= \frac{1}{4} P^- (\sqrt{-1} \partial \Lambda \partial) (\alpha^* \beta) \\
 &= \frac{\sqrt{-1}}{4} P^- \partial \{ \Lambda (\overline{\partial}_a \alpha)^* \wedge \beta + \Lambda \alpha^* \partial \beta \} \\
 (6.8) \quad &= -\frac{\sqrt{-1}}{4} P^- \partial \{ -\Lambda (\overline{\partial}_a^* \beta)^* \wedge \beta + \sqrt{-1} (\alpha^* \overline{\partial}_a^* \beta) \} \\
 &= -\frac{\sqrt{-1}}{4} P^- \partial \{ -\Lambda (\overline{\partial}_a^* \beta)^* \wedge \beta - \sqrt{-1} \alpha^* (\overline{\partial}_a \alpha) \}
 \end{aligned}$$

where we have used (3.1.1) and the Kähler identities. Applying to (6.8) the identity  $\sqrt{-1} \Lambda (\beta \wedge (\overline{\partial}_a^* \beta)^*) \equiv \beta (\partial_{\tilde{\nabla}_a} \beta)^*$  which also derives from the Kähler identities, we obtain

$$(6.9.1) \quad P^- d^* d(-\frac{1}{4} \alpha^* \beta) = -\frac{1}{4} P^- \partial \{ -\langle \beta, \partial_{\tilde{\nabla}_a} \beta \rangle_{L \otimes \overline{K}} + \langle \overline{\partial}_a \alpha, \alpha \rangle_L \}.$$

Taking its complex conjugate, we obtain the equality

$$(6.9.2) \quad P^- d^* d(\frac{1}{4} \alpha \beta^*) = -\frac{1}{4} P^- \overline{\partial} \{ \langle \partial_{\tilde{\nabla}_a} \beta, \beta \rangle_{L \otimes \overline{K}} - \langle \alpha, \overline{\partial}_a \alpha \rangle_L \}.$$

Therefore, by applying the identities

$$(6.10.1) \quad P^- F_a \equiv P^- (\overline{\partial}_a \partial_a + \partial_a \overline{\partial}_a),$$

$$(6.10.2) \quad P^- (F_a + F_{\overline{K}}) \equiv P^- (\overline{\partial}_{\tilde{\nabla}_a} \partial_{\tilde{\nabla}_a} + \partial_{\tilde{\nabla}_a} \overline{\partial}_{\tilde{\nabla}_a})$$

to (6.4), (6.7) and (6.9) and summing up the result, it follows that

$$\begin{aligned}
 (6.11) \quad \frac{1}{2} \nabla \nabla^* F_a^- + \mathcal{R} F_a^- &= -\frac{r}{4} \langle P^- F_a \alpha, \alpha \rangle_L - \frac{r}{4} \langle P^- (F_a + F_{\overline{K}}) \beta, \beta \rangle_{L \otimes \overline{K}} \\
 &\quad + \frac{r}{4} P^- \{ \langle d_a \alpha, d_a \alpha \rangle_L + \langle d_{\tilde{\nabla}_a} \beta, d_{\tilde{\nabla}_a} \beta \rangle_{L \otimes \overline{K}} \} + \frac{1}{4} P^- (d^* d \eta).
 \end{aligned}$$

By taking the Hermitian inner product of this with  $P^- F_a$ , and making use of the inequality  $(\Delta |F|) |F| \leq \langle \nabla^* \nabla F, F \rangle$  for an arbitrary non-vanishing real 2-form  $F$ , we obtain the inequality

$$\begin{aligned}
 (6.12) \quad (\frac{1}{2} \Delta + \frac{r}{4} |\alpha|^2) t &\leq |\mathcal{R}| t + \frac{r}{4} |\beta|^2 |F_{\overline{K}}| + \frac{r}{4} |P^- \{ \langle d_a \alpha, d_a \alpha \rangle_L + \langle d_{\tilde{\nabla}_a} \beta, d_{\tilde{\nabla}_a} \beta \rangle_{L \otimes \overline{K}} \}| \\
 &\quad + \frac{1}{4} |P^- d^* d \eta| \\
 &\leq |\mathcal{R}| t + C r |\beta|^2 + \frac{r}{4\sqrt{2}} (|\nabla_a \alpha|^2 + |\tilde{\nabla}_a \beta|^2) + |h_\eta|.
 \end{aligned}$$

□

**Step (1).** We will introduce a comparison function  $q_0$ .

Take a sufficiently large  $\kappa > 1$  and define the function  $q_0$  by

$$(6.13) \quad q_0 := \frac{r}{4\sqrt{2}} (1 + \frac{2\kappa_1}{r^{\frac{1}{2}}}) (1 - |\alpha|^2 - r^{\frac{1}{2}} |\beta|^2 + \frac{\kappa}{r^{\frac{3}{2}}}).$$

Lemma 6.2, Proposition 5.1 and Proposition 5.3 imply that if  $r$  is sufficiently large,  $q_0$  is positive and obeys

$$(6.14) \quad \left(\frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2\right)(t - q_0) \leq R_0 \cdot t + |h_\eta|,$$

where  $R_0$  denotes  $\sup_X |\mathcal{R}|$ .

Here we applied Proposition 5.1 to  $p=\frac{1}{2}$  and  $\zeta=1$  and used the fact that  $1 \leq (1 - \frac{\kappa_1}{r^{\frac{1}{2}}})(1 + \frac{2\kappa_1}{r^{\frac{1}{2}}})$  if  $r$  is sufficiently large. We included the constant term  $\kappa r^{-\frac{3}{2}}$  in the definition of  $q_0$  in order to compensate for the last term of the R.H.S. in the inequality of Proposition 5.1, whose existence derives from the Nijenhuis tensor of  $J$ .

**Step (2).** We will define a good comparison function  $q \in W_0^{2,2}(X)$  so that  $q$  obeys  $t \leq q_0 + q$ .

**Lemma 6.3.** *The operator  $(\frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2) : C_0^\infty(X) \mapsto C_0^\infty(X)$  extends to a self-adjoint operator  $\tilde{L}$  over  $L^2(X)$  with  $\text{Dom}(\tilde{L}) = W_0^{2,2}(X)$ . Further,  $\tilde{L}$  is surjective.*

Here  $W_0^{2,2}(X)$  denotes the completion of  $C_0^\infty(X)$  with respect to the Sobolev norm  $\|*\|_{W^{2,2}(X)}$  defined by  $\|f\|_{W^{2,2}(X)} := \|f\|_{L^2(X)} + \|\nabla f\|_{L^2(X)} + \|\nabla \nabla f\|_{L^2(X)}$ . We will give its proof in Appendix.

Define  $q \in W_0^{2,2}(X)$  to be the unique solution of the equation

$$(6.15) \quad \left(\frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2\right)q = R_0 \cdot t + h.$$

Here  $h$  denotes  $|h_\eta|$ . By the construction,  $h$  obeys  $h \leq Ce^{-r}$  and decays like the function  $e^{-\epsilon_1 \sigma}$ .

**Lemma 6.4.**  *$q$  obeys the following:*

1.  $q$  tends to zero uniformly at the end of  $X$ .
2.  $q \in C^{2, \frac{1}{2}}(X)$ .
3.  $q \geq 0$ .

Then a maximum principle applied to (6.14) and (6.15) implies that

$$(6.16) \quad t \leq q_0 + q.$$

*Proof of Lemma 6.4.* Let  $l > 0$  denote  $\min_X \sigma$ . The property (A) of the Riemannian metric  $g_J$  means that the geometries  $\{g_J|_{B(x,l)}\}_{x \in X}$  are bounded. Thus it follows from the standard  $L^p$ -theory of elliptic operators and the Sobolev embedding theorem that there exists a positive constant  $C$  such that

$$(6.17) \quad \|q\|_{C^0(B(x, \frac{l}{2}))} \leq C \left( \|q\|_{L^2(B(x,l))} + \|R_0 \cdot t + h\|_{C^0(B(x,l))} \right).$$

Then the first assertion follows from the fact that  $\|R_0 \cdot t + h\|_{C^0(B(x,l))}$  tends to zero uniformly when  $\sigma(x)$  tends to infinity. The second assertion follows from the standard Hölder theory of elliptic operators since the right-hand side of (6.15) is in  $C^{\frac{1}{2}}(X)$ . Then a maximum principle verifies the third assertion.  $\square$

**Step (3).** We will estimate  $\|q\|_{L^2(X)}$  in terms of  $\sup_X q$ .

Taking the multiple of (6.15) with  $q$  and adding  $\frac{r}{4}(1 - |\alpha|^2)q^2$  to both sides of it, we obtain the equality

$$(6.18) \quad \frac{1}{4}\Delta(q^2) + \frac{1}{2}|\nabla q|^2 + \frac{r}{4}|q|^2 = (R_0 \cdot t + h)q + \frac{r}{4}(1 - |\alpha|^2)q^2.$$

Applying Hölder's inequality to the both terms of the right-hand side, we see that it is no more than  $\frac{3}{r}(R_0 \cdot t + h)^2 + \frac{3}{16}r(1 - |\alpha|^2)^2(\sup_X q)^2 + \frac{r}{6}|q|^2$ . Thus it follows that

$$(6.19) \quad \frac{1}{4}\Delta(q^2) + \frac{r}{12}|q|^2 \leq \frac{6}{r}(R_0^2 \cdot t^2 + h^2) + \frac{3}{16}r(1 - |\alpha|^2)^2(\sup_X q)^2.$$

The very definition of  $W_0^{2,2}(X)$  immediately implies that  $\int_X \Delta(q^2) d\text{vol}_X = 0$ . Thus by integrating (6.19) over  $X$ , we obtain the following inequality:

$$(6.20) \quad \int_X |q|^2 \leq C \left\{ \frac{1}{r^2} \left( \int_X t^2 \right) + \frac{e^{-2r}}{r^2} + \frac{1}{r} (\sup_X q)^2 \left( \int_X r(1 - |\alpha|^2)^2 \right) \right\}.$$

On the other hand, Proposition 3.1 implies that

$$(6.21) \quad \int_X r(1 - |\alpha|^2)^2 \leq C.$$

Further, it holds that

$$(6.22) \quad \int_X t^2 \leq Cr + C.$$

In fact, the Chern-Weil theory implies that

$$(6.23) \quad \int_X t^2 = \int_X |F_a^-|^2 = \int_X |F_a^+|^2 - \langle c_1^2(L), [X] \rangle.$$

On the other hand, (3.1.2), (3.1.3) and Proposition 3.1 imply that

$$(6.24) \quad \int_X |F_a^+|^2 = \frac{r}{32} \int_X r \{ (1 - |\alpha|^2 - |\beta|^2)^2 + 2|\beta|^2 \} \leq Cr.$$

Combining (6.20), (6.21) and (6.22) together, we obtain

$$(6.25) \quad \|q\|_{L^2(X)} \leq \frac{C}{r^{\frac{1}{2}}} + \frac{C}{r^{\frac{1}{2}}} \sup_X q.$$

**Step (4).** We will introduce a comparison function  $p \in W_0^{2,2}(X)$ .

Define  $p \in W_0^{2,2}(X)$  to be the unique solution of the equation

$$(6.26) \quad \left( \frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2 \right) p = (R_0 \cdot t + h)|\alpha|^2.$$

Exactly the same way as in the case of Lemma 6.4, we can prove

**Lemma 6.5.** *p obeys the following:*

- (1) *p tends to zero uniformly at the end of X.*
- (2)  *$p \in C^{2, \frac{1}{2}}(X)$ .*
- (3)  *$p \geq 0$ .*

Applying a maximum principle to  $p - \frac{4}{r}(R_0 \sup_X t + \sup_X h)$ , it follows that

$$(6.27) \quad \sup_X p \leq \frac{4}{r}(R_0 \sup_X t + \sup_X h).$$

**Step (5).** We will estimate the upper bounds of both  $q$  and  $t$ .

We consider the equality below obtained from (6.15) and (6.26):

$$(6.28) \quad \left( \frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2 \right) (q - p) = (R_0 \cdot t + h)(1 - |\alpha|^2).$$

We will apply to it the maximum principle of Gilbarg and Trudinger derived from Alexandrov and Bakel'man. See Theorem 9.20 in [G-T].

**Theorem 6.6** ([G-T]). *Let  $D := -a^{ij} \frac{\partial}{\partial x_i} \frac{\partial}{\partial x_j} + b^j \frac{\partial}{\partial x_j} + c$  be an elliptic operator defined on the unit ball in  $\mathbb{R}^4$  and satisfy the conditions below:*

- (1) *Let  $A$  be a symmetric matrix  $[a^{ij}]_{ij}$ . There exist positive constants  $\lambda_1 \geq \lambda_2$  such that  $\lambda_1 |\xi|^2 \geq \xi^t A \xi \geq \lambda_2 |\xi|^2$  for all  $\xi \in \mathbb{R}^4$ . In other words,  $A$  is uniformly positive definite.*
- (2) *There exists  $\lambda_3 \geq 0$  such that  $|b| \leq \lambda_3$  and  $c \geq -\lambda_3$ .*

*Then there exists a positive constant  $C'$  which depends only on  $\lambda_1, \lambda_2$  and  $\lambda_3$  and has the following significance: If  $f \in C^2(\overline{B_1})$  obeys the differential inequality  $Df \leq g$  on  $B_1^+ \subset B_1$ , then it follows that*

$$\sup_{B_{\frac{1}{2}}^+} f \leq C' (\|f\|_{L^2(B_1^+)} + \|g\|_{L^4(B_1^+)}),$$

where  $B_1^+$  denotes the subset  $\{x \in B_1 \mid f(x) \geq 0\}$ .

The important point is that the coefficient  $c$  is required to be bounded only from below. In [G-T], the assumption of the statement requires that  $|c| \leq \lambda_3$ . But looking closely at the proof, we can easily see that this condition can be relaxed as above.

Let  $x \in X$  attain the maximum of  $q - p$ . Theorem 6.6 is applied to (6.28) to show that

$$(6.29) \quad \begin{aligned} \sup_X (q - p) = \sup_{B_{\frac{1}{2}}(x)} (q - p) &\leq C \left\{ \|q\|_{L^2(X)} + \|p\|_{L^2(B_1(x_n))} \right. \\ &\quad \left. + (\sup_X t + e^{-r}) \|1 - |\alpha|^2\|_{L^4(X)} \right\}. \end{aligned}$$

The right-hand side of (6.29) is bounded from above by

$$C \left( \frac{1}{r^{\frac{1}{2}}} + \frac{1}{r^{\frac{1}{2}}} \sup_X q + \frac{1}{r^{\frac{1}{4}}} \sup_X t \right),$$

because of (6.25), (6.27) and the inequality

$$\int_X (1 - |\alpha|^2)^4 \leq C \int_X (1 - |\alpha|^2)^2 \leq \frac{C}{r}$$

that derives from Proposition 5.2 and (6.21). On the other hand, the left-hand side of (6.29) is bounded from below by

$$\sup_X q - \frac{C}{r} \sup_X t - C \frac{e^{-r}}{r}.$$

Therefore, for sufficiently large  $r$ , it follows that

$$(6.30) \quad \sup_X q \leq C \left( \frac{1}{r^{\frac{1}{2}}} + \frac{1}{r^{\frac{1}{4}}} \sup_X t \right).$$

Applying it to (6.16), it follows that

$$(6.31) \quad \sup_X t \leq Cr,$$



which is applied back to (6.27) and (6.30) to prove that

$$(6.32.1) \quad \sup_X q \leq Cr^{\frac{3}{4}},$$

$$(6.32.2) \quad \sup_X p \leq C.$$

**Step (6).** We will derive a good comparison function which bounds  $q - p$  from above in order to refine the estimate of  $t$ .

**Lemma 6.7.** *There exists a constant  $\delta > 0$  such that if  $r$  is sufficiently large, the function  $v := 1 - |\alpha|^2 - |\beta|^2 + \frac{3\delta}{r}$  obeys the following:*

- (1)  $v \geq 1 - |\alpha|^2 + \frac{2\delta}{r} \geq \frac{\delta}{r}$ .
- (2)  $(\frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2)v \geq 0$ .

The lemma above follows easily from Propositions 5.1 and 5.3.

**Lemma 6.8.** *Define  $v_1$  by  $v_1 := v^{1-r^{-\frac{3}{4}}}$ . Then it obeys the following:*

- (1)  $2v \geq v_1 \geq \frac{1}{2}v$ .
- (2)  $(\frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2)v_1 \geq \frac{r^{\frac{1}{4}}}{2}|\alpha|^2(1 - |\alpha|^2 + \frac{2\delta}{r})$ .

Define the function  $v_2$  by

$$(6.33) \quad v_2 := \left\{ \frac{8}{r^{\frac{1}{4}}} (R_0 \sup_X t + \sup_X h) + 4 \sup_X q \right\} v_1.$$

The right-hand side of (6.28) is bounded from above by

$$(6.34) \quad \begin{aligned} & (R_0 t + h) \left( 1 - |\alpha|^2 + \frac{C_M}{r^2} \right) \\ & \leq (R_0 \sup_X t + \sup_X h) \left( 1 - |\alpha|^2 + \frac{C_M}{r^2} \right), \end{aligned}$$

where we have used Proposition 5.3. Thus (2) of Lemma 6.8 implies that, if  $r$  is sufficiently large,

$$(6.35) \quad \left( \frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2 \right) (q - p - v_2) \leq 0$$

on the domain  $\Omega_{\frac{1}{2}} \{x \in X \mid |\alpha|_x^2 \geq \frac{1}{2}\}$ . On the other hand, (1) of Lemma 6.7 and (1) of Lemma 6.8 imply that, if  $r$  is sufficiently large,

$$(6.36) \quad v_2 \geq \sup_X q \text{ on } X \setminus \Omega_{\frac{1}{2}}.$$

Therefore, a maximum principle is applied to prove that, if  $r$  is sufficiently large,

$$(6.37) \quad q - p \leq v_2 \leq Cr^{\frac{3}{4}}(1 - |\alpha|^2) + \frac{C}{r^{\frac{1}{4}}}.$$

Then (6.16), (6.32.2) and Proposition 5.3 imply that

$$(6.38) \quad t \leq Cr(1 - |\alpha|^2) + C.$$

*Proof Lemma 6.8.* The first assertion follows from the inequality

$$\left( 1 + \frac{3\delta}{r} \right)^{(r^{-\frac{3}{4}})} \geq \frac{v}{v_1} \geq \left( \frac{\delta}{r} \right)^{(r^{-\frac{3}{4}})},$$

where the left-most and right-most sides tend to 1 when  $r$  tends to infinity. The second assertion follows from (1) of Lemma 6.7, (1) of Lemma 6.8 and the fact that

$\Delta(f^b) \geq b(\Delta f)f^{b-1}$  for a smooth positive function  $f$  and a constant  $b$  that obeys  $1 \geq b > 0$ .  $\square$

**Step (7).** We will derive a good comparison function which bounds  $p$  from above and verify the required estimate of  $t$ .

(6.26), (6.38) and (2) of Lemma 6.8 imply that there exists a positive constant  $\kappa_3$  which depends only on  $(\omega, J, c_1(L_\varrho))$  such that the function  $f := p - \kappa_3 r^{\frac{3}{4}} v_1$  obeys  $(\frac{1}{2}\Delta + \frac{r}{4}|\alpha|^2)f \leq 0$ . Since  $f$  is negative outside some compact set, it follows from a maximum principle that

$$(6.39) \quad p \leq \kappa_3 r^{\frac{3}{4}} v_1 \leq C r^{\frac{3}{4}} (1 - |\alpha|^2) + \frac{C}{r^{\frac{1}{4}}}.$$

Combining (6.37) with (6.39), it follows that

$$(6.40) \quad q \leq C r^{\frac{3}{4}} (1 - |\alpha|^2) + \frac{C}{r^{\frac{1}{4}}}.$$

Then (6.16) verifies the required estimate in the statement of Proposition 6.1.

## 7. AN A PRIORI ESTIMATE FOR THE TOTAL ENERGY INTEGRAL

**Proposition 7.1.** *There exists a positive constant  $C$  which depends only on  $(\omega, J, c_1(L_\varrho))$  and has the following significance: Let  $(\alpha, \beta, a)$  be a solution of equations (3.1) with  $r \geq 1$ . Then it holds that*

$$(7.1) \quad |\nabla_a \alpha|^2 + r |\tilde{\nabla}_a \beta|^2 \leq C \{r(1 - |\alpha|^2) + 1\}.$$

This corresponds to Proposition 2.8 of [T1] and can be proved exactly in the same way by making use of the estimates in Section 5 and Proposition 6.1. See [T1] for the proof.

With this in hand, we will devote the latter part of this section to prove

**Proposition 7.2.** *There exists a positive constant  $C_e$  which depends only on  $(\omega, J, c_1(L_\varrho))$  and has the following significance: Let  $(\alpha, \beta, a)$  be a solution of equations (3.1) with  $r \geq 1$ . Then it holds that*

$$(7.2) \quad \int_X \frac{r}{4} |1 - |\alpha|^2| \leq C_e.$$

Of course, the pointwise a priori estimates of the integrand that we have obtained in Section 5 do not directly imply the estimate above since a noncompact weakly convex manifold has infinite volume.

*Proof.* The proof is divided into 3 steps.

**Step (1).** Let  $X_{\frac{1}{2}}$  denote the set  $\{x \in X \mid |\alpha|_x^2 \leq \frac{1}{2}\}$ . We will introduce good subsets  $X^1, X^2 \subset X$ .

**Lemma 7.3.** *There exists a positive constant  $C_v$  which depends only on  $(\omega, J, c_1(L_\varrho))$  and has the following significance: Let  $V$  be a finite subset  $\{x_i\}_{1 \leq i \leq k} \subset X_{\frac{1}{2}}$  such that  $B(x_i, r^{-\frac{1}{2}})$  are mutually disjoint, where  $B(x_i, r^{-\frac{1}{2}})$  denotes the geodesic ball of radius  $r^{-\frac{1}{2}}$  with center  $x_i$ . Then  $\#V \leq C_v r$ .*

Let  $V_M$  be one of the sets described in Lemma 7.3 and suppose it is maximal among such sets. Define  $X^1$  and  $X^2$  as  $\bigcup_{x \in V_M} B(x_i, 2r^{-\frac{1}{2}})$  and  $\bigcup_{x \in V_M} B(x_i, 4r^{-\frac{1}{2}})$ , respectively. Then the following properties hold:

- (1)  $X_{\frac{1}{2}} \subset X^1 \subset X^2$ .

- (2)  $\text{dist}(X^1, X \setminus X^2) \geq r^{-\frac{1}{2}}$ .  
 (3)  $\text{Vol}(X^2) \leq \frac{C}{r}$ .

In fact, the first property follows from the maximality of  $V_M$ . The second property is obvious. The third property follows from the bound of  $\#V_M$  given in Lemma 7.3.

*Proof of Lemma 7.3.* Proposition 7.1 and the inequality  $|\nabla|\alpha|^2| \leq 2|\nabla_a \alpha| \cdot |\alpha|$  imply that there exists a positive constant  $C$  such that if  $\text{dist}(y, X_{\frac{1}{2}}) \leq \frac{1}{C}r^{-\frac{1}{2}}$ , then  $1 - |\alpha|_y^2 \geq \frac{1}{4}$ . Thus there exists a positive constant  $C$  such that

$$(7.3) \quad \int_{B(x, r^{-\frac{1}{2}})} r(1 - |\alpha|^2)^2 \geq \frac{1}{C}r^{-1},$$

if  $x \in X_{\frac{1}{2}}$ . On the other hand, the integral  $\int_X r(1 - |\alpha|^2)^2$  is bounded from above by a constant which does not depend on  $r$ . See Proposition 3.1. Thus we are done.  $\square$

**Step (2).** We will introduce a comparison function  $\phi$  on  $X$  which obeys

$$(7.4) \quad |1 - |\alpha|^2| \leq \phi \text{ on } X \setminus X^2.$$

Let  $C_g$  be a constant determined later such that  $0 < C_g \leq 1$  and such that it depends only on the weakly convex almost Kähler structure.

**Lemma 7.4.** *There exist positive constants  $C_c$  and  $\epsilon_c$  which depend only on  $(\omega, J, c_1(L_\varrho))$  and have the following significance: Let  $y \in X \setminus X^2$ .*

- (1) *If  $\text{dist}(y, X^1) \geq C_g \sigma(y)$ , then*

$$(7.5.1) \quad |1 - |\alpha|_y^2| \leq C_c \exp^{-\epsilon_c \sqrt{r}\{C_g \sigma(y)\}}.$$

- (2) *If  $\text{dist}(y, X^1) \leq C_g \sigma(y)$ , then*

$$(7.5.2) \quad |1 - |\alpha|_y^2| \leq C_c \max_{x_i \in V_M} \left\{ \exp^{-\epsilon_c \sqrt{r}\{\text{dist}(y, x_i) - 2r^{-\frac{1}{2}}\}} \right\}.$$

*Proof of Lemma 7.4.* Define  $d_y$  by

$$d_y := \min \left\{ C_g \sigma(y), \min_{x_i \in V_M} \text{dist}(y, x_i) - 2r^{-\frac{1}{2}} \right\}.$$

Then we see that  $|\alpha|^2 \geq \frac{1}{2}$  on the geodesic ball  $B(y, d_y)$ . Let  $\tilde{e}_y: TX_y \mapsto X$  be the map defined by  $\tilde{e}_y(v) := \exp_y(d_y v)$ . Then the pull-back  $(\underline{\alpha}, \underline{\beta}, \underline{a}) := \tilde{e}_y^*(\alpha, \beta, a)$  is a solution of (3.1) with rescaling parameter  $rd_y^2$ . The pull-back Riemannian metric and symplectic form are a priori bounded in the sense of Definition 1.1. Then the assertion is an immediate consequence of the following:

**Proposition 7.5** ([K-M2]). *Let  $(\alpha, \beta, a)$  be a solution of equations (3.1) defined on the unit ball  $B_1$  with rescaling parameter  $r_1 > 0$  and suppose that  $\eta$  obeys  $\|\eta\|_{C^1(B_1)} \leq e^{-\delta_1 r_1}$  for a positive constant  $\delta_1$ . Then there exist positive constants  $C_u$  and  $\epsilon_1$  that depend only on  $\delta_1$  and on the Riemannian metric and the symplectic form of the unit ball and have the following significance: If  $|\alpha|^2 \geq \frac{1}{2}$  on  $B_1$ , then*

$$(7.6) \quad \sup_{B_{\frac{1}{2}}} \left\{ |1 - |\alpha|^2| + |\beta|^2 + |\nabla_a \alpha|^2 + |\tilde{\nabla}_a \beta|^2 + |F_a| \right\} \leq C_u \exp^{-\epsilon_1 \sqrt{r_1}}.$$

See Proposition 3.22 in [K-M2] for the proof.

We may define the comparison function  $\phi$  by

$$(7.7) \quad \phi(y) := C_c \left\{ \sum_{x_i \in V_M} \Omega_{B(x_i, \sigma(x_i))}(y) \cdot \exp^{-\epsilon_c \sqrt{r} \{ \text{dist}(y, x_i) - 2r^{-\frac{1}{2}} \}} + \exp^{-\epsilon_c \sqrt{r} \{ C_g \sigma(y) \}} \right\},$$

where the symbol  $\Omega_A$  for  $A \subset X$  denotes the characteristic function of  $A$ . Then Lemma 7.4 and the following lemma verify (7.4).

**Lemma 7.6.** *There exists a constant  $C_g$  such that  $0 < C_g \leq 1$  and such that it depends only on  $(\omega, J, \sigma)$  and has the following significance: If  $r$  is sufficiently large and if  $\text{dist}(y, x) \leq C_g \sigma(y) + 2r^{-\frac{1}{2}}$ , then  $\text{dist}(y, x) \leq \sigma(x)$ .*

The lemma above follows from (4) of Property (A) in Definition 1.1.  $\square$

**Step (3).** We will verify the required estimate for the energy integral.

We will estimate first the integral over  $X \setminus X^2$ . It follows from (7.4) that

$$(7.8) \quad \begin{aligned} \int_{X \setminus X^2} |1 - |\alpha|^2| &\leq \int_X \phi \\ &\leq C_c \sum_{x_i \in V_M} \int_{B(x_i, \sigma(x_i))} \exp^{-\epsilon_c \sqrt{r} \{ \text{dist}(y, x_i) - 2r^{-\frac{1}{2}} \}} \\ &\quad + C_c \int_X \exp^{-\epsilon_c \sqrt{r} \{ C_g \sigma(y) \}}. \end{aligned}$$

Lemma 7.3 implies that the first term of the right-hand side of (7.8) is bounded from above by

$$(7.9) \quad C_c(C_v r) C_l \int_{\mathbb{R}^4} \exp^{-\epsilon_c \sqrt{r} (|y| - 2r^{-\frac{1}{2}})}.$$

Here  $C_l$  is a positive constant which depends only on  $(\omega, J)$  and has the following significance: Let  $x \in X$ . Fix an isometry  $P_x : \mathbb{R}^4 \mapsto TX_x$  and define the Riemannian metric  $g_x$  on  $B(0, \sigma(x)) \subset \mathbb{R}^4$  by  $g_x := (\exp_x \circ P_x)^*(g_J)$  where  $g_J$  is the Riemannian metric of  $X$ . Then it follows that  $\int_{\partial B(0, R)} i_{\frac{\partial}{\partial R}} d \text{vol}_{g_i} \leq 4\pi^2 C_l R^3$  where the coordinate  $R$  stands for the distance from the origin. The existence of  $C_l$  is assured by the weak convexity of  $X$ .

The integrand in (7.9) is no more than  $\exp^{2\epsilon_c} \cdot \exp^{-\epsilon_c \sqrt{r} |y|}$ . Thus (7.9) is no more than

$$(7.10) \quad \begin{aligned} &Cr \int_0^\infty dR R^3 \exp^{-\epsilon_c \sqrt{r} R} \\ &= \frac{C}{r} \int_0^\infty dQ Q^3 \exp^{-\epsilon_c Q} \\ &\leq \frac{C}{r}. \end{aligned}$$

The second term of the right-hand side of (7.8) is estimated as follows (see Property (B) of the Definition 1.1):

$$(7.11) \quad \begin{aligned} C_c \int_{\mathbb{R}} dy g_\sigma \exp^{-\epsilon_c \sqrt{r} \{ C_g y \}} &\leq C \int_{\mathbb{R}} dy y^{\max(1, \epsilon_0)} \exp^{-\epsilon_c \sqrt{r} \{ C_g y \}} \\ &\leq Cr^{-\max(1, \frac{1+\epsilon_0}{2})}. \end{aligned}$$

Thus we have obtained

$$(7.12) \quad \int_{X \setminus X^2} |1 - |\alpha|^2| \leq Cr^{-1}.$$

On the other hand, the third property of  $X^2$  and Proposition 5.2 imply that

$$(7.13) \quad \int_{X^2} |1 - |\alpha|^2| \leq C \cdot \text{Vol}(X^2) \leq Cr^{-1}.$$

Therefore, the required estimate (7.2) is verified.  $\square$

## 8. A MONOTONICITY FORMULA FOR LOCAL ENERGY INTEGRALS

We will prove in this section a monotonicity formula for local energy integral, the formula which is a slightly refined version of Proposition 3.2 in [T1].

**Proposition 8.1.** *Let  $(\alpha, \beta, a)$  be a solution of equations (3.1) with  $r \geq 1$ . For  $x \in X$  define the function  $\mathcal{E}_x$  by  $\mathcal{E}_x(R) := \int_{B(x,R)} \frac{r}{4} |1 - |\alpha|^2|$ . There exist positive constants  $\mu_7, \mu_8, \mu_9$  and  $\rho_0$  with  $1 \geq \rho_0 > 0$  which depend only on  $(\omega, J, c_1(L_\varrho))$  and have the following significance:*

*If  $\rho_0 \sigma(x) \geq R \geq 0$ , then*

$$(8.1) \quad \mathcal{E}_x(R) \leq \frac{R}{2} \left( 1 + \mu_7 \frac{R}{\sigma(x)} \right) \left( 1 + \frac{\mu_8}{r^{\frac{1}{4}}} \right) \frac{d}{dR} \mathcal{E}_x(R) + \frac{\mu_9}{r^{\frac{1}{4}}} R^4.$$

We omit the proof since it is exactly the same as that of Proposition 3.2 in [T1]. But it is essential to make use of Proposition 6.1. Our formula is different from the one in [T1] in that it has  $r^{-\frac{1}{4}}$  factor in the second term of the right-hand side, which is due to the existence of the  $r^{-\frac{1}{4}}$  factor in the second term of the right-hand side of (6.1) (and that of the  $r^{-1}$  factor in the second term of the R.H.S. of (5.7)).

By making use of it, we will prove

**Proposition 8.2.** *There exist a positive constant  $C_f$  and a positive function  $\pi_0$  which depend only on  $(\omega, J, C_1(L_\varrho))$  and have the following significance: If  $|\alpha|_x^2 \leq \frac{1}{2}$  and if  $R$  obeys  $r \geq \pi_0(R)$  and  $\rho_0 \sigma(x) \geq R$ , then*

$$(8.2) \quad \mathcal{E}_x(R) \geq \frac{1}{C_f} R^2.$$

*Proof of Proposition 8.2.* We will mimic the proof of Proposition 3.1 in [T1].

Define the function  $f_x$  by

$$f_x(R) := -2 \left( 1 + \frac{\mu_8}{r^{\frac{1}{4}}} \right)^{-1} \log \left( \frac{R}{1 + \mu_7 \frac{R}{\sigma(x)}} \right).$$

It follows from (8.1) that if  $\rho_0 \sigma(x) \geq R \geq 0$ , then

$$(8.3) \quad \frac{d}{dR} (\exp^f \mathcal{E}_x) \geq -Cr^{-\frac{1}{4}} R^3 \exp^f.$$

Fix a positive constant  $R_0$ . Let  $x \in X$  satisfy the condition that  $\rho_0 \sigma(x) \geq R_0$ . There exists a positive constant  $C_{R_0}$  which depends only on  $R_0$  and on  $(\omega, J, c_1(L_\varrho))$  and has the following significance: If  $r^{-\frac{1}{2}} \leq R \leq R_0$ , then  $\exp^{f(R)} \leq C_{R_0} R^{-2}$ . Here we have used the fact that  $\lim_{r \rightarrow \infty} r^{(r^{-\frac{1}{4}})} = 0$ . This implies that if  $r^{-\frac{1}{2}} \leq R \leq R_0$ , then

$$(8.4) \quad \frac{d}{dR} (\exp^f \mathcal{E}_x) \geq -C \cdot C_{R_0} r^{-\frac{1}{4}} R.$$

Integrating (8.4) over  $[r^{-\frac{1}{2}}, R_0]$ , we obtain

$$(8.5) \quad \mathcal{E}_x(R_0) \geq \exp\left(f_x(r^{-\frac{1}{2}}) - f_x(R_0)\right) \left\{ \mathcal{E}_x(r^{-\frac{1}{2}}) - CC_{R_0} r^{-\frac{1}{4}} \cdot \exp^{-f_x(r^{-\frac{1}{2}})} \cdot R_0^2 \right\}.$$

We will estimate the right-hand side of (8.5);

**Lemma 8.3.** *If  $|\alpha|_x^2 \leq \frac{1}{2}$ , then  $\mathcal{E}_x(r^{-\frac{1}{2}}) \geq \zeta_0 r^{-1}$  for some positive constant  $\zeta_0$  that depends only on  $(\omega, J, c_1(L_\varrho))$ .*

On the other hand, we can easily check that

$$\begin{aligned} \lim_{r \rightarrow \infty} r^{-1} \left( \exp\left(f_x(r^{-\frac{1}{2}}) - f_x(R_0)\right) \right) &= \left(1 + \mu_7 \frac{R_0}{\sigma(x)}\right)^{-2} R_0^2, \\ \lim_{r \rightarrow \infty} r \left( C_{R_0} r^{-\frac{1}{4}} \cdot \exp^{-f_x(r^{-\frac{1}{2}})} \cdot R_0^2 \right) &= 0. \end{aligned}$$

Therefore, if  $r$  is no less than a sufficiently large constant  $\pi_0(R_0)$  that depends only on  $R_0$ , we see that

$$(8.6) \quad \mathcal{E}_x(R_0) \geq \frac{1}{2} \left(1 + \mu_7 \frac{R_0}{\sigma(x)}\right)^{-2} \zeta_0 R_0^2.$$

The assumption that  $\rho_0 \sigma(x) \geq R_0$  implies that the coefficient of  $\zeta_0 R_0^2$  in the right-hand side is no less than a positive constant that depends only on  $(\omega, J, c_1(L_\varrho))$ . Thus we are done.  $\square$

*Proof of Lemma 8.3.* Proposition 7.1 implies that there exists a positive constant  $C$  such that  $1 - |\alpha|^2 \geq \frac{1}{4}$  on the geodesic ball  $B(x, \frac{1}{C} r^{-\frac{1}{2}})$  if  $|\alpha|_x^2 \leq \frac{1}{2}$ . Thus it follows that

$$\mathcal{E}_x(r^{\frac{1}{2}}) = \frac{r}{4} \int_{B(x, r^{\frac{1}{2}})} |1 - |\alpha|^2| \geq \frac{r}{16} \text{Vol}\{B(x, r^{-\frac{1}{2}} \cdot \min(1, C^{-1}))\} \geq \frac{1}{Cr}.$$

$\square$

## 9. FINAL ARGUMENTS FOR MAIN THEOREM

Recall that  $C_e$ ,  $\rho_0$  and  $C_f$  are the constants defined in Proposition 7.2, 8.1 and 8.2 respectively and that  $\pi_0$  is the function defined in Proposition 8.2.

**Proposition 9.1.** *There exists a constant  $R_M$  which depends only on  $(\omega, J, c_1(L_\varrho))$  and has the following significance: Let  $(\alpha, \beta, a)$  be a solution of (3.1). If  $x \in X$  obeys  $|\alpha|_x^2 \leq \frac{1}{2}$  and if  $r \geq \pi_0(R_M)$ , then  $\rho_0 \sigma(x) < R_M$ .*

*Proof.* We may take  $R_M$  to be  $\sqrt{2C_e C_f}$ . In fact, suppose that  $|\alpha|_x^2 \leq \frac{1}{2}$ ,  $r \geq \pi_0(R_M)$  and  $\rho_0 \sigma(x) \geq R_M$ . Then Proposition 8.2 implies that

$$\mathcal{E}_x(R_M) \geq \frac{1}{C_f} R_M^2 = 2C_e,$$

which contradicts the assertion of Proposition 7.2 that  $C_e \geq \mathcal{E}_x(R_M)$ .  $\square$

**Corollary 9.2.** *There exist a compact set  $K_M \subset X$  and a positive constant  $C$  which depend only on  $(\omega, J, c_1(L_\varrho))$  and have the following significance: Any solution  $(\alpha, \beta, a)$  of (3.1) with  $r \geq 1$  obeys  $|\alpha|^2 \geq \frac{1}{2}$  on  $X \setminus K_M$ .*

Combining Corollary 9.2 with Proposition 7.5, we immediately obtain

**Proposition 9.3.** *There exist positive constants  $\epsilon_2$  and  $C_d$  which depend only on  $(\omega, J, c_1(L_\varrho))$  and have the following significance: Let  $(\alpha, \beta, a)$  be a solution of (3.1) with  $r \geq 1$ . Then it obeys*

$$|1 - |\alpha|^2| + |\beta|^2 + |\nabla_a \alpha|^2 + |\tilde{\nabla}_a \beta|^2 + |F_a| \leq C_d \exp^{-\epsilon_2 \sqrt{r} \sigma} \quad \text{on } X \setminus K_M.$$

Once Proposition 9.3 is achieved, the arguments in [T1] can be applied to our case almost directly to prove Theorem 4.1. But we need the following two minor modifications to complete the proof:

The first one is in Lemma 3.5 of [T1]. Let  $\Omega_M$  be the domain  $\{x \in X \mid \sigma(x) < 2 \max_{K_M} \sigma\}$ . Then  $\overline{\Omega}_M$  is compact. We may assume that  $\partial\Omega_M$  is smooth. With this understood, we require the function  $u$  in the statement not to be an element of  $C^\infty(X)$  but that of  $C^\infty(\Omega_M) \cap C^0(\overline{\Omega}_M)$  with  $u|_{\partial\Omega_M} \equiv 0$ . Accordingly, we replace the Green function  $G$  in the proof by the fundamental solution of the Dirichlet problem of  $d^*d$  with domain  $\Omega_M$ . Then by making use of the modified  $u$  with Proposition 9.4, we can prove exactly the same result as in (d) of Section 3 in [T1].

The second modification is in Part (1) of the proof of Lemma 4.3 in [T1], where we must bound the function  $|P^+ F_a|^2 - |P^- F_a|^2$  from below by a function  $f$  that obeys  $\|f\|_{L^1(X)} \leq C$  for a constant  $C$  which depends only on  $(\omega, J, c_1(L_\varrho))$ . For this purpose, we may define  $f$  by

$$f := - \begin{cases} \kappa \{r|1 - |\alpha|^2| + 1\} & \text{on } \Omega_M, \\ \kappa C_d^2 \exp^{-2\epsilon_2 \sqrt{r} \sigma} & \text{on } X \setminus \Omega_M, \end{cases}$$

where the constant  $\kappa$  is chosen sufficiently large.

## 10. AN APPLICATION

Let  $\Gamma$  be a discrete subgroup of  $\mathrm{SU}(2)$ . The classification of such groups is well known. They are in one-to-one correspondence with the Dynkin diagrams of type  $A_n$ ,  $D_n$  and  $E_6, E_7, E_8$ .

Let  $Y_1, Y_2$  and  $Y_3$  be the standard basis of the Lie algebra  $\mathfrak{su}(2)$  which we regard as right invariant vector fields. Define the contact 2-plane field  $\zeta$  to be the span of  $Y_1$  and  $Y_2$ , which is called the standard contact structure of the 3-sphere.  $\zeta$  drops to the quotient space  $M_\Gamma := \mathrm{SU}(2)/\Gamma$  as a contact structure denoted by  $\zeta_\Gamma$ .

**Theorem 10.1.** *Let  $(X_0, \omega)$  be a symplectic filling of  $(M_\Gamma, \zeta_\Gamma)$  such that it is minimal.*

- (1) *The intersection form of  $X_0$  is negative definite.*
- (2) *The trivialization of the canonical line bundle  $K_{X_0}$  given over  $\partial X_0$  by  $Y_1$  extends to the interior of  $X_0$ . In particular,  $X_0$  must be a spin manifold.*

Notice that, if we regard  $\zeta_\Gamma$  as a complex line bundle, it is canonically isomorphic to  $K_{X_0}|_{\partial X_0}$ .

*Remark 10.2.* (1) Ohta and Ono [O-O] proved this theorem in the case where the Dynkin diagram of  $\Gamma$  is  $E_8$ , that is,  $M_\Gamma$  is the Poincaré homology of the 3-sphere.

(2) Combining our result with that in [F], we get a good estimate of  $b_2(X_0)$ . In particular, if  $\Gamma$  corresponds to  $E_8$ , the intersection matrix of  $X_0$  must be  $-E_8$ .

*Proof of (2) of Theorem 10.1.* Denote by  $Y_1^*, Y_2^*, Y_3^*$  the standard dual basis of  $\mathfrak{su}(2)^*$  which we regard as right-invariant 1-forms of  $\mathrm{SU}(2)$ . Proposition 1.3 implies that there exists a weakly convex almost Kähler manifold  $(X, \tilde{\omega}, J)$  obeying:

- (1)  $X \cong X_0 \cup_{id} M_\Gamma \times [1, \infty)$  as smooth manifolds.
- (2)  $\tilde{\omega}|_{X_0} = \omega$ .
- (3) There exists a positive constant  $l \geq 1$  such that  $\tilde{\omega}|_{M_\Gamma \times [l, \infty)} = d(t^2 Y_3^*)$  and such that  $J|_{M_\Gamma \times [l, \infty)}$  obeys the formulae  $J(Y_1) = Y_2, J(Y_2) = -Y_1, J(\partial_t) = Y_3, J(Y_3) = -\partial_t$ .

For simplicity, we may assume that  $l = 1$ .

All through the later arguments, we regard  $X_0$  as a subset of  $X$  and denote by  $X^+$  the conical end  $M_\Gamma \times [1, \infty)$ .

Define the 1-parameter family of symplectic forms  $\{\omega_\nu\}_{0 \leq \nu \leq 1}$  on  $X^+$  by

$$\omega_\nu := d\{t^2((\cos \pi\nu)Y_3^* + (\sin \pi\nu)Y_1^*)\}$$

These  $\omega_\nu$  are self-dual 2-forms of length  $\sqrt{2}$  with respect to  $g_J|_{X^+}$ . Hence, for each  $\nu$  there exists a unique almost complex structure  $J_\nu$  compatible with  $\omega_\nu$  such that the associated metric  $\omega_\nu(*, J*)$  coincides with  $g_J|_{X^+}$ . Then we see that  $(\tilde{\omega}, J)|_{X^+} = (\omega_0, J_0) = (-\omega_1, -J_1)$ .

*Remark.* These  $J_\nu$  are integrable. In fact,  $g_J|_{X^+}$  is a hyper-Kähler metric.

For the time being, we fix an element  $(s, \varrho) \in S(X, \omega)$ .

Let  $\mathbb{I}$  be the unit length section of  $W_s|_{X^+}$  given as the pull-back of  $(1, 0) \in \Gamma(W_{s_{\tilde{\omega}}}) \equiv \Gamma(\mathbb{C} \oplus K)$  through the identification map  $\varrho: W_s|_{X^+} \mapsto W_{s_{\tilde{\omega}}}|_{X^+}$ . Notice that  $\rho(\tilde{\omega})\mathbb{I} = -2\sqrt{-1}\mathbb{I}$ . We see that there exists a smooth 1-parameter family of unit length sections  $\{\mathbb{I}_\nu\}_{0 \leq \nu \leq 1}$  which obeys the equation  $\rho(\omega_\nu)\mathbb{I}_\nu = -2\sqrt{-1}\mathbb{I}_\nu$  and the initial condition  $\mathbb{I}_0 = \mathbb{I}$ . This induces a smooth 1-parameter family of the isomorphisms  $\varrho_{\mathbb{I}_\nu}: W_s|_{X^+} \mapsto W_{s_{\tilde{\omega}}}|_{X^+}$  by imposing the condition  $\varrho_{\mathbb{I}_\nu}(\mathbb{I}_\nu) = (1, 0)$ . Notice that  $\varrho_{\mathbb{I}_0} = \varrho$ . We can easily show that the family  $\{(\omega_\nu, J_\nu, \varrho_{\mathbb{I}_\nu})\}_{0 \leq \nu \leq 1}$  satisfies the assumption in the statement of the first property of the monopole invariant  $SW$  (see Section 1). Thus it follows that

$$SW(X, \tilde{\omega}, J, s, \varrho) = \pm SW(X, -\tilde{\omega}, -J, s, \varrho_{\mathbb{I}_1}).$$

Let  $-s$  be the  $\text{Spin}^c$  structure obtained from  $s$  by changing the sign of the complex structure. Then  $W_{-s}$  is canonically isomorphic to  $W_s$  as a real vector bundle. Denote by  $\overline{\mathbb{I}}_1$  the section of  $W_{-s}$  that corresponds to  $\mathbb{I}_1$ . Then  $\rho(\tilde{\omega})\overline{\mathbb{I}}_1 = -2\sqrt{-1}\overline{\mathbb{I}}_1$  since  $\rho(\tilde{\omega})\mathbb{I}_1 = 2\sqrt{-1}\mathbb{I}_1$ . Since the change of the sign of the complex structure does not affect the underlying equation, we obtain

$$SW(X, -\tilde{\omega}, -J, s, \varrho_{\mathbb{I}_1}) = \pm SW(X, \tilde{\omega}, J, -s, \varrho_{\overline{\mathbb{I}}_1}).$$

Now suppose that the element  $(s, \varrho)$  that we have fixed so far to be  $(s_{\tilde{\omega}}, \text{id})$ . Since  $SW(X, \tilde{\omega}, J, s_{\tilde{\omega}}, \text{id}) = 1$  (see Section 1), by combining the two formulae above, we have

$$SW(X, \tilde{\omega}, J, -s_{\tilde{\omega}}, \varrho_{\overline{\mathbb{I}}_1}) = \pm 1.$$

The corresponding line bundle to  $-s_{\tilde{\omega}}$  is  $K$  since  $W_{-s_{\tilde{\omega}}} = K \oplus \mathbb{C} = (\mathbb{C} \oplus \overline{K}) \otimes K$ . Hence, we have only to show that  $(K, \varrho_{\overline{\mathbb{I}}_1})$  coincides with  $(\mathbb{C}, \text{id})$ .

Assume to the contrary. Then applying Theorem 4.1, we obtain a non-empty, compact  $J$ -holomorphic curve  $D \subset X$  such that  $P.D.[D] = c_1(K, \varrho_{\overline{\mathbb{I}}_1}) \in H_{\text{cpt}}^2(X, \mathbb{Z})$ . Taking multiplicities into account,  $D$  is written as

$$D = \sum_{i=1}^k n_i D_i$$



where the  $D_i$ 's are mutually distinct and non-multiple such that each  $D_i$  is the image of a non-constant  $J$ -holomorphic map from a connected compact Riemann surface.  $n_i$  is a positive integer that represents the multiplicity of  $D_i$  in  $D$ . The minimality of  $(X_0, \omega)$  means that  $(X, \tilde{\omega}, J)$  contains no embedded  $J$ -holomorphic rational curve whose self-intersection number is  $-1$ . Thus the argument in the proof of Proposition 7.1 in [T1] shows that, if  $J$  is generic, each  $D_i$  is a smooth submanifold,  $D_i \cap D_j = \emptyset$  for  $i \neq j$  and  $D_i \cdot D_i \geq 0$ . Since  $\partial X_0$  is a rational homology 3-sphere, the intersection form of  $X$  is non-degenerate. Further,  $[D_i] \in H_2(X, \mathbb{Z})$  is not zero since  $\int_{D_i} \tilde{\omega} > 0$ . Thus it follows that  $b_2^+(X_0) \geq 1$ , which contradicts assertion (1).  $\square$

*Remark 10.3.* In general, a compact  $J$ -holomorphic curve  $D$  in a weakly convex almost Kähler manifold  $(X, \omega, J)$  is contained in a compact set  $K_{[D]}$  which is determined a priori by the value  $\langle [D], [\omega] \rangle$  due to the monotonicity formula of energy density. Thus to have  $J$  generic, it is sufficient to consider the space  $\mathfrak{J}$  of compatible complex structures which agree with a fixed almost complex structure outside a fixed compact set. In fact, we can show that there exists a Baire subset of  $\mathfrak{J}$  whose elements have the needed genericity for all  $J$ -holomorphic curves of a fixed homology class.

*Proof of (1) of Theorem 10.1.* This follows from the standard necking argument in gauge theory, which is well-known by the experts. Hence, we will give here only the sketch of the proof. See [M-S-T]) and [F] for the details.

We will derive a contradiction by assuming that  $b_2^+(X_0) > 0$ .

Let  $X$  and  $X^+$  be as defined in the proof of assertion (1). Perturb the Riemannian metric of  $X$  only near  $\partial X_0 (= -\partial X^+)$  so that some regular neighborhood of  $\partial X_0$  is isometric to the Cartesian product of  $(M_\Gamma, g_M)$  with a small open interval and so that  $\partial X_0$  is totally geodesic. Here  $g_M$  is the standard Riemannian metric of  $M_\Gamma$ . Splitting  $X$  along  $\partial X_0$  into pieces and gluing back  $(X_R^m, g_R^m) := (M_\Gamma \times [-R, R], g_M + dt^2)$  with  $R > 1$  between them, we obtain the new Riemannian manifold  $(X_R, g_R)$  with no boundary.

Fix an element  $(s, \varrho) \in S(X, \tilde{\omega})$  and consider the following monopole equation on  $X_R$  with variables  $(\Phi, B) \in \Gamma(W_s^+) \times \mathcal{A}(\det W_s^+)$ :

$$\mathcal{D}_a \Phi = 0,$$

$$F_a^+ = \rho^{-1}(\Phi \Phi^*)_0 - \frac{\sqrt{-1}}{2} \tau^+ \cdot \tilde{\omega} + \tau_m \cdot \text{Pr}^+ \{ \pi^*(\mu_m) \} + \mu_1 + \mu_2.$$

We will explain the notation in order:

1.  $\rho^{-1}: \sqrt{-1} \text{SkewEnd}(W_s^+) \mapsto \Lambda^+$  is the inverse of the bundle isomorphism  $\rho$ .
2.  $\tau^+ \in C^\infty(X_R)$ ,  $\text{supp}(\tau^+) \subset X^+$  and  $\tau_+ \equiv 1$  outside some compact set.
3.  $\tau_m \in C^\infty(X_R)$ ,  $\text{supp}(\tau_m) \subset X_R^m$  and  $\tau_m \equiv 1$  on  $M_\Gamma \times [-R+1, R-1] \subset X_R^m$ .
4.  $\mu_m$  is an exact 2-form of  $M_\Gamma$ .
5.  $\pi: M_\Gamma \times [-R, R] \mapsto M_\Gamma$  is the natural projection.
6.  $\mu_1$  is a self-dual 2-form such that  $\text{supp}(\mu_1) \subset X_0$ .
7.  $\mu_2$  is a self-dual 2-form such that  $\text{supp}(\mu_2) \subset X^+$  and such that  $\|\mu_2\|_{\mathcal{N}(X^+)} < \infty$ .

Notice that this equation can be written in the form of (3.1) with  $r = 1$  when restricted to the complement of a sufficiently large compact set. We impose the

same boundary condition as in Section 3 to  $(\Phi, B)$  and define the moduli space  $\mathcal{M}_R$  to be the gauge equivalence classes of the solutions.

**Assertion 10.4.** *For the generic choice  $(\mu_m, \mu_1, \mu_2)$  (which can be taken smooth), the equation is transverse, that is, the linearized equation at any solution is surjective.*

The important feature of this equation is that it restricts to  $X_R^m \cong M_\Gamma \times [-R, R]$  as the gradient flow equation of the perturbed Chern-Simons-Dirac functional  $C.S.D_{\mu_m}$  when adopting temporal gauge. In general, a stationary point of  $C.S.D_{\mu_m}$  is a solution of the following, reduced-to-3-dimensional monopole equation:

$$\begin{aligned}\mathcal{D}_B \Phi &= 0, \\ \rho(F_B)|_{W_{s'}^+} &= (\Phi \Phi^*)_0 + \mu_m.\end{aligned}$$

Here the  $\text{Spin}^c$  structure  $s'$  on  $M_\Gamma$  is such that  $\pi^*(s') \cong s|_{X_R^m}$ . In general, the same argument that derives the a priori  $C^0$  estimate of monopole equations shows that if  $\|\mu_m\|_{C^0}$  is sufficiently small and if the Riemannian metric has positive scalar curvature, there are only reducible solutions, that is,  $\Phi \equiv 0$ . Thus we have

**Lemma 10.5.** *The stationary point of  $C.S.D_{\mu_m}$  on  $M_\Gamma$  is reducible and unique up to gauge equivalence if  $\mu_m$  is sufficiently small. Further, for generic  $\mu_m$  the Hessian of  $C.S.D_{\mu_m}$  at the stationary point  $(0, B_0)$  is non-degenerate when restricted to the orthogonal complement of the tangent space of the gauge orbit containing  $(0, B_0)$ .*

Gluing the half cylinder  $(M_\Gamma \times [0, \infty), g_M + dt^2)$  along the boundary to  $X_0$  and  $X^+$  respectively, we get two Riemannian manifolds with no boundary denoted by  $X_1$  and  $X_2$  respectively. Assume that  $\mathcal{M}_R$  be non-empty for all sufficiently large  $R$ . Then taking  $R$  to infinity and following the standard necking argument (see [M-S-T]), we obtain on each  $X_i$  a solution  $(\Phi_i, B_i)$  described as follows:

(1)  $(\Phi_1, B_1) \in \Gamma(W_{s^+}^+|_{X_1}) \times \mathcal{A}(\det W_{s^+}^+|_{X_1})$  obeys

$$\begin{aligned}\mathcal{D}_{B_1} \Phi_1 &= 0, \\ F_{B_1}^+ &= \rho^{-1}(\Phi \Phi^*)_0 + \mu_1 + \tau \cdot \text{Pr}^+ \{ \pi^*(\mu_m) \}, \\ \|(\Phi_1, B_1)|_{M_\Gamma \times [0, \infty)} - \pi^*(0, B_0)\|_{W^{k,2}(M_\Gamma \times [0, \infty))} &< \infty.\end{aligned}$$

(2)  $(\Phi_2, B_2) \in \Gamma(W_{s^+}^+|_{X_2}) \times \mathcal{A}(\det W_{s^+}^+|_{X_2})$  obeys

$$\begin{aligned}\mathcal{D}_{B_2} \Phi_2 &= 0, \\ F_{B_2}^+ &= \rho^{-1}(\Phi \Phi^*)_0 - \frac{\sqrt{-1}}{2} \tau^+ \cdot \tilde{\omega} + \mu_2 + \tau \cdot \text{Pr}^+ \{ \pi^*(\mu_m) \}, \\ \|(\Phi_2, B_2)|_{M_\Gamma \times [0, \infty)} - \pi^*(0, B_0)\|_{W^{k,2}(M_\Gamma \times [0, \infty))} &< \infty, \\ \|(\Phi_2, B_2)|_{X^+} - (\mathbb{I}, B_d)\|_{W^{k,2}(X^+)} &< \infty.\end{aligned}$$

Here  $k$  is fixed sufficiently large and  $\tau \in C_0^\infty(M_\Gamma \times [0, \infty))$  such that  $\tau \equiv 1$  on  $M_\Gamma \times [1, \infty)$ .  $B_d$  in (2) is the connection of  $\det(W_{s_\omega})$  that corresponds to the trivial connection of the trivial line bundle. (See Section 2.)

Now we have to define suitable moduli spaces  $\mathcal{M}_i$  in which the gauge equivalence classes of the solutions  $(\Phi_i, B_i)$  should live respectively. The usual Sobolev norm is not suitable for this purpose since the “boundary value”  $(0, B_0)$  is reducible. We have to adopt a weighted Sobolev norm  $\|*\|_{W_\delta^{2,k}}$  which is in the form  $\|f\|_{W_\delta^{2,k}} :=$

$\sum_{i=1}^k \{\int e^{\delta t} |\nabla^k f|^2\}^{\frac{1}{2}}$  in the cylindrical end and agrees with the usual one in the complement. Here  $\delta$  is a sufficiently small positive constant and  $f$  stands for a 1-form, a section of the spinor bundle and so on. Anyway, we can construct the moduli spaces  $\mathcal{M}_i$  according to the standard procedure.

Any solution  $(\Phi_2, B_2)$  is irreducible since  $\Phi_2$  approaches asymptotically to the unit length section  $\mathbb{I}$  at the infinity of the conical end. As for  $(\Phi_1, B_1)$ , its irreducibility is assured by the assumption that  $b_2^+(X_1) > 0$ . In fact, we can choose  $\mu_1$  from the complement of the affine subspace

$$F_{B_b}^+ - \tau \cdot \text{Pr}^+\{\pi^*(\mu_m)\} + \text{Image}\{d^+ : \Omega_{W_\delta^{k,2}}^1(X_1) \mapsto \Omega_{W_\delta^{k-1,2}}^+(X_1)\}$$

because the codimension is no less than 1. Here  $B_b$  is a fixed base point of connections such that  $F_{B_b}^+|_{M_\Gamma \times [1, \infty)} = \text{Pr}^+\{\pi^*(\mu_m)\}$ . The second line of the equation prevents  $\Phi_1$  from vanishing identically.

Thus we have

**Proposition 10.6.** 1. If  $\mathcal{M}_R$  are non-empty for all sufficiently large  $R$ , then neither  $\mathcal{M}_1$  nor  $\mathcal{M}_2$  are empty.  
2. If  $b_2^+(X_0) > 0$  and if the perturbation is generic,  $\mathcal{M}_i$  are finite dimensional smooth manifolds such that  $\dim \mathcal{M} = \dim \mathcal{M}_1 + \dim \mathcal{M}_2 + 1$ .

*Remark 10.7.* The term 1 in the right-hand side of the formula in (2) is the dimension of  $U(1)$ , which is the isotropy subgroup at  $(0, B_0)$  of the gauge group.

Now suppose  $(s, \varrho)$  to be  $(s_{\tilde{\omega}}, \text{id})$ . Then  $\dim \mathcal{M}_R = 0$  and  $\mathcal{M}_R$  is non-empty (see Section 1). Then the first assertion of Proposition 10.6 implies that  $\mathcal{M}_1$  and  $\mathcal{M}_2$  are both non-empty. But the second assertion of Proposition 10.6 implies that either  $\mathcal{M}_1$  or  $\mathcal{M}_2$  is a negative dimensional manifold and thus must be empty. This is a contradiction.  $\square$

## 11. APPENDIX

We will prove Lemma 6.3.

Denote by  $L$  the operator  $\Delta + \frac{r}{2}|\alpha|^2$  acting on  $C_0^\infty(X)$ .

**Lemma 11.1.** *There exist a compact set  $K \subset X$  and a positive constant  $C$  such that, for an arbitrary  $f \in C_0^\infty(X)$ , it holds that*

$$(11.0) \quad \|f\|_{W_0^{2,2}(X)} \leq C \left( \|Lf\|_{L^2(X)} + \|f\|_{L^2(K)} \right).$$

*Proof.* A short calculation shows that

$$\nabla^* \nabla \nabla f = \nabla \Delta f + \text{Ric}(\nabla f, *)$$

where  $\text{Ric}$  means the Ricci curvature of the Riemannian metric. Taking the inner product of both sides with  $\nabla f$ , integrating the result over  $X$  and using an integration by parts, we obtain

$$(11.1) \quad \int_X |\nabla \nabla f|^2 = \int_X |\Delta f|^2 + \int_X \text{Ric}(\nabla f, \nabla f).$$

On the other hand, the very definition of  $L$  and an integration by parts imply that

$$(11.2) \quad \int_X |\nabla f|^2 = \int_X \langle Lf, f \rangle - \int_X \phi |f|^2$$

where  $\phi$  denotes the function  $\frac{r}{2}|\alpha|^2$ . Choose a compact set  $K$  sufficiently large so that  $\phi \geq \frac{1}{2}$  on the complement of  $K$ . (See Proposition 3.2.) Then Hölder's inequality implies that

$$(11.3) \quad \int_X |\nabla f|^2 \leq \frac{1}{2} \int_X |Lf|^2 + \frac{1}{2} \int_K |f|^2.$$

Further, integrating the identity

$$|\Delta f|^2 = |Lf|^2 - 2\langle \Delta f, \phi f \rangle - \phi^2 |f|^2$$

over  $X$  and making use of an integration by parts and Schwarz' inequality, we obtain

$$\int_X |\Delta f|^2 \leq \int_X |Lf|^2 - \int_X (2\phi |\nabla f|^2 + \phi^2 |f|^2) + 2 \int_X |f| \cdot |\nabla f| \cdot |\nabla \phi|.$$

By choosing  $K$  sufficiently large, we may assume that  $|\nabla \phi| \leq \frac{1}{4}$  on the complement of  $K$ . (See Proposition 3.2.) Thus we obtain

$$(11.4) \quad \int_X |\Delta f|^2 \leq \int_X |Lf|^2 - \frac{1}{8} \int_{X-K} (|\nabla f|^2 + |f|^2) + C \int_K (|\nabla f|^2 + |f|^2).$$

The weak convexity implies that  $|\text{Ric}|$  is bounded over  $X$ . Thus combining the inequalities (11.1), (11.3) and (11.4), we get the required inequality.  $\square$

*Proof of Lemma 6.3.* It is trivial that  $L$  extends uniquely to the symmetric operator  $\tilde{L}$  over  $L^2(X)$  with domain  $W_0^{2,2}(X)$ .

First, we will show that  $\tilde{L}$  is self-adjoint. Suppose that  $u, v \in L^2(X)$  satisfy

$$(11.5) \quad \langle u, \tilde{L}f \rangle = \langle v, f \rangle$$

for an arbitrary  $f \in W_0^{2,2}(X)$ . Let  $\{u_n\}_{n \in \mathbb{N}} \subset C_0^\infty(X)$  converges strongly to  $u$  with respect to the  $L^2(X)$  norm. It follows by using an integration by parts that

$$\langle u_n, \tilde{L}f \rangle = \langle Lu_n, f \rangle$$

which implies that  $\{Lu_n\}_{n \in \mathbb{N}}$  converges weakly to  $v$  with respect to the  $L^2(X)$  norm. Thus  $\{\|Lu_n\|_{L^2(X)}\}_{n \in \mathbb{N}}$  is bounded. Then Lemma 11.1 implies  $\{\|u_n\|_{W^{2,2}(X)}\}_{n \in \mathbb{N}}$  is also bounded. By passing to a suitable subsequence, we may assume that  $\{u_n\}_{n \in \mathbb{N}}$  converges weakly to an element  $u_0 \in W_0^{2,2}(X)$  with respect to the  $W^{2,2}(X)$  norm. It follows from (11.5) that  $u_0 = u$  almost everywhere, that is,  $u \in W_0^{2,2}(X)$ .

Second, we will show

**Lemma 11.2.** *If a sequence  $\{u_n\}_{n \in \mathbb{N}} \subset W_0^{2,2}(X)$  satisfies that  $\lim_{n \rightarrow \infty} \|\tilde{L}u_n\|_{L^2(X)} = 0$  and  $\|u_n\|_{L^2(X)} = 1$ , then  $\{u_n\}_{n \in \mathbb{N}}$  converges strongly to an element  $u_0 \in \text{Ker } \tilde{L}$  with respect to the  $W^{2,2}(X)$  norm.*

It is easy to check that this lemma derives immediately the closedness of  $\text{Im } \tilde{L}$  in  $L^2(X)$  and the finite dimensionality of  $\text{Ker } \tilde{L}$ .

The inequality (11.0) implies that  $\{u_n\}_{n \in \mathbb{N}}$  is bounded in  $W_0^{2,2}(X)$ . Thus by passing to a suitable subsequence, we may assume that  $\{u_n\}_{n \in \mathbb{N}}$  converges weakly to an element  $u_0$  with respect to the  $W_0^{2,2}(X)$  norm. Then the Sobolev embedding theorem implies that  $\{u_n|_K\}_{n \in \mathbb{N}}$  converges strongly to  $u_0|_K$  in  $L^2(X)$ . (It is this part where we have to use the compactness of  $K$ .) Applying (11.0) to  $\{u_n - u_0\}_{n \in \mathbb{N}}$ , we can show that  $\{u_n\}_{n \in \mathbb{N}}$  converge strongly to  $u_0$  in  $W_0^{2,2}$ .

Third, we will show that  $\text{Ker } \tilde{L} = \{0\}$ , which in the same time implies that  $\text{Coker } \tilde{L} = \{0\}$ . Suppose  $\tilde{L}u = 0$ . The local elliptic regularity means that  $u$  is smooth. On the other hand, (11.2) implies that  $\int_X |\nabla u|^2 = 0$ . Thus  $u$  is a constant function. Since  $u \in L^2(X)$ ,  $u \equiv 0$ .  $\square$

*Remark 11.3.* Due to the weak convexity, we can show that  $W_0^{2,2}(X)$  coincides with the function space  $W^{2,2}(X)$  that consists of  $L^2$ -functions whose distributional derivatives of order 1 and 2 are realized as  $L^2$ -functions. (See [K-M2].)

## REFERENCES

- [E] Y.Eliashberg, *Contact 3-manifolds, twenty years since Martinet's work*, Ann. Inst. Fourier. **42** (1992), 165-192. MR **93k**:57029
- [F] K.Fløyshov, *The Seiberg-Witten equations and 4-manifolds with boundary*, Math. Res. Lett. **3** (1996), 373-390. MR **97i**:57037
- [Gr] M.Gromov, *Pseudo-holomorphic curves in symplectic manifolds*, Invent. Math. **82** (1985), 307-347. MR **87j**:53053
- [G-T] D.Gilbarg and N.S.Trudinger, *Elliptic partial differential equalities of second order*, 2nd ed., Springer-Verlag, 1983. MR **86c**:35035
- [K-M1] P.B.Kronheimer and T.S.Mrowka, *The genus of embedded surfaces in the projective plane*, Math. Res. Lett. **1** (1994), 797-808. MR **96a**:57073
- [K-M2] ———, *Monopoles and contact structures*, Inv. Math. **130** (1997), 209-255. MR **98h**:57058
- [Ko] D.Kotschick, *The Seiberg-Witten invariants of symplectic four-manifolds [after C.H.Taubes]*, Séminaire Bourbaki 48ème année n°812 (1995-1996). MR **98h**:57057
- [Ma] R.Mandelbaum, *Irrational connected sums*, Trans. Amer. Math. Soc. **247** (1979), 137-156. MR **80e**:57023
- [M-S-T] J.W.Morgan, Z.Szabó and C.H.Taubes, *A product formula for the Seiberg-Witten invariants and the generalized Thom conjecture*, Jour. Diff. Geom. **44** (1996), 706-788. MR **97m**:57052
- [O-O] H.Ohta and K.Ono, *Simple singularities and topology of symplectically filling 4-manifold*, Comment. Math. Helv. **74** (1999), 575-590. CMP 2000:06
- [T1] C.H.Taubes, *SW  $\Rightarrow$  Gr: From the Seiberg-Witten equations to pseudo-holomorphic curves*, J. Amer. Math. Soc. **9** (1996), 845-918. MR **97a**:57053
- [T2] ———, *Gr  $\Rightarrow$  SW: From pseudo-holomorphic curves to Seiberg-Witten solutions*, J. Differential Geom. **51** (1999), 203-334. MR **2000i**:53123
- [T3] ———, *Counting pseudo-holomorphic curves in dimension 4*, preprint.
- [W] E.Witten, *Monopoles and four-manifolds*, Math. Res. Lett. **1** (1994), 769-796. MR **96d**:57035

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO 060-0810, JAPAN  
*E-mail address:* `kanda@math.sci.hokudai.ac.jp`