

METRIC PROPERTIES OF THE GROUP OF AREA PRESERVING DIFFEOMORPHISMS

MICHEL BENAÏM AND JEAN-MARC GAMBAUDO

ABSTRACT. Area preserving diffeomorphisms of the 2-disk which are identity near the boundary form a group \mathcal{D}_2 which can be equipped, using the L^2 -norm on its Lie algebra, with a right invariant metric. With this metric the diameter of \mathcal{D}_2 is infinite. In this paper we show that \mathcal{D}_2 contains quasi-isometric embeddings of any finitely generated free group and any finitely generated abelian free group.

1. INTRODUCTION

Let \mathcal{D}_2 be the group of smooth area preserving diffeomorphisms of the 2-disk which are identity near the boundary of the disk. A tangent vector to \mathcal{D}_2 at a point ϕ is a divergence free vector field X_ϕ (see [1] and [5]) whose L^2 -norm is defined by

$$\|X_\phi\|_2 = \left(\int_{\mathbf{D}^2} \|X_\phi(x)\|^2 dx \right)^{1/2},$$

where $\|\cdot\|$ stands for the standard Euclidean norm. Consider a path $\{\phi_t\}_{t \in [0,1]}$ in \mathcal{D}_2 connecting two maps ϕ_0 and ϕ_1 . The length of the path $\{\phi_t\}$ is given by the formula

$$l_2(\{\phi_t\}) = \int_0^1 \left\| \frac{d\phi_t}{dt} \right\|_2 dt.$$

Any two maps in \mathcal{D}_2 being connected by a path with finite length l_2 , induces a distance on \mathcal{D}_2 defined by $d_2(\phi_0, \phi_1) = \inf l_2(\{\phi_t\})$, where the infimum is taken over all paths joining ϕ_0 to ϕ_1 . Notice that this distance is right invariant:

$$d_2(\phi_0 \circ \phi, \phi_1 \circ \phi) = d_2(\phi_0, \phi_1).$$

Equipped with this distance, \mathcal{D}_2 has an infinite diameter (see [13], [6] and [1]). The group of area preserving diffeomorphisms is important in hydrodynamics; it is the configuration space of an incompressible 2-dimensional fluid (see [1]). The Euler-Lagrange equation associated to the metric d_2 corresponds to the equation of the motion of a perfect incompressible flow (Euler equation).

Finitely generated groups, i.e. groups generated with a finite number of generators obeying a finite number of relations, are another standard class of groups equipped with a right invariant metric. Let \mathcal{G} be such a group, and \mathcal{S} a system of generators of \mathcal{G} . The *length* $l_{\mathcal{S}}(g)$ of an element g in \mathcal{G} is the minimal number of

Received by the editors April 11, 2000 and, in revised form, October 30, 2000.

1991 *Mathematics Subject Classification*. Primary 20F36, 58B05, 58B25, 76A02.

Key words and phrases. Area preserving diffeomorphisms, braids, free groups, quasi-isometry.

generators in \mathcal{S} and their inverses needed to write g . It defines a right invariant natural metric on \mathcal{G} :

$$d_{\mathcal{S}}(g_1, g_2) = l_{\mathcal{S}}(g_1 g_2^{-1}).$$

The free group \mathbf{F}_n and the abelian free group \mathbf{Z}^n with n generators are classical examples of finitely generated groups.

Let (\mathcal{G}_1, d_1) and (\mathcal{G}_2, d_2) be two metric groups, a morphism $m : \mathcal{G}_1 \rightarrow \mathcal{G}_2$ is a *quasi-expanding embedding* if:

- m is injective;
- there exists a pair of positive constants $k, k' > 0$ such that for any pair g_1, g'_1 in \mathcal{G}_1 :

$$d_2(m(g_1), m(g'_1)) \geq k d_1(g_1, g'_1) - k'.$$

An isomorphism m between two metric groups is a *quasi-isometry* if m and its inverse are quasi-expanding embeddings. A *quasi-isometric embedding* is a morphism which is a quasi-isometry onto its image. We refer the reader to [10] for more details and results concerning quasi-isometries.

In the case of a finitely generated group \mathcal{G} equipped with 2 systems of generators \mathcal{S} and \mathcal{S}' , we have for any g in \mathcal{G} :

$$(1) \quad \frac{1}{k(\mathcal{S}', \mathcal{S})} l_{\mathcal{S}}(g) \leq l_{\mathcal{S}'}(g) \leq k(\mathcal{S}, \mathcal{S}') l_{\mathcal{S}}(g),$$

where $k(\mathcal{S}, \mathcal{S}')$ stands for the maximum over all the generators s in \mathcal{S} of the length $l_{\mathcal{S}'}(s)$. Thus the identity map is a quasi-isometric isomorphism from $(\mathcal{G}, d_{\mathcal{S}})$ to $(\mathcal{G}, d_{\mathcal{S}'})$ where $d_{\mathcal{S}}$ and $d_{\mathcal{S}'}$ stand respectively for the right invariant metrics induced by \mathcal{S} and \mathcal{S}' . It follows that for a morphism of metric groups with values in (or taking values in) a finitely generated group $(\mathcal{G}, d_{\mathcal{S}})$ (where $d_{\mathcal{S}}$ is the right invariant metric induced by a system of generators \mathcal{S}) the property of being a quasi-isometric embedding does not depend on the set of generators.

This note is an attempt to understand the anatomy of the group \mathcal{D}_2 seen as a metric space. Since closed balls are not compact in \mathcal{D}_2 , a direct approach using growth of balls seems pointless. Our point of view here is to show the complexity of \mathcal{D}_2 by showing that very different finitely generated groups can be embedded by quasi-isometric morphisms in \mathcal{D}_2 . We prove the following result:

Theorem 1. *Any finitely generated free group and any finitely generated abelian free group is quasi-isometrically embedded in \mathcal{D}_2 .*

The keys for this theorem rely on the following two facts that we will prove and make more explicit in the sequel:

- any finitely generated free group and any finitely generated abelian free group can be mapped by a quasi-isometric embedding in a braid group;
- the braid subgroups obtained this way can be in turn mapped by a quasi-isometric embedding in the group \mathcal{D}_2 .

Finally, we conclude with three remarks related to Theorem 1 and its possible generalizations.

2. THE FREE GROUPS AND THE ABELIAN FREE GROUPS

For any integer $n > 0$ we denote by \mathbf{F}_n (resp. \mathbf{Z}^n) the free group (resp. the abelian free group) with n generators e_1, \dots, e_n and d_n the corresponding right

invariant metric. For any pair of integers $2 \leq r \leq s$, there exists a quasi-isometric embedding from the (abelian) free group with r generators to the (abelian) free group with s generators. In the case of the free groups the converse is also true, more precisely we have the following standard result:

Lemma 1. *For any $n \geq 2$, there exists a morphism from the free group with n generators \mathbf{F}_n to the free group with 2 generators \mathbf{F}_2 which is a quasi-isometric embedding.*

Proof. For a given n , one can choose an integer r large enough, and a system of n elements in \mathbf{F}_2 , g_1, \dots, g_n , such that

- these words can be written using only the generators e_1 and e_2 of \mathbf{F}_2 (and not their inverses);
- they have a same length equal to $10r$;
- for any pair i, j in $1, \dots, n$ the word $g_i g_j^{-1}$ has length at least $18r$.

It follows that the morphism $\mu : \mathbf{F}_n \rightarrow \mathbf{F}_2$ which associates to each generator e_i of \mathbf{F}_n the element g_i satisfies

$$8rd_n(f_1, f_2) \leq d_2(\mu(f_1), \mu(f_2)) \leq 10rd_n(f_1, f_2),$$

for any pair of elements f_1, f_2 in \mathbf{F}_n .

The second inequality shows that μ is an injective morphism; combined with the first one, we get that it is a quasi-isometric embedding. \square

3. THE BRAID GROUP AND ITS SUBGROUPS

The *Artin Braid group* \mathbf{B}_n is a group given by the set of generators $\sigma_1, \sigma_2, \dots, \sigma_{n-1}$ and the relations

$$\sigma_i \sigma_j = \sigma_j \sigma_i,$$

for all i, j in $\{1, \dots, n-1\}$ with $|i-j| \geq 2$ and

$$\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$$

for all i in $\{1, \dots, n-2\}$. An element of this group is called a *braid*.

A geometrical way to represent braids (see for instance [3] and [12]) consists in fixing the $2n$ points $P_i = (i/(n-1) - 1/2, 0, 1)$ and $Q_i = (i/(n-1) - 1/2, 0, 0)$ for $i = 0, \dots, n-1$ in the solid cylinder $\mathbf{D}^2 \times [0, 1]$. A braid β can be seen as the isotopy class of a system of n non-intersecting arcs, joining each point P_i to a point $Q_{\tau_\beta(i)}$, where τ_β is a permutation on $\{0, \dots, n-1\}$, and such that the intersection of any of these arcs with any disk $\mathbf{D}^2 \times \{t\}$, t in $[0, 1]$, consists in a unique point. With this representation, it is easy to understand what the group law, the generators and the relations mean.

There exists a classical surjective representation ρ of the braid group \mathbf{B}_3 in the group $\mathbf{PSL}(2, \mathbf{Z})$ which is defined by

$$\rho(\sigma_1) = s_1 \quad \text{and} \quad \rho(\sigma_2) = s_2,$$

where s_1 and s_2 are respectively the representatives in $\mathbf{PSL}(2, \mathbf{Z})$ of the following two elements in $\mathbf{SL}(2, \mathbf{Z})$:

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}.$$

It is a representation since the only relation in \mathbf{B}_3

$$\sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2$$

is preserved by ρ :

$$s_1 s_2 s_1 = s_2 s_1 s_2.$$

It is surjective since $\mathbf{PSL}(2, \mathbf{Z})$ is generated by the two elements s_1 and s_2 . The two elements in $\mathbf{PSL}(2, \mathbf{Z})$,

$$a = s_1 s_2 \text{ and } b = s_1 s_2 s_1,$$

satisfy $a^3 = b^2 = Id$ and generate $\mathbf{PSL}(2, \mathbf{Z})$ which actually is isomorphic to the free product of $\mathbf{Z}/2\mathbf{Z}$ and $\mathbf{Z}/3\mathbf{Z}$:

$$\mathbf{PSL}(2, \mathbf{Z}) = \langle a, b; a^3, b^2 \rangle.$$

The generators s_1 and s_2 can be written using a and b :

$$s_1 = a^{-1}b \text{ and } s_2 = ba^{-1}.$$

In the particular case of the group $\mathbf{PSL}(2, \mathbf{Z})$ equipped with the two systems of generators $\{s_1, s_2\}$ and $\{a, b\}$, equation (1) reads

$$(2) \quad \frac{1}{3}l_{\{s_1, s_2\}} \leq l_{\{a, b\}} \leq 2l_{\{s_1, s_2\}}.$$

The following proposition is a classical result. For the sake of completeness, we give it below followed by a short proof.

Proposition 1. *The morphism m from the free group \mathbf{F}_2 to $\mathbf{PSL}(2, \mathbf{Z})$ defined by*

$$m(e_1) = s_1^2 = a^{-1}ba^{-1}b \text{ and } m(e_2) = s_2^2 = ba^{-1}ba^{-1},$$

is a quasi-isometric embedding.

Proof. Consider a word f in \mathbf{F}_2 , written in a reduced form (i.e. with the smallest possible length) and let f^n be the word in \mathbf{F}_2 with length $n < l_{\{e_1, e_2\}}(f)$ which coincides with f truncated after its n^{th} symbol. It is easy to check that

$$l_{\{a, b\}}(m(f^{n+1})) \geq l_{\{a, b\}}(m(f^n)) + 1.$$

It follows that

$$(3) \quad l_{\{a, b\}}(m(f)) \geq l_{\{e_1, e_2\}}(f) - 1 + l_{\{a, b\}}(m(f^1)) = l_{\{e_1, e_2\}}(f) + 3 \geq l_{\{e_1, e_2\}}(f).$$

On the other hand, for any element f in \mathbf{F}_2 we have

$$l_{\{a, b\}}(m(f)) \leq 4l_{\{s_1^2, s_2^2\}}(m(f)) = 4l_{\{e_1, e_2\}}(f).$$

This series of inequalities show that the morphism m is a quasi-isometric embedding. \square

This quasi-isometric embedding can be easily visualized on the Caley graph of the group $\mathbf{PSL}(2, \mathbf{Z})$ (see this graph in [10]).

Corollary 1. *The morphism \bar{m} from the free group \mathbf{F}_2 to \mathbf{B}_3 defined by*

$$\bar{m}(e_1) = \sigma_1^2 \text{ and } \bar{m}(e_2) = \sigma_2^2,$$

is a quasi-isometric embedding.

Proof. On one hand, we obviously have

$$l_{\{\sigma_1, \sigma_2\}}(\bar{m}(f)) \leq 2l_{\{e_1, e_2\}}(f).$$

On the other hand, we get from the relation

$$m = \rho \circ \bar{m}$$

that for any element f in \mathbf{F}_2 ,

$$l_{\{s_1, s_2\}}(m(f)) \leq l_{\{\sigma_1, \sigma_2\}}(\bar{m}(f)).$$

Combining Equations (2) and (3) we get

$$\frac{1}{2}l_{\{e_1, e_2\}}(f) \leq \frac{1}{2}l_{\{a, b\}}(m(f)) \leq l_{\{s_1, s_2\}}(m(f)).$$

Thus,

$$\frac{1}{2}l_{\{e_1, e_2\}}(f) \leq l_{\{\sigma_1, \sigma_2\}}(\bar{m}(f)).$$

These inequalities prove the corollary. \square

Remark 1. The embedding of the free group \mathbf{F}_2 in the braid group \mathbf{B}_3 maps \mathbf{F}_2 onto the subgroup of \mathbf{B}_3 which consists in all equivalence classes of systems of 3 arcs, the first one and the last one being vertical and the second one turning around the other two. This interpretation allows us to identify $\bar{m}(\mathbf{F}_2)$, hence \mathbf{F}_2 , with the fundamental group of the 2-punctured disk.

From Lemma 1 and Corollary 1 we get

Corollary 2. *Any finitely generated free group can be embedded in \mathbf{B}_3 by a quasi-isometric embedding.*

Consider now embeddings of the abelian free groups in the braid groups.

Proposition 2. *For any $n \geq 1$, the morphism \tilde{m}_n from the group \mathbf{Z}^n to the braid group \mathbf{B}_{2n} defined for $i = 1, \dots, n$ by*

$$\tilde{m}_n(e_i) = (\sigma_{2i-1})^2,$$

*is a quasi-isometric embedding.*¹

Proof. Since two generators σ_i and σ_j of the braid group \mathbf{B}_n commute when then are not consecutive ($|i - j| \neq 1$), the morphism \tilde{m}_n is well defined. Any element z in $\tilde{m}_n(\mathbf{Z}^n)$ can be written in a unique way as the product:

$$z = \sigma_1^{\alpha_1(z)} \sigma_3^{\alpha_3(z)} \sigma_5^{\alpha_5(z)} \dots \sigma_{2n-1}^{\alpha_{2n-1}(z)},$$

where the $\alpha_j(z)$'s are even integers.

For $i = 0, \dots, n-1$, the map $z \rightarrow \alpha_i(z)$ is a morphism. By identifying \mathbf{B}_2 with \mathbf{Z} , this morphism corresponds to the forgetful map: $\tau_i : \mathbf{B}_{2n} \rightarrow \mathbf{B}_2$ which consists in forgetting all the strands of a representative of an element in \mathbf{B}_n but the $(2i-1)^{th}$ and $(2i)^{th}$ ones. This yields the equality:

$$l_{\{\sigma_1, \dots, \sigma_{2n-1}\}}(z) = \sum_{i=0}^{i=n-1} |\alpha_i(z)|.$$

¹It is only for technical reasons that will be clear in the next section that we define $\tilde{m}_n(e_i) = (\sigma_{2i-1})^2$ and not $\tilde{m}_n(e_i) = (\sigma_{2i-1})$.

Thus for any f in \mathbf{Z}^n ,

$$l_{\{\sigma_1, \dots, \sigma_{2n-1}\}}(\tilde{m}_n(f)) = 2l_{\{e_1, \dots, e_n\}}(z).$$

□

4. AREA PRESERVING DIFFEOMORPHISMS

4.1. A lower bound for the distance between two maps. We recall in this paragraph some results that are developed in [9].

Let us fix again an element ϕ in \mathcal{D}_2 and an isotopy $\{\phi_t\}_{t \in [0,1]}$ in \mathcal{D}_2 connecting identity to ϕ . To any pair of distinct points x, y in \mathbf{D}^2 and to every $t \in [0, 1]$, we associate the unit vector

$$u(t, x, y) = \frac{\phi_t(y) - \phi_t(x)}{\|\phi_t(y) - \phi_t(x)\|}.$$

Lemma 2 ([9]). *The integral*

$$\mathcal{G}(\{\phi_t\}) = \frac{1}{2\pi} \int_0^1 \int_{\mathbf{D}^2 \times \mathbf{D}^2} \left\| \frac{du}{dt}(t, x, y) \right\| dx dy dt$$

is well defined.

Proof. An easy calculation yields

$$\left\| \frac{du}{dt}(t, x, y) \right\| = \frac{\|(\dot{\phi}_t(y) - \dot{\phi}_t(x)) \wedge (\phi_t(y) - \phi_t(x))\|}{\|\phi_t(y) - \phi_t(x)\|^2},$$

where \wedge is the wedge product and $\dot{\phi}_t = \frac{\partial \phi_t}{\partial t}$. It follows that

$$\left\| \frac{du}{dt}(t, x, y) \right\| \leq \frac{2}{\|\phi_t(y) - \phi_t(x)\|} \sup_{(t,x) \in [0,1] \times \mathbf{D}^2} \|\dot{\phi}_t(x)\|.$$

The quantity $\sup_{(t,x) \in [0,1] \times \mathbf{D}^2} \|\dot{\phi}_t(x)\|$ is bounded. Hence it is enough to show that since ϕ_t is area preserving for every t in $[0, 1]$,

$$\int_{\mathbf{D}^2 \times \mathbf{D}^2} \frac{dx dy}{\|\phi_t(y) - \phi_t(x)\|} = \int_{\mathbf{D}^2 \times \mathbf{D}^2} \frac{dx dy}{\|y - x\|}$$

and that this last integral converges. □

We now introduce the quantity

$$\mathcal{G}(\phi) = \inf \mathcal{G}(\{\phi_t\}),$$

where the infimum is taken over all isotopies in \mathcal{D}_2 joining identity to ϕ .

Lemma 3 ([9]). *There exists a constant $K > 0$ such that for any map ϕ in \mathcal{D}_2*

$$\mathcal{G}(\phi) \leq K l_2(\phi).$$

Proof. The quantity $\mathcal{G}(\{\phi_t\})$ satisfies

$$\mathcal{G}(\{\phi_t\}) \leq \frac{1}{\pi} \int_0^1 \int_{\mathbf{D}^2 \times \mathbf{D}^2} \frac{\|\dot{\phi}_t(x)\|}{\|\phi_t(y) - \phi_t(x)\|} dx dy dt.$$

The Cauchy-Schwarz inequality gives

$$\mathcal{G}(\{\phi_t\}) \leq \frac{1}{\pi} \int_0^1 \|\dot{\phi}_t\|_2 \|I_t\|_2 dt,$$

where

$$I_t(x) = \int_{\mathbf{D}^2} \frac{dy}{\|\phi_t(y) - \phi_t(x)\|}.$$

For every t in $[0, 1]$ the map ϕ_t is area preserving; consequently, the L_2 -norm of I_t satisfies

$$\|I_t\|_2 = \left(\int_{\mathbf{D}^2} \left(\int_{\mathbf{D}^2} \frac{dy}{\|x - y\|} \right)^2 dx \right)^{1/2} < +\infty.$$

Thus, for $K = \frac{1}{\pi} \|I_t\|_2$ we get

$$\mathcal{G}(\{\phi_t\}) \leq Kl_2(\{\phi_t\}).$$

It follows that

$$\mathcal{G}(\phi) \leq Kl_2(\phi).$$

□

The quantity $\mathcal{G}(\phi)$ can be interpreted as follows. For an isotopy $\{\phi_t\}_{t \in [0,1]}$ connecting identity to ϕ and a pair of distinct points x, y in \mathbf{D}^2 , we consider the map

$$\begin{aligned} u_{x,y} : [0, 1] &\rightarrow \mathbf{S}^1 \\ t &\mapsto u(t, x, y). \end{aligned}$$

The change of variables induced by the map $u_{x,y}$ leads to the equality

$$\int_0^1 \left\| \frac{du_{x,y}}{dt}(t) \right\| dt = \int_{\mathbf{S}^1} \# \{u_{x,y}^{-1}(\omega)\} d\omega,$$

where $\#$ stands for cardinality. By integrating we get

$$\mathcal{G}(\{\phi_t\}) = \frac{1}{2\pi} \int_{\mathbf{D}^2 \times \mathbf{D}^2} \int_{\mathbf{S}^1} \# \{u_{x,y}^{-1}(\omega)\} d\omega dx dy.$$

Consider the 2 arcs $\gamma_x : t \mapsto (\phi_t(x), t)$ and $\gamma_y : t \mapsto (\phi_t(y), t)$ in the cylinder $\mathbf{D}^2 \times [0, 1]$ and choose a direction ω in \mathbf{S}^1 tangent to \mathbf{D}^2 . When projected in the direction of ω onto a plane orthogonal to ω , γ_y overcrosses $\# \{u_{x,y}^{-1}(\omega)\}$ times γ_x . The integral

$$Cr_{\{\phi_t\}}(x, y) = \frac{1}{2\pi} \int_{\mathbf{S}^1} \# \{u_{x,y}^{-1}(\omega)\} d\omega$$

is the averaged number of times the arc γ_y overcrosses γ_x over all directions ω tangent to \mathbf{D}^2 . $\mathcal{G}(\{\phi_t\})$ is then the spatial average $\int_{\mathbf{D}^2 \times \mathbf{D}^2} Cr_{\{\phi_t\}}(x, y) dx dy$.

Changing the isotopy $\{\phi_t\}$ may reduce the averaged quantity of overcrossings $\mathcal{G}(\{\phi_t\})$; the infimum is given by $\mathcal{G}(\phi)$.

4.2. A subgroup of \mathcal{D}_2 . For r small enough, consider, for $i = 0, \dots, n-1$, the n disks D_i in \mathbf{D}^2 , with radius r and centered at $P_i = (i/(n-1) - 1/2, 0)$. We define the subgroup \mathcal{B}_n which consists in these maps whose restriction to the disks D_i is identity for $i = 0, \dots, n-1$. There is a natural morphism μ_n from the group \mathcal{B}_n to the braid group \mathbf{B}_n which is defined as follows:

- choose a map ϕ in \mathcal{B}_n and an isotopy $\{\phi_t\}_{t \in [0,1]}$ from identity to ϕ ;

- a representative of the braid $\mu_n(\phi)$ consists in the system of arcs in $\mathbf{D}^2 \times [0, 1]$:

$$\bigcup_{i=0}^{i=n-1} \{(\phi_t(P_i), t), t \in [0, 1]\}.$$

Lemma 4. *Let ϕ be a map in \mathcal{B}_n . There exists a quantity $C(r) \geq 0$ which depends only on r and goes to zero with r such that*

$$l_2(\phi) \geq \frac{1}{K} l_{\{\sigma_1, \dots, \sigma_{n-1}\}}(\mu_n(\phi))(1 - C(r)) \times (\text{area } D_0)^2.$$

Proof. Let $\{\phi_t\}_{t \in [0, 1]}$ in \mathcal{D}_2 be an isotopy connecting identity to ϕ . The integral $\mathcal{G}(\{\phi_t\})$ reads

$$\mathcal{G}(\{\phi_t\}) = \int_{\mathbf{D}^2 \times \mathbf{D}^2} Cr_{\{\phi_t\}}(x, y) dx dy,$$

and consequently

$$\mathcal{G}(\{\phi_t\}) \geq \sum_{i \neq j} \int_{D_i \times D_j} Cr_{\{\phi_t\}}(x, y) dx dy = \int_{D_0 \times D_0} \left(\sum_{i \neq j} Cr_{\{\phi_t\}}(T_i(x), T_j(y)) \right) dx dy,$$

where T_i stands for the translation which maps D_0 onto D_i .

For $(x, y, \omega) \in D_0 \times D_0 \times \mathbf{S}^1$, let $\mathcal{U}_\phi(x, y, \omega)$ be the minimum over all isotopies in \mathcal{D}_2 connecting identity to ϕ of the quantity

$$\sum_{i \neq j} \#\{u_{T_i(x), T_j(y)}^{-1}(\omega)\},$$

We have

$$\mathcal{G}(\phi) \geq \frac{1}{2\pi} \int_{\mathcal{S}^1} \int_{D_0 \times D_0} \mathcal{U}_\phi(x, y, \omega) d\omega dx dy.$$

Let $M(r)$ be the set of ω 's for which the orthogonal projections (in the direction ω) of the disks D_i overlap. We denote by $2\pi C(r)$ the measure of the set $M(r)$. One can easily check that $C(r)$ goes to zero with r . It is also clear that for ω in $\mathcal{S}^1 \setminus M(r)$,

$$\mathcal{U}_\phi(x, y, \omega) \geq l_{\{\sigma_1, \dots, \sigma_{n-1}\}}(\mu_n(\phi)).$$

By integrating over $\mathcal{S}^1 \setminus M(r)$ we get

$$\mathcal{G}(\phi) \geq l_{\{\sigma_1, \dots, \sigma_{n-1}\}}(\mu_n(\phi))(1 - C(r)) \times (\text{area } D_0)^2.$$

□

5. PROOF OF THEOREM 1

Consider in \mathcal{B}_3 two maps Θ_1 and Θ_2 such that

$$\mu_3(\Theta_1) = \sigma_1^2 \quad \text{and} \quad \mu_3(\Theta_2) = \sigma_2^2.$$

The morphism $\pi : \mathbf{F}_2 \rightarrow \mathcal{D}_2$ defined by

$$\pi(e_1) = \Theta_1 \quad \text{and} \quad \pi(e_2) = \Theta_2,$$

is injective since $\mu_3 \circ \pi = \bar{m}$ is injective. For any f in \mathbf{F}_2 , it also satisfies

$$\begin{aligned} l_2(\pi(f)) &\geq \frac{1}{K} l_{\{\sigma_1, \sigma_2\}} \mu_3(\pi(f))(1 - C(r)) \times (\text{area } D_0)^2 \\ &\geq \frac{1}{2K} l_{\{e_1, e_2\}}(f)(1 - C(r)) \times (\text{area } D_0)^2; \end{aligned}$$

the left inequality coming from Lemma 4, the right one from the second inequality in the proof of Corollary 1.

On the other hand, for

$$f = e_{\epsilon_1}^{\alpha_1} e_{\epsilon_2}^{\alpha_2} \dots e_{\epsilon_n}^{\alpha_n},$$

where $\epsilon_i = 1$ or 2 , we have

$$l_2(\pi(f)) \leq \sum_{i=1}^n |\alpha_i| l_2(\pi(e_{\epsilon_i})) \leq l_{e_1, e_2}(f) \times \max_{i \in \{1, 2\}} l_2(\pi(e_i)).$$

Using Lemma 1, we also get a quasi-isometric embedding of \mathbf{F}_n in \mathcal{D}_2 for any $n \geq 2$. This proves Theorem 1 for finitely generated free groups.

Consider now n maps $\Omega_1, \dots, \Omega_n$ in \mathcal{B}_{2n} such that

- $\mu_{2n}(\Omega_i) = \sigma_{2i-1}^2$ for $i = 1, \dots, n$;
- the maps Ω_i are different from identity on disjoint supports, thus they commute.

Since the maps Ω_i commute, the morphism $\tilde{\pi} : \mathbf{Z}^n \rightarrow \mathcal{D}_2$ defined for $i = 1, \dots, n$, by

$$\tilde{\pi}(e_i) = \Omega_i$$

is well defined. It is injective since $\mu_{2n} \circ \tilde{\pi} = \tilde{m}_n$ is injective. For any f in \mathbf{Z}^n , it also satisfies

$$\begin{aligned} l_2(\tilde{\pi}(f)) &\geq \frac{1}{K} l_{\{\sigma_1, \dots, \sigma_{2n-1}\}} \mu_{2n}(\tilde{\pi}(f)) (1 - C(r)) \times (\text{area } D_0)^2 \\ &= \frac{2}{K} l_{\{e_1, \dots, e_n\}}(f) (1 - C(r)) \times (\text{area } D_0)^2; \end{aligned}$$

the left inequality coming from Lemma 4, the right one from the last inequality in the proof of Proposition 2.

On the other hand, given

$$f = e_1^{\alpha_1} e_2^{\alpha_2} \dots e_n^{\alpha_n},$$

we have

$$l_2(\tilde{\pi}(f)) \leq \sum_{i=1}^n |\alpha_i| l_2(\tilde{\pi}(e_i)) \leq l_{e_1, \dots, e_n}(f) \times \max_{i \in \{1, \dots, n\}} l_2(\tilde{\pi}(e_i)).$$

Thus for any $n \geq 1$, we get a quasi-isometric embedding of \mathbf{Z}^n in \mathcal{D}_2 . This achieves the proof of Theorem 1.

6. FINAL REMARKS

6.1. Random walks on \mathcal{D}_2 . Let G be a group equipped with a right invariant metric d . Consider the Borel σ -field of G , $\mathcal{B}(G)$, and let $\mu : \mathcal{B}(G) \mapsto [0, 1]$ be a Borel probability measure on G . We can associate to μ the random walk on \mathcal{D}_2 $X = \{X^n\}_{n \in \mathbf{Z}^+}$ defined by

$$X^0 = Id,$$

$$X^n = g_n \cdot g_{n-1} \dots g_1 \quad \text{for } n \geq 1,$$

where $\{g_i\}_{i \in \mathbf{Z}^+}$ is a sequence of independent random variables defined on some probability space (Ω, \mathcal{F}, P) and identically distributed according to μ .

For any random variable $X : \Omega \mapsto \mathbf{R}$ and $p \geq 1$ we let $\|X\|_p = (\int_{\Omega} |X|^p dP)^{1/p}$. $L^p(P)$ is the set of random variables X for which $\|X\|_p < \infty$. For $X \in L^1(P)$ we let $E(X) = \int_{\Omega} X dP$ denote the mathematical expectation of X .

Proposition 6.1. *Suppose $E(d(X^1, Id)) = \int_{\mathbf{R}} d(g, Id) \mu(dg) < \infty$, and let*

$$\gamma_1 = \inf_{n \in \mathbf{Z}^+} \frac{1}{n} E(d(X^n, Id)).$$

Then

$$\lim_{n \rightarrow \infty} \frac{l_2(d(X^n, Id))}{n} = \gamma_1$$

P -almost surely.

Proof. The proof is standard. We can always suppose that (Ω, \mathcal{F}, P) is the canonical space where $\Omega = G^{\mathbf{Z}^+}$, $\mathcal{F} = \mathcal{B}(G^{\mathbf{Z}^+})$ and $P = \mu^{\otimes \mathbf{Z}^+}$ is the product measure. Let $\Theta : \Omega \mapsto \Omega$ denote the shift defined by $\Theta(\mathbf{g})_i = g_{i+1}$ where $\mathbf{g} = \{g_i\}_{i \in \mathbf{Z}^+}$ and let $\tau_n(\mathbf{g}) = d(g_n \cdot g_{n-1} \dots g_1, Id)$. The right invariance of d implies that

$$\begin{aligned} \tau_{n+m}(\mathbf{g}) &= d(g_{n+m} \dots g_1, Id) \\ &= d(g_{n+m} \dots g_{n+1}, (g_n \dots g_1)^{-1}) \\ &\leq \tau_n(\mathbf{g}) + \tau_m(\Theta^n(\mathbf{g})). \end{aligned}$$

Hence Proposition 6.1 follows from Kingman's subadditive Ergodic Theorem [11]. \square

The quantity γ_1 is called *the linear escape rate* of the random walk.

Examples. • Consider the free group \mathbf{F}_d with generators $\mathcal{S} = \{e_1, \dots, e_d\}$ and equipped with the right invariant metric induced by the word length. Charge the generators and their inverse with an equidistributed measure ($\mu(e_i) = \mu(e_i^{-1}) = \frac{1}{d}$) and consider the random walk on this group which consists in starting from an element chosen at random in $\mathcal{S} \cup \mathcal{S}^{-1}$ and multiplying at each step on the left by an element chosen at random in $\mathcal{S} \cup \mathcal{S}^{-1}$. It is easy to check that the linear rate of escape in this particular case satisfies

$$\gamma_1 = 1 - \frac{1}{d}.$$

- For a similar random walk on the abelian free group \mathbf{Z}^d it is also known that the linear escape rate vanishes ($\gamma_1 = 0$).
- From Theorem 1 we get that for any positive integer d , there exist d maps $\Theta_1, \dots, \Theta_d$ in \mathcal{D}_2 such that the group generated by $\Theta_1, \dots, \Theta_d$ is quasi-isometric to the free group. It implies that the associated random walk on this group has a strictly positive linear escape rate.
- From Theorem 1 we also get that for any positive integer d , there exist d maps $\Omega_1, \dots, \Omega_d$ in \mathcal{D}_2 such that the group generated by $\Omega_1, \dots, \Omega_d$ is quasi-isometric to the abelian free group \mathbf{Z}^d . It implies that the associated random walk on this group has a linear escape rate which vanishes.

Question 1. For any $d > 1$ and for any generic choice of maps ϕ_1, \dots, ϕ_d in \mathcal{D}_2 , consider the subgroup $\mathcal{D}_2(\phi_1, \dots, \phi_d)$ of \mathcal{D}_2 generated by $\{\phi_1, \dots, \phi_d\}$. Is this subgroup a quasi-isometric embedding of the free group \mathbf{F}_d ? More generally, does the random walk which gives an equal probability to each generator and their inverse, have a non-zero linear rate of escape?

6.2. Embeddings in the kernel of the Calabi morphism. Consider a map ϕ in \mathcal{D}_2 and a 1-form α on \mathbf{D}^2 which is a primitive of the area 2-form. Since ϕ is area preserving, the form $\phi^*\alpha - \alpha$ is closed and vanishes near the boundary of \mathbf{D}^2 . Thus, there exists a unique function $H(\phi, \alpha) : \mathbf{D}^2 \rightarrow \mathbf{R}$, which vanishes near the boundary of \mathbf{D}^2 and such that

$$dH(\phi, \alpha) = \phi^*\alpha - \alpha.$$

The Calabi invariant [4] of ϕ is defined by

$$\mathcal{C}(\phi) = \int_{\mathbf{D}^2} H(\phi, \alpha).$$

It is easy to verify that this integral does not depend on the choice of the primitive α and that the Calabi invariant is actually a morphism from \mathcal{D}_2 to \mathbf{R} :

$$\mathcal{C}(\phi_0 \circ \phi_1) = \mathcal{C}(\phi_0) + \mathcal{C}(\phi_1).$$

A. Banyaga [2] proved that the kernel of this morphism, $\ker \mathcal{C}$, is a simple group and Y. Eliashberg and T. Ratiu [6] showed that its subgroup has an infinite diameter (see also [9]).

There is a second definition of the Calabi invariant which is due to A. Fathi [7] (see also [8]) and which can be seen as an estimate in terms of braiding of pairs of orbits. Consider an isotopy $\{\phi_t\}_{t \in [0,1]}$ in \mathcal{D}_2 connecting identity to ϕ . The map

$$Ang_\phi : \mathbf{D}^2 \times \mathbf{D}^2 \setminus \Delta \rightarrow \mathbf{R},$$

(where Δ stands for the diagonal) which associates to any pair of points $x \neq y$ in \mathbf{D}^2 the angular variation of the vector $\overrightarrow{\phi_t(x)\phi_t(y)}$ when t goes from 0 to 1, does not depend on the choice of the isotopy and is bounded where it is defined. The Calabi invariant is the integral of this function, i.e.:

$$\mathcal{C}(\phi) = \int_{\mathbf{D}^2 \times \mathbf{D}^2} Ang_\phi(x, y) dx dy.$$

With this second definition, $\text{Ker } \mathcal{C}$ is the subgroup of area preserving maps whose angular variation of a pair of points vanishes in average. Using this second geometric interpretation, Theorem 1 can be made more precise:

Theorem 2. *Any finitely generated free group and any finitely generated abelian free group is quasi-isometrically embedded in $\text{Ker } \mathcal{C}$.*

We only give here the main lines of the proof and insist on its differences with the proof of Theorem 1.

Consider the morphism $m' : \mathbf{F}_2 \rightarrow \mathbf{B}_5$ which maps the generator e_1 on $\sigma_1^2 \sigma_3^{-2}$ and e_2 on $\sigma_2^2 \sigma_4^{-2}$. Using the same technics as the ones we used in the proof of Theorem 1 we get that m' is a quasi-isometric embedding. It is possible to find two maps in \mathcal{B}_5 , Θ'_1 and Θ'_2 such that

- $\mu_6(\Theta'_1) = \sigma_1^2 \sigma_3^{-2}$ and $\mu_6(\Theta'_2) = \sigma_2^2 \sigma_4^{-2}$;
- Θ'_1 and Θ'_2 are in $\text{Ker } \mathcal{C}$.

Similarly to the proof of Theorem 1, we prove that the morphism $\pi' : \mathbf{F}_2 \rightarrow \text{Ker } \mathcal{C}$ defined by $\pi'(e_1) = \Theta'_1$ and $\pi'(e_2) = \Theta'_2$ is a quasi-isometric embedding. Using Corollary 1, we get quasi-isometric embeddings of any finitely generated free group in $\text{Ker } \mathcal{C}$. A similar construction works for finitely generated abelian free groups.

6.3. Symplectomorphisms in higher dimensions. Let $n > 0$ and consider the group \mathcal{D}_{2n} of diffeomorphisms of the unit ball in the $2n$ -dimensional space B^{2n} which preserve the standard symplectic 2-form $\omega = \sum_{i=1}^n dx_i \wedge dy_i$, and are identity near the boundary ∂B^{2n} . This group, like \mathcal{D}_2 , can be equipped with a right invariant metric. With this metric, \mathcal{D}_{2n} also has an infinite diameter [6].

Question 2. Does a quasi-isometric embedding of any finitely generated free group and any finitely generated abelian free group in \mathcal{D}_{2n} exist for all $n > 0$?

ACKNOWLEDGMENTS

It is a pleasure to thank J. González-Meneses and L. Paris for helpful discussions on braid groups.

REFERENCES

- [1] ARNOLD, V. AND KHEVIN, B. *Topological methods in hydrodynamics*. Applied Mathematical Sciences, **125**, (1998). MR **99b**:58002
- [2] BANYAGA, A.: On the group of diffeomorphisms preserving an exact symplectic form, in *Differential topology*, Varenna (1976), 5-9. MR **83g**:58006
- [3] BIRMAN, J. *Braids, Links and Mapping Class Groups*. Annals of Math. Studies **82**, Princeton University Press, (1974) Erratum, 1975. MR **51**:11477; MR **54**:13894
- [4] CALABI, E.: On the group of automorphisms of a symplectic manifold, in *Problems in analysis*, Symposium in honor of S. Bochner, R. C. Gunning, Ed., Princeton Univ. Press, Princeton (1970), 1-26. MR **50**:3268
- [5] EBIN, D. G. AND MARSDEN, J.: Groups of diffeomorphisms and the notion of an incompressible fluid, *Ann. of Math.* **92**, (1970), 102-163.
- [6] ELIASBERG, Y. AND RATIU, T.: The diameter of the symplectomorphism group is infinite, *Invent. Math.* **103**, (1991), 327-340. MR **92a**:58018
- [7] FATHI, A.: Transformations et homéomorphismes préservant la mesure. Systèmes dynamiques minimaux., Thèse Orsay (1980).
- [8] GAMBAUDO, J.-M. AND GHYS, É.: Enlacements asymptotiques, *Topology* **36**, (1997), 1355-1379. MR **98f**:57050
- [9] GAMBAUDO, J.-M. AND LAGRANGE, M.: Topological lower bounds on the distance between area preserving diffeomorphisms, *Bol. Soc. Brasil. Mat.* **31**, (2000), 1-19. CMP 2000:12
- [10] GHYS, É. AND DE LA HARPE, P. *Sur les groupes hyperboliques d'après Mikhael Gromov*, Ghys, É and de la Harpe, eds., Progress in Mathematics **83**, Birkhauser, (1990). MR **92f**:53050
- [11] KINGMAN, J. F. C.: The ergodic theory of subadditive stochastic processes, *J. Royal Stat. Soc.*, **30**, (1968), 499-510. MR **40**:8114
- [12] MURASUGI, K. *Knot theory and its applications*. Translated from the 1993 Japanese original by Bohdan Kurpita. Birkhäuser Boston, Inc., Boston, (1996). MR **97g**:57011
- [13] SHNIRELMAN, A.: The geometry of the group of diffeomorphisms and the dynamics of an ideal incompressible fluid, *Matem. Sbornik* **128**, (1985), 82-109; English transl: Math. USSR, Sbornik **56** (1987), 79-105. MR **87d**:58034

UNIVERSITÉ DE CERGY PONTOISE, LABORATOIRE DE MATHÉMATIQUES, 2, AVENUE ADOLPHE CHAUVIN, 95302 CERGY-PONTOISE CEDEX, FRANCE

E-mail address: benaim@math.u-cergy.fr

UNIVERSITÉ DE BOURGOGNE, LABORATOIRE DE TOPOLOGIE, UMR CNRS 5584, B.P. 47870-21078-DIJON CEDEX, FRANCE