

SCATTERING POLES FOR ASYMPTOTICALLY HYPERBOLIC MANIFOLDS

DAVID BORTHWICK AND PETER PERRY

ABSTRACT. For a class of manifolds X that includes quotients of real hyperbolic $(n + 1)$ -dimensional space by a convex co-compact discrete group, we show that the resonances of the meromorphically continued resolvent kernel for the Laplacian on X coincide, with multiplicities, with the poles of the meromorphically continued scattering operator for X . In order to carry out the proof, we use Shmuel Agmon's perturbation theory of resonances to show that both resolvent resonances and scattering poles are simple for generic potential perturbations.

1. INTRODUCTION

The purpose of this paper is to show the equivalence of two possible notions of ‘scattering resonances’ for the Laplacian on asymptotically hyperbolic manifolds, i.e., complete Riemannian manifolds of infinite volume with ‘constant curvature at infinity’. On the one hand, it is very natural to define scattering resonances with respect to the meromorphically continued resolvent of the Laplace operator. This point of view has been very fruitful and has led to a large body of results on the distribution of scattering resonances in the complex plane; see [29, 31] for surveys and [5, 6, 7, 30] for results on the class of manifolds studied here. On the other hand, it is also reasonable, by analogy with scattering theory for the Schrödinger or wave equations in Euclidean space (see e.g. [14, 26]), to define scattering resonances as poles of the scattering operator for the Laplacian. For Schrödinger scattering on Euclidean space, the equivalence of scattering resonances and resolvent resonances is well known [8, 9, 10]. The analogous result for hyperbolic manifolds is of interest since the poles of the scattering operator have a geometric and dynamical interpretation: they are among the poles of Selberg's zeta function for geodesic flow on the manifold [22, 23, 25]. Thus the scattering resonances serve, in a sense, as discrete data similar in character to the eigenvalues of a compact surface. The methods used to prove equivalence of resonances in the Euclidean setting do not appear to work in the hyperbolic case.

For noncompact Riemann surfaces and certain metric perturbations, Guillopé and Zworski showed that the set of resolvent resonances and the set of scattering poles coincide with multiplicities ([7], Proposition 2.11); here we will show that,

Received by the editors March 19, 1999 and, in revised form, June 28, 2001.

2000 *Mathematics Subject Classification*. Primary 58J50, 35P25; Secondary 47A40.

Key words and phrases. Scattering resonances, hyperbolic manifolds.

Supported in part by NSF grant DMS-9796195 and by an NSF Postdoctoral Fellowship.

Supported in part by NSF grant DMS-9707051.

with some restrictions, the same equivalence holds for asymptotically hyperbolic manifolds in higher dimension.

To describe our results, we first recall that an asymptotically hyperbolic manifold is a compact manifold \overline{X} with boundary, endowed with a Riemannian metric of a special form. A *defining function* for the boundary of a compact manifold \overline{X} is a nonnegative C^∞ function on \overline{X} with $x^{-1}\{0\} = \partial\overline{X}$ and $dx|_{\partial\overline{X}}$ nowhere vanishing. The metric g then takes the form $x^{-2}h$, where x is a defining function and h is a nondegenerate smooth metric on \overline{X} such that $|dx|_h \rightarrow 1$ as $x \rightarrow 0$. Note that this metric puts $\partial\overline{X}$ ‘at infinity’ and makes X , the interior of \overline{X} , a complete Riemannian manifold of infinite volume. The condition on $|dx|_h$ insures that the sectional curvatures approach -1 at metric infinity. If Δ_g denotes the positive Laplace-Beltrami operator on (X, g) and X has dimension $n+1$, it is known that the spectrum of Δ_g consists of at most finitely many L^2 eigenvalues of finite multiplicity in the interval $[0, n^2/4)$, and absolutely continuous spectrum in $[n^2/4, \infty)$ with no embedded eigenvalues (see [3, 15, 16] for quotients of hyperbolic space and [17, 19] for asymptotically hyperbolic manifolds). Thus the resolvent $\mathcal{R}_g(z) = (\Delta_g - z)^{-1}$ is a meromorphic operator-valued function on the cut plane $\mathbb{C} \setminus [n^2/4, \infty)$ with poles at the L^2 eigenvalues having finite-rank residues. It is convenient to introduce a uniformizing parameter ζ and set $R_g(\zeta) = (\Delta_g - \zeta(n - \zeta))^{-1}$, which is then a meromorphic operator-valued function on the half-plane $\Re(\zeta) > n/2$. The operator $R_g(\zeta)$ has first-order poles at points ζ_0 whenever $\zeta_0(n - \zeta_0)$ is an $L^2(X)$ -eigenvalue of Δ_g . We denote by Z_p the (finite and possibly empty) set of all such ζ_0 . The *multiplicity* of $\zeta_0 \in Z_p$ is the dimension, m_{ζ_0} , of the eigenspace of Δ_g with eigenvalue $\zeta_0(n - \zeta_0)$. Equivalently,

$$(1.1) \quad m_{\zeta_0} = \text{rank} \left(\int_{\gamma_{\zeta_0}} R_g(\zeta) d\zeta \right),$$

where γ_{ζ_0} is a simple closed contour surrounding ζ_0 and no other pole of $R_g(\zeta)$.

First, we define the resolvent resonance set of Δ_g . Let $\dot{C}^\infty(X)$ denote the smooth functions on \overline{X} vanishing to all orders at $\partial\overline{X}$. Viewed as a map from $\dot{C}^\infty(X)$ to $C^\infty(X)$, the resolvent operator $R_g(\zeta)$ admits a meromorphic continuation to $\mathbb{C} \setminus \frac{1}{2}(n - \mathbb{N})$, as was shown in [20]. The set \mathcal{R} of *resolvent resonances* consists of the poles of this meromorphic continuation in the half-plane $\Re(\zeta) < n/2$, excluding the set $\frac{1}{2}(n - \mathbb{N})$. Lower bounds on resolvent resonances proven in [27] show that this set is always nontrivial for constant curvature spaces, and explicit examples (see, for instance, section 3 of [5]) show that resolvent resonances can form an infinite lattice in the half-plane $\Re(\zeta) < n/2$. If ζ_0 is a resolvent resonance with $\Re(\zeta_0) < n/2$, the *multiplicity* of ζ_0 is the number m_{ζ_0} defined by (1.1), where again γ_{ζ_0} is a simple closed curve that encloses ζ_0 and no other pole of $R_g(\zeta)$. The point ζ_0 is a *semi-simple resonance* if $R_g(\zeta)$ has a first-order pole at ζ_0 . The point ζ_0 is a *simple resonance* if, in addition, the residue of the pole has rank one.

Next, we define the scattering operator and the scattering resonance set of Δ_g . For $\zeta \in \mathbb{C}$ with $\Re(\zeta) = n/2$ and $\zeta \neq n/2$, and each $f_- \in C^\infty(\partial\overline{X})$, there is a unique smooth solution of the eigenvalue equation $(\Delta_g - \zeta(n - \zeta))u = 0$ having the asymptotic form

$$u = x^\zeta f_+ + x^{n-\zeta} f_- + O(x^{n/2+1}),$$

where $f_+ \in C^\infty(\partial\overline{X})$. (This is the definition from [21]; for a proof see [11].) It follows that f_+ is uniquely determined and that there is a linear map $S(\zeta) : C^\infty(\partial\overline{X}) \rightarrow C^\infty(\partial\overline{X})$ with $S(\zeta)f_- = f_+$; moreover it is clear that $S(\zeta)S(n-\zeta) = I$. It can be shown that $S(\zeta)$ extends to a meromorphic family of operators on \mathbb{C} (see [11]); these operators will have infinite-rank poles at $\zeta \in \frac{1}{2}n + \mathbb{N}$, and infinite-rank zeros at $\zeta \in \frac{1}{2}n - \mathbb{N}$. The set \mathcal{S} of *scattering poles* consists of the poles of the meromorphic continuation of $S(\zeta)$ in the half-plane $\Re(\zeta) < n/2$, excluding the set $\frac{1}{2}(n - \mathbb{N})$. For $\zeta_0 \notin \frac{1}{2}(n - \mathbb{N})$, the *multiplicity* of a scattering pole ζ_0 is the integer

$$(1.2) \quad \nu_{\zeta_0} = \frac{1}{2\pi i} \operatorname{Tr} \left(\int_{\gamma_{\zeta_0, \varepsilon}} S(\zeta)^{-1} S'(\zeta) d\zeta \right)$$

(compare [7, 23] where similar definitions are made). Results of [4] show that ν_{ζ_0} is an integer equal to the ‘null multiplicity’ of $S(\zeta_0)$ minus the ‘null multiplicity’ of $S^{-1}(\zeta_0)$, where null multiplicity must be defined with some care since neither operator may exist at ζ_0 (see Section 3). We will say that ζ_0 is semi-simple if the pole of $S(\zeta)$ at ζ_0 is of first order, and simple if the residue is rank-one.

We would like to show a correspondence, with multiplicities, between the sets \mathcal{R} and \mathcal{S} . A direct method (see for example [7], where the case $n = 1$ is treated) would compare the Laurent expansion of the meromorphically continued resolvent at $\zeta_0 \in \mathcal{R}$ to the Laurent expansion of the scattering operator at ζ_0 , using the fact that the Schwarz kernel of the scattering operator can be recovered from that of the resolvent kernel. This direct method works easily when the resolvent resonance is simple but is somewhat complicated for nonsimple resonances. For this reason, we will perturb the operator Δ_g with a potential $V \in \dot{C}^\infty(X)$ which, as we will show, can be chosen so that the meromorphically continued resolvent,

$$R_V(\zeta) = (\Delta_g + V - \zeta(n - \zeta))^{-1},$$

has only simple resonances. The perturbation will split each resonance of multiplicity m into m resonances of multiplicity one localized near the unperturbed resonance, and similarly each eigenvalue of multiplicity m into m eigenvalues of multiplicity one. This result, which is of some independent interest, will allow us to count multiplicities properly but avoid technicalities associated with nonsimple resonances.

Our analysis of generic potential perturbations is inspired by Klopp and Zworski’s analysis of resonances in potential scattering [13]. To carry out the analysis, we will use Shmuel Agmon’s perturbation theory of resonances [1] in which the resonances are realized as eigenvalues of a non-self-adjoint operator on a cleverly constructed Banach space; this replaces the complex scaling used in [13]. Standard Kato-Rellich perturbation theory [12] can then be used to study how the resonances move under perturbation.

Our first result is:

Theorem 1.1. *Let (X, g) be an asymptotically hyperbolic manifold, and let \mathcal{R} and \mathcal{S} be respectively the resolvent resonance set and scattering resonance set for the Laplacian Δ_g (with the points $\frac{1}{2}(n - \mathbb{N})$ excluded). Then $\mathcal{R} = \mathcal{S}$, and the relationship*

$$\nu_{\zeta_0} = m_{n-\zeta_0} - m_{\zeta_0}$$

holds at any $\zeta_0 \in \mathcal{R}$.

Note that $m_{n-\zeta_0}$ is nonzero only for the finitely many ζ_0 with $n - \zeta_0 \in Z_p$; for all other $\zeta_0 \notin \frac{1}{2}(n - \mathbb{N})$ the scattering resonances and resolvent resonances coincide with multiplicities. If $n - \zeta_0 \in Z_p$, then we could have $\nu_{\zeta_0} = 0$ even though $\zeta_0 \in \mathcal{S}$.

We can make a stronger statement if (X, g) has even dimension constant and curvature ‘near’ infinity, i.e., if there is a compact subset K of X so that g has constant negative curvature -1 on $X \setminus K$. In [6] it was shown that in this case $R_g(\zeta)$ is meromorphic with finite rank poles on all of \mathbb{C} . This class includes the convex co-compact hyperbolic manifolds. Recall that a geometrically finite group Γ of isometries of real hyperbolic $(n+1)$ -dimensional space \mathbb{H}^{n+1} is called convex co-compact if the orbit space $\Gamma \backslash \mathbb{H}^{n+1}$ has infinite volume and Γ contains no parabolic elements. If Γ is torsion-free (which we can insure by passing to a subgroup of finite index), the orbit space $X = \Gamma \backslash \mathbb{H}^{n+1}$ is a complete Riemannian manifold when given the induced hyperbolic metric g .

The full meromorphic continuation of the resolvent allows us to drop the restriction $\zeta \notin \frac{1}{2}(n - \mathbb{N})$ and define an enlarged resonance set $\tilde{\mathcal{R}}$ that includes all resolvent poles in the region $\Re(\zeta) < n/2$. The definition of the extended set $\tilde{\mathcal{S}}$ requires a bit more care, because at $\zeta \in \frac{1}{2}n - \mathbb{N}$, the inverse $S(\zeta)^{-1} = S(n - \zeta)$ has infinite-rank poles, which means that the formula (1.2) cannot be used to define the multiplicity at such points. Following [7], we can use a gamma-function to remove the infinite-rank zeroes at these problem points. Define the modified scattering operator

$$\tilde{S}(\zeta) = \Gamma(\zeta - n/2 + 1)S(\zeta),$$

and let $\tilde{\mathcal{S}}$ be the set of poles of $\tilde{S}(\zeta)$ for $\Re(\zeta) < n/2$. The multiplicity is now defined by

$$\nu_{\zeta_0} = \frac{1}{2\pi i} \operatorname{Tr} \left(\int_{\gamma_{\zeta_0, \varepsilon}} \tilde{S}(\zeta)^{-1} \tilde{S}'(\zeta) d\zeta \right).$$

We will show that, in even dimension, S_ζ vanishes at the points $\frac{1}{2}n - \mathbb{N}$, which implies that $R_g(\zeta)$ and $\tilde{S}(\zeta)$ are both holomorphic at these points.

Theorem 1.2. *Let (X, g) have even dimension and constant curvature near infinity, and let $\tilde{\mathcal{R}}$ and $\tilde{\mathcal{S}}$ denote the enlarged resolvent resonance and scattering resonance sets for Δ_g . Then $\tilde{\mathcal{R}} = \tilde{\mathcal{S}}$ and the relation*

$$\nu_{\zeta_0} = m_{n-\zeta_0} - m_{\zeta_0}$$

holds for all $\zeta_0 \in \tilde{\mathcal{R}}$.

This paper is organized as follows. In Section 2 we review the Mazzeo-Melrose construction of the resolvent and study its behavior near resolvent resonances. In Section 3 we recall how the scattering operator can be recovered from the resolvent and discuss its behavior near scattering poles. In Section 4 we study the perturbation behavior of resonances when the operator Δ_g is perturbed by a potential $V \in \dot{C}^\infty(X)$. In Section 5 we show that the operator $P_V = \Delta_g + V$ has only simple resonances ζ_0 for $\zeta_0 \notin \frac{1}{2}(n - \mathbb{N})$ for potentials V in a dense open subset of $\dot{C}^\infty(X)$. Finally, in Section 6, we prove Theorems 1.1 and 1.2.

In what follows, $x^N L^2(X)$ denotes the space of locally square-integrable functions v on X with $v = x^N u$ for a function $u \in L^2(X)$ and a fixed real number N . For a fixed, given N , we denote by B_0 the Banach space $x^N L^2(X)$, and by B_1 the Banach

space $x^{-N}L^2(X)$. If \mathcal{X} and \mathcal{Y} are Banach spaces, $\mathcal{L}(\mathcal{X}, \mathcal{Y})$ denotes the Banach space of bounded operators from \mathcal{X} to \mathcal{Y} . We metrize $\dot{C}^\infty(X)$ by introducing the seminorms

$$(1.3) \quad d_{\alpha,n}(V) = \sup_{x \in X} |x^{-n} D^\alpha V(x)|,$$

for nonnegative integers n and multi-indices α , and we denote by $d(\cdot, \cdot)$ the associated metric on $\dot{C}^\infty(X)$.

2. THE RESOLVENT OF $\Delta_g + V$ AND ITS MEROMORPHIC CONTINUATION

The resolvent of the operator $P_V = \Delta_g + V$ has a distribution kernel, with respect to the Riemannian density on X , which is smooth away from the diagonal Λ of $X \times X$. To describe its singularities, it is useful to introduce the blow up of $\overline{X} \times \overline{X}$ along $\partial\overline{X} \times \partial\overline{X}$, the stretched product $\overline{X} \times_0 \overline{X}$. This amounts to introducing polar coordinates at the diagonal in the corner of $\overline{X} \times \overline{X}$ where Λ intersects the ‘top’ boundary face $\overline{X} \times \partial\overline{X}$ and the ‘bottom’ boundary face $\partial\overline{X} \times \overline{X}$; globally one replaces $\partial\Lambda$ with the doubly inward-pointing spherical normal bundle of $\partial\Lambda$. If (x, y) and (x', y') are local coordinates on \overline{X} in a neighborhood of the boundary, $\partial\Lambda$ is given by $x = x' = y - y' = 0$; local coordinates for $\overline{X} \times_0 \overline{X}$ near the boundary are then given by $(r, \eta, \eta', \theta, y)$ where

$$r = \sqrt{x^2 + (x')^2 + |y - y'|^2},$$

$$(\eta, \eta', \theta) = (x/r, x'/r, (y - y')/r).$$

We denote by β the ‘blow-down map’ $\beta : \overline{X} \times_0 \overline{X} \rightarrow \overline{X} \times \overline{X}$; in the local coordinates described above, $\beta(r, \eta, \eta', \theta, y) = (r\eta, y, r\eta', y - r\theta)$.

The following theorem summarizes the Mazzeo-Melrose [20] construction of the resolvent. Although Mazzeo and Melrose did not treat potential perturbations, potentials in the class $\dot{C}^\infty(X)$ may be accommodated without difficulty (see [11], Theorem 3.1 and its proof). We denote by G_ζ the integral kernel of the resolvent operator $R_V(\zeta) = (P_V - \zeta(n - \zeta))^{-1}$ with respect to Riemannian measure on X , initially a meromorphic function of ζ with $\Re(\zeta) > n/2$.

Theorem 2.1. *Let (X, g) be an asymptotically hyperbolic manifold, and let $V \in \dot{C}^\infty(X)$. The resolvent kernel G_ζ has a meromorphic continuation to \mathbb{C} with*

$$\beta^* G_\zeta = A_\zeta + B_\zeta + C_\zeta$$

where

$$A_\zeta \in I^{-2}(\overline{X} \times_0 \overline{X}),$$

$$B_\zeta \in (\eta\eta')^\zeta C^\infty(\overline{X} \times_0 \overline{X}),$$

and

$$C_\zeta \in \beta^* [(xx')^\zeta C^\infty(\overline{X} \times \overline{X})].$$

Moreover A_ζ is an entire function of ζ , B_ζ is holomorphic in $\mathbb{C} \setminus \frac{1}{2}(n - \mathbb{N})$, and C_ζ is meromorphic in $\mathbb{C} \setminus \frac{1}{2}(n - \mathbb{N})$.

Sketch of the proof. Let $P_\zeta = \Delta_g + V - \zeta(n - \zeta)$. Given an operator A on $C^\infty(X)$, we will denote by $\kappa(A)$ the lift of the kernel of A (with respect to the Riemannian density on X) to $\overline{X} \times_0 \overline{X}$. The construction in [20] may be broken into three pieces. First, we construct an operator A_ζ to cancel the conormal singularity of $\kappa(P_\zeta)$ on the lifted diagonal. The family A_ζ is entire and has the property that $\kappa(I - P_\zeta A_\zeta) \in C^\infty(\overline{X} \times_0 \overline{X})$. This remainder does not yet correspond to the integral kernel of a compact operator on the original space.

To improve the error term, one uses the model resolvent. A second operator B_ζ is constructed so that $E_\zeta = I - P_\zeta(A_\zeta + B_\zeta)$ has

$$\kappa(E_\zeta) \in \eta^\zeta(\eta' r)^\infty C^\infty(\overline{X} \times_0 \overline{X}).$$

The operator B_ζ is holomorphic in $\mathbb{C} \setminus \frac{1}{2}(n - \mathbb{N})$; the operator E_ζ is a compact operator on the weighted L^2 space $x^N L^2(X)$ for all ζ with $\Re(\zeta) > n/2 - N$, and is also holomorphic in $\mathbb{C} \setminus \frac{1}{2}(n - \mathbb{N})$.

Finally, one inverts $(I - E_\zeta)$ using analytic Fredholm theory. Composition theorems of [18] show that if

$$(I + F_\zeta) = (I - E_\zeta)^{-1},$$

then $\kappa(F_\zeta)$ also lies in $\eta^\zeta(\eta' r)^\infty C^\infty(\overline{X} \times_0 \overline{X})$. This in turn implies that $C_\zeta = (A_\zeta + B_\zeta)F_\zeta$ belongs to $\beta^*[(xx')^\zeta C^\infty(\overline{X} \times \overline{X})]$, and therefore β^*G_ζ has the claimed form. \square

Remark 2.2. It follows from the form of the resolvent kernel that $R_V(\zeta)$ is a continuous mapping from $\dot{C}^\infty(X)$ to $C^\infty(X)$ when defined, and extends to a bounded mapping from $x^N L^2(X)$ to $x^{-N} L^2(X)$ for $\Re(\zeta) > n/2 - N$.

Remark 2.3. The Mazzeo-Melrose construction does not rule out the possibility of poles at $\zeta \in \frac{1}{2}(n - \mathbb{N})$, possibly of infinite rank. If (X, g) has constant negative curvature in a neighborhood of infinity, the operator $A_\zeta + B_\zeta$ may be replaced by the model resolvent (the resolvent of the Laplacian on the covering space \mathbb{H}^{n+1}), which is entire if n is even and has finite-rank poles at $\zeta = -k$ if n is odd (see for example the explicit formulas in [6], section 2). In either case, these terms contain only poles of finite rank, and the last step of the construction, involving the meromorphic Fredholm theorem, gives at most poles with finite-rank residues. A detailed construction of the resolvent in this case is given in [6], section 3. This observation plays a crucial role in the proof of Theorem 1.2.

Theorem 2.1 and standard arguments (see [7], Lemma 2.4) enable us to characterize the polar part of $R_V(\zeta)$ at a resolvent resonance $\zeta_0 \notin \frac{1}{2}(n - \mathbb{N})$. We will view the meromorphically continued resolvent as a mapping from the space $B_0 = x^N L^2(X)$ to $B_1 = x^{-N} L^2(X)$ as in Remark 2.2, where N is chosen so that $\Re(\zeta_0) > n/2 - N$. Introduce the nondegenerate form

$$\langle u, v \rangle = \int_X uv \, dx$$

(no complex conjugation), which can be used to pair elements in B_0 and B_1 . The resolvent operator is symmetric with respect to this form.

Proposition 2.4. *Let $\zeta_0 \in \mathbb{C} \setminus \frac{1}{2}(n - \mathbb{N})$ be a pole of $R_V(\zeta)$. Then*

$$R_V(\zeta) = \sum_{j=-k}^{-1} (\zeta(n - \zeta) - \zeta_0(n - \zeta_0))^j A_j + H(\zeta),$$

where $H(\zeta)$ is a holomorphic $\mathcal{L}(B_0, B_1)$ -valued function near $\zeta = \zeta_0$ and the A_j are finite-rank operators in $\mathcal{L}(B_0, B_1)$ with

$$A_{-j} = (\Delta_g + V - \zeta_0(n - \zeta_0))^{j-1} A_{-1}$$

for $j \geq 2$. The operator A_{-1} commutes with $\Delta_g + V - \zeta_0(n - \zeta_0)$. Moreover, there is a basis $\{\psi_i\}_{i=1}^{m_{\zeta_0}}$ for $\text{Ran}(A_{-1})$ so that

$$A_{-1}f = \sum_i \langle f, \psi_i \rangle \psi_i,$$

and the operator $(\Delta_g + V - \zeta_0(n - \zeta_0))$ is represented on $\text{Ran}(A_{-1})$ by a matrix M with $M^{k-1} \neq 0$ but $M^k = 0$.

Remark 2.5. If (X, g) has constant curvature in a neighborhood of infinity, the same result holds for any resolvent resonance ζ_0 including those $\zeta_0 \in \frac{1}{2}(n - \mathbb{N})$.

3. THE SCATTERING OPERATOR FOR P_V

Let $S_V(\zeta)$ denote the scattering operator for $P_V = \Delta_g + V$. To describe the scattering operator and its singularities, we recall its connection with the resolvent kernel. First, we blow up $\partial\overline{X} \times \partial\overline{X}$ along the diagonal Λ_∞ to obtain the space $\partial\overline{X} \times_0 \partial\overline{X}$. The map $\beta_\partial : \partial\overline{X} \times_0 \partial\overline{X} \rightarrow \partial\overline{X} \times \partial\overline{X}$ is the ‘blow-down map’ for this resolution. If (y, y') are local coordinates for $\partial\overline{X} \times \partial\overline{X}$, $r = |y - y'|$, and $\theta = (y - y')/r$, then (r, θ, y) give local coordinates for $\partial\overline{X} \times_0 \partial\overline{X}$, and $\beta_\partial(r, \theta, y) = (y, y + r\theta)$. The kernel of the scattering operator is recovered as an asymptotic limit of the resolvent kernel. Let $\kappa(A)$ denote the lift of the integral kernel of A (with respect to the measure on $\partial\overline{X}$ induced by the metric $h|_{\partial\overline{X}}$) to $\partial\overline{X} \times_0 \partial\overline{X}$. Then [2, 11]

$$(3.1) \quad \kappa(S_V(\zeta)) = \beta_\partial^* ((xx')^{-\zeta} G_\zeta)|_{T \cap B},$$

where $T \cap B$ is the intersection of the top and bottom faces of $\overline{X} \times_0 \overline{X}$. From this formula and Theorem 2.1, we easily obtain:

Theorem 3.1. *The decomposition*

$$\kappa(S_V(\zeta)) = r^{-2\zeta} F_\zeta + \beta_\partial^*(K_\zeta)$$

holds, where F_ζ and K_ζ are meromorphic maps respectively into $C^\infty(\partial\overline{X} \times_0 \partial\overline{X})$ and $C^\infty(\partial\overline{X} \times \partial\overline{X})$, and F_ζ is holomorphic in $\mathbb{C} \setminus \frac{1}{2}\mathbb{Z}$. At poles $\zeta_0 \notin \mathbb{C} \setminus \frac{1}{2}\mathbb{Z}$, the kernel K_ζ has polar part with finite Laurent series and coefficients in $C^\infty(\partial\overline{X} \times \partial\overline{X})$. The set of such poles is contained in the set of poles of $R_V(\zeta)$. For $\zeta_0 \in Z_p$ with $\zeta_0 \notin \frac{1}{2}n + \mathbb{Z}$, $S_V(\zeta)$ has a semi-simple pole at ζ_0 whose residue has rank m_{ζ_0} .

Note that the distribution $r^{-2\zeta}$ has poles at $\zeta \in \frac{1}{2}n + \mathbb{N}$, giving rise to infinite-rank first-order poles of $S_V(\zeta)$. The statement about poles at $\zeta_0 \in Z_p$ follows from the fact that the resolvent has a semi-simple pole at each such ζ_0 and that the residue is a finite-rank projection onto eigenfunctions of the form $x^{\zeta_0}\psi$ for $\psi \in C^\infty(\overline{X})$ with $\psi|_{\partial\overline{X}} \neq 0$. Thus from (3.1) it follows that $S_V(\zeta)$ has a semi-simple pole with residue $\sum_{i=1}^{m_{\zeta_0}} \langle \varphi_i, \cdot \rangle \varphi_i$ where $\varphi_i = x^{-\zeta_0} \psi_i|_{\partial\overline{X}}$.

The scattering operator $S_V(\zeta)$ is a family of Fredholm operators meromorphic in $\mathbb{C} \setminus \frac{1}{2}\mathbb{Z}$, satisfying

$$S_V(\zeta)S_V(n - \zeta) = I,$$

with finite polar parts and finite-rank residues at each pole $\zeta_0 \in \mathbb{C} \setminus \frac{1}{2}\mathbb{Z}$. For any $\zeta_0 \notin \frac{1}{2}\mathbb{Z}$, the multiplicity is

$$\nu_{\zeta_0} = \operatorname{Tr} \left(\frac{1}{2\pi i} \int_{\gamma_{\zeta_0, \varepsilon}} S_V(\zeta)^{-1} S'_V(\zeta) d\zeta \right).$$

In [4] Gohberg-Sigal gave a general method for computing the integer ν_{ζ_0} . Suppose $A(\zeta)$ is a meromorphic family of Fredholm operators on the Banach space \mathcal{B} , holomorphic for $\zeta \neq \zeta_0$ in some neighborhood of ζ_0 , such that the nonsingular part of A at ζ_0 has index zero. Assume also that there is a meromorphic inverse $B(\zeta)$ such that $A(\zeta)B(\zeta) = B(\zeta)A(\zeta) = I$. A *root function* of A is a holomorphic \mathcal{B} -valued function such that $\phi(\zeta_0) \neq 0$ but

$$\lim_{\zeta \rightarrow \zeta_0} A(\zeta)\phi(\zeta) = 0.$$

Then $\phi_0 = \phi(\zeta_0)$, which plays the role of a null vector of $A(\zeta_0)$, is called a *root vector*. As an illustration, consider

$$A(z) = \begin{pmatrix} z & 0 \\ 0 & 1 \end{pmatrix}, \quad C(z) = \begin{pmatrix} 1 & z^{-1} \\ 0 & 1 \end{pmatrix}.$$

Then A , C , and C^{-1} have a root vector $\phi_0 = (1, 0)$ at $z = 0$, while A^{-1} has no root vector.

The rank of ϕ_0 is the maximal order of vanishing of $A(\zeta)\phi(\zeta)$ at ζ_0 . Gohberg-Sigal define a number $N_{\zeta_0}(A)$ which is essentially the total rank of the space of null vectors of A at ζ_0 . Since we will be splitting into simple poles, we need only worry about the simplest possible cases: $N_{\zeta_0}(A) = 0$ if no root vector exists and $N_{\zeta_0}(A) = 1$ if the space of root vectors is 1-dimensional with $A(\zeta)\phi(\zeta)$ vanishing to first order. Theorem 2.1 of [4] says

$$\nu_{\zeta_0}(A) = N_{\zeta_0}(A) - N_{\zeta_0}(B),$$

where, in this setting, $\nu_{\zeta_0}(A)$ is defined by the equation

$$\nu_{\zeta_0}(A) = \operatorname{Tr} \frac{1}{2\pi i} \left(\int_{\gamma_{\zeta_0, \varepsilon}} B(\zeta) A'(\zeta) d\zeta \right)$$

for $\gamma_{\zeta_0, \varepsilon}$ a simple closed contour enclosing ζ_0 and no other singularity of $A(\zeta)$ or $B(\zeta)$.

Now we can prove equality of multiplicities in the special case when all resonances are simple. In the next two sections we shall show that all resonances are simple for ‘generic’ V , from which the general result will follow.

Proposition 3.2. *$R_V(\zeta)$ has a simple resonance at $\zeta_0 \in \mathbb{C} \setminus \frac{1}{2}\mathbb{Z}$ iff $S_V(\zeta)$ has a simple pole at ζ_0 . Moreover assuming that ζ_0 and $n - \zeta_0$ are at most simple poles of R_V , we have*

$$\nu_{\zeta_0} = m_{n-\zeta_0} - m_{\zeta_0}.$$

Proof. Assume $\zeta_0 \in \mathbb{C} \setminus \frac{1}{2}\mathbb{Z}$ is a simple resonance of $R_V(\zeta)$. The polar part of $R_V(\zeta)$ is

$$(\zeta(n - \zeta) - \zeta_0(n - \zeta_0))^{-1} \langle \psi_0, \cdot \rangle \psi_0$$

where $\psi_0 \in x^{\zeta_0} C^\infty(\overline{X})$ solves the eigenvalue equation $(\Delta_g + V - \zeta_0(n - \zeta_0))\psi_0 = 0$. We claim that $\varphi_0 = x^{-\zeta_0}\psi_0|_{\partial\overline{X}}$ is a nonzero element of $C^\infty(\partial\overline{X})$. If not, then

introducing local coordinates (x, y) on X where x is a defining function for $\partial\overline{X}$, we have $\psi_0 \in x^{\zeta_0+1}C^\infty$, and a power series argument using the eigenvalue equation shows that, in fact, $\psi_0 \in \dot{C}^\infty(X)$, so that ψ_0 is an L^2 -eigenfunction, a contradiction. It now follows from (3.1) that $S_V(\zeta)$ has a first-order pole at $\zeta = \zeta_0$ with rank-one residue $(\varphi_0, \cdot)\varphi_0$, where (\cdot, \cdot) is the real inner product on functions in $C^\infty(\partial\overline{X})$ with Riemannian measure induced by the metric $h|_{\partial\overline{X}}$.

Thus $S_V(\zeta)$ can be factored as

$$S_V(\zeta) = E(\zeta)[(\zeta - \zeta_0)^{-1}P_0 + H(\zeta)]F(\zeta),$$

where E and F are holomorphically invertible, H is holomorphic, and P_0 is the orthogonal projection onto the span of ϕ_0 . By [4] we can drop the E and F in computing N_{ζ_0} . Let

$$\begin{aligned}\phi(\zeta) &= (\zeta - \zeta_0)S_V(\zeta)\phi_0 \\ &= \phi_0 + (\zeta - \zeta_0)H(\zeta)\phi_0.\end{aligned}$$

Since

$$S_V^{-1}(\zeta)\phi(\zeta) = (\zeta - \zeta_0)\phi_0,$$

ϕ_0 is a rank one root vector for S_V^{-1} , and it is unique up to a scalar multiple. Thus (under the simplicity assumption)

$$N_{\zeta_0}(S_V^{-1}) = m_{\zeta_0}.$$

Applying the same argument at $n - \zeta_0$ gives

$$N_{\zeta_0}(S_V) = N_{n-\zeta_0}(S_V^{-1}) = m_{n-\zeta_0}.$$

So the result follows from the Gohberg-Sigal formula for the multiplicity. \square

A careful analysis of the resolvent parametrix construction (see for example [2]) shows that the map

$$\left(\mathbb{C} \setminus \frac{1}{2}\mathbb{Z}\right) \times \dot{C}^\infty(X) \ni (\zeta, V) \longmapsto S_V(\zeta)$$

is a continuous mapping away from poles of $S_V(\zeta)$. The following continuity result for scattering poles is a simple consequence.

Proposition 3.3. *Let $\zeta_0 \in \mathcal{S}$ with $\zeta_0 \in \mathbb{C} \setminus \frac{1}{2}(n - \mathbb{N})$. For sufficiently small $\varepsilon > 0$ there is a $\delta > 0$ so that for all $V \in \dot{C}^\infty(X)$ with $d(V, 0) < \delta$, the multiplicity*

$$\nu_{\zeta_0}(V) = \operatorname{Tr} \left(\frac{1}{2\pi i} \int_{\gamma_{\zeta_0, \varepsilon}} S_V^{-1}(\zeta) S'_V(\zeta) d\zeta \right)$$

is continuous as a map from $\dot{C}^\infty(X)$ to \mathbb{Z} . Hence

$$\nu_{\zeta_0}(V) = \nu_{\zeta_0}(0)$$

for such V .

4. PERTURBATION THEORY OF RESONANCES

Now we apply Agmon's perturbation theory of resonances [1] to study the behavior of resonances under potential perturbations. For a fixed, asymptotically hyperbolic manifold (X, g) we consider the family of operators P_V where V ranges over complex-valued $\dot{C}^\infty(X)$ functions. For any such V , Theorem 2.1 guarantees that $R_V(\zeta) = (P_V - \zeta(n - \zeta))^{-1}$ admits a meromorphic continuation to any half-plane $\Re(\zeta) > n/2 - N$, N a positive integer, as a mapping from $B_0 = x^N L^2(X)$ to $B_1 = x^{-N} L^2(X)$. We will set $\mathcal{R}_V(z) = (P_V - z)^{-1}$ with the understanding that z lies on the second sheet of the Riemann surface for the inverse function of $f(\zeta) = \zeta(n - \zeta)$, so that $\mathcal{R}_V(z)$ is the meromorphic continuation of the resolvent to the second sheet. Using the fact that $P_V : \dot{C}^\infty(X) \rightarrow \dot{C}^\infty(X)$ together with Theorem 2.1, it is not difficult to check that the operator P_V satisfies the hypotheses of Agmon's abstract theory.

To study the perturbation of a resonance z_0 , Agmon introduces auxiliary operators and Banach spaces associated to an open connected domain Δ containing z_0 with C^1 boundary Γ . Let B_Γ be the subset of B_1 consisting of functions of the form

$$u = f + \int_\Gamma \mathcal{R}_V(w) \Phi(w) dw,$$

where $f \in B_0$ and $\Phi \in C(\Gamma; B_0)$, the continuous functions on Γ with values in B_0 . Finally, let Y be the closed subspace of $B_0 \times C(\Gamma, B_0)$ consisting of those (g, Φ) with

$$0 = g + \int_\Gamma \mathcal{R}_V(w) \Phi(w) dw.$$

The space B_Γ is a Banach space as the quotient of $B_0 \times C(\Gamma, B_0)$ by the closed subspace Y when equipped with the quotient norm

$$\|u\|_{B_\Gamma} = \inf \left\{ \|f\|_{B_0} + \|\Phi\|_{C(\Gamma; B_0)} : u = f + \int_\Gamma \mathcal{R}_V(w) \Phi(w) dw \right\}.$$

The space B_Γ satisfies $B_0 \subset B_\Gamma \subset B_1$, where the canonical injections are continuous.

The theory of [1], which implies that there is a closed operator $P_V^\Gamma : \mathcal{D}(P_V^\Gamma) \rightarrow B_\Gamma$ which is a restriction of P_V , in a sense we will make precise, and whose eigenvalues in Δ are exactly the resonances of $\mathcal{R}_V(z)$ in Δ . In fact, let \overline{P}_V be the closure of P_V as a densely defined operator from B_1 to itself. Then $P_V^\Gamma u = \overline{P}_V u$ for all $u \in \mathcal{D}(P_V^\Gamma)$. The Laurent expansion of $\mathcal{R}_V^\Gamma(z) = (P_V^\Gamma - z)^{-1}$ near a resonance $z_0 \in \Delta$ takes the form

$$\sum_{j=-k}^{-1} (z - z_0)^j A_j^\Gamma + H^\Gamma(z),$$

where $H^\Gamma(z)$ is a holomorphic $\mathcal{L}(B_\Gamma)$ -valued function in a neighborhood of z_0 and the A_j^Γ are finite-rank operators belonging to $\mathcal{L}(B_\Gamma, \mathcal{D}(P_V^\Gamma))$. For $f \in B_0$ we have

$$A_j^\Gamma f = A_j f,$$

where A_j are the corresponding Laurent coefficients for $\mathcal{R}_V(z)$.

Now fix $V \in \dot{C}^\infty(X)$ and set $P(t) = P_V + tW$ for another potential $W \in \dot{C}^\infty(X)$. The operators $P(t)$ have resolvents which admit meromorphic continuation to $\mathcal{L}(B_0, B_1)$ -valued meromorphic functions in $\Re(s) > n/2 - N$ for any fixed $N > 0$.

Moreover, for t small and a fixed region Δ , the spaces $B_\Gamma(t)$ corresponding to $P(t)$ are equal as sets and carry equivalent norms, and the operators $P_\Gamma(t)$ form an analytic family of type (A) in the sense of Kato [12]. Let $\mathcal{R}(t, z) = (P(t) - z)^{-1}$ and $\mathcal{R}^\Gamma(t, z) = (P^\Gamma(t) - z)^{-1}$. Theorem 7.7 of [1] shows that, for small t , the resolvents $\mathcal{R}^\Gamma(t, z)$ and $\mathcal{R}(t, z)$ coincide on B_0 , possess the same set of poles for each fixed t , and $\text{Ran}(A_j^\Gamma(t)) = \text{Ran}(A_j(t))$, where $A_j(t)$ and $A_j^\Gamma(t)$ are the respective Laurent coefficients of $\mathcal{R}(t, z)$ and $\mathcal{R}^\Gamma(t, z)$ at a given pole in Δ .

5. GENERIC SIMPLICITY OF RESONANCES

For L^2 eigenvalues of the Laplacian and its perturbations, it has long been known that ‘generic’ potential perturbations split degenerate eigenvalues so that a single eigenvalue of multiplicity m becomes m simple eigenvalues, localized near the original eigenvalue (see for example Uhlenbeck [28], where generic simplicity is proved for the Laplacian on compact manifolds, and Kato [12] for the background in perturbation theory of linear operators; Uhlenbeck’s methods adapt without difficulty to eigenvalues below the continuous spectrum). The purpose of this section is to show that the same is true of the resolvent resonances.

Theorem 5.1. *The set E of potentials $V \in \dot{C}^\infty(X)$ for which all eigenvalues of $\Delta_g + V$ and all resonances of $\Delta_g + V$ in $\mathbb{C} \setminus \frac{1}{2}(n - \mathbb{N})$ are simple is open and dense in $\dot{C}^\infty(X)$.*

We will follow rather closely the argument of [13] except that Agmon’s perturbation theory replaces the exterior complex scaling used there. Since generic simplicity results for eigenvalues are well-known we will only prove that generic simplicity holds for the resonances.

For positive integers N and real numbers $r > 0$, we define

$$\mathcal{R}_N^r = \left\{ \zeta \in \mathcal{R} : |\zeta| < r, \text{ dist}(\zeta, \tfrac{1}{2}(n - \mathbb{N})) > 1/N \right\},$$

and let

$$E_N^r = \left\{ V \in \dot{C}^\infty(X) : \text{each } \zeta \text{ in } \mathcal{R}_N^r \text{ is simple} \right\}.$$

We set

$$E = \bigcap_{n=1}^{\infty} \bigcap_{N=1}^{\infty} E_N^n,$$

and we define

$$F = \dot{C}^\infty(X) \setminus E.$$

We wish to show that E is dense in $\dot{C}^\infty(X)$, i.e., that F has empty interior. By the Baire category theorem, it suffices to show that $F_N^n = \dot{C}^\infty(X) \setminus E_N^n$ is nowhere dense for each n and N . By the discreteness of the resonance set in $\mathbb{C} \setminus \frac{1}{2}(n - \mathbb{N})$, it suffices to show that for any $V \in F_N^n$, any nonsimple pole ζ_0 , and any $\varepsilon > 0$, there is a W with $x(W, 0) < \varepsilon$ so that $V + W$ has only simple poles in a neighborhood of ζ_0 . As in Section 4, it will be convenient to work with the meromorphically continued resolvent $\mathcal{R}(z)$, and we will denote by z_0 the point on the second sheet of the Riemann surface for $\mathcal{R}(z)$ corresponding to ζ_0 .

Consider the family of operators $P_{V+W} = \Delta_g + V + W$, where $d(W, 0) < \varepsilon_0$, and $\varepsilon_0 > 0$ is to be chosen (here $d(\cdot, \cdot)$ is the metric on $\dot{C}^\infty(X)$ defined in (1.3)). Let

Γ be a contour enclosing z_0 and no other pole of $\mathcal{R}(z)$. For ε_0 small enough, the operators P_{V+W}^Γ defined in Agmon's abstract theory can be considered to act on a single Banach space B_Γ , and the associated projection

$$\Pi_{V+W}^\Gamma = \frac{1}{2\pi i} \int_\Gamma (P_{V+W}^\Gamma - w)^{-1} dw$$

is analytic in W with $d(W, 0) < \varepsilon_0$, and of constant rank, say m . As in [13], we note that either

- (1) for each $\varepsilon > 0$, there is a W with $x(W, 0) < \varepsilon$ so that P_{V+W}^Γ has at least two distinct eigenvalues, or
- (2) there is an $\varepsilon > 0$ so that for all W with $x(W, 0) < \varepsilon$, P_{V+W}^Γ has a single eigenvalue $z(W)$ and there is an integer $k(W)$, $1 \leq k(W) \leq m$, so that

$$(P_{V+W}^\Gamma - z(W))^{k(W)} \Pi_{V+W}^\Gamma = 0 \quad (P_{V+W}^\Gamma - z(W))^{k(W)-1} \Pi_{V+W}^\Gamma \neq 0.$$

If case (2) does not occur, we can split resonances repeatedly by small perturbations. Thus we will suppose that case (2) does occur and obtain a contradiction.

First, note that $k(W)$ is locally constant, so by taking ε_0 small enough we may assume that $k(W)$ is constant for W with $x(W, 0) < \varepsilon_0$. As in [13] we consider in turn the possibilities $k(W) = 1$ (the semi-simple case) and $k(W) \geq 2$.

First suppose that $k(W) = 1$, that $z(W)$ is an eigenvalue of P_{V+W}^Γ of multiplicity m , and let $\{\psi_i\}_{i=1}^m$ be a basis for $\text{Ran}(A_{-1}^\Gamma)$, where A_{-1}^Γ occurs in the Laurent expansion for $(P_V^\Gamma - z)^{-1}$ at $z = z_0$. The vectors $\{\psi_j\}_{j=1}^m$ belong to $B_\Gamma \subset B_1$ and may be chosen to diagonalize A_{-1}^Γ as in Proposition 2.4. Let $\{f_j\}_{j=1}^m$ be a set of vectors in B_0 with $\langle \psi_i, f_j \rangle = \delta_{ij}$. Finally, for fixed W , let $L(t) = P_{V+tW}^\Gamma$, let $\Pi_t = \Pi_{V+tW}^\Gamma$, let $\psi_i(t) = \Pi_t \psi_i$, and let $z(t) = z(tW)$. By differentiating the eigenvalue equation

$$(L(t) - z(t))\psi_i(t) = 0$$

at $t = 0$, we recover the identity

$$(W - z'(0))\psi_i + (L(0) - z(0))\psi'_i(0) = 0.$$

We now apply the projection Π_0 to both sides, pair with f_j , and use the fact that $(L(0) - z(0))\Pi_0 = \Pi_0(L(0) - z(0)) = 0$ to conclude that

$$\langle f_j, \Pi_0 W \psi_i \rangle = z'(0) \delta_{ij}.$$

From the choice of $\{f_i\}$ and the fact that $\Pi_0 = \sum_i \langle \psi_i, \cdot \rangle \psi_i$ it now follows that

$$\langle \psi_i, W \psi_j \rangle = z'(0) \delta_{ij}.$$

Since this must hold for any $W \in \dot{C}^\infty(X)$ (in particular for all $W \in C_0^\infty(U)$ with U an open subset of X), it follows that at least one of the ψ_i vanishes on U , and hence on X by unique continuation. This gives a contradiction.

Now suppose that $z(W)$ is not semi-simple, but that there is a fixed $k \geq 2$ so that

$$(L(t) - z(t))^k \Pi_t = 0, \quad (L(t) - z(t))^{k-1} \Pi_t \neq 0.$$

Choose a vector $h \in B$ with $\psi(t) = (L(t) - z(t))^{k-1} \Pi_t h \neq 0$, so that

$$(L(t) - z(t))\psi(t) = 0.$$

Let $\psi = \psi(0)$. A perturbation calculation again leads to

$$(5.1) \quad (W - z'(0))\psi + (L(0) - z(0))\psi'(0) = 0.$$

If we apply the projection Π_0 to both sides of (5.1) and pair with a vector $f \in B_0$ with $\Pi_0 f = \psi$, we obtain

$$\langle \psi, W\psi \rangle = z'(0) \langle f, \Pi_0 \psi \rangle.$$

We have used the fact that Π_0 is symmetric with respect to the pairing $\langle \cdot, \cdot \rangle$. To evaluate the right-hand side, we use the fact that $L(0) - z(0)$ preserves B_0 to write

$$\begin{aligned} \langle f, \Pi_0 \psi \rangle &= \langle f, \Pi_0 (L(0) - z(0))^{k-1} \Pi_0 h \rangle \\ &= \langle \psi, \Pi_0 (L(0) - z(0))^{k-1} \Pi_0 h \rangle \\ &= \langle (L(0) - z(0))^{k-1} \Pi_0 h, (L(0) - z(0))^{k-1} \Pi_0 h \rangle \\ &= \langle (L(0) - z(0))^{k-2} \Pi_0 h, (L(0) - z(0))^k \Pi_0 h \rangle \\ &= 0, \end{aligned}$$

so that $\langle \psi, W\psi \rangle = 0$ for all $W \in \dot{C}^\infty(X)$. It follows that ψ vanishes on X , a contradiction.

We have now shown that for each $\varepsilon > 0$, there is a W with $d(W, 0) < \varepsilon$ so that P_{V+W}^Γ has at least two distinct eigenvalues. It follows that any resonance can be split by an arbitrarily small perturbation $W \in \dot{C}^\infty(X)$. This fact implies that the set E of potentials V for which $\Delta_g + V$ has only simple resonances in $\mathbb{C} \setminus \frac{1}{2}(n - \mathbb{N})$ is open and dense in $\dot{C}^\infty(X)$, and Theorem 5.1 is proved.

In case (X, g) has constant curvature near infinity, this result can be improved if we work with the class $C_0^\infty(U)$ for a fixed open subset U of X . In this case the methods of [6] can be used to show that the resolvent of $\Delta_g + V$ has a meromorphic continuation with only finite-rank poles, including any poles at $\zeta_0 \in \frac{1}{2}(n - \mathbb{N})$. One can then apply the above arguments without essential changes to prove:

Theorem 5.2. *Let U be a fixed open subset of X with compact closure. The set E of potentials $V \in C_0^\infty(U)$ for which all eigenvalues and all resonances of $\Delta_g + V$ are simple is open and dense in $C_0^\infty(U)$.*

6. RESOLVENT RESONANCES AND SCATTERING POLES

Finally, we give the proofs of Theorems 1.1 and 1.2.

To prove Theorem 1.1, we choose a $\dot{C}^\infty(X)$ potential V so that all of the eigenvalues and resonances of $\Delta_g + V$ are simple. We further choose V small enough that, for a given point $\zeta_0 \in \mathcal{R}$ with $\zeta_0 \notin \mathbb{C} \setminus \frac{1}{2}(n - \mathbb{N})$, some $\varepsilon > 0$, and any $t \in (0, 1)$, no resonances of $\Delta_g + tV$ cross the circle $\gamma_{\zeta_0, \varepsilon}$ of radius ε about ζ_0 , and the projection

$$\Pi_{tV} = \frac{1}{2\pi i} \int_{\gamma_{\zeta_0, \varepsilon}} (2\zeta - n)(\Delta_g + tV - \zeta(n - \zeta))^{-1} d\zeta$$

(as well as its analogue for $n - \zeta_0$ if $\zeta_0 \in Z_p$) is continuous in t . It follows from Kato-Rellich perturbation theory for small t , the rank of Π_{tV} is continuous, so that

$$m(t) = \text{rank}(\Pi_{tV}) = m_{\zeta_0}$$

is constant for t small. On the other hand, we can also assume that for some $t \neq 0$ and small, the resonances of $\Delta_g + tV$ are all simple. Thus we can apply Proposition 3.2 to conclude

$$\text{Tr} \left(\frac{1}{2\pi i} \int_{\gamma_{\zeta_0, \varepsilon}} S_{tV}^{-1}(\zeta) S'_{tV}(\zeta) d\zeta \right) = m_{n-\zeta_0} - m_{\zeta_0}.$$

Finally, by Proposition 3.3, we have

$$\mathrm{Tr} \left(\frac{1}{2\pi i} \int_{\gamma_{\zeta_0, \varepsilon}} S_{tV}^{-1}(\zeta) S'_{tV}(\zeta) d\zeta \right) = \nu_{\zeta_0},$$

where ν_{ζ_0} is the multiplicity of the scattering pole ζ_0 of $S(\zeta)$, the scattering operator for the unperturbed Δ_g . This proves Theorem 1.1.

For poles $\zeta_0 \in \frac{1}{2}(n - \mathbb{N})$, the proof breaks down for several reasons: (i) the resolvent may have infinite rank poles at these points, making m_{ζ_0} undefined, (ii) the scattering operator may have infinite rank poles at the points $\frac{1}{2}n + \mathbb{N}$, making ν_{ζ_0} undefined at $\frac{1}{2}n - \mathbb{N}$. If (X, g) has constant curvature in a neighborhood of infinity, then as remarked in the introduction, problem (i) is avoided. Moreover, Theorem 5.1 on generic simplicity of resonances holds at all $\Re(\zeta) < n/2$ in this case because all poles have finite rank.

If in addition to constant curvature near infinity X has even dimension (i.e. n is odd), then problem (ii) can be bypassed because of very particular behavior of the resolvent and scattering operators at the points $\frac{1}{2}n - \mathbb{N}$.

Lemma 6.1. *Suppose that X has constant curvature near infinity, V is compactly supported, and $\dim X$ is even. Then for $\zeta \in \frac{1}{2}n - \mathbb{N}$, $S_V(\zeta) \equiv 0$.*

Proof. The proof is a direct generalization of [7], Lemma 2.5. Let $G_0(\zeta; w, w')$ be the integral kernel of $R_0(\zeta)$ with respect to Riemannian measure on \mathbb{H}^{n+1} , where $w = (x, y)$, $w' = (x', y')$ for $y, y' \in \mathbb{R}^n$, $x, x' > 0$. In general one has $G_0(\zeta) \in (xx')^\zeta C^\infty(\mathbb{H}^{n+1} \times \mathbb{H}^{n+1} \setminus \Delta)$, where Δ denotes the diagonal. However, at the special points $\frac{1}{2}n - \mathbb{N}$ we actually have

$$(6.1) \quad G_0(n/2 - k) \in (xx')^{\frac{n}{2}+k} C^\infty(\mathbb{H}^{n+1} \times \mathbb{H}^{n+1} \setminus \Delta).$$

This can be checked by writing down the resolvent explicitly or by using the formula

$$G_0(\zeta; w, w') - G_{0, n-\zeta}(w, w') = \int_{\mathbb{R}^n} e_0(\zeta; w, y'') e_0(n - \zeta; w, y'') dy'',$$

where

$$e_0(\zeta; w, y'') = \pi^{-n/2} \frac{\Gamma(\zeta)}{\Gamma(\zeta - n/2)} \frac{x^\zeta}{(|y - y''|^2 + x^2)^\zeta}.$$

The point is that $e_0(\zeta)$ vanishes at $\zeta = \frac{1}{2}n - \mathbb{N}$ if $\dim X = n + 1$ is even.

Now we want to make use of the iterative parametrix construction in [6]. Since g has constant curvature near infinity, a neighborhood of $\partial\overline{X}$ can be covered with finitely many neighborhoods each of which is isometric to a hemisphere bordering the real axis in \mathbb{H}^{n+1} . Guillope-Zworski construct a boundary model $M_0(\zeta)$ for the resolvent by pulling back G_0 into these neighborhoods. So $\beta^* M_0(\zeta) \in (\eta\eta')^\zeta C^\infty(\overline{X} \times_0 \overline{X})$, in the notation of §2, but has the extra decay as in (6.1) for $\zeta \in \frac{1}{2}n - \mathbb{N}$.

For each $N > 0$ one can construct a correction $M'(\zeta)$ such that

$$[\Delta_g + V - \zeta(n - \zeta)](M_0(\zeta) - M'(\zeta)) = I - K(\zeta),$$

where M' and K are meromorphic with finite rank poles, and the kernels satisfy

$$\begin{aligned} \beta^* \kappa(M') &\in \eta^{\zeta+2} \eta'^\zeta r^2 C^\infty(\overline{X} \times_0 \overline{X}), \\ \beta^* \kappa(K) &\in \eta^{\zeta+N} \eta'^\zeta r^2 C^\infty(\overline{X} \times_0 \overline{X}). \end{aligned}$$

This implies that K is compact on $x^N L^2(X)$ for $\Re \zeta > \frac{1}{2}n - N$, so the analytic Fredholm theory can be applied to give

$$R_V(\zeta) = (M_0(\zeta) - M'(\zeta))(I - K(\zeta))^{-1}.$$

Using the general composition properties of the kernels (see [18, 2]), one can show that $(I - K(\zeta))^{-1} = I + K'$ where K' has a similar structure to K , but possibly with extra logarithms at lower orders. We then have

$$(6.2) \quad R_V(\zeta) = (M_0(\zeta) - M'(\zeta))(I + K'(\zeta)).$$

Recall the characterization (3.1) of the scattering operator as a boundary limit of the resolvent. From this we see that $M'(I + K')$ never contributes to the scattering operator. The extra decay of $M_0(\zeta)$ at the special points $\zeta \in \frac{1}{2}n - \mathbb{N}$ shows that $(xx')^{-\zeta} M_0(I + K')$ vanishes in the limit $x \rightarrow 0$ at these points. So the scattering operator is zero. \square

Now we are ready to prove Theorem 1.2. By the perturbation argument as above we can assume that R_V has only simple resonances, and away from $\frac{1}{2}n - \mathbb{N}$ no new argument is needed.

Fix $k \in \mathbb{N}$ and suppose that R_V had a pole at $\zeta_0 = \frac{n}{2} - k$. Arguing as in Proposition 3.2, the polar part of $R_V(\zeta)$ is

$$(\zeta(n - \zeta) - \zeta_0(n - \zeta_0))^{-1} \langle \psi_0, \cdot \rangle \psi_0,$$

where $\psi_0 \in x^{\frac{n}{2}-k} C^\infty(\overline{X})$ solves the eigenvalue equation $(\Delta_g + V - \zeta_0(n - \zeta_0))\psi_0 = 0$. Suppose that $\varphi_0 = x^{-\frac{n}{2}+k} \psi_0|_{\partial \overline{X}} = 0$. Then $\psi_0 \in x^{\frac{n}{2}-k+1} C^\infty(\overline{X})$. The power series argument can be applied at these points to show that $\psi_0 \in x^{\frac{n}{2}+k} C^\infty(\overline{X})$. This would be L^2 and so gives contradiction. Therefore ϕ_0 would have to be nonzero, implying that $S_V(\zeta)$ would have a pole at ζ_0 . Since $S_V(\zeta_0) = 0$ we conclude that R_V is holomorphic at $\zeta_0 = \frac{n}{2} - k$ and $m_{\zeta_0} = 0$.

Now consider $\tilde{S}_V(\zeta) = \Gamma(\zeta - \frac{n}{2} + 1) S_V(\zeta)$ as in the introduction. By Lemma 6.1, \tilde{S}_V is holomorphic at $\zeta_0 = \frac{n}{2} - k$, so $N_{\zeta_0}(\tilde{S}_V^{-1}) = 0 = m_{\zeta_0}(V)$.

It remains to compute $N_{\zeta_0}(\tilde{S}_V)$ by considering the behavior of \tilde{S}_V^{-1} at ζ_0 . We can appeal to (6.2) to argue that

$$\begin{aligned} \kappa(\tilde{S}_V^{-1}(\zeta)) &= \frac{1}{\Gamma(\zeta - \frac{n}{2} + 1)} \kappa(S_V(n - \zeta)) \\ &= \frac{1}{\Gamma(\zeta - \frac{n}{2} + 1)} \beta_\partial^* \left((xx')^{-(n-\zeta)} G_{n-\zeta} \right) \Big|_{T \cap B} \end{aligned}$$

for ζ in a neighborhood of ζ_0 . Here the infinite-rank pole in S_V at $n - \zeta_0 = \frac{n}{2} + k$, which comes from the model term M_0 in $G_{n-\zeta}$, is cancelled by dividing out the gamma function.

The rest of the argument proceeds as before. We conclude that $N_{\zeta_0}(\tilde{S}_V) = 1$ if and only if $R_V(\zeta)$ has a simple pole at $\frac{n}{2} + k$. We thus have

$$\nu_{\zeta_0}(V) = N_{\zeta_0}(\tilde{S}_V) - N_{\zeta_0}(\tilde{S}_V^{-1}) = m_{n-\zeta_0}(V) - m_{\zeta_0}(V),$$

and the theorem follows.

REFERENCES

1. S. Agmon, A perturbation theory for resonances. *Comm. Pure Appl. Math.* **51** (1998), 1255–1309. MR **99i**:47023; erratum MR **2000g**:47010
2. D. Borthwick, Scattering theory and deformations of asymptotically hyperbolic metrics. Preprint, 1997.
3. R. Froese, P. Hislop, P. Perry, A Mourre estimate and related bounds for hyperbolic manifolds with cusps of non-maximal rank. *J. Funct. Anal.* **98** (1991), 292–310. MR **92h**:58198
4. I. C. Gohberg, E. I. Sigal, An operator generalization of the logarithmic residue theorem and the theorem of Rouché. *Math. U. S. S. R. Sbornik* **84** (1971), 607–629. MR **47**:2409
5. L. Guillopé, M. Zworski, Upper bounds on the number of resonances for non-compact Riemann surfaces. *J. Funct. Anal.* **129** (1995), 364–389. MR **96b**:58116
6. L. Guillopé, M. Zworski, Polynomial bounds on the number of resonances for some complete spaces of constant curvature. *Asymptotic Anal.* **11** (1995), 1–22. MR **96h**:58172
7. L. Guillopé, M. Zworski, Scattering asymptotics for Riemann surfaces. *Ann. Math.* **145** (1997), 597–660. MR **98g**:58181
8. G. Hagedorn, Link between scattering resonances and dilation-analytic resonances in few-body quantum mechanics. *Commun. Math. Phys.* **65** (1979), 181–201. MR **80f**:81093
9. A. Jensen, Local distortion technique, resonances, and poles of the S -matrix. *J. Math. Anal. Appl.* **59** (1977), 505–513. MR **55**:14017
10. A. Jensen, Resonances in an abstract analytic scattering theory. *Ann. Inst. Henri Poincaré* **33** (1980), 209–223. MR **82b**:47007
11. M. Joshi, A. Sá Barreto, Inverse scattering on asymptotically hyperbolic manifolds. *Acta Math.* **184** (2000), 41–86. CMP 2000:12
12. T. Kato, *Perturbation Theory for Linear Operators*. Berlin, Heidelberg, New York: Springer-Verlag, 1976. MR **53**:11389
13. F. Klopp, M. Zworski, Generic simplicity of resonances. *Helv. Phys. Acta* **68** (1995), 531–538. MR **97j**:81067
14. P. Lax, R. S. Phillips, *Scattering Theory*. New York: Academic Press, 1967. MR **36**:530
15. P. Lax, R. S. Phillips, The asymptotic distribution of lattice points in Euclidean and non-Euclidean spaces. *J. Funct. Anal.* **46**, 280–350 (1982). MR **83j**:10057
16. P. Lax, R. S. Phillips, Translation representation for automorphic solutions of the non-Euclidean wave equation I, II, III. *Comm. Pure. Appl. Math.* **37** (1984), 303–328, **37** (1984), 779–813, and **38** (1985), 179–208. MR **86c**:58148; MR **86h**:58140; MR **86j**:58150
17. R. Mazzeo, The Hodge cohomology of a conformally compact metric. *J. Diff. Geom.* **28** (1988), 309–339. MR **89i**:58005
18. R. Mazzeo, Elliptic theory of edge operators. *Comm. P. D. E.* **16** (1991), 1615–1664. MR **93d**:58152
19. R. Mazzeo, Unique continuation at infinity and embedded eigenvalues for asymptotically hyperbolic manifolds. *American J. Math.* **113** (1991), 25–56. MR **92f**:58187
20. R. Mazzeo, R. Melrose, Meromorphic extension of the resolvent on complete spaces with asymptotically constant negative curvature. *J. Funct. Anal.* **75** (1987), 260–310. MR **89c**:58133
21. R. B. Melrose, *Geometric Scattering Theory*. New York, Melbourne: Cambridge University Press, 1995. MR **96k**:35129
22. S. J. Patterson, The Selberg zeta-function of a Kleinian group. In *Number Theory, Trace Formulas, and Discrete Groups: Symposium in honor of Atle Selberg*, Oslo, Norway, July 14–21, 1987, New York, Academic Press, 1989, pp. 409–442. MR **91c**:11029
23. S. J. Patterson, P. A. Perry, Divisor of Selberg’s zeta function for Kleinian groups. *Duke Math. J.* **106** (2001), 321–390. CMP 2001:08
24. P. A. Perry, The Laplace operator on a hyperbolic manifold, II. Eisenstein series and the scattering matrix. *J. Reine Angew. Math.* **398** (1989), 67–91. MR **90g**:58138
25. P. A. Perry, The Selberg zeta function and a local trace formula for Kleinian groups. *J. Reine Angew. Math.* **410** (1990), 116–152. MR **92e**:11057
26. M. Reed, B. Simon, *Methods of Modern Mathematical Physics, III. Scattering Theory*. New York, Academic Press, 1979. MR **80m**:81085
27. J. Sjöstrand, M. Zworski, Lower bounds on the number of scattering poles. *Comm. P. D. E.* **18** (1993), 847–857. MR **94h**:35198

- 28. K. Uhlenbeck, Generic properties of eigenfunctions. *American J. Math.* **98** (1976), 1059–1078. MR **57**:4264
- 29. M. Zworski, Counting scattering poles. In *Proceedings of the Taniguchi International Workshop, Spectral and Scattering Theory*, M. Ikawa, ed., Marcel Dekker, New York, Basel, Hong Kong, 1994. MR **95i**:35210
- 30. M. Zworski, Dimension of the limit set and density of resonances for convex co-compact hyperbolic quotients. *Inventiones Math.* **136** (1999), 353–409. CMP 99:12
- 31. M. Zworski, Resonances in physics and geometry. *Notices Amer. Math. Soc.* **46** (1999), no. 3, 319–328. MR **2000d**:58051

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, EMORY UNIVERSITY, ATLANTA, GEORGIA 30322

E-mail address: davidb@mathcs.emory.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KENTUCKY, LEXINGTON, KENTUCKY 40506–0027

E-mail address: perry@ms.uky.edu