

ON DEGREES OF IRREDUCIBLE BRAUER CHARACTERS

W. WILLEMS

ABSTRACT. Based on a large amount of examples, which we have checked so far, we conjecture that $|G|_{p'} \leq \sum_{\varphi} \varphi(1)^2$ where p is a prime and the sum runs through the set of irreducible Brauer characters in characteristic p of the finite group G . We prove the conjecture simultaneously for p -solvable groups and groups of Lie type in the defining characteristic. In non-defining characteristics we give asymptotically an affirmative answer in many cases.

1. INTRODUCTION

Let G be a finite group and let $\text{IBr}_p(G)$, resp. $\text{IBr}_p(B)$, be the set of irreducible p -Brauer characters of G , resp. of a p -block B . For $\varphi \in \text{IBr}_p(G)$ let Φ_{φ} denote the projective indecomposable character corresponding to φ . Due to a result of Brauer and Nesbitt [6] the term $|G| - \frac{|G|}{\Phi_1(1)}$, where Φ_1 denotes the projective character corresponding to the trivial character, is an upper bound for the dimension of the Jacobson radical of the p -modular group algebra of G . An obvious reformulation of this result leads to

$$(1) \quad \frac{|G|}{\Phi_1(1)} \leq \sum_{\varphi \in \text{IBr}_p(G)} \varphi(1)^2.$$

Moreover, equality holds if and only if G has a normal Sylow p -subgroup. This was proved by Wallace in [18] for p -solvable groups, and in full generality by Brockhaus in [7] using the classification of finite simple groups.

In case $p = 2$ the classification can be avoided to show the normality of a Sylow p -subgroup if equality holds in (1). Since Φ_{φ} is a constituent of $\varphi\Phi_1$, equality in (1) implies that $\Phi_{\varphi} = \varphi\Phi_1$. Now, if $p \mid \varphi(1)$, the trivial character has multiplicity at least 2 in $\varphi\overline{\varphi}$, and a contradiction follows by considering the scalar product relation

$$1 = (\varphi, \Phi_{\varphi}) = (\varphi, \varphi\Phi_1) = (\varphi\overline{\varphi}, \phi_1) \geq 2$$

(see [13], VII, 8.5 d)). Thus p does not divide any irreducible Brauer character degree, and a nice argument of Okuyama (see [16], Theorem 2.33) implies that for $p = 2$ the Sylow p -subgroup of G is normal.

The lower bound in (1) has been improved in [14], replacing $\Phi_1(1)$ by the spectral radius $\rho(C)$ of the Cartan matrix C , i.e. the maximum value $|\lambda|$ where λ runs through the set of complex eigenvalues of C .

However, for obvious reasons we are interested in a lower bound which only depends on terms of G . This is a weaker question than the one Brauer asked in

Received by the editors January 9, 2003 and, in revised form, October 29, 2003.
 2000 *Mathematics Subject Classification*. Primary 20C20, 20G40.

Problem 15 of his famous list [5]. He actually wanted to have a characterization of the dimension of the Jacobson radical by group-theoretical properties.

In the case of p -solvable groups (the existence of a p -complement suffices) we have $\Phi_1(1) = |G|_p$ (see [11], Chap. X, 3.2). Hence the desired bound is

$$(2) \quad |G|_{p'} \leq \sum_{\varphi \in \text{IBr}_p(G)} \varphi(1)^2.$$

But for non- p -solvable groups, $\Phi_1(1)$ (and $\rho(C)$) may differ extremely from $|G|_p$. For instance, let $G = R(q)$ with $q = 3^{2m+1}$ ($m \geq 1$) be a Ree group and let $p = 2$. Then $\Phi_1(1) = 2(q^3 + 1) \gg 8$ and $\rho(C) \approx 16.38$ (see [15] for the Cartan matrix), but $|G|_2 = 8$ and moreover

$$\frac{|G|}{8} = |G|_{2'} \leq \sum_{\varphi \in \text{IBr}_p(G)} \varphi(1)^2.$$

(In order to prove the inequality it is enough to consider the $\frac{q-3m}{6}$ characters of defect zero of degree $(q^2 - 1)(q + 1 + 3m)$, see [19].) Nevertheless for the sporadic groups that we have checked so far, the bound in (2) holds true. The same bound turns out to be true if the Sylow p -subgroups of G are cyclic [14]. So for any finite group G we are led to the

Conjecture. *We always have*

$$(3) \quad |G|_{p'} \leq \sum_{\varphi \in \text{IBr}_p(G)} \varphi(1)^2,$$

and equality holds if and only if G has a normal Sylow p -subgroup.

Note that the *if part* is trivial, since a normal p -subgroup is always contained in the kernel of an irreducible representation in characteristic p .

Suppose that the conjecture has an affirmative answer. Thus in the extreme case $|G|_{p'} = \sum_{\varphi \in \text{IBr}_p(G)} \varphi(1)^2$ the group G has a normal Sylow p -subgroup, say P . By Lemma 4.26 of ([11], Chapter IV), we know that

$$\Phi_\varphi(x) = |C_P(x)|\varphi(x)$$

for p' -elements $x \in G$. On the other hand, by Lemma 3.8 of ([11], Chapter IV), we have

$$\sum_{\varphi \in \text{IBr}_p(G)} \Phi_\varphi(x) \overline{\Phi_\varphi(y)} = \begin{cases} 0 & \text{if } x \not\sim y, \\ |C_G(x)| & \text{if } x \sim y, \end{cases}$$

for p' -elements $x, y \in G$ where \sim denotes conjugation in G . Thus we get the relation

$$\sum_{\varphi \in \text{IBr}_p(G)} \varphi(x) \overline{\varphi(y)} = \begin{cases} 0 & \text{if } x \not\sim y, \\ \frac{|C_G(x)|}{|C_P(x)|} & \text{if } x \sim y. \end{cases}$$

So we may ask the

Question. Is it even true that for p' -elements $x \in G$ we always have

$$(4) \quad \frac{|C_G(x)|}{|G|_p} \leq \sum_{\varphi \in \text{IBr}_p(G)} |\varphi(x)|^2 ?$$

If so, the bound may be sharp for $x \neq 1$ even if the Sylow p -subgroup is not normal. For instance, if $p = 5$ and x is an element of order 3 in $G = L_2(16)$, then

$$3 = \frac{15}{5} = \frac{|C_G(x)|}{|G|_5} = \sum_{\varphi \in \text{IBr}_5(G)} \varphi(x) \overline{\varphi(x)} = 1^2 + 1^2 + (-1)^2 = 3.$$

Note that the inequality (4) reduces to (3) for $x = 1$.

We are furthermore tempted to ask a p -local version of (3). In this case we may replace

$$|G|_{p'} = \frac{|G|}{p^a} = \frac{\dim KG}{p^a}$$

by $\frac{\dim B}{p^d}$ where K is a splitting field of G of characteristic p and B is a p -block of defect d . Thus we ask whether

$$(5) \quad \frac{\dim B}{p^d} \leq \sum_{\varphi \in \text{IBr}_p(B)} \varphi(1)^2?$$

An affirmative answer to (5) was given by Kiyota and Wada in [14] in case G is p -solvable or B is a p -block with cyclic defect group. In the latter case equality holds if and only if the Brauer tree is a star and all irreducible Brauer characters have the same degree.

Unfortunately the principal 2-block B_0 of the alternating group A_5 shows that inequality (5) fails to be true in general, because

$$11 = \frac{\dim B_0}{2^d} > \sum_{\varphi \in \text{IBr}_p(B)} \varphi(1)^2 = 1 + 2^2 + 2^2 = 9.$$

A more advanced counterexample is the non-principal 3-block of maximal defect of 6.A7. This example was brought to my attention by Thomas Breuer.

2. ON THE CONJECTURE

In this section we prove the conjecture for groups which have a projective character with properties similar to those of the Steinberg character for groups of Lie type. We start with a result due to Alperin. Let Φ be any projective character of a finite group G and let $\text{IBr}(G) = \{\varphi_1, \varphi_2, \dots, \varphi_s\}$. Then Φ uniquely determines a matrix $A = (a_{ij})$ by the equations

$$\varphi_i \Phi = \sum_{j=1}^s a_{ij} \Phi_j \quad (i = 1, \dots, s).$$

If x_1, \dots, x_s are representatives of the p' -conjugacy classes of G and if C denotes the Cartan matrix of G , then we have

Lemma 2.1 ([1]).

$$\det A = \frac{\prod_{i=1}^s \Phi(x_i)}{\det C}.$$

Proof. For $C = (c_{ij})$, we have

$$\varphi_i \Phi = \sum_{j,k=1}^s a_{ij} c_{jk} \varphi_k.$$

Therefore we get the matrix equation

$$(\varphi_i(x_l)) \begin{pmatrix} \Phi(x_1) & 0 & \cdots & 0 \\ 0 & \Phi(x_2) & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & \Phi(x_s) \end{pmatrix} = AC(\varphi_i(x_l)).$$

Since $(\varphi_i(x_l))$ and C are invertible ([11], Chap. IV, 3.6), the assertion follows.

Theorem 2.2. *Let G be a finite group. Suppose that G has a projective character Φ satisfying the following two conditions.*

- (a) $\Phi(1) = |G|_p$.
- (b) $\Phi(x) \neq 0$ for all p' -elements $x \in G$.

Then

$$|G|_{p'} \leq \sum_{i=1}^s \varphi_i(1)^2,$$

and equality holds if and only if G has normal Sylow p -subgroup.

Proof. Condition (b) implies that $\det A \neq 0$ in the above lemma. Thus there exists a permutation π of $\{1, \dots, s\}$ such that

$$\prod_{i=1}^s a_{\pi i, i} \neq 0.$$

This means $\varphi_{\pi i} \Phi = \Phi_i + \cdots$. If ρ denotes the regular character of G , then

$$\sum_{i=1}^s \varphi_i(1) \varphi_{\pi i} \Phi = \sum_{i=1}^s \varphi_i(1) \Phi_i + \cdots = \rho + \cdots.$$

In particular we have

$$(6) \quad \sum_{i=1}^s \varphi_i(1) \varphi_{\pi i}(1) \Phi(1) \geq \rho(1) = |G|,$$

and by condition (a)

$$(7) \quad \sum_{i=1}^s \varphi_i(1) \varphi_{\pi i}(1) \geq |G|_{p'}.$$

Applying the Cauchy-Schwarz inequality yields

$$(8) \quad \left(\sum_{i=1}^s \varphi_i(1) \varphi_{\pi i}(1) \right)^2 \leq \sum_{i=1}^s \varphi_i(1)^2 \sum_{i=1}^s \varphi_{\pi i}(1)^2,$$

and hence

$$(9) \quad \sum_{i=1}^s \varphi_i(1) \varphi_{\pi i}(1) \leq \sum_{i=1}^s \varphi_i(1)^2.$$

Putting (7) and (9) together, we obtain the desired assertion

$$|G|_{p'} \leq \sum_{i=1}^s \varphi_i(1)^2.$$

So it remains to prove that $|G|_{p'} = \sum_{i=1}^s \varphi_i(1)^2$ implies that a Sylow p -subgroup of G is normal.

By (6) we get $\varphi_{\pi i}\Phi = \Phi_i$ for all i . In particular we have $\varphi_{\pi 1}\Phi = \Phi_1$ for the trivial character φ_1 . The equality in the Cauchy-Schwarz inequality (see (8)) forces $\varphi_{\pi 1}(1) = \varphi_1(1) = 1$. Thus we obtain $\Phi_1(1) = |G|_p$ and by a result of Brockhaus [7] the group G has a normal Sylow p -subgroup.

The properties we required on the projective character in 2.2 seem to be very strong. However there are interesting examples for which the theorem applies.

Examples 2.3. a) Let G be a finite group with a split BN -pair of characteristic p and satisfying the commutator relations (see [9]). For Φ we take the Steinberg character. Then conditions a) and b) are satisfied, since

$$\Phi(x) = |C_G(x)|_p$$

for all p' -elements x of G (see [9], 6.4.7). Note that for Chevalley groups (twisted or non-twisted) defined in characteristic p , the square of the degree of the Steinberg character already dominates the p' -part of the order of G .

b) Let G be a finite group with a p -complement. Now we may take for Φ the projective indecomposable character corresponding to the trivial character. Observe that Φ is the trivial character of a p -complement induced to G and satisfies the assumptions of 2.2.

c) More subtle is the following. Let N be a normal p -solvable subgroup of G , and suppose that $H = G/N$ possess a projective character satisfying conditions a) and b). We prove that G has a projective character with the same properties. So let $M \leq N$ be a minimal normal subgroup of G . By the inductive hypothesis the group G/M has a projective character, say Ψ , satisfying a) and b). If M is a p' -group, we may take for Φ the inflation $\Phi = \text{infl}_G(\Psi)$ of Ψ . Thus suppose that M is a p -group. Let λ be the Brauer character of the conjugation action of G on M . Now we put

$$\Phi = \lambda \text{infl}_G(\Psi).$$

That Φ is projective follows from a result of Alperin, Collins and Sibley (see [2]). Clearly,

$$\Phi(1) = \lambda(1) \text{infl}_G(\Psi)(1) = |M||H| = |G|_p$$

and

$$\Phi(x) = \lambda(x) \text{infl}_G(\Psi)(x) = |C_M(x)| \text{infl}_G(\Psi)(x) \neq 0$$

for all p' -elements $x \in G$.

3. ASYMPTOTIC RESULTS FOR GROUPS OF LIE TYPE

To give more evidence for inequality (3) we will look in this section asymptotically on groups of Lie type in non-defining characteristics. For a detailed introduction to the character theory of groups of Lie type, the reader is referred to [9].

Let G be a simple linear algebraic group over an algebraically closed field K of positive characteristic $r \neq p$, and let $G^F = G(q)$ (where $q = r^f$) denote the group of fixed points under a Frobenius map F . Furthermore let $T_w = T^F$ be the maximal torus in G^F corresponding to $w \in W = W(T)$ where W denotes the Weyl group of an F -stable maximal torus $T \subseteq G$.

Now we look at irreducible Deligne-Lusztig characters $R_{T,\theta}$ (up to a sign) where $\theta \in \hat{T}^F = \text{Hom}(T^F, \mathbb{C}^*)$ is an irreducible character of $T_w = T^F$ and θ is in general

position. Let $\mathcal{R}(q)$ denote the set of all such $R_{T,\theta}$'s. Note that $R_{T,\theta} = R_{T,\theta'}$ if and only if $\theta = {}^{w'}\theta'$ for some

$$w' \in W(T)^F = C_{w,F} = \{w' \in W \mid w'^{-1}wF(w') = w\}$$

(see [9], p. 219). Thus the number $|\mathcal{R}(q)|$ of different $R_{T,\theta}$'s is equal to the number of $W(T)^F$ -orbits of characters θ in general position. By a result of Veldkamp ([17]) this number is equal to the number of $W(T)^F$ -conjugacy classes of elements in general position in the fixed point group T^{*F^*} of the dual torus $T^* \subseteq G^*$.

For simplicity let us suppose that G^* is simply connected. In this case the regular elements in T^{*F^*} coincide with the elements in general position. (Note that $t \in G$ is regular (resp. in general position) if $C_G(t)^\circ = T$ (resp. $C_G(t) = T$).) Let l denote the Lie rank of G . Applying 3.1 and 3.2 in [10] (see also [12]), we get

$$|\mathcal{R}(q)| = \{R_{T,\theta} \mid \theta \text{ in general position}\} = \frac{q^l}{|C_{w,F}|} + O(q^{l-1}).$$

Note that $R_{T,\theta}$ is of degree

$$R_{T,\theta}(1) = \frac{|G(q)|_{r'}}{|T_w|}.$$

Thus we obtain

$$(10) \quad \sum_{R_{T,\theta} \in \mathcal{R}(q)} R_{T,\theta}(1)^2 = \left(\frac{q^l}{|C_{w,F}|} + O(q^{l-1}) \right) \frac{|G(q)|_{r'}^2}{|T_w|^2}.$$

Let N denote the number of positive roots. As polynomials in q we have

- $|T_w| = q^l + \text{lower terms in } q$,
- $|G(q)|_{r'} = q^{l+N} + \text{lower terms in } q$,
- $|G(q)| = q^{l+2N} + \text{lower terms in } q$.

Inserting these equations in (10), we obtain

$$(11) \quad \begin{aligned} \sum_{R_{T,\theta} \in \mathcal{R}(q)} R_{T,\theta}(1)^2 &= \frac{1}{|C_{w,F}|} q^{l+2N} + O(q^{l+2N-1}) \\ &= |G(q)|_{p'} + \left(\frac{1}{|C_{w,F}|} - \frac{1}{|G(q)|_p} \right) q^{l+2N} + O(q^{l+2N-1}). \end{aligned}$$

We may assume that $p \mid |G(q)|$. If we exclude the Steinberg triality ${}^3D_4(q^2)$ for a moment, then

$$|G(q)|_{r'} = \prod_{i=1}^l (q^{d_i} - \epsilon_i)$$

where $\epsilon_i \in \{1, -1\}$ and the d_i are the exponents of the underlying group (see [9], p. 75). Thus $p \mid q^{2d_i} - 1$. Note that

$$(q^{2d_i} - 1)_p < (q^{2pd_i} - 1)_p$$

and $|W|$ is independent of q (see [3], Planche II). Thus there exists an $s \in \mathbb{N}$ (for instance s a suitable power of p) such that

$$|C_{w,F}| \leq |W| < |G(q^s)|_p,$$

and therefore

$$|C_{w,F}| < |G(q^{st})|_p$$

for all $t \in \mathbb{N}$ and all $w \in W$. This is also true for ${}^3D_4(q^3)$, as a similar argument shows. By (11) we obtain

$$\sum_{R_{T,\theta} \in \mathcal{R}(q^{st})} R_{T,\theta}(1)^2 \geq |G(q^{st})|_{p'}$$

for all $t \geq t_0$. Now, if there exists a torus $T_w = T_w(q^{st}) \subseteq G(q^{st})$ with $p \nmid |T_w|$ for $t \geq t_0$, then all characters $R_{T,\theta}$ with θ in general position are irreducible and of p -defect zero. In particular they are irreducible Brauer characters for the prime p and we have the desired inequality

$$\sum_{\varphi \in \text{IBr}_p(G(q^{st}))} \varphi(1)^2 \geq |G(q^{st})|_{p'}$$

for all $t \geq t_0$. Thus we have proved

Theorem 3.1. *Let $G(q) = G^F$ be a finite group of Lie type where G is a linear simple algebraic group and G^* is simply connected. Let p be a prime with $p \nmid q$ and $p \mid |G(q)|$.*

a) *Then there exists an $s \in \mathbb{N}$ such that*

$$\sum_{R_{T,\theta} \in \mathcal{R}(q^{st})} R_{T,\theta}(1)^2 \geq |G(q^{st})|_{p'}$$

for all $t \geq t_0$.

b) *If $T_w = T_w(q^{st})$ is a torus in $G(q^{st})$ for some $w \in W$ and $p \nmid |T_w|$ for all $t \geq t_0$, then*

$$\sum_{\varphi \in \text{IBr}_p(G(q^{st}))} \varphi(1)^2 \geq |G(q^{st})|_{p'}$$

for all $t \geq t_0$.

The critical point in the whole process is the assumption $p \nmid |T_w| = |T_w(q^{st})|$ for a suitable $w \in W$ and all $t \geq t_0$. The author would like to thank an anonymous referee for pointing out the following argument.

If we replace s by $s(p-1)$, then we have $q^s \equiv 1 \pmod{p}$. Note that the order of $T_w(q)$ can be written as

$$|T_w(q)| = \Phi_{n_1}(q) \cdots \Phi_{n_r}(q)$$

where Φ_n denotes the n -th cyclotomic polynomial. Now we choose $w \in W$ in such a way that $1 < n_j$ and $p \nmid n_j$ for all $j = 1, \dots, r$. Since

$$\begin{aligned} \Phi_{n_j}(q^{st}) \mid \frac{q^{stn_j}-1}{q^{st}-1} &= 1 + q^{st} + \dots + q^{st(n_j-1)} \\ &\equiv 1 + 1 + \dots + 1 \pmod{p} \\ &\equiv n_j \pmod{p}, \end{aligned}$$

we get $p \nmid \Phi_{n_j}(q^{st})$ for all t ; hence $p \nmid |T_w(q^{st})|$ for all t .

The following example may illustrate the above. In particular, in the case where $G(q) = \text{PGL}(2, q)$ we are not able to find a torus as required for the prime $p = 2$. Thus our method fails for $p = 2$.

Example. Let $G^F = G(q) = \text{PGL}(l+1, q)$ be the finite adjoint group of type A_l for $l \geq 2$ with q a power of r . Note that $G^{*F^*} = \text{SL}(l+1, q)$ is the fixed point group of a simply connected group. Let $w \in W$ be a Coxeter element, i.e. w is a cycle of length $l+1$, and $C_{w,F} = C_{S_{l+1}}(w) = \langle w \rangle$ where S_n is the symmetric group on

n letters. Let p be a prime with $p > l + 1$. Thus we either have $p \nmid |G(q)|$ and (3) holds with equality, or $|C_{w,F}| < |G(q)|_p$ and we have $\sum_{R_{T,\theta} \in \mathcal{R}} R_{T,\theta}(1)^2 \geq |G(q)|_{p'}$ for $q \geq q_0$ and $p \mid |G(q)|$. Let such a q be given. Now if $p \nmid |T_w| = 1 + q + \dots + q^l$, then

$$\sum_{\varphi \in \text{IBr}_p(G(q))} \varphi(1)^2 \geq |G(q)|_{p'}.$$

The condition $p \nmid |T_w|$ is satisfied for instance if $p \nmid q - 1$ and $p \mid q^s - 1$ where $2 < s < l + 1$ and $\gcd(s, l + 1) = 1$.

Let us specialize to $G(q) = \text{PGL}(2, q)$ and let $2 < p \neq r$, hence $2 = |W| < p$. Since $|G(q)| = q(q+1)(q-1)$, we have $p \nmid |T_w| = (q+1)$ or $p \nmid |T_1| = q-1$ where T_w is the Coxeter and T_1 the split torus. Thus for all q large enough and all odd primes inequality (3) holds true for $\text{PGL}(2, q)$. However in this small case we get (3) for any q . If $p = r$ we may apply 2.3 a). In case $2 \neq p \neq r$ the Sylow p -subgroups of $G(q)$ are cyclic. Thus the inequality (5) holds true for every p -block by [14], which obviously implies (3). For $p = 2$ one easily checks inequality (3) using the results of Section VIII in [8].

ADDED IN PROOF

There is an analogue of the Conjecture for p' -class lengths, namely $|G|_{p'} \leq \sum_x |G : C_G(x)|$ where the sum runs through a set of representatives of the p' -conjugacy classes, and with equality if and only if G has a normal p -complement. In contrast to characters this can easily be proved using the Frobenius Conjecture which we know to be true. We shall discuss this and similar questions in a forthcoming paper.

REFERENCES

- [1] J. ALPERIN. Projective modules and tensor products. *J. Pure and Appl. Algebra* 8 (1976), 235-241. MR53:5712
- [2] J.L. ALPERIN, M.J. COLLINS AND D.A. SIBLEY. Projective modules, filtrations and Cartan invariants. *Bull. London Math. Soc.* (3) 16 (1984), 416-420. MR85m:20009
- [3] N. BOURBAKI. *Groupes et algèbre de Lie. Chapitres 4,5 et 6.* Masson, Paris 1981. MR83g:17001
- [4] R. BRAUER. Notes on representations of finite groups. *J. London Math. Soc.* (2), 13 (1976), 162-166. MR53:3091
- [5] R. BRAUER. Representations of finite groups. *Lectures on Modern Mathematics*, Vol. 1, Wiley, New York, 1963, 133-175. MR31:2314
- [6] R. BRAUER AND C.J. NESBITT. On the modular characters of groups, *Ann. Math.*(2) 42 (1941), 556-590. MR2:309c
- [7] P. BROCKHAUS. On the radical of a group algebra. *J. Algebra* 95 (1985), 454-472. MR87h:20021
- [8] R. BURKHARDT. Die Zerlegungsmatrizen der Gruppen $\text{PSL}(2, p^f)$. *J. Algebra* 40 (1976), 75-96. MR58:864
- [9] R.W. CARTER. *Finite groups of Lie type, conjugacy classes and complex characters.* Wiley, Chichester 1985. MR87d:20060
- [10] D.I. DERIZIOTIS. On the number of conjugacy classes in finite groups of Lie type. *Comm. Algebra* 13 (1985), 1019-1045. MR86i:20067
- [11] W. FEIT. *The representation theory of finite groups.* North Holland, Amsterdam 1982. MR83g:20001
- [12] P. FLEISCHMANN AND I. JANISZCZAK. The number of regular semisimple elements for Chevalley groups of classical type. *J. Algebra* 155 (1993), 482-528. MR94f:20090
- [13] B. HUPPERT AND N. BLACKBURN. *Finite groups II.* Springer-Verlag, Berlin/Heidelberg/New York 1982. MR84i:20001a

- [14] M. KIYOTA AND T. WADA. Some remarks on the eigenvalues of the Cartan matrix in finite groups. *Commun. Algebra* 21, No.11 (1993), 3839-3860. MR94i:20021
- [15] P. LANDROCK AND G.O. MICHLER. Principal 2-blocks of simple groups of Lie type. *Trans. Am. Math. Soc.* 260 (1980), 83-111. MR81h:20013
- [16] G. NAVARRO. Characters and blocks of finite groups. *London Math. Soc., Lecture Notes Series* 250, Cambridge Univ. Press, Cambridge, 1998. MR2000a:20018
- [17] F.D. VELDKAMP. Regular characters and regular elements. *Comm. Algebra* 5(12) (1977), 1259-1273. MR58:16857
- [18] D.A.R. WALLACE. On the radical of a group algebra, *Proc. Am. Math. Soc.* 12 (1961), 133-137. MR22:12146
- [19] H.N. WARD. On Ree's series of simple groups. *Trans. Amer. Math. Soc.* 121 (1966), 62-89. MR33:5752

INSTITUT FÜR ALGEBRA UND GEOMETRIE, FAKULTÄT FÜR MATHEMATIK, OTTO-VON-GUERICKE-UNIVERSITÄT, 39016 MAGDEBURG, GERMANY