

ON THE EXTENSION OF A THEOREM OF POINCARÉ FOR DIFFERENCE-EQUATIONS*

BY

EDWARD B. VAN VLECK

One of the most useful theorems relating to difference equations is the well-known theorem of POINCARÉ† for the equation

$$(1) \quad u(n+r) + P_1(n)u(n+r-1) + \cdots + P_r(n)u(n) = 0,$$

in which the coefficients $P_i(n)$ possess limits A_i for $n = +\infty$. POINCARÉ showed, namely, that if the roots of the equation

$$(2) \quad y^r + A_1 y^{r-1} + \cdots + A_r = 0$$

are no two of equal modulus, the ratio $u(n)/u(n-1)$ has a limit for $n = +\infty$, and this limit is a root of (2), in general the root of greatest modulus.‡

* Presented to the Society at the Chicago meeting, April 29, 1911.

† *American Journal of Mathematics*, vol. 7 (1885), p. 213. For careful analysis and further development of the theorem, see PERRON, *Crelle's Journal*, vol. 136 (1909), p. 17 and vol. 137, p. 6.

‡ In 1904, in an article on continued fractions in these *Transactions*, p. 253, I gave a proof of POINCARÉ's theorem for $r = 2$ which was suitable for application to the class of continued fractions there under consideration. This proof, and consequently its immediate applicability, was not fully understood by PRINGSHEIM and called forth an interesting memoir from him (*Sitzungsberichte der Bayerischen Akademie, math.-physikalischer Klasse*, 1910, p. 3) devoted in large part to rederiving my results. Later after oral and written communication between us he has recognized the legitimacy of my proof (*Ibid.*, 1911, p. 64). My argument was somewhat condensed, but not more so, it seemed to me, than is current in articles of an investigational character. A few additional lines, which it is worth while to add now, would have removed any difficulty.

When Pringsheim reaches the following theorem (p. 46), he inadvertently fails to attribute it to me, though the attribution is intended in the introduction:

If in the continued fraction

$$A(z) = \frac{1}{1 + \frac{a_2 z}{1} + \frac{a_3 z}{1} + \cdots}$$

we have $\lim a_n = k$ for $n = \infty$, the continued fraction will converge over the entire plane except (1) along the whole or a part of a rectilinear cut drawn from $z = -\frac{1}{k}$ to $z = \infty$ with an argument equal to that of the vector from the origin to $z = -\frac{1}{k}$, and except possibly (2) at certain isolated points p_1, p_2, \dots . Within the plane so cut the continued fraction is holomorphic except at the points p_1, p_2, p_3, \dots , which are poles.

* See, for example, pages 72–73 of HORN'S *Gewöhnliche Differentialgleichungen beliebiger Ordnung*.

definite combinations,

$$(11) \quad v_i(n) = \sum_{j=1}^r c_{ji}(n) y_j(n-1) \quad (i = 1, \dots, r),$$

which possess the following property: When transformed by the linear substitution (A), they undergo the multiplicative transformation

$$\sum_{j=1}^r c_{ji}(n) y_j(n) = \rho_i(n) \sum_{j=1}^r c_{ji}(n) y_j(n-1).$$

For the limiting form of (11) we have

$$(12) \quad V_i(n) = \sum_{j=1}^r C_{ji} Y_j(n-1) \quad (i = 1, \dots, r).$$

In the theory of linear substitutions* it is shown that the determinant $|c_{ji}(n)|$ can not vanish when the roots of (6) are distinct. We may therefore solve (11) for the $y_j(n-1)$ in terms of the $v_i(n)$. Similarly the determinant $|C_{ji}| \neq 0$, and equations (12) may be solved for the $Y_j(n-1)$. The solutions will be denoted by

$$(13) \quad y_i(n-1) = \sum_{j=1}^r d_{ji}(n) v_j(n) \quad (i = 1, \dots, r),$$

$$(14) \quad Y_i(n-1) = \sum_{j=1}^r D_{ji} V_j(n),$$

in which D_{ji} is obviously the limit of $d_{ji}(n)$ for $n = \infty$.

The ratios of the D_{ji} can be most easily determined by considering the linear system of difference equations with constant coefficients

$$(15) \quad Y_i(n) = \sum_{j=1}^r A_{ij} Y_j(n-1) \quad (i = 1, \dots, r),$$

to which the limiting relations (12) and (14) may be attached. For this system we have

$$V_i(n+1) = R_i V_i(n).$$

Hence

$$(16) \quad Y_i(n) = \sum_{j=1}^r D_{ji} V_j(n+1) = \sum_{j=1}^r D_{ji} R_j V_j(n).$$

If the expressions for $Y_i(n-1)$ and $Y_i(n)$ which are given in (14) and (16) are substituted in (15), and the coefficients of each $V_j(n)$ on the two sides of the equation are then equated, we obtain the following r sets of relations

* Cf. HORN, loc. cit., p. 73.

and consequently for all succeeding values. *This can happen only if the determinant of the coefficients $a_{ij}(n)$ vanishes for at least one value of n .* We shall assume in §2 that at least one of the $v_i(n)$ does not vanish for an infinity of values of n .

§2. Extension of Poincaré's theorem.

It will be supposed henceforth that

$$(19) \quad |R_1| > |R_2| > \cdots > |R_r|.$$

When n increases indefinitely, some one of the r moduli $|v_i(n)|$ will be as large or larger than the remaining $r - 1$ for an infinite number of values of n . I will show that this $|v_i(n)|$ will become and remain dominant over all the other $r - 1$ moduli.

Suppose first that for an infinite set of values of n we have

$$(20) \quad |v_1(n)| \geq |v_i(n)| \quad (i = 2, 3, \dots, r),$$

and denote this set by $\{n'\}$. The division of the first of equations (B) by $v_1(n)$ and of the succeeding equations by $v_1(n+1)$ gives

$$(21) \quad \frac{v_1(n+1)}{v_1(n)} = R_1 + e_1(n)$$

and

$$(22) \quad \frac{v_i(n+1)}{v_1(n+1)} = \frac{R_i}{R_1 + e_1(n)} \cdot \frac{v_i(n)}{v_1(n)} + \frac{e_i(n)}{R_1 + e_1(n)} \quad (i = 2, \dots, r),$$

in which the $e(n)$ satisfy for $n = n'$ the inequalities

$$(23) \quad |e_1(n')| \leq \bar{r}\bar{e}(n'), \quad |e_i(n')| \leq \bar{r}\bar{e}(n'),$$

where $\bar{e}(n')$ denotes the largest of the $|e_{ij}(n')|$ in (B). Since $\lim \bar{e}(n') = 0$ for $n' = \infty$, it follows that for a sufficiently large value of n' we have

$$\frac{|v_i(n'+1)|}{|v_1(n'+1)|} \leq \frac{|R_i|}{|R_1|} + a,$$

where a is an arbitrarily prescribed positive quantity subject to the condition $|R_i|/|R_1| + a < 1$. Consequently $n' + 1$ will be included with n' in the set $\{n'\}$, provided only n' is taken sufficiently large. It follows that equations (20) to (23) will hold without exception from and after some sufficiently large value of n .

From (21) we now obtain

$$(24) \quad \lim_{n \rightarrow \infty} \frac{v_1(n+1)}{v_1(n)} = R_1.$$

since $\lim \bar{e}(n) = 0$. It may next be shown that

$$(25) \quad \lim_{n \rightarrow \infty} \frac{v_i(n)}{v_1(n)} = 0. \quad (i = 2, 3, \dots, r).$$

For if this is not true, there will be at least some one value of i and some fixed magnitude $M > 0$ such that for an infinity of values of n one has

$$\frac{|v_i(n)|}{|v_1(n)|} \geq M.$$

Call this set of values $\{n''\}$. Then since $\lim e_1(n) = \lim e_i(n) = 0$, equation (22) for $n = n''$ may be written

$$\frac{v_i(n'' + 1)}{v_1(n'' + 1)} = \left(\frac{R_i}{R_1} + e'_i(n'') \right) \frac{v_i(n'')}{v_1(n'')},$$

where $\lim e'_i(n'') = 0$ for $n'' = \infty$. Since $|R_i| < |R_1|$, this shows that when n'' is sufficiently large, the ratio $|v_i(n'' + j)| / |v_1(n'' + j)|$ for $j = 1, 2, \dots$ will decrease in approximately the ratio $|R_i| : |R_1|$ until a value of j is reached for which the ratio falls below M . Thus if $|v_i(n)| / |v_1(n)|$ is equal to or greater than M for an infinity of values of n , there must also be an infinity of such values for which it falls below M . But when it falls below M , equation (22) gives

$$\frac{|v_i(n + 1)|}{|v_1(n + 1)|} < \frac{|R_i|}{|R_1 + e_1(n)|} M + \frac{|e_i(n)|}{|R_1 + e_1(n)|}.$$

Since now $\lim e_1(n) = \lim e_i(n) = 0$ for $n = \infty$, n may be chosen so large in the last inequality that

$$\frac{|v_i(n + 1)|}{|v_1(n + 1)|} < M.$$

Thus if $|v_i(n)| / |v_1(n)| < M$ for a sufficiently large value of n , it must remain so for all larger values of n . But this contradicts the hypothesis that it exceeds M for an infinity of values of n . Consequently (25) must hold, as was to be proved.

We shall next establish like results in case the $r - 1$ inequalities

$$(26) \quad |v_2(n)| \geq |v_i(n)| \quad (i = 2).$$

hold simultaneously for an infinity of values of n . Denote this particular set of values by $\{n'\}$. Then on dividing equations (B) for $i = 2$ by $v_2(n)$ and for $i \neq 2$ by $v_2(n + 1)$ we obtain the following equations:

$$(27) \quad \frac{v_2(n + 1)}{v_2(n)} = R_2 + e_2(n),$$

$$(28) \quad \frac{v_1(n+1)}{v_2(n+1)} = \frac{R_1}{R_2 + e_2(n)} \cdot \frac{v_1(n)}{v_2(n)} + \frac{e_1(n)}{R_2 + e_2(n)},$$

$$(29) \quad \frac{v_i(n+1)}{v_2(n+1)} = \frac{R_i}{R_2 + e_2(n)} \cdot \frac{v_i(n)}{v_2(n)} + \frac{e_i(n)}{R_2 + e_2(n)} \quad (i = 3, 4, \dots, r),$$

in which the $e(n)$ for $n = n'$ satisfy the inequalities

$$(30) \quad |e_i(n')| \leq r\bar{e}(n') \quad (i = 1, 2, \dots, r),$$

where $\bar{e}(n')$ denotes the largest of the $|e_{ij}(n')|$ in (B).

Equations (29) show that for a sufficiently large value of n' we have

$$(31) \quad |v_i(n'+1)| < |v_2(n'+1)| \quad (i = 3, 4, \dots, r)$$

since

$$|R_i| < |R_2|, \quad |v_i(n')| \leq |v_2(n')|, \quad \lim_{n' \rightarrow \infty} \bar{e}(n') = 0.$$

Furthermore, we must also have for a sufficiently large value of n'

$$(32) \quad |v_1(n'+1)| < |v_2(n'+1)|,$$

for otherwise the inequality

$$|v_1(n'+1)| \geq |v_i(n'+1)| \quad (i = 1, 3, \dots, r)$$

would hold for an infinity of values of n' . But the latter is impossible, since it has been shown that $|v_1(n)|$ would then ultimately become and remain dominant over all the other $|v_i(n)|$, which contradicts the hypothesis that (26) is to hold for an infinity of values of n . Now (31) and (32) together show that $n' + 1$ will be included simultaneously with n' in the set $\{n'\}$, provided only n' is sufficiently large. It follows that (26)–(30) will all hold without exception from and after some fixed but large enough value of n .

From (27) we see next that

$$(33) \quad \lim_{n \rightarrow \infty} \frac{v_2(n+1)}{v_2(n)} = R_2.$$

From (29) it follows also that

$$(34) \quad \lim_{n \rightarrow \infty} \frac{v_i(n+1)}{v_2(n+1)} = 0 \quad (i = 3, \dots, r),$$

the reasoning by which this is established being exactly parallel to that by which (25) was established on our former hypothesis. Lastly, it remains to be proved that

$$(35) \quad \lim_{n \rightarrow \infty} \frac{v_1(n)}{v_2(n)} = 0.$$

Suppose, if possible, this not to be true. Then there must be some magnitude

$M > 0$ which $|v_1(n)|/|v_2(n)|$ will exceed for an infinity of values of n . Denote this set of values by $\{n''\}$. For these values (28) may be written

$$\frac{v_1(n''+1)}{v_2(n''+1)} = \left(\frac{R_1}{R_2} + e'(n'') \right) \frac{v_1(n'')}{v_2(n'')},$$

where $\lim e'(n'') = 0$ for $n'' = \infty$. Since $|R_1| > |R_2|$, this shows that ultimately $n''+1$ will be included with n'' in the set $\{n''\}$. Then as i increases through the natural number system, the ratio $|v_1(n''+i)|/|v_2(n''+i)|$ will increase successively by factors approximating closer and closer to $|R_1|/|R_2|$, and hence finally we will have $|v_1(n''+i)| > |v_2(n''+i)|$ from and after some fixed value of i . But this contradicts our hypothesis (26). Hence (35) must hold, as stated.

In similar manner we may treat successively for $j = 3, 4, \dots, r$ the hypothesis that the $r-1$ inequalities

$$|v_j(n)| \geq |v_i(n)| \quad (i \neq j)$$

hold for an infinity of values of n . On dividing (B) by $v_j(n)$ and $v_j(n+1)$ we obtain three sets of inequalities similar to (27), (28), (29), which furnish like results. We reach thus the following conclusion:

THEOREM 1. *If in the system of differential equations (B) we have*

$$\lim_{n=\infty} e_{ij}(n) = 0 \quad (i, j = 1, 2, \dots, r)$$

and we suppose

$$|R_1| > |R_2| > \dots > |R_r|,$$

then as n increases indefinitely, either by exception the r quantities $v_i(n)$ will all vanish from and after some fixed value of n , or some one of them will ultimately dominate over the other $r-1$. If this be $v_k(n)$, then

$$\lim_{n=\infty} \frac{v_{k+1}(n)}{v_k(n)} = R_k$$

and

$$\lim_{n=\infty} \frac{v_i(n)}{v_k(n)} = 0 \quad (i \neq k).$$

We return now to our initial equations (A). We have seen that in the relations

$$(13) \quad y_i(n-1) = \sum_{j=1}^r d_{ji}(n) v_j(n) \quad (i = 1, \dots, r)$$

the coefficients tend toward limiting values D_{ji} . The determinant $|D_{ji}|$ can not vanish since it is the reciprocal of $|C_{ji}| \neq 0$. Consequently for each value of j not all of the D_{ji} ($i = 1, \dots, r$) can vanish. Consider now

the equation

$$(36) \quad \frac{y_i(n)}{y_i(n-1)} = \frac{\sum_{j=1}^r d_{ji}(n+1) v_j(n+1)}{\sum_{j=1}^r d_{ji}(n) v_j(n)}.$$

If, as in Theorem 1, $v_k(n)$ denotes that one of the $v_i(n)$ which ultimately becomes dominant, and if D_{kl} denotes one of the D_{kj} ($j = 1, \dots, r$) which does not vanish, it follows by application of the theorem to (36) that

$$\lim_{n=\infty} \frac{y_l(n-1)}{y_l(n)} = \lim \frac{d_{kl}(n+1) v_k(n+1)}{d_{kl}(n) v_k(n)} = \lim \frac{v_k(n+1)}{v_k(n)} = R_k.$$

Also we have

$$\lim_{n=\infty} \frac{y_i(n)}{y_l(n)} = \lim \frac{d_{ki}(n+1) v_k(n+1)}{d_{kl}(n+1) v_k(n+1)} = \frac{D_{ki}}{D_{kl}}. \quad (i \neq l).$$

This shows that the point $y_i(n)$ ($i = 1, \dots, r$) for $n = \infty$ approaches as its limit one of the invariant points of (15). This result may be restated as follows:

THEOREM 2. *Let (A) be a system of difference equations in which the coefficients $a_{ij}(n)$ have limiting values A_{ij} for $n = \infty$. If the roots R_i of the limiting characteristic equation (7) are no two of equal modulus, then unless by exception the $y_i(n)$ all vanish from and after some fixed value of n , there is at least one function $y_l(n)$ in the solution $y_j(n)$, ($j = 1, \dots, r$), for which we have*

$$\lim_{n=\infty} \frac{y_l(n)}{y_l(n-1)} = R_k,$$

where R_k is one of the roots of (7).

If also (A) is a sequence of non-singular collineations in r homogeneous variables, which have for $n = \infty$ a limiting form, then any initial point $y_1(0), \dots, y_r(0)$ by the successive application of the collineations will approach as a limit one of the invariant points of the limiting collineation (15).

The theorem of POINCARÉ is an immediate consequence of the first part of this theorem, for by the very construction of (3) we have

$$\lim_{n=\infty} \frac{u_i(n)}{u_i(n+1)} = \lim_{n=\infty} \frac{u_{i-1}(n)}{u_{i-1}(n-1)} \quad (i = 2, \dots, r).$$

To guard against misconception I will add in conclusion that not all of the r ratios $y_i(n)/y_i(n-1)$ need have limits for $n = \infty$, and even if they have, not all of the limits need to be roots of (7). The truth of the latter

statement will be obvious from a consideration of the special system,

$$v_1(n) = R_1 v_1(n-1),$$

$$v_2(n) = e(n) v_1(n-1) + R_2 v_2(n-1)$$

in which $e(n)$ is an infinitesimal having the value

$$(S_1 - R_2) \frac{\bar{v}_2(n-1)}{v_1(n-1)} \quad (0 < R_2 < S_1 < R_1).$$

Here the limit of $v_2(n)/v_1(n-1)$ for the particular integral $v_i(n) = \bar{v}_i(n)$ is an arbitrarily chosen constant S_1 .

UNIVERSITY OF WISCONSIN.
