

ON MATRICES WHOSE COEFFICIENTS ARE FUNCTIONS OF A SINGLE VARIABLE

BY

J. H. M. WEDDERBURN

1. The methods usually* employed in reducing to its normal form a matrix whose coefficients are polynomials in a variable λ are of such a nature that it is not at all obvious how they can be extended when polynomials are replaced by analytic functions.† The object of this note is to show that the main theorems on elementary factors are not restricted to matrix polynomials but apply without appreciable modification to analytic matrix functions.

As vectors are employed freely in the sequel, a short explanation of the notation used is necessary. A *vector*, $x = (\xi_1, \xi_2, \dots, \xi_n)$, is an ordered set of n coefficients: two vectors are equal if, and only if, their corresponding coefficients are equal. The *sum* of two vectors, $x = (\xi_1, \xi_2, \dots, \xi_n)$ and $y = (\eta_1, \eta_2, \dots, \eta_n)$ is defined as

$$x + y = (\xi_1 + \eta_1, \xi_2 + \eta_2, \dots, \xi_n + \eta_n).$$

The *product* of any number ρ into a vector x is defined as

$$\rho x = (\rho \xi_1, \rho \xi_2, \dots, \rho \xi_n).$$

Except in this elementary case, multiplication of vectors will not be used. A set of n linearly independent vectors is called a *basis*. Any vector can be expressed linearly in terms of the elements of a basis and, in particular, if e_i is the vector for which $\xi_i = 1$, $\xi_j = 0$ ($j \neq i$), we have $x = \sum_{i=1}^n \xi_i e_i$.

If A is a matrix (a_{rs}) and $x = \sum \xi_i e_i$, the vector $\sum_{i=1}^n (\sum_{j=1}^n a_{ij} \xi_j) e_i$ is denoted by Ax . We evidently have

$$A(x_1 + x_2) = Ax_1 + Ax_2, \quad (A + B)x = Ax + Bx, \quad A \cdot Bx = (AB)x.$$

A matrix is completely determined when its action on n linearly independent vectors is given. If the determinant of a matrix is not zero, it transforms a set of linearly independent vectors into a linearly independent set; while if its rank is r ($r < n$), there is a unique complex of vectors every element of

* See for instance Bôcher, *Introduction to Higher Algebra*, p. 262.

† To avoid frequent repetition, a matrix whose coefficients are functions of a variable, λ , will be called a *matrix function* of λ .

which is annihilated by the matrix. The following lemma is an immediate consequence of this.

LEMMA I. *If x_1, x_2, \dots, x_n is a set of vector functions which are holomorphic and linearly independent for all values of λ lying in a given region R , the determinant of the matrix defined by*

$$x_r = P(\lambda) e_r \quad (r = 1, 2, \dots, n)$$

does not vanish for any value of λ in R .

It may be remarked that the matrix $P(\lambda)$ is holomorphic in R and, since $|P(\lambda)|^*$ does not vanish, $P^{-1}(\lambda)$ is also holomorphic in that region. Also, since

$$e_r = P^{-1}(\lambda) x_r, \quad (r = 1, 2, \dots, n),$$

the fixed basis e_1, e_2, \dots, e_n may be replaced by any set of vector functions which are holomorphic and linearly independent throughout R .

2. In the reduction of a matrix to its normal form, we require the following extension of a well-known algebraic theorem.

LEMMA II. *If $f(\lambda)$ and $g(\lambda)$ are two functions which are holomorphic in R and have no factor† in common in that region, there exist two functions, holomorphic in R , such that*

$$p(\lambda)f(\lambda) + q(\lambda)g(\lambda) \equiv 1.$$

If we expand $1/[f(\lambda)g(\lambda)]$ in terms of its principal parts in a Mittag-Leffler series‡ we get

$$(1) \quad \frac{1}{f(\lambda)g(\lambda)} = F(\lambda) + G(\lambda) + \phi(\lambda),$$

where $F(\lambda)$ and $G(\lambda)$ are the parts of the series arising from the zeros of $f(\lambda)$ and $g(\lambda)$, respectively, which lie in R , and $\phi(\lambda)$ is a function which is holomorphic in R . If we now put

$$p(\lambda) = g(\lambda)[G(\lambda) + \phi(\lambda)], \quad q(\lambda) = f(\lambda)F(\lambda),$$

both $p(\lambda)$ and $q(\lambda)$ are holomorphic in R and, on multiplying both sides of (1) by $f(\lambda)g(\lambda)$, we have

$$\begin{aligned} 1 &= f(\lambda) \cdot g(\lambda)[G(\lambda) + \phi(\lambda)] + g(\lambda) \cdot f(\lambda)F(\lambda) \\ &= f(\lambda)p(\lambda) + g(\lambda)q(\lambda), \end{aligned}$$

as required by the Lemma.

* The determinant of the matrix $P(\lambda)$ is denoted by $|P(\lambda)|$.

† I. e., the two functions have no common zeros in R . Functions which are holomorphic and nowhere zero in R play the same rôle as constants do in the algebraic theory and will therefore not be regarded as factors in this paper.

‡ The series used here is a special case of the "generalized Mittag-Leffler theorem." See Osgood, *Funktionentheorie* vol. 1 (second edition), p. 540, or Mittag-Leffler, *Acta Mathematica*, vol. 4 (1884), p. 8.

3. LEMMA III. If $p_1(\lambda), p_2(\lambda), \dots, p_n(\lambda)$ are n functions which are holomorphic and have no factor common to all in R , and x_1, x_2, \dots, x_n is a set of vector functions which are holomorphic and linearly independent for all values of λ in R , there exists a matrix function $P(\lambda)$, holomorphic in R , for which (i) $|P(\lambda)| \neq 0$ for any λ in R , and (ii)

$$P(\lambda) x_1 = p_1 x_1 + p_2 x_2 + \dots + p_n x_n.$$

Assume that the lemma is true for bases of order less than n : then, if l is the H.C.F. of p_1, p_2, \dots, p_{n-1} and $p'_r = p_r/l$, ($r = 1, 2, \dots, n-1$), there exists a matrix function, Q_0 , relative to the basis x_1, x_2, \dots, x_{n-1} which is holomorphic and never singular in R and is such that

$$Q_0 x_1 = p'_1 x_1 + p'_2 x_2 + \dots + p'_{n-1} x_{n-1}.$$

Let Q_1 be the matrix defined relatively to the basis x_1, x_2, \dots, x_n by

$$Q_1 x_n = x_n, \quad Q_1 x_r = Q_0 x_r \quad (r = 1, 2, \dots, n-1),$$

and set $x'_r = Q_1 x_r$ ($r = 1, 2, \dots, n$). Since l and p_n have no common factor in R , we can, as in Lemma II, find two functions, α and β , which are holomorphic in R and are such that* $\alpha l + \beta p_n \equiv 1$. If therefore Q_2 is the matrix defined by

$$Q_2 x'_1 = l x'_1 + p_n x'_n, \quad Q_2 x'_n = -\beta x'_2 + \alpha x'_n, \quad Q_2 x'_r = x'_r \\ (r = 2, \dots, n-1),$$

we have $|Q_2| = 1$, and $P(\lambda) = Q_2 Q_1$ satisfies the conditions of the lemma since

$$P x_1 = Q_2 Q_1 x_1 = Q_2 x'_1 = l x'_1 + p_n x'_n = \sum_1^n p_r x_r.$$

Since the lemma is obviously true for $n = 1$, the required result follows immediately by induction.

4. THEOREM. If $A(\lambda)$ is a matrix function of rank r which is holomorphic in a region R , there exist two matrix functions, $P(\lambda)$ and $Q(\lambda)$, which are holomorphic and non-singular in R , and are such that

$$P(\lambda) A(\lambda) Q(\lambda) = \left\| \begin{array}{ccccccc} E_1(\lambda) & & & & & & \\ & E_2(\lambda) & & & & & \\ & & \cdot & & & & \\ & & & \cdot & & & \\ & & & & \cdot & & \\ & & & & & E_r(\lambda) & \\ & & & & & & 0 \\ & & & & & & \cdot \\ & & & & & & & \cdot \\ & & & & & & & & 0 \end{array} \right\|,$$

* If $p_n = 0$, then $l = 1 = \alpha$.

where $E_1(\lambda), \dots, E_r(\lambda)$ are functions of λ which are holomorphic in R and are such that E_s is a factor of E_t when $s < t$ ($s, t = 1, 2, \dots, r$).

This theorem is obviously true when $n = 1$ so that we may make its proof depend on induction. We assume therefore that it is true for matrices of order less than n .

If the rank of A is less than n , there is at least one vector function x , holomorphic in R , for which $Ax \equiv 0$ in R ; and we can insure that x itself does not vanish by removing any common factor from its coefficients when expressed in terms of e_1, e_2, \dots, e_n . By Lemma III, there is a matrix function, Q_0 , holomorphic and non-singular in R , for which $x = Q_0 e_n$, whence $AQ_0 e_n \equiv 0$: the elements in the last column of AQ_0 are therefore all zero. Considering now the conjugate matrix $Q'_0 A'$, we see in the same manner that there is a matrix function P'_0 holomorphic and non-singular in R , which is such that the coefficients in the last column of $Q'_0 A'$ are also all zero. It follows then that $P_0 AQ_0$ has zeros both in the last column and in the last row, and may therefore be regarded as a matrix of order $n - 1$ relative to the basis e_1, e_2, \dots, e_{n-1} . There are therefore by hypothesis two matrix functions, P_1 and Q_1 , of order $n - 1$ which are holomorphic and non-singular in R and are such that $P_1 P_0 AQ_0 Q_1$ has the desired normal form with regard to the basis e_1, e_2, \dots, e_{n-1} : and we have only to extend P_1 and Q_1 to the original basis, e_1, e_2, \dots, e_n , by adding the conditions $P_1 e_n = e_n = Q_1 e_n$ in order to have A in the required normal form with regard to this basis.

Assume now that the rank of A is n . If the coefficients of A have a H.C.F., $f(\lambda)$, we may write $A(\lambda) = f(\lambda) A_1(\lambda)$: therefore, since multiplying $P(\lambda) A(\lambda) Q(\lambda)$ by a scalar factor still leaves it in the normal form, we may, without loss of generality, assume that there is no common factor.

If $x = (\xi_1, \xi_2, \dots, \xi_n)$ is a constant vector, Ax can only vanish for values of λ for which $|A| = 0$: but the values of λ for which $Ax = 0$ are necessarily continuous functions of the ξ 's and, since the coefficients of A have no common factor in R , there are therefore no values of λ in R for which $Ax = 0$ for every constant vector x ; hence there is some constant vector, x , for which $Ax \neq 0$ for any value of λ in R . Let X be a constant matrix whose first column consists of the coördinates of this vector, the remaining coefficients being so chosen as to make the determinant of X not zero. The coefficients in the first column of AX are then the coefficients of Ax and therefore have no factor in common.

Let $AXe_1 = \Sigma a_{r1}(\lambda) e_r$. By Lemma III, we can find a matrix function $P_1^{-1}(\lambda)$, holomorphic and non-singular in R , which is such that

$$P_1^{-1} e_1 = \sum a_{r1}(\lambda) e_r,$$

whence $P_1 AXe_1 = e_1$. If, therefore,

$$P_1 AX e_r = c_{r1} e_1 + \sum_{s=2}^n c_{rs} e_s \quad (r = 2, 3, \dots, n),$$

and Q_1 is the matrix defined by

$$e_1 = Q_1 e_1, \quad e_r - c_{r1} e_1 = Q_1 e_r \quad (r = 2, 3, \dots, n),$$

then Q_1 is a matrix function, holomorphic in R , for which $|Q_1| = 1$ and

$$P_1 AX Q_1 e_1 = e_1, \quad P_1 AX Q_1 e_r = P_1 AX (e_r - c_{r1} e_1) = \sum_{s=2}^n c_{rs} e_s \quad (r = 2, 3, \dots, n).$$

All the coefficients in the first row and column of $P_1 AX Q_1$ are therefore zero, except the first, which is 1. Striking out this row and column, we have a matrix of order $n - 1$, and, as we have assumed the theorem true for matrices of order less than n , this matrix can be reduced to the required form by means of two matrix functions P_2 and Q_2 which are defined relative to the basis e_2, e_3, \dots, e_n and are holomorphic and non-singular in R . If we extend these matrices to the basis e_1, e_2, \dots, e_n as above by adding the conditions $P_2 e_1 = e_1 = Q_2 e_1$, the matrix $P_2 P_1 AX Q_1 Q_2$ has the form required by the theorem since in multiplying by P_2 and Q_2 the first row and column of $P_1 AX Q_1$ remain unaltered; the induction is therefore complete.
