

ON THE WEDDLE SURFACE AND ANALOGOUS LOCI*

BY

ARNOLD EMCH

I. INTRODUCTION

Geometrically, the well known quartic, the Weddle surface, may be defined as the locus of the vertices of the quadric cones in the linear three-dimensional system of quadrics through six independent points in space. In a similar manner, the locus of the vertices of cones through seven points, or through the eight base points of the net of quadrics determined by the seven points, is a sextic of genus three, which however is not the most general sextic with this genus.

It is the purpose of this paper to establish analogous loci for cubic cones, and, subsequently, for general n -ic cones. The base points through which the cones pass shall be denoted by $\mathfrak{A}_1, \mathfrak{A}_2, \dots, \mathfrak{A}_i$, the vertices of the coördinate tetrahedron by A_1, A_2, A_3, A_4 , the join of any two base points \mathfrak{A}_i and \mathfrak{A}_k by g_{ik} or simply by g ; a surface locus of vertices of cones by W , a curve locus by S .

In case of n -ic cones there is $i = N+1$ ($N = n(n+3)/2$) for a W , $i = N+2$, for an S .

Before taking up the cubic cones it is perhaps well to state briefly some of the properties of the locus of vertices of quadric cones through six points, and through the eight base points of a net of quadrics.

II. THE WEDDLE SURFACE

When $n = 2$, $N+1 = 6$, the order of W is $m = 4$, and we have the Weddle surface as the locus of quadric cones through six points. The number of lines g is 15 which in sets of 5 pass through the six points, the nodes of W . The six points can be arranged in 10 double-triples, each of which determines two planes through the six points, or a degenerate cone whose axis lies on W . Thus, in addition to the 15 g 's, there are 10 other lines on W , so that the total number of lines on W is 25, as is well known.

The order of the curve S of vertices of quadric cones through seven points is $s = 6$. But in the case of $n = 2$ we have the circumstance that all quadric cones through seven points pass through an eighth point, so that the S_8 appears as the locus of vertices of quadric cones in a net of quadrics.

* Presented to the Society, December 29, 1923.

The S_6 cuts each of the 28 lines joining every pair of the base points in 2 points. Every sextuple of the base points determines a Weddle surface, so that there are 28 Weddle surfaces through the S_6 . The S_6 is also the base curve of the involutory cubic Cremona transformation connected with the eight base points, and the surface of the eighth order formed by the trisecants of the S_6 is the base surface of the transformation. The cubic surfaces corresponding to any two planes pass through S_6 , and intersect moreover in a cubic rest curve which corresponds to the line of intersection of the two planes. The S_6 is therefore of genus 3. But this S_6 is not the most general sextic of this genus, since the system of general sextics of genus 3 has the dimension 24 that of our system being $3 \cdot 7 = 21$. For 7 points determine such a sextic and there are $\infty^{3 \cdot 7}$ sextuples. What characterizes our sextic is the fact that it is uniquely determined by the 8 base points of a net of quadrics. The number 21 as the dimension of the system of sextics is also verified as follows. There are ∞^{18} sextuples of points and consequently the same manifold of Weddle surfaces. With a definite Weddle surface fixed, every seventh point determines an S_6 uniquely. Hence on every Weddle surface there are ∞^3 sextics, so that their manifold is ∞^{21} .

In the involutory cubic transformation determined by the octuple of base points, bisecants of the S_6 are transformed into bisecants of S_6 , a quartic surface through S_6 into a quartic which passes singly through S_6 , because every trisecant of S_6 cuts the original quartic in one point only (outside of S_6). The 28 Weddle surfaces through S_6 are therefore invariant in the transformation.

Now with every Weddle surface is also uniquely associated a Geiser transformation of order 7. As is well known this involutory Cremona transformation may be defined by the net of quadrics through the six nodes of the surface and a seventh point P . Then all quadrics of this net will pass through an eighth point P' . Thus to every point P corresponds a point P' and conversely to P' corresponds in the same transformation P . The Weddle surface is pointwise invariant in this transformation and passes through a twisted cubic on the six nodes. The join of P and P' is a bisecant s of this cubic, so that corresponding pairs on s form an involution with the intersection of s and the Weddle surface (outside of the cubic) as double points. Summing up, we have

THEOREM 1. *There are 28 Weddle surfaces through a sextic S_6 determined by the octuple of base points of a net of quadrics. Each of the W 's is invariant in the involutory cubic transformation determined by the octuple. There are also 28 involutory Geiser transformations each leaving one of the W 's, and consequently the S_6 , pointwise invariant.*

The 28 Geiser transformations determine an infinite discontinuous group G which leaves the sextic pointwise invariant.

Denoting by T_{ik} the Geiser transformation associated with the Weddle surface W_{ik} which has the six points obtained from the octuple A_h ($h = 1, 2, \dots, 8$) by leaving out A_i and A_k ($h = i, h = k$), T_{ik} has the property that it permutes A_i and A_k and leaves the remaining 6 points of the octuple unchanged. Thus the 28 T_{ik} 's leave the configuration of the octuple unchanged. Any permutation of the 8 points may be obtained as a product of Geiser transformations T_{ik} . For example, $A_1 A_2 A_3 A_4 A_5 A_6 A_7 A_8$ is transformed into $A_5 A_3 A_1 A_8 A_4 A_2 A_7 A_6$ by the transformation $T_{68} T_{56} T_{45} T_{36} T_{26} T_{13}$. This is obvious since every substitution may be obtained as a product of transpositions. The transformations of G therefore transform the octuple in $8! = 40,320$ different ways.

To prove that the transformations of G are infinite in number, consider for example the repeated product $T_{78} T_{12} T_{78} T_{12} \dots$ and the effect which it has on the Weddle surface W_{12} . If the repeated product at some stage could lead to an identity, W_{12} , which in turn is transformed by $T_{78}, T_{12}, T_{78}, T_{12}, \dots$ into surfaces of definite orders, would accordingly be transformed into itself. Now, as a matter of fact, T_{78} transforms W_{12} into a surface W'_{12} of order 12 with double points at A_7 and A_8 , six-fold points at A_3, A_4, A_5, A_6 and eight-fold points at A_1 and A_2 , as can easily be ascertained. T_{12} transforms W'_{12} into a surface W''_{12} of order 28 with eight-fold points at A_1 and A_2 , 14-fold points at A_3, A_4, A_5, A_6 , 18-fold points at A_7, A_8 . Next, T_{78} transforms W''_{12} into W'''_{12} , of order 52; again T_{12} transforms W'''_{12} into $W^{(4)}_{12}$ of order 84, T_{78} transforms $W^{(4)}_{12}$ into $W^{(5)}_{12}$ of order 124, etc. Thus the succession of $1, 2, 3, 4, \dots, m, \dots$ transformations (including 1 as the identical one) leads consecutively to surfaces of order $4, 12, 28, 52, 84, 124, \dots, 4(m^2 - m + 1), \dots$. From this follows that the repeated products of T_{78} and T_{12} all represent different transformations; and that their number, and hence also the number of transformations of the G , is infinite.

III. W -SURFACE OF VERTICES OF CUBIC CONES

Let $\mathcal{U}_i (a_i b_i c_i d_i)$, $i = 1, 2, \dots, 10$, be the ten base points and $P(x_1, x_2, x_3, x_4)$ any point in space distinct from any of the \mathcal{U}_i 's. Assume the coördinate tetrahedron in a general position, and project the points \mathcal{U}_i from P upon $x_4 = 0$. The coördinates of the projections \mathcal{U}'_i are

$$\varrho X_i = a_i x_4 - d_i x_1, \quad \varrho Y_i = b_i x_4 - d_i x_2, \quad \varrho Z_i = c_i x_4 - d_i x_3,$$

or, writing cartesian coördinates in space

$$x = \frac{x_1}{x_4}, \quad y = \frac{x_2}{x_4}, \quad z = \frac{x_3}{x_4},$$

and putting $d_i = 1$,

$$(1) \quad \sigma X_i = x - a_i, \quad \sigma Y_i = y - b_i, \quad \sigma Z_i = z - c_i.$$

The ten projections \mathfrak{A}'_i lie on a cubic in $x_4 = 0$, when the determinant

$$(2) \quad \begin{vmatrix} X_1^3, & Y_1^3, & Z_1^3, & X_1^2 Y_1, & X_1^2 Z_1, & Y_1^2 X_1, & Y_1^2 Z_1, & Z_1^2 X_1, & Z_1^2 Y_1, & X_1 Y_1 Z_1 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ X_{10}^3, & Y_{10}^3, & Z_{10}^3, & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & X_{10} Y_{10} Z_{10} \end{vmatrix} = 0.$$

The expansion of this gives an equation of degree 30 in x, y, z for the locus of vertices (x, y, z) of cubic cones through the ten given points. But this equation is reducible, as is seen by expanding the X, Y , and Z terms in (2), and by subtracting the first row in succession from the 9 remaining rows. In this manner all cubic terms in the 9 rows vanish, so that the degree of the expanded determinant now becomes 21. By repeated division by proper numerical factors and subsequent subtraction the degree may be reduced still further. But the degree of the final irreducible equation may be established by a much less laborious method. The order of the W -surface is equal to the number of its intersections with a generic line. Such a line is $y = 0, z = 0$. Substituting this in (2) the first column contains only cubic terms in x , the second and third columns only constants, the fourth and fifth only quadratic terms in x , the sixth only linear terms in x , the seventh constant terms, the eighth linear terms in x , the ninth constant terms, and the tenth linear terms in x . The determinant consequently expands into a polynomial of degree $1 \cdot 3 + 2(3 - 1) + 3(3 - 2) = 10$ in x . Consequently the line $y = 0, z = 0$ cuts the W -surface in 10 points. When the determinant (2) is expanded by multiplying the elements of the k th row by their corresponding minors, a homogeneous polynomial of degree 3 in $x - a_k, y - b_k, z - c_k$ is obtained. From this follows that the point (a_k, b_k, c_k) is a triple point of the surface, and that the cubic cone with this point as a vertex, and through the remaining nine points, is the tangential cone to the surface at this point.

Consider any point V of a line g joining any two of the ten points \mathfrak{A}_i . This line g and the 9 lines joining V to the 8 remaining points determine a cubic cone uniquely. This is a cubic cone through the 10 points, consequently V , and the line g , lie on the W -surface. These results may be summed up in

THEOREM 2. *The locus of vertices of cubic cones through 10 fixed points in space is a surface of order 10 with these points as triple-points and with the 45 joins of the 10 points as single lines on the surface.*

IV. S -CURVE OF VERTICES OF CUBIC CONES

In addition to the 10 fixed points $\mathfrak{A}_1, \dots, \mathfrak{A}_{10}$ of the W -surface of order 10 consider another fixed point \mathfrak{A}_{11} . The locus of the vertices of cubic cones through these 11 points is a curve S which may be obtained from the intersection of any two W -surfaces associated with two sets of ten points each chosen from the 11 points. Thus, S may be obtained from the intersection of the two surfaces W_{10} and W_{11} associated with $\mathfrak{A}_1 \dots \mathfrak{A}_9 \mathfrak{A}_{10}$ and $\mathfrak{A}_1 \dots \mathfrak{A}_9 \mathfrak{A}_{11}$, respectively. Now consider the line l joining \mathfrak{A}_{10} and \mathfrak{A}_{11} ; it will cut W_{10} in 7 points V_1, \dots, V_7 outside of \mathfrak{A}_{10} which is a triple point of W_{10} . Hence V_1, \dots, V_7 are vertices of cubic cones which pass through all 11 points and are therefore points of S . Thus, each of the 55 joins l of the 11 points cuts the curve S in 7 points. S itself does not pass through the 11 points $\mathfrak{A}_1, \dots, \mathfrak{A}_{11}$. This fact may be immediately verified analytically by choosing \mathfrak{A}_{10} at $(a_{10} = 0, b_{10} = 0, c_{10} = 0)$, and $y = 0, z = 0$ as the line l . Then (2) reduces to an equation of degree 7 in x .

As W_{10} and W_{11} have the 36 joins g of $\mathfrak{A}_1, \dots, \mathfrak{A}_9$ in common, their curve of intersection outside of these lines is of order 64. But it is reducible. To show this take the plane p determined by any triple of points out of the nine common to W_{10} and W_{11} , say by $\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{A}_3$. Let g_1, g_2, g_3 be the joins of $\mathfrak{A}_2 \mathfrak{A}_3, \mathfrak{A}_3 \mathfrak{A}_1, \mathfrak{A}_1 \mathfrak{A}_2$, and B_1, B_2, \dots, B_{15} the intersections with p of the 15 joins g of $\mathfrak{A}_4, \mathfrak{A}_5, \dots, \mathfrak{A}_9$. The plane p cuts W_{10} and W_{11} each in g_1, g_2, g_3 and a curve of order 7 which passes through the 15 B 's and singly through $\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{A}_3$, since these are triple points. Hence the two curves of order 7, outside of the B 's and \mathfrak{A} 's intersect in $49 - 15 - 3 = 31$ residual points, which lie on the curve S . But S cuts each g_1, g_2, g_3 in 7 points. Consequently p intersects S in $31 + 21 = 52$ points, which is the order of S .

As the curve of intersections of W_{10} and W_{11} outside of the 36 g 's is of order 64, of which S is a part, there is a residual curve R of order 12. Since each of the 9 \mathfrak{A} 's common to W_{10} and W_{11} is a triple point for both surfaces at which 8 g 's, common to W_{10} and W_{11} , are concurrent, the R passes through each of the 9 points. Thus, in the plane p , R passes singly through $\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{A}_3$, and must cut g_1, g_2, g_3 each in 3 additional points. The curve R does not belong to the locus of vertices of cubic cones through 11 points. To ascertain the geometric meaning of R , consider a pencil $\Phi_1 + \lambda \Phi_2 = 0$ of cubic cones through 8 of the 9 points, say $\mathfrak{A}_1, \dots, \mathfrak{A}_8$, with $V(x, y, z)$ as a vertex. All cones of the pencil pass through the 8 lines $V\mathfrak{A}_1, \dots, V\mathfrak{A}_8$, and a ninth fixed line g_9 . Intersect this pencil with $x_4 = 0$, as before, and denote the projections by the same letters with accents. Then in order that \mathfrak{A}'_9 may coincide with the intersections of g_9 with $x_4 = 0$, on substituting the coördinates of \mathfrak{A}_9 in

$\phi'_1 + \lambda \phi'_2 = 0$, there must be simultaneously $\phi'_1 = 0$, $\phi'_2 = 0$. The irreducible part of the intersection of the two surfaces is precisely the curve R . From every point of R the nine points are projected into 9 points of a pencil of cubics. Hence there is a cubic of the pencil through \mathfrak{A}'_{10} and another cubic through \mathfrak{A}'_{11} . Consequently every point of R is a vertex of two cubic cones through $\mathfrak{A}_1, \dots, \mathfrak{A}_9, \mathfrak{A}_{10}$ and $\mathfrak{A}_1, \dots, \mathfrak{A}_9, \mathfrak{A}_{11}$. The result may be stated in

THEOREM 3. *The locus of vertices of cubic cones through 11 points is a curve S of order 52, which cuts each of the 55 joins in 7 points.*

There are 55 W -surfaces through S . Any two surfaces of the set intersect in a residual curve R of order 12, uniquely associated with the 9 points \mathfrak{A} common to the two W 's. The curve R is the locus of points from which the nine points are projected into the base points of a pencil of cubics.

The curve S has no effective singularities, so that its genus may be determined by evaluating the number of its apparent double points from any generic point O , and by subtracting this number from the maximum number of double points of its projection. The sextic cone C through S with O as a vertex cuts W_{11} , outside of S , in a residual curve S^* of order $10 \cdot 52 - 52 = 468$. Now S' cuts W_{10} in 4680 points. Among these points are the $9 \cdot 52 = 468$ points in which the first polar of O with respect to W_{11} cuts S . At such a point the tangents to S and S' at this point lie in the tangent plane to W_{11} at this point. Hence the projection of a point does not result in a double point. The cone C cuts each join g of the nine points \mathfrak{A} , associated with S , in 52 points, among which are the 7 intersections of g with S , which do not lie on S^* . The remaining 45 points on each g lie on S^* , and on W_{10} , but not on S . Thus there are $36 \cdot 45 = 1620$ such points, which also are not projected into double points of the projections of S' of S from O . The remaining $4680 - 468 - 1620 = 2592$ intersections of S^* with W_{10} arrange themselves into 1296 couples P, P' which lie simultaneously on S and S^* and on lines through O . In fact, such a point of intersection P^* of S^* with W_{10} lies on W_{11} and W_{10} , and consequently on S . But P^* is the projection from O upon W_{11} of a point P and S , so that P^* is projected back into P , and P also lies on S^* .

The genus of S is therefore $\frac{1}{2} \cdot 51 \cdot 50 - 1296 = 1254$.

As a verification, the same method applied to the curve S in case of the Weddle surface yields the genus 3, as is well known.

V. W -SURFACES OF VERTICES OF n -IC CONES

Following the same method and notation as in the case of cubic cones it is easy to generalize the results for n -ic cones, and it is not necessary to restate the arguments in detail.

The determinant (2) is now of order $N+1$ and expands into a polynomial of degree $n(N+1)$ in x, y, z . If we write the equation of the n -ic in $x_4 = 0$ in the form

$$X^n + X^{n-1} \phi_1 + X^{n-2} \phi_2 + \dots + \phi_n = 0,$$

in which ϕ_i is a homogeneous polynomial of degree i in Y and Z , so that ϕ_i contains $i+1$ terms, the determinant

$$(3) \quad \begin{vmatrix} (x-a_1)^n, & (x-a_1)^{n-1}(y-b_1), & (x-a_1)^{n-1}(z-c_1) & \dots, & (z-c_1)^n \\ \cdot & \cdot & \cdot & \cdot & \cdot, \dots, \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot, \dots, \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot, \dots, \cdot \\ (x-a_{N+1})^n, & (x-a_{N+1})^{n-1}(y-b_{N+1}), & (x-a_{N+1})^{n-1}(z-c_{N+1}), & \dots, & (z-c_{N+1})^n \end{vmatrix} = 0$$

contains x^n as the highest power in the first column only, x^{n-1} in the next two, x^{n-2} in the next three, x^{n-k} in $k+1$ consecutive columns. The reduced form of (3) gives the equation of the W -surface. It is evident from (3) that the generic line $y = 0, z = 0$ gives for its intersection with W an equation of degree

$$(4) \quad m = 1 \cdot n + 2(n-1) + 3(n-2) + \dots + (n-1)2 + n \cdot 1 \\ = \frac{n(n+1)(n+2)}{6}$$

in x . Hence m is the order of the W -surface. Moreover if we write (3) in the form (2) and expand from the elements of the k th row, it is apparent that \mathfrak{A}_k is an n -fold point of W . Again, every point of a join g of two of the $N+1$ points is a vertex of an n -ic cone through the $N+1$ points. Hence

THEOREM 4. *The locus of vertices of n -ic cones through $\frac{1}{2}n(n+3)+1$ points \mathfrak{A}_i is a W -surface of order $\frac{1}{6}n(n+1)(n+2)$ with the \mathfrak{A}_i 's as n -fold points and the joins g as single lines. The tangential cones at the \mathfrak{A}_i 's are determined by the N joins g through each \mathfrak{A}_i .*

VI. S -CURVES

The locus of vertices of n -ic cones through $N+2$ points is an S -curve, through which all the $\frac{1}{2}(N+2)(N+1)$ W -surfaces associated with the $(N+1)$ -tuples of the $N+2$ points pass. Let W_{N+1} and W_{N+2} be two of these surfaces with the group of common points $(\mathfrak{A}) = (\mathfrak{A}_1, \mathfrak{A}_2, \dots, \mathfrak{A}_N)$ and with \mathfrak{A}_{N+1} and \mathfrak{A}_{N+2} individualizing the two surfaces respectively. Their full intersection, to which S belongs, is of order $m^2 = \frac{1}{96}n^2(n+1)^2(n+2)^2$,

from which split off the $\frac{1}{2}N(N-1)$ joins g of (\mathfrak{A}) . The intersection proper of W_{N+1} and W_{N+2} is therefore a curve T of order

$$t = m^2 - \frac{1}{2}N(N-1) = \frac{1}{72}(2n^6 + 12n^5 + 17n^4 - 30n^3 - 55n^2 + 54n).$$

This curve is composed of S and a residual curve R . To determine the order of these curves, the number of points in which S cuts a join g must first be established. Let g be the join of \mathfrak{A}_{N+1} and \mathfrak{A}_{N+2} . It cuts W_{N+1} in $m-n$ points V , outside of \mathfrak{A}_{N+1} . The n -ic cones with the V 's as vertices and through (\mathfrak{A}) and \mathfrak{A}_{N+1} also pass through \mathfrak{A}_{N+2} ; consequently the same V 's are also vertices of n -ic cones through (\mathfrak{A}) and \mathfrak{A}_{N+2} , and lie on W_{N+2} , and hence on S . Thus, S cuts each g in $m-n$ points. Next cut both W_{N+1} and W_{N+2} by the plane p through $\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{A}_3$. Neglecting the joins g_1, g_2, g_3 , which are common to both surfaces, the residual intersections are curves C_1 and C_2 of order $m-3$, which both pass through the $\frac{1}{2}N(N-1)-3$ intersections B of the joins g of the group (\mathfrak{A}) (minus the three lines g_1, g_2, g_3). As W_{N+1} and W_{N+2} have n -tuple points at $\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{A}_3$, and at each of these have $N-1$ g 's in common, the curve T has multiplicities of order $n^2-N+1 = \frac{1}{2}(n-1)(n-2)$ at each $\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{A}_3$. The multiplicity of each C_1 and C_2 at these points is $n-2$. Hence these points and the points B absorb $\frac{1}{2}N(N-1)-3+3(n-2)^2$ of the $(m-3)^2$ intersections of C_1 and C_2 . The residual set of intersections, that of S with p , outside of g_1, g_2, g_3 , consists therefore of

$$\begin{aligned}\mu &= (m-3)^2 - \frac{1}{2}(N-3)(N-4) - 3(n-2)^2 \\ &= \frac{1}{72}(2n^6 + 12n^5 + 17n^4 - 102n^3 - 379n^2 + 1098n - 648).\end{aligned}$$

To this number must be added the $3(m-n) = \frac{1}{4}(n^3 + 3n^2 - 4n)$ intersections of S with g_1, g_2, g_3 . Hence the order of S is

$$s = \frac{1}{72}(2n^6 + 12n^5 + 17n^4 - 66n^3 - 271n^2 + 954n - 648).$$

The order of R , the residual curve of intersection of W_{N+1} and W_{N+2} , is $r = t - s$, or

$$r = \frac{1}{2}(n^3 + 6n^2 - 25n + 18) = \frac{1}{2}(n-1)(n-2)(n+9).$$

R is the locus of points from which the N joins of the group (\mathfrak{A}) are projected into the base points of a pencil of n -ics.

Thus may be stated

THEOREM 5. *The locus of vertices of n -ic cones through $\frac{1}{2}n(n+3)+2$ points \mathfrak{A} in space is an S -curve of order*

$$s = \frac{1}{72} (2n^6 + 12n^5 + 17n^4 - 66n^3 - 271n^2 + 954n - 648),$$

which cuts each of the $\frac{1}{2}(N+1)(N+2)$ joins of each two of the \mathfrak{A} 's in $\frac{1}{2}n(n+1)(n+2)-n$ points, and does not pass through the \mathfrak{A} 's. Any two W -surfaces each associated with $N+1$ points of the set \mathfrak{A} pass through S and intersect moreover in a residual curve R of order

$$r = \frac{1}{2} (n-1)(n-2)(n+9)$$

which has multiple points of order $\frac{1}{2}(n-1)(n-2)$ at the points of the group (\mathfrak{A}) associated with R . R is the locus of points from which the group (\mathfrak{A}) is projected into the base of a pencil of n -ics.

The genus of S (and also of R) could easily be determined as in the case of cubic cones and, for the sake of brevity, will be omitted.

UNIVERSITY OF ILLINOIS,
URBANA, ILL.
