

AN EXTENSION OF SCHWARZ'S LEMMA*

BY

LARS V. AHLFORS

I. THE FUNDAMENTAL INEQUALITY

1. To every neighborhood on a Riemann surface there is given a map onto a region of the complex plane. For any two overlapping neighborhoods the corresponding maps are directly conformal.† We agree to denote points on the surface by w , corresponding values of the local complex parameter by z .

We introduce a Riemannian metric of the form

$$(1) \quad ds = \lambda |dz|,$$

where the positive function λ is supposed to depend on the particular parameter chosen, in such a way that ds becomes invariant. The metric is regular if λ is of class C_2 . In this paper we shall, without mentioning it further, allow λ to become zero, although such points are of course singularities of the metric.

It is well known that the Gaussian curvature of the metric (1) is given by

$$(2) \quad K = -\lambda^{-2} \Delta \log \lambda,$$

and that this expression remains invariant under conformal mappings of the w -plane. We are interested in the case of a metric with negative curvature, bounded away from zero. It is convenient to choose the upper bound of the curvature equal to -4 . From (2) it follows that the corresponding λ satisfies the condition

$$(3) \quad \Delta \log \lambda \geq 4\lambda^2.$$

When we set $u = \log \lambda$ this is equivalent to

$$(4) \quad \Delta u \geq 4e^{2u}.$$

The hyperbolic metric of the unit circle $|z| < 1$ is defined by

$$(5) \quad d\sigma = (1 - |z|^2)^{-1} |dz|$$

and has the constant curvature -4 .

2. Consider now an analytic function $w=f(z)$ from the circle $|z| < 1$ to a Riemann surface W . The analyticity is expressed by the fact that every local parameter w is an analytic function of z . To a differential element dz corresponds an element dw whose length does not depend on the direction of dz . The corresponding value of $ds = \lambda |dw| = \lambda_z |dz|$ is therefore uniquely de-

* Presented to the Society, September 8, 1937; received by the editors April 1, 1937.

† For the definition of a Riemann surface see T. Radó, *Über den Begriff der Riemannschen Fläche*, Acta Szeged, vol. 2 (1925).

terminated, and we have $\lambda_z = \lambda |w'(z)|$. It is also seen that $u = \log \lambda_z$ satisfies the condition (4) whenever the given metric has a curvature ≤ -4 . An exception has to be made for the possible zeros of λ_z , corresponding to the zeros of λ and $w'(z)$.

THEOREM A. *If the function $w = f(z)$ is analytic in $|z| < 1$, and if the metric (1) of W has a negative curvature ≤ -4 at every point, then the inequality*

$$(6) \quad ds \leq d\sigma$$

will hold throughout the circle.

Proof: Choose an arbitrary $R < 1$ and set $v = \log R(R^2 - |z|^2)^{-1}$ for $|z| < R$. We note that $\Delta v = 4e^{2v}$ and consequently

$$(7) \quad \Delta(u - v) \geq 4(e^{2u} - e^{2v}).$$

Let us denote by E the open point set in $|z| < R$ for which $u > v$. It is clear that E cannot contain any zeros of λ_z . Hence (7) is valid and shows that $u - v$ is subharmonic in E . It follows that $u - v$ can have no maximum in E and must approach its least upper bound on a sequence tending to the boundary of E . But E can have no boundary points on $|z| = R$, for v becomes positively infinite as z tends to that circle, and at interior boundary points we must have $u - v = 0$, by continuity. A contradiction is thus obtained, unless E is vacuous. The inequality $u \leq v$ consequently subsists for all points with $|z| < R$, and letting R tend to 1 we find $u \leq -\log(1 - |z|^2)$ at all points. This is equivalent to (6).

If W is the unit circle and ds its hyperbolic metric, Theorem A is simply the differential form of Schwarz's lemma given by Pick.*

3. Several generalizations of the theorem just proved suggest themselves at once. Since the only thing we need is to prevent the function $u - v$ from having a maximum in E , it is obvious that the assumptions on λ can be considerably weakened, without affecting the validity of the argument. We shall give below two such generalizations which are found to be particularly useful for the applications.

THEOREM A1. *Let λ be continuous and such that at every point, either (a) the second derivatives of $u = \log \lambda$ are continuous and satisfy (4), or (b) it is possible to find two opposite directions n', n'' for which $\partial u / \partial n' + \partial u / \partial n'' > 0$. Then the statement of the previous theorem is still true.*

Opposite directions in the w -plane correspond to opposite directions in the z -plane. At a maximum of $u - v$ we have $\partial u / \partial n \leq \partial v / \partial n$ in any direction, when-

* An account of all questions related to Schwarz's lemma will be found in R. Nevanlinna, *Eindeutige analytische Funktionen*, Springer, 1936, pp. 45-58.

ever the directional derivative exists. For opposite directions $\partial v/\partial n' + \partial v/\partial n'' = 0$; hence $\partial u/\partial n' + \partial u/\partial n'' \leq 0$ in case of a maximum. It follows that no maximum can be attained in points satisfying condition (b).

We shall call $ds' = \lambda' |dw|$ a supporting metric of $ds = \lambda |dw|$ at the point w_0 if: (1) $\lambda' = \lambda$ at w_0 , (2) λ' is defined and $\leq \lambda$ in a neighborhood of w_0 .

THEOREM A2. *Suppose that λ is continuous, and that it is possible to find a supporting metric, satisfying (4), at every point of W . Then the inequality (6) still holds.*

If $u - v > 0$ at z_0 , then $u' - v$ will also be positive, and consequently subharmonic, in a neighborhood of z_0 .^{*} A maximum of $u - v$ will *a fortiori* be a maximum of $u' - v$. Hence $u - v$ can have no maximum in E .

II. SCHOTTKY'S THEOREM

4. As a first application we prove Schottky's theorem with definite numerical bounds.

THEOREM B. *If $f(z)$ is analytic and different from 0 and 1 in $|z| < 1$, then*

$$(8) \quad \log |f(z)| < \frac{1 + \theta}{1 - \theta} (7 + \log |f(0)|)$$

for $|z| \leq \theta < 1$.[†]

Let $\zeta_1 = \zeta_1(w)$ map the region outside of the segment $(0, 1)$ onto the exterior of the unit circle, so that $w = \infty$ corresponds to $\zeta_1 = \infty$, $w = 1$ to $\zeta_1 = 1$, and $w = 0$ to $\zeta_1 = -1$. We also set $\zeta_2(w) = \zeta_1(w^{-1})$ and $\zeta_3(w) = \zeta_2(1 - w)$. Clearly these functions define similar maps of the regions outside of the segments $(1, \infty)$ and $(-\infty, 0)$. Explicitly, $\zeta_1(w)$ is obtained from the equation

$$(9) \quad \zeta_1 + \zeta_1^{-1} = 4w - 2.$$

We introduce the coordinates $\rho_1 = |w|$, $\rho_2 = |w - 1|$ and divide the plane into regions Ω_1 : $\rho_1 \geq 1$, $\rho_2 \geq 1$; Ω_2 : $\rho_1 \leq 1$, $\rho_1 \leq \rho_2$; Ω_3 : $\rho_2 \leq 1$, $\rho_2 \leq \rho_1$. The metric

$$(10) \quad ds_i = \frac{|d \log \zeta_i|}{2(4 + \log |\zeta_i|)} = \lambda_i |dw|$$

^{*} u' corresponds to λ' as u to λ .

[†] Schottky's original theorem was purely qualitative. Numerical relations have been studied at great length, notably by Ostrowski (*Studien über den Schottky'schen Satz*, Basel, 1931, and *Asymptotische Abschätzung des absoluten Betrags einer Funktion, die die Werte 0 und 1 nicht annimmt*, *Commentarii Mathematici Helvetici*, vol. 5 (1933)), but no simple inequality comparable with (8) has ever been proved.

Added in proof: Numerical bounds of the same order of magnitude are found by A. Pfluger, *Über numerische Schranken im Schottky'schen Satz*, *Commentarii Mathematici Helvetici*, vol. 7 (1935). His proof depends on the use of modular functions, while ours is strictly elementary.

is readily recognized as the hyperbolic metric of a half-plane with the constant curvature -4 . Computing the derivatives $\zeta'_i(w)$ we find

$$(11) \quad \begin{aligned} \lambda_1^{-1} &= 2(\rho_1\rho_2)^{1/2}(4 + \log |\zeta_1|), \\ \lambda_2^{-1} &= 2\rho_1\rho_2^{1/2}(4 + \log |\zeta_2|), \\ \lambda_3^{-1} &= 2\rho_2\rho_1^{1/2}(4 + \log |\zeta_3|). \end{aligned}$$

We now set $ds = \lambda|dw|$ with $\lambda = \lambda_i$ in Ω_i . This metric is regular and satisfies condition (3) except at the singular points $0, 1, \infty$ and on the lines separating the regions Ω_i . On these lines λ is still continuous, as seen from (11) and the relations between ζ_1, ζ_2 , and ζ_3 .

Next we wish to show that condition (b) in Theorem A1 holds on the singular lines. We consider the arc $\rho_1 = 1, \rho_2 > 1$ and choose n', n'' as the outer and inner normals of the circle. The required condition is

$$\frac{\partial}{\partial n'} \log \lambda_1 + \frac{\partial}{\partial n''} \log \lambda_2 = \frac{\partial}{\partial n'} \log \frac{\lambda_1}{\lambda_2} > 0.$$

From (11) we obtain

$$\frac{\partial}{\partial n'} \log \frac{\lambda_1}{\lambda_2} = \frac{1}{2} - \frac{\frac{\partial}{\partial n'} \log \left| \frac{\zeta_1}{\zeta_2} \right|}{4 + \log |\zeta_1|},$$

which is also equal to

$$\frac{1}{2} - 2(4 + \log |\zeta_1|)^{-1} \frac{\partial \Phi_1}{\partial \phi},$$

where $\Phi_1 = \arg \zeta_1, \phi = \arg w$. For Φ_1 we have the simple relation $\cos \Phi_1 = \rho_1 - \rho_2$, which for $\rho_1 = 1$ becomes $\cos \Phi_1 = 1 - 2 \sin \phi/2$. Differentiating we find

$$\frac{\partial \Phi_1}{\partial \phi} = \frac{1}{2} \left(1 + \csc \frac{\phi}{2} \right)^{1/2},$$

and by use of the inequalities $\pi/3 \leq \phi \leq 5\pi/3, |\zeta_1| > 1$, we are finally led to the desired result,

$$\frac{\partial}{\partial n'} \log \frac{\lambda_1}{\lambda_2} > \frac{1}{2} - \frac{3^{1/2}}{4} > 0.$$

By symmetry, the same must be true for the arc $\rho_2 = 1, \rho_1 > 1$. The transformation $w' = (1 - w)^{-1}$ takes Ω_1 into Ω_2 and Ω_2 into Ω_3 . Since the function λ is invariant under the transformation we conclude at once that condition (b) will hold also on the line separating Ω_2 and Ω_3 .

From Theorem A1 we can now conclude that $w = f(z)$ satisfies the differ-

ential inequality $\lambda |dw| \leq (1 - |z|^2)^{-1} |dz|$. Integrating, we find that the shortest distance between the points $f(0)$ and $f(z)$, $|z| = \theta$, measured in the metric $ds = \lambda |dw|$, cannot exceed $[\log (1+\theta)/(1-\theta)]/2$.

The shortest path between the circles $\rho_1 = m$ and $\rho_1 = M$, where $M > m \geq 2$, is a segment of the negative real axis, whose length is found to be

$$\frac{1}{2} \log \frac{4 + \log |\zeta_1(-M)|}{4 + \log |\zeta_1(-m)|}.$$

To simplify we introduce the lower and upper bounds $|\zeta_1(-M)| \geq 4M$, $|\zeta_1(-m)| \leq 5m$. Setting $M = |f(z)|$ and m equal to the greater of the numbers $|f(0)|$ and 2 we obtain

$$4 + \log 4M \leq \frac{1 + \theta}{1 - \theta} (4 + \log 5m).$$

Here $\log 5m \leq \log 10 + \log^+ |f(0)| < 3 + \log^+ |f(0)|$ and we find

$$4 + \log 4M < \frac{1 + \theta}{1 - \theta} (7 + \log^+ |f(0)|)$$

which is stronger than (8).

III. BLOCH'S THEOREM

5. Let $w = f(z)$ be analytic in $|z| < 1$ with $|f'(0)| = 1$. Let $B' = B'(f)$ be the l.u.b. of the radii of all simple (*schlicht*) circles contained in the Riemann surface W generated by $f(z)$. Bloch's theorem is $B = \min B' > 0$. Landau has proved $B > .396$.* Grunsky and Ahlfors proved in a recent paper $B < .472$.†

We show that the method developed in this paper gives an immediate proof of Bloch's theorem with a better lower bound for B . For an arbitrary point \mathfrak{w} on W let $\rho(\mathfrak{w})$ denote the radius of the largest simple circle of center \mathfrak{w} contained in W . It is clear that $\rho(\mathfrak{w})$ is continuous, and equal to zero only at the branch-points. We introduce the metric $ds = \lambda |dw|$ with

$$(12) \quad \lambda = \frac{A}{2\rho^{1/2}(A^2 - \rho)} \quad (\rho = \rho(\mathfrak{w}))$$

and w denoting the variable of the function plane (not the uniformizing variable). A is a constant satisfying the preliminary condition $A^2 > B'$.

In the neighborhood of a branch-point α we have $\rho = |w - \alpha|$. Let n be the multiplicity of α ; then $w_1 = (w - \alpha)^{1/n}$ is a uniformizing variable, and

* E. Landau, *Über den Blochschen Satz und zwei verwandte Weltkonstanten*, Mathematische Zeitschrift, vol. 30 (1929).

† L. V. Ahlfors and H. Grunsky, *Über die Blochsche Konstante*, Mathematische Zeitschrift, vol. 42 (1937). The result was found independently by R. M. Robertson.

the corresponding λ_1 is determined from $\lambda_1 |dw_1| = \lambda |dw|$. We obtain $\lambda_1 = n\rho^{1/2-1/n}/2(A^2-\rho)$, and it is seen at once that the metric is regular in case $n=2$ and that λ_1 becomes zero in case $n>2$.

We wish to apply Theorem A2 and therefore look for a supporting metric satisfying the requirements of that theorem. For a regular point w_0 the surrounding circle of radius $\rho(w_0)$ must pass through at least one singularity b which is either a branch-point or a boundary point for the surface. We set $\rho' = |w-b|$ and define $\lambda' = A/[2\rho'^{1/2}(A^2-\rho')]$. This metric has the curvature -4 for it is obtained from the hyperbolic metric of a circle by means of the transformation $w' = w^{1/2}$. In all points of our circle we have $\rho \leq \rho'$ by the definition of ρ . The inequality $\lambda' \leq \lambda$ is therefore satisfied in a neighborhood of w_0 if the function $t^{1/2}(A^2-t)$ increases for $t \leq \rho(w_0)$. Under this condition λ' will be a supporting function of λ , for at the center w_0 we have $\lambda' = \lambda$. The function $t^{1/2}(A^2-t)$ is increasing as long as $t < A^2/3$. Consequently all the conditions in Theorem A2 are fulfilled if we suppose that $A^2 > 3B'$.

Apply the theorem with $z=0$. Using the condition $|dw/dz|_{z=0}=1$ we get

$$(13) \quad A \leq 2\rho_0^{1/2}(A^2 - \rho_0),$$

where ρ_0 is the radius of the largest simple circle with center at the image of $z=0$. The function in the right member of (13) is increasing, and we can replace ρ_0 by B' obtaining $A \leq 2B'^{1/2}(A^2 - B')$. Letting A tend to $(3B')^{1/2}$ we finally get $B' \geq 3^{1/2}/4$. This implies that Bloch's constant $B \geq 3^{1/2}/4 > .433$.

On the other side, if we insert $A^2 = (3B')^{1/2}$ in (13), lower and upper bounds for ρ_0 in terms of B' can be found.

6. Landau has considered a closely related constant L . Let $L' = L'(f)$ be the l.u.b. of the radii of all circles in the w -plane contained in the projection of W , that is, whose values are taken by the function $w=f(z)$, $|f'(0)|=1$. L is defined as the minimum of all such L' . Clearly, $L \geq B$.

The method employed above is immediately applicable if we choose $\lambda = (2\rho \log C/\rho)^{-1}$. This metric is regular at all branch-points, and when we replace ρ by the distance ρ' from a fixed boundary point, the curvature becomes -4 . In order that the function λ' thus obtained be a supporting function it is sufficient that $t \log C/t$ is increasing. This is true for $t < Ce^{-1}$. We therefore choose $C > eL'$, obtaining the inequality $1 \leq 2L' \log C/L'$ as above. Letting C tend to eL' we find $L' \geq 1/2$ and hence $L \geq 1/2$.

This lower bound is the best known. It shows in particular that $L > B$.*

HARVARD UNIVERSITY,
CAMBRIDGE, MASS.

* In the other direction R. M. Robinson has proved $L < .544$. This result has not been published.