

ONE-PARAMETER SEMIGROUPS IN A SEMIGROUP⁽¹⁾

BY

PAUL S. MOSTERT AND ALLEN L. SHIELDS

1. Introduction. A topological semigroup is a Hausdorff space S with a continuous, associative multiplication. If there is an identity, 1 , then $H(1)$ will denote the maximal subgroup of S containing 1 ; in other words, $H(1)$ is the set of all elements with two-sided inverses. A *one-parameter semigroup* in S is a continuous, one-to-one function $\sigma: [0, 1] \rightarrow S$ such that $\sigma(0) = 1$, and $\sigma(a+b) = \sigma(a)\sigma(b)$ for all $a, b \in [0, 1]$ for which $a+b \in [0, 1]$.

In this paper we obtain the following result on the existence of one-parameter semigroups.

THEOREM 1. *Let S be a compact semigroup with identity, and assume that $H(1)$ is not an open set in S . Let there be a neighborhood V of the identity containing no other idempotents. Then S contains a one-parameter semigroup σ such that $\sigma(a) \notin H(1)$ for $0 < a \leq 1$. Moreover, $\sigma(a) = \sigma(b)g$, $g \in H(1)$, implies $a = b$ and $g = 1$.*

This is a generalization of a previous result of the authors [4, Theorem A] in which it is assumed that $H(1)$ is a Lie group. Our proof will actually establish the following more general result.

THEOREM 2. *The conclusion of the previous theorem is correct if S is merely assumed to be locally compact, provided $H(1)$ contains a compact subgroup G open in $H(1)$ but not open in S , and provided there is a neighborhood V of G containing no idempotents other than the identity.*

In the last part of the paper it is shown that if S is a compact semigroup with zero and identity and no other idempotents, and if $H(1)$ is not an open set in S , then S is arcwise connected and S contains a subsemigroup that is an arc with 0 and 1 as its endpoints. Examples are given to show that this result need not be true if there are additional idempotents. Finally, a theorem is stated giving conditions when a semigroup can be locally imbedded in a Lie group.

2. Preliminaries. The topological closure of a set A will be denoted by A^- . The set-theoretic difference A minus B will be denoted by $A \setminus B$. For the basic properties of topological semigroups see the survey by A. D. Wallace [5].

Let V be the neighborhood of the identity mentioned in the hypotheses of Theorems 1 and 2; without loss of generality we may assume that V has

Presented to the Society, November 30, 1957; received by the editors August 17, 1959.

(¹) This work was supported in part by Tulane University's National Science Foundation contract.

compact closure and that V^- contains no idempotents other than the identity. Let N denote the index set of a fundamental system of neighborhoods of the identity, partially ordered by inclusion. N is thus a directed set and we give it the discrete topology. Let $\{x_\alpha\}$ ($\alpha \in N$) denote a net in V^- (for the properties of nets and subnets see [2, Chapter 2]).

Let N^* denote the Stone-Čech compactification of N . Any continuous function f from N to a compact space X admits a unique extension to a continuous function $f^*: N^* \rightarrow X$. In particular, define $f: N \rightarrow V^-$ by $f(\alpha) = x_\alpha$. Choose a point $b \in N^* \setminus N$ such that b is a limit point of the directed set N , and hold this point b fixed throughout the discussion. A generalized limit is then defined by

$$\text{Lim } x_\alpha = f^*(b).$$

If the ordinary limit exists it will be denoted by $\lim x_\alpha$. If $\{y_\alpha\}$ is another set in V^- indexed by N we define $\text{Lim } y_\alpha$ in a similar manner, using the same point b . The generalized limit has the following properties (Gleason [1]).

- (i) $\text{Lim } x_\alpha$ exists for every set in V^- indexed by N .
- (ii) $\text{Lim } x_\alpha = \lim x_\alpha$ whenever $\lim x_\alpha$ exists.
- (iii) $\text{Lim } x_\alpha y_\alpha = (\text{Lim } x_\alpha)(\text{Lim } y_\alpha)$.
- (iv) If $x_\alpha \in C$, then $\text{Lim } x_\alpha \in C^-$.

A *local semigroup* is a Hausdorff space R with a distinguished element 1 , called the identity, a neighborhood V of 1 , and a continuous function $m: V \times V \rightarrow R$ which is associative whenever the triple products are defined, and which has the property that $m(x, 1) = m(1, x) = x$. We shall write xy instead of $m(x, y)$.

Let G denote the set of all elements of V having inverses in V , in other words, $x \in G$ if and only if $x \in V$ and there is an element $y \in V$ such that $xy = yx = 1$. R is said to be an *adequate local semigroup* if G is a group (not merely a local group). We shall need the following result (see [4, Theorem A]).

Let R be a compact adequate local semigroup whose maximal subgroup G is a Lie group. If G is not an open set in R , and if there exists a neighborhood of the identity containing no other idempotents, then there exists a one-parameter semigroup σ such that $\sigma(a) \notin G$ for $a > 0$. Moreover, if $\sigma(a) = \sigma(b)g$ for some $g \in G$, then $a = b$ and $g = 1$.

(The last sentence was not included in the statement of Theorem A but may be found in §4.4.4 of [4].)

3. Proof of Theorem 1. Throughout this section we assume the hypotheses of the theorem. As remarked in §2, we may assume that V is open and has compact closure, and that V^- contains no idempotents other than 1 . Let $x_\alpha \rightarrow 1$ ($x_\alpha \in V \setminus H(1)$, $\alpha \in N$).

LEMMA 1. *For each $\alpha \in N$ there is a positive integer $n(\alpha)$ such that*

$$(1) \quad \begin{aligned} x_\alpha^k &\in V, & (1 \leq k \leq n(\alpha)), \\ x_\alpha^{n(\alpha)+1} &\notin V. \end{aligned}$$

Proof. Let $x \in V \setminus H(1)$ and assume that $x^n \in V$ ($n=1, 2, \dots$). Then $\Gamma(x) = \{x, x^2, x^3, \dots\} \subset V^-$. But $\Gamma(x)$ is a compact semigroup and hence contains an idempotent. This can only be 1, and therefore $\Gamma(x)$ is a group and $x \in H(1)$, which is a contradiction. (See [3] and [5] for the properties of $\Gamma(x)$.)

Let $\tau: [0, 1] \rightarrow V^-$ be defined by

$$\tau(a) = \lim x_\alpha^{[an(\alpha)]}$$

where $[s]$ denotes the greatest integer less than or equal to s . (x^0 is defined to be 1 for all $x \in V$.)

LEMMA 2. *The function τ has the following properties.*

- (i) $\tau(0) = 1$,
- (ii) $\tau(1) \in V \setminus V$,
- (iii) $\tau(a+b) = \tau(a)\tau(b)$ for $a, b, a+b \in [0, 1]$.

Proof. (i) is obvious.

(ii) Since $x_\alpha \rightarrow 1$ we have $\tau(1) = \lim x^{[n(\alpha)]} = \lim x^{[n(\alpha)]+1}$; the result now follows from (1).

(iii) Fix a and b and let $c(\alpha) = [an(\alpha) + bn(\alpha)]$. Then $c(\alpha) = [an(\alpha)] + [bn(\alpha)] + \epsilon(\alpha)$, where $\epsilon(\alpha) = 0$ or 1 . Thus $\tau(a+b) = \lim x_\alpha^{c(\alpha)} = \lim x_\alpha^{[an(\alpha)]} \lim x_\alpha^{[bn(\alpha)]} \lim x_\alpha^{\epsilon(\alpha)}$ from which the result follows.

The function τ is not necessarily continuous. However, in the local semigroup $R^* = \{\tau([0, 1])\}^-$ we shall be able to construct a one-parameter semigroup. Notice that R^* is abelian and that $R^* \subset V^-$ (and therefore R^* contains no idempotents except the identity).

LEMMA 3. *Let $Q = \bigcap_{a>0} \{\tau([0, a])\}^-$. Then Q is a compact abelian subgroup of $H(1)$.*

Proof. By (iii) of Lemma 2, $Q^2 \subset Q$, so Q is a compact abelian semigroup. Furthermore, $1 \in Q$ and Q contains no other idempotents (since $Q \subset V^-$). Thus Q is a group, and therefore $Q \subset H(1)$.

LEMMA 4. *If $a_n \rightarrow a$ ($a_n, a \in [0, 1]$), then all cluster points of $\{\tau(a_n)\}$ are contained in the set $\tau(a)Q$.*

Proof. We may assume $a_n > a$ for all n , or $a_n < a$ for all n . In the first case, $\tau(a_n) = \tau(a_n - a)\tau(a)$ and the result follows since $a_n - a \rightarrow 0$ and therefore $\tau(a_n - a)$ clusters in Q . In the second case $\tau(a) = \tau(a_n)\tau(a - a_n)$. Let x be a cluster point of $\{\tau(a_n)\}$. By taking subnets we may assume that $\tau(a_n) \rightarrow x$, and that $\tau(a - a_n)$ is convergent, say to an element y (necessarily in Q). Then $\tau(a) = xy$ or $x = \tau(a)y^{-1}$.

LEMMA 5. If $a > 0 (a \in [0, 1])$, then $\tau(a) \in H(1)$.

Proof. Assume the statement were false, so $\tau(a) = g \in H(1)$ for some $a > 0$. Let $x = \tau(a/2)$, so that $x^2 = g$. Let $y = xg^{-1}$, $z = g^{-1}x$. Then $xy = 1$, $zx = 1$ and so x has both a left and a right inverse and therefore $x \in H(1)$. Similarly, $\tau(a/2^n) \in H(1)$ ($n = 0, 1, 2, \dots$).

Choose dyadic rationals r_n such that $r_n a \rightarrow 1^-$. Then by Lemma 4, $\tau(r_n a)$ clusters in $\tau(1)Q$. But since $\tau(r_n a) \in H(1)$ for all n this implies that $\tau(1) \in H(1)$, contrary to Lemma 2 (ii).

LEMMA 6. There exists $a_0 > 0$ such that τ is one-to-one on $[0, a_0]$. Moreover, $\tau(a) = \tau(b)g$ with $a \leq a_0$ and $g \in Q$ implies $a = b$.

Proof. Let $c_0 = \inf \{c \in [0, 1] : \tau(c) = \tau(b)g \text{ for some } b > c, g \in Q\}$. It will be sufficient to prove that $c_0 > 0$, for then a_0 can be taken to be any positive number smaller than c_0 .

Assume $c_0 = 0$. Then there exist $c_n < b_n$, $c_n \rightarrow 0$, $g_n \in Q$ such that $\tau(c_n) = \tau(b_n)g_n$. Let $d_n = b_n - c_n$. Then

$$(2) \quad \tau(c_n) = \tau(c_n)\tau(d_n)g_n = \tau(c_n)\tau(d_n)^m g_n^m = \tau(c_n)\tau(md_n)g_n^m$$

for every positive integer m such that $md_n \leq 1$. There exist integers $m(n)$ such that $m(n)d_n$ converges to a number d , $1/2 \leq d \leq 1$. By taking subnets we may assume

$$\begin{aligned} \tau(c_n) &\rightarrow h \in Q, \\ \tau(m(n)d_n) &\rightarrow z \in \tau(d)Q \text{ (Lemma 4),} \\ g_n^{m(n)} &\rightarrow g \in Q. \end{aligned}$$

From (2) we have $h = hzg$ or $z \in Q$, which is a contradiction to Lemma 5.

Let $R = \tau([0, a_0])Q$. R is a closed set by Lemma 4, and therefore R is a compact abelian adequate local semigroup. Also by Lemma 4, Q is not an open set in R . If H is a closed subgroup of Q then the factor semigroup R/H is an adequate local semigroup whose maximal subgroup is Q/H .

Consider the family F of all pairs (H, σ) where H is a closed subgroup of Q and σ is a one-parameter semigroup in R/H , $\sigma: [0, a_0] \rightarrow R/H$, such that

$$(3) \quad \sigma(t) \in \pi(\tau(t)Q), \quad (0 \leq t \leq a_0),$$

and

$$(4) \quad \sigma(a) = \sigma(b)g \quad (g \in Q/H) \text{ implies } a = b \text{ and } g = 1.$$

Here π denotes the natural projection homomorphism from R to R/H . We shall consider all one-parameter semigroups to be defined on the interval $[0, a_0]$. This is no restriction since we may always change parameters and pass to the interval $[0, 1]$. Notice that condition (3) implies that

$$(5) \quad \sigma(t) \notin Q/H, \quad (t > 0).$$

The family F is not empty since we may take $H=Q$ in which case R/Q is just the arc $[0, a_0]$, in other words, R/Q is itself a one-parameter semigroup.

We introduce a partial order into F by saying that $(H, \sigma) \leq (H_1, \sigma_1)$ if

$$(a) \quad H_1 \subset H,$$

$$(b) \quad p\sigma_1(t) = \sigma(t), \quad (0 \leq t \leq a_0),$$

where p is the natural projection from R/H_1 to R/H . Notice that $p^{-1}(1)$ is isomorphic to H/H_1 . (The symbol 1 will be used indiscriminately to denote the identity of any semigroup with which we are dealing.)

Let $(H_\alpha, \sigma_\alpha)$ be a linearly ordered subfamily, and let $K = \bigcap_\alpha H_\alpha$. One sees easily that there is a one-parameter semigroup σ in R/K such that $(H_\alpha, \sigma_\alpha) \leq (K, \sigma)$ for all $(H_\alpha, \sigma_\alpha)$ in the subfamily. Indeed, let $R_\alpha = R/H_\alpha$. Then R/K is the inverse limit of the R_α , and σ may be taken to be the inverse limit of the $\sigma_\alpha(t)$. It is easily verified that (3) and (4) hold for σ . Now let $(H_\alpha, \sigma_\alpha)$ be a maximal linearly ordered subfamily. The proof of Theorem 1 will be complete if we can show that in this case $K = \{1\}$. This will follow from the following lemma.

LEMMA 7. *Let $(K, \sigma) \in F$. Let $K_1 \subset K$ be a closed subgroup such that K/K_1 is a Lie group. Then there is a one-parameter semigroup σ_1 in R/K_1 such that $(K, \sigma) \leq (K_1, \sigma_1)$.*

Proof. Let p denote the natural projection from R/K_1 to R/K . Let $R' = p^{-1}(\sigma([0, a_0]))$. Then R' is an adequate local semigroup contained in R/K_1 , and the maximal group of R' is isomorphic to K/K_1 . We may apply the result stated at the end of §2 to conclude that there is a one-parameter semigroup $\gamma: [0, 1] \rightarrow R'$ such that if $\gamma(a) = \gamma(b)g$ ($g \in p^{-1}(1)$), then $a = b$ and $g = 1$. Then $\gamma(1) \in \pi_1(\tau(a_1)Q)$ for some a_1 ($0 < a_1 \leq a_0$), where π_1 denotes the natural projection from R to R/K_1 .

Define $\sigma_1(t): [0, a_0] \rightarrow R'$ by

$$\sigma_1(t) = \gamma(t/a_1), \quad (0 \leq t \leq a_1),$$

$$\sigma_1(t) = (\sigma_1(a_1))^k \sigma_1(r), \quad (t \leq a_0, t = ka_1 + r).$$

Then $p\sigma_1(a_1) = \sigma(a_1)$. From this it follows easily that $p\sigma_1(t) = \sigma(t)$ ($0 \leq t \leq a_0$).

It only remains to show that σ_1 satisfies (4). Assume that $\sigma_1(a) = \sigma_1(b)g$ for some $g \in Q/K_1$. Applying p we have $\sigma(a) = \sigma(b)p(g)$. But (4) holds for σ and therefore $a = b$ and $p(g) = 1$. This means that $g \in p^{-1}(1)$, but γ was so chosen that this implies $g = 1$. This completes the proof.

4. Applications. An (I) -semigroup is a semigroup on an arc such that one endpoint is a zero and the other is an identity. The general structure is given in [4, Theorem B]. In particular, if an (I) -semigroup has no idempotents except the two endpoints, then it is isomorphic to one of the following:

- (1) $[0, 1]$ with the usual multiplication of real numbers,

(2) $[1/2, 1]$ with the product $x \cdot y$ defined by $x \cdot y = \max(1/2, xy)$, where xy denotes the ordinary multiplication of real numbers.

THEOREM 3. *Let S be a compact semigroup with zero and identity and no other idempotents, and assume that $H(1)$ is not an open set. Then S contains an (I) -subsemigroup J , and S is arcwise connected.*

For the proof of this theorem we shall need the following lemma.

LEMMA 8. *Let S be a compact semigroup with zero and identity and no other idempotents. Let $y \notin H(1)$. Then $yz^n \rightarrow 0$, and if $yz = y$ for any $z \in H(1)$ then $y = 0$.*

Proof. Let $\Gamma(y) = \{y, y^2, y^3, \dots\}$. This is a compact semigroup and therefore contains an idempotent. As in the proof of Lemma 1, this idempotent cannot be the identity. But it is known [3, Theorem 1] that if $0 \in \Gamma(y)$ then $yz^n \rightarrow 0$.

If $yz = y$, then $yz^2 = yz = y$, and so $yz^n = y$. But $z^n \rightarrow 0$ and so $y = 0$.

We do not know whether $yz = y$ is possible for some $z \neq 1$ in $H(1)$.

Proof of Theorem 3. Let σ be the one-parameter semigroup of Theorem 1. Define $\sigma(t)$ for $t \in [1, 2]$ by

$$(6) \quad \sigma(t) = \sigma(1)\sigma(t-1), \quad (1 \leq t \leq 2).$$

It is easy to see that if $a, b, a+b \in [0, 2]$ then $\sigma(a+b) = \sigma(a)\sigma(b)$, and that $\sigma(t) \in H(1)$ for $t > 0$.

Assume that $\sigma(a) = \sigma(b)$ ($a < b$). Then $\sigma(a) = \sigma(a)\sigma(b-a)$, and so by Lemma 8 $\sigma(a) = 0$. Thus either σ is one-to-one on $[0, 2]$, or $\sigma(a) = 0$ for some a .

We may successively extend σ into the intervals $[2, 3]$, $[3, 4]$, etc., so that σ is defined for $0 \leq t < \infty$. By Lemma 8, $\lim \sigma(t) = 0$ ($t \rightarrow \infty$). Thus, either σ is one-to-one on $[0, \infty]$, or there is a finite number a such that $\sigma(a) = 0$ and then $\sigma(t) = 0$ for all $t \geq a$. If we let a' be the smallest such a , then σ is one-to-one on $[0, a']$. In any case, $J = \sigma([0, \infty])$ is the required (I) -semigroup.

To show that S is arcwise connected we observe that if $x \neq 0$ then xJ is an arc from x to 0.

We conjecture that J is a homomorphic retract of S . This would imply that S has a "character" (a homomorphism onto an (I) -semigroup).

If S contains an additional idempotent the result need not be true, even when S is connected. For example, let S be the points of the form $(1 + 1/(1+x)) \exp(2\pi ix)$ ($0 \leq x < \infty$) together with the unit disc in the complex plane. We multiply as follows:

$$\begin{aligned} \left(1 + \frac{1}{1+x}\right) e^{2\pi ix} \cdot \left(1 + \frac{1}{1+y}\right) e^{2\pi iy} &= \left(1 + \frac{1}{1+x+y}\right) e^{2\pi i(x+y)}, \\ \rho_1 e^{2\pi ix} \cdot \rho_2 e^{2\pi iy} &= \rho_1 \rho_2 e^{2\pi i(x+y)}, & (\rho_1, \rho_2 \leq 1), \\ \left(1 + \frac{1}{1+x}\right) e^{2\pi ix} \cdot \rho e^{2\pi iy} &= \rho e^{2\pi i(x+y)}, \end{aligned}$$

and take multiplication to be commutative. This is a ray winding asymptotically about the unit disc. The identity ($2e^{2\pi i0}=2$) of the ray is an identity for S , while the zero of the unit disc is a zero for S . In this case there is no (I) -semigroup, in fact no arc at all, from zero to the identity, although a one-parameter semigroup does exist.

We now give another example to illustrate the possibilities when there are additional idempotents. In this example S is the Cantor set endowed with a multiplication which will be described presently. The endpoint 0 will be a zero, 1 will be an identity, and every neighborhood of the identity will contain additional idempotents.

Let $x \in S$. Then x admits a triadic expansion in which only the digits 0 and 2 are used:

$$(7) \quad x = \sum_1^{\infty} \epsilon_n / 3^n, \quad (\epsilon_n = 0 \text{ or } 2).$$

Let the indices in which two occurs be $n_1 < n_2 < \dots$, and let $J^* = \{1, 2, 3, \dots, \infty\}$. Define a function $f_x: J^* \rightarrow J^*$ by:

$$(8) \quad \begin{aligned} f_x(j) &= n_j, & (j \neq \infty), \\ f_x(\infty) &= \infty. \end{aligned}$$

This definition is not complete if the expansion (7) has only a finite number of two's. In this case the indices for which a two occurs have a greatest member, n_N . We complete the definition of f_x by

$$(8') \quad f_x(j) = \infty, \quad (j > N).$$

In any case the function f_x has the following property:

(P) f_x is strictly increasing as long as it is finite-valued; once it takes the value ∞ it is constantly equal to ∞ thereafter.

Conversely, let $g: J^* \rightarrow J^*$ satisfy (P), and let $x = \sum 2/3^{g(j)}$. Then $x \in S$ and $g = f_x$. So there is a one-to-one correspondence between the Cantor set S and the set of functions satisfying (P). But the functions form a semigroup under composition, and we take this as the multiplication in S (continuity is easily checked).

The idempotents are precisely the elements 1 and $e_n = 2/3 + 2/3^2 + \dots + 2/3^n$; they converge to 1. The multiplication is order-preserving on one side only: $x < y$ implies $cx \leq cy$ for all $c \in S$.

Another corollary to Theorems 1 and 2 is the following result (whose proof we omit).

THEOREM 4. *Let S be a locally compact semigroup with identity and assume S is an n -dimensional manifold with regular boundary in some neighborhood of 1. If the hypotheses of Theorem 2 are satisfied and if $\dim H(1) = n - 1$, then S can be locally imbedded in a Lie group.*

This result need not be true if $\dim H(1) < n-1$, as the following example shows.

Let I_1 be $[0, 1]$ with ordinary multiplication, and let $I_2 = [0, 1]$ with the multiplication $x \cdot y = \min(x, y)$. Let $S = \{(x, y) : x \in I_1, y \in I_2, y \geq x\}$ and give S the coordinatwise multiplication. Then S satisfies the hypotheses of Theorem 4 except that $\dim H(1) = 0$. However, every neighborhood of 1 contains elements for which the cancellation law fails, and so no neighborhood of the identity can be imbedded in a group.

5. Concluding remarks. The hypotheses of Theorem 1 are not necessary and sufficient for the existence of a one-parameter semigroup. Let us agree to say that an element x is *Archimedean* with respect to a neighborhood V of the identity ($x \in V$) if some power of x is not in V . We originally conjectured that if S is a compact semigroup with identity then a necessary and sufficient condition for the existence of a one-parameter semigroup (with $\sigma(t) \in H(1)$ for $t > 0$) is the existence of a neighborhood V of 1 and a net x_a in V , $x_a \rightarrow 1$, such that each element x_a is Archimedean with respect to V . The main difficulty in proving this seems to be in showing that Q is a group (Lemma 3). A. M. Gleason has given us an example of a semigroup with an Archimedean collection for which Q is not a group. Although this example does not necessarily kill the conjecture, it certainly throws doubt on it.

We further conjecture that if S is a locally compact connected semigroup with identity, and if there is a neighborhood of 1 containing no other idempotents, then S contains a one-parameter semigroup.

Finally, we note that R. J. Koch [6] has an unpublished result showing that every compact connected semigroup with identity contains a generalized arc, but an example of R. P. Hunter shows that there need not be any arc having the identity as one endpoint. These results will appear soon.

REFERENCES

1. A. M. Gleason, *Groups without small subgroups*, Ann. of Math. vol. 56 (1952) pp. 193-212.
2. J. L. Kelley, *General topology*, New York, Van Nostrand, 1955.
3. R. J. Koch, *On monothetic semigroups*, Proc. Amer. Math. Soc. vol. 8 (1957) pp. 397-401.
4. P. S. Mostert and A. L. Shields, *On the structure of semigroups on a compact manifold with boundary*, Ann. of Math. vol. 65 (1957) pp. 117-143.
5. A. D. Wallace, *The structure of topological semigroups*, Bull. Amer. Math. Soc. vol. 61 (1955) pp. 95-112.
6. R. J. Koch, *Arcs in partially ordered spaces*, Pacific J. Math. vol. 9 (1959) pp. 723-728.

TULANE UNIVERSITY,
NEW ORLEANS, LOUISIANA
UNIVERSITY OF MICHIGAN,
ANN ARBOR, MICHIGAN