

ON DIFFERENCE METHODS FOR THE SOLUTION OF A TRICOMI PROBLEM⁽¹⁾

BY

HAJIMU OGAWA

1. Introduction. The purpose of this paper is the investigation of approximate solutions to a Tricomi problem for the partial differential equation

$$(1.1) \quad Lu = K(y)u_{xx} + u_{yy} + a(x, y)u_x + b(x, y)u_y + c(x, y)u = f(x, y),$$

where K is a continuous function which is positive for y positive and negative for y negative. Thus the equation is elliptic, parabolic or hyperbolic according as $y > 0$, $y = 0$ or $y < 0$.

For $y < 0$ the equation (1.1) has real characteristics given by the two families

$$(1.2a) \quad dy/dx = (-K)^{-1/2},$$

$$(1.2b) \quad dy/dx = -(-K)^{-1/2}.$$

Let A and B be two points on the x -axis with $x_A < x_B$. By D we denote the open domain bounded by the characteristic Γ_1 of the family (1.2b) passing through the point A , the characteristic Γ_2 of the family (1.2a) passing through the point B , and by the simple arc Γ^+ in the upper half-plane with endpoints at A and B . (We will consider only bounded domains.)

The Tricomi problem for the equation (1.1) on D is the problem of finding a function u , continuous in $\bar{D}$, which satisfies the equation (1.1) in D , and which takes on the boundary values

$$(1.3) \quad u = \phi_1 \text{ on } \Gamma^+, \quad u = \phi_2 \text{ on } \Gamma_1,$$

where ϕ_1 and ϕ_2 are given functions. Such a problem was first solved by Tricomi [1] for the case $K(y) = y$, $a \equiv b \equiv c \equiv f \equiv 0$.

We approximate (1.1) by a difference equation

$$(1.4) \quad L_h U = f$$

in a function U which is defined on a mesh region D_h depending on the original domain D . The problem of solving the differential equation (1.1) with the boundary conditions (1.3) is replaced by the problem of solving the difference equation (1.4) with suitable boundary conditions, and investigating the behavior of U as the mesh size tends to zero. Filippov [2] proved that if the Tricomi problem for the equation

Presented to the Society, April 22, 1961; received by the editors December 29, 1960.

⁽¹⁾ This paper is part of a doctoral dissertation presented at the University of California, Berkeley. The author is indebted to his thesis director, Professor M. H. Protter, for his valuable advice and encouragement.

$$yu_{xx} + u_{yy} = f(x, y)$$

has a solution u in D which is sufficiently smooth, then the solution to an appropriate difference equation converges to u as the mesh size tends to zero. The techniques used in this paper, as in [2], depend essentially on determining under what conditions the solution of the difference equation satisfies a maximum principle, in analogy with the maximum principle for the differential equation. The latter may be stated in the following form: if $Lu \geq 0$ in D and u is a nondecreasing function of y on Γ_1 , then the maximum of u on $\bar{D}$, if non-negative, is attained on Γ^+ . Such a principle was first discovered by Germain and Bader [3] for the Tricomi equation

$$yu_{xx} + u_{yy} = 0,$$

and was extended by Agmon, Nirenberg and Protter [4] to the equation (1.1) with the coefficients satisfying a complicated set of inequalities. In the present paper, we find that solutions to the difference equation have the maximum property provided that the coefficients of the difference equation satisfy certain inequalities which are consequences of the conditions of [4] away from the x -axis. However, near the x -axis, conditions in addition to those of [4] are necessary in order that the difference equation have the maximum property.

Using the maximum principle we prove that, for a sufficiently fine mesh, the difference equation has a unique solution U for arbitrary functions f, ϕ_1 and ϕ_2 , and that U converges to the solution u of the differential equation (1.1) with boundary conditions (1.3), provided that u exists and is twice continuously differentiable in $\bar{D}$. In §6 we prove that under certain restrictions on K near the x -axis, and on the domain D^+ , the part of D for which $y \geq 0$, the regularity conditions on the derivatives of u at the boundary of the region may be weakened.

2. The difference problem. We assume that K is of class $C^3(\bar{D})$ with $K(y) > 0$ for $y > 0$ and $K(y) < 0$ and $K'(y) > 0$ for $y < 0$. Furthermore, we suppose that a and b are of class $C^1(\bar{D})$ and c and f are of class $C(\bar{D})$.

Integrating the equations (1.2) with respect to y , we obtain the characteristics of the differential equation in the form $x - x_0 = \pm G(y)$, where

$$(2.1) \quad G(y) = \int_y^0 [-K(\eta)]^{1/2} d\eta; \quad y \leq 0.$$

We divide the segment AB into N equal parts, each of length h , and through each of the points $x_k = x_A + kh$ ($k = 1, 2, \dots, N-1$) we draw the characteristics

$$x - x_k = \pm G(y).$$

These characteristics, together with the characteristics Γ_1 and Γ_2 , intersect at the points

$$(2.2) \quad \left(x_A + \frac{nh}{2} + kh, -y_n\right); \quad k = 0, 1, \dots, N-n; \quad n = 1, 2, \dots, N,$$

with the ordinates satisfying

$$G(-y_n) = nh/2; \quad n = 1, 2, \dots, N.$$

Taking for $y < 0$ the points given by (2.2), and for $y \geq 0$ the points of the form $(x_A + kh, mh)$ (k, m integers) which lie in $\bar{D}$, we obtain a mesh region $\bar{D}_h$. For $y > 0$, we define the neighbors of the mesh point (x, y) to be the four mesh points $(x+h, y)$, $(x-h, y)$, $(x, y+h)$ and $(x, y-h)$. We call the boundary Γ_h of $\bar{D}_h$ those points of $\bar{D}_h$ in the upper half-plane for which not all four neighboring points belong to $\bar{D}_h$, together with all points of $\bar{D}_h$ which lie on Γ_1 , and the point B . The totality of points of $\bar{D}_h$ which are not boundary points we call the interior mesh region D_h . Let D_h^- and D_h^+ consist of all points of D_h for which $y < 0$ and $y > 0$, respectively, and let Γ_h^- and Γ_h^+ be the points of Γ_h for which $y < 0$ and $y \geq 0$, respectively. Finally we let γ_h be the points in D_h for which $y = 0$.

We now introduce a difference operator L_h which acts on any function U defined on $\bar{D}_h$. For any point $(x, -y_n)$ of D_h^- we define

$$\begin{aligned} L_h U(x, -y_n) = & \frac{1}{\lambda_n \lambda_{n+1}} \left\{ \frac{2\lambda_{n+1}}{\lambda_n + \lambda_{n+1}} U\left(x - \frac{h}{2}, -y_{n-1}\right) \right. \\ & + \frac{2\lambda_n}{\lambda_n + \lambda_{n+1}} U\left(x - \frac{h}{2}, -y_{n+1}\right) - U(x-h, -y_n) - U(x, -y_n) \Big\} \\ (2.3) \quad & + a(x, -y_n) \frac{1}{h} \{ U(x, -y_n) - U(x-h, -y_n) \} \\ & + b(x, -y_n) \frac{1}{\lambda_n + \lambda_{n+1}} \left\{ U\left(x - \frac{h}{2}, -y_{n-1}\right) - U\left(x - \frac{h}{2}, -y_{n+1}\right) \right\} \\ & + c(x, -y_n) U(x, -y_n), \end{aligned}$$

where $\lambda_n = y_n - y_{n-1}$ and $y_0 = 0$. At a point (x, y) of D_h^+ we let

$$\begin{aligned} L_h U(x, y) = & K(y) \frac{1}{h^2} \{ U(x-h, y) - 2U(x, y) + U(x+h, y) \} \\ & + \frac{1}{h^2} \{ U(x, y-h) - 2U(x, y) + U(x, y+h) \} \\ (2.4) \quad & + a(x, y) \frac{1}{h} \{ U(x, y) - U(x-h, y) \} \\ & + b(x, y) \frac{1}{2h} \{ U(x, y+h) - U(x, y-h) \} + c(x, y) U(x, y), \end{aligned}$$

and at a point $(x, 0)$ of γ_h we define

$$\begin{aligned}
 L_h U(x, 0) = & \frac{2}{hy_2} \left\{ \frac{y_2}{h + y_2} U(x, h) + \frac{h}{h + y_2} U(x, -y_2) - U(x, 0) \right\} \\
 (2.5) \quad & + a(x, 0) \frac{1}{h} \{ U(x, 0) - U(x - h, 0) \} \\
 & + b(x, 0) \frac{1}{h + y_2} \{ U(x, h) - U(x, -y_2) \} + c(x, 0) U(x, 0).
 \end{aligned}$$

The problem of finding the solution u of the differential equation (1.1) in D subject to the boundary conditions (1.3) is replaced by the problem of finding the solution U of the difference equation

$$(2.6) \quad L_h U = f$$

on the region D_h , which satisfies the boundary conditions

$$\begin{aligned}
 (2.7) \quad & U = \phi_1 \text{ on } \Gamma_h^+, \\
 & U = \phi_2 \text{ on } \Gamma_h^-.
 \end{aligned}$$

We have assumed here that ϕ_1 is a function which is defined and continuous on the domain $\bar{D}^+ = \{(x, y) \in \bar{D} \mid y \geq 0\}$. The equations (2.6) and (2.7) form a system of linear algebraic equations in the unknown values of U at the points of D_h , in which the number of equations is equal to the number of unknowns.

The following result establishes a relation between the difference operator L_h and the differential operator L for functions of class $C^2(\bar{D})$.

THEOREM 2.1. *Let u be of class $C^2(\bar{D})$. Then on D_h , $L_h u \rightarrow Lu$ uniformly as $h \rightarrow 0$.*

Proof. Using Taylor's theorem in a neighborhood of the point $(x, -y_n)$ of D_h^- , we find from (1.1) and (2.3) that for $y < 0$,

$$\begin{aligned}
 |Lu - L_h u| \leq & \left| K(-y_n) + \frac{h^2}{4\lambda_n \lambda_{n+1}} \right| |u_{xx}| + \frac{h^2}{2\lambda_n \lambda_{n+1}} \epsilon_1 + \frac{2h}{\lambda_n + \lambda_{n+1}} \epsilon_2 + \epsilon_3 \\
 (2.8) \quad & + \frac{h}{2} \left\{ |a| + \frac{h}{2(\lambda_n + \lambda_{n+1})} |b| \right\} |\bar{u}_{xx}| + \frac{h}{2} |b| |\bar{u}_{xy}| \\
 & + \frac{\lambda_n^2 + \lambda_{n+1}^2}{2(\lambda_n + \lambda_{n+1})} |b| |\bar{u}_{yy}|,
 \end{aligned}$$

where all unbarred functions are evaluated at $(x, -y_n)$, and the barred partial derivatives denote values of the functions at points $(\bar{x}, \bar{y})$ which satisfy $x - h < \bar{x} < x$, $-y_{n+1} < \bar{y} < -y_{n-1}$. The quantities ϵ_1 , ϵ_2 and ϵ_3 are the moduli of

continuity of u_{xx} , u_{xy} and u_{yy} , respectively. In the same way we obtain for $y > 0$, the estimate

$$(2.9) \quad |Lu - L_h u| \leq K\epsilon_1 + \left(1 + \frac{h}{4} |b|\right) \epsilon_3 + \frac{h}{2} |a| |\bar{u}_{xx}|$$

and for $y = 0$,

$$(2.10) \quad |Lu - L_h u| \leq \epsilon_3 + \frac{h}{2} |a| |\bar{u}_{xx}| + \frac{1}{2} |b| (h + y_2) |\bar{u}_{yy}|.$$

By hypothesis, K , a , b , u_{xx} and u_{yy} are uniformly bounded on $\bar{D}$. Moreover, as $h \rightarrow 0$, y_2 , ϵ_1 and ϵ_3 tend to zero, uniformly on $\bar{D}$. Hence (2.9) and (2.10) imply that $L_h u$ tends uniformly to Lu as $h \rightarrow 0$, for $y \geq 0$. For $y < 0$ we must evaluate the coefficients appearing in the estimate (2.8). For this it will be convenient to introduce the inverse function to G . More precisely, if $x = G(y)$ is the equation of the characteristic of the family (1.2b) passing through the origin, then we can express it also as

$$(2.11) \quad y = -H(x).$$

The relation between H and K can be obtained by differentiating (2.11) to obtain

$$(2.12) \quad H'(x) = [-K(y)]^{-1/2}$$

at each point (x, y) on the curve (2.11). Hence H , which is defined for $0 \leq x \leq (x_B - x_A)/2$, is an increasing function with $H(0) = 0$, and H' is a decreasing function with $H'(0) = +\infty$. That is, $H(x) > 0$, $H'(x) > 0$ and $H''(x) < 0$ for $x > 0$. Moreover, since we assume that K has three continuous derivatives, H will have four continuous derivatives for $x > 0$.

Let us now return to (2.8). First we note that

$$(2.13) \quad \lambda_n = y_n - y_{n-1} = H\left(\frac{nh}{2}\right) - H\left(\frac{nh-h}{2}\right) = \frac{h}{2} H'\left(\frac{nh-\theta h}{2}\right),$$

with $0 < \theta < 1$. Since $H(x) \rightarrow 0$ as $x \rightarrow 0$ we may choose $\delta > 0$ so small that $H(\delta) < \epsilon/2$. Then for all n and h such that $nh/2 \leq \delta$, we have $\lambda_n < \epsilon$. On the other hand, for $x \geq \delta/2$, $H'(x)$ is uniformly bounded. Therefore, we may choose h so small that for $nh/2 > \delta$, $\lambda_n < \epsilon$. Hence for all n , $\lambda_n \rightarrow 0$ uniformly as $h \rightarrow 0$. Using the relations (2.12) and (2.13) we find that

$$K(-y_n) + \frac{h^2}{4\lambda_n\lambda_{n+1}} = K(-y_n) + [-K(-y_{n-\alpha})]^{1/2}[-K(-y_{n+\beta})]^{1/2}$$

where $y_{n-1} < y_{n-\alpha} < y_n < y_{n+\beta} < y_{n+1}$. Therefore since K is continuous, the expression $h^2/2\lambda_n\lambda_{n+1}$ is uniformly bounded as $h \rightarrow 0$ and

$$K(-y_n) + \frac{h^2}{4\lambda_n\lambda_{n+1}} \rightarrow 0$$

as $h \rightarrow 0$. Also, since $\lambda_n > \lambda_{n+1}$, we have

$$\frac{\lambda_n^2 + \lambda_{n+1}^2}{2(\lambda_n + \lambda_{n+1})} < \frac{\lambda_n}{2}.$$

Hence, each term on the right side of (2.8) can be made arbitrarily small by choosing h sufficiently small. This completes the proof of the theorem.

3. Maximum principles. To establish a maximum principle for the difference equation on the region D_h , we first prove maximum principles on each of the regions D_h^- and D_h^+ separately. Let us take as the boundary of D_h^- the set $\Gamma_h^- \cup \gamma_h$, and as the boundary of D_h^+ the set $\Gamma_h^+ \cup \gamma_h$. We then denote by $\bar{D}_h^-$ the set D_h^- plus its boundary, and by $\bar{D}_h^+$ the set D_h^+ plus its boundary.

From (2.2) we note that each point $(x, -y_n)$ of $\bar{D}_h^-$ may be uniquely represented by a pair of indices (k, n) , where the first denotes the negatively sloping characteristic and the second the ordinate corresponding to the point. In this notation the difference operator (2.3) takes the form

$$(3.1) \quad L_h U_{k,n} = \frac{1}{\lambda_n \lambda_{n+1}} [(1 - A_{k,n}) U_{k,n-1} + (1 + A_{k,n}) U_{k-1,n+1} \\ - (1 + \alpha_{k,n}) U_{k-1,n} - (1 - \alpha_{k,n} - \gamma_{k,n}) U_{k,n}],$$

in which we have defined

$$(3.2) \quad A_{k,n} = \frac{\lambda_n - \lambda_{n+1}}{\lambda_n + \lambda_{n+1}} - \frac{\lambda_n \lambda_{n+1}}{\lambda_n + \lambda_{n+1}} b(x, -y_n),$$

$$(3.3) \quad \alpha_{k,n} = \frac{\lambda_n \lambda_{n+1}}{h} a(x, -y_n),$$

$$(3.4) \quad \gamma_{k,n} = \lambda_n \lambda_{n+1} c(x, -y_n).$$

THEOREM 3.1. Let $L_h U \geq 0$ on D_h^- and

$$(3.5) \quad U_{0,n+1} \leq U_{0,n}; \quad n \geq 1.$$

Assume that the conditions

$$(3.6) \quad \gamma_{k,n} \leq 0,$$

$$(3.7) \quad 1 + \alpha_{k,n} > 0,$$

$$(3.8) \quad A_{k,n} \leq 1,$$

$$(3.9) \quad A_{k,n} - \alpha_{k,n} > 0,$$

$$(3.10) \quad A_{k+1,n-1} - A_{k,n} - A_{k,n} A_{k+1,n-1} - \alpha_{k+1,n-1} + \alpha_{k,n} + \alpha_{k,n} \alpha_{k+1,n-1} \\ + (1 + \alpha_{k+1,n-1}) \gamma_{k,n} \geq 0,$$

are satisfied at all points (k, n) of D_h^- . Then the maximum of U on $\bar{D}_h^-$, if non-negative, is attained on the boundary. If $\gamma_{k,n} \equiv 0$, then the result holds without the requirement that the maximum be non-negative.

Proof. Assume that the maximum of U occurs at an interior point (k, n) , and that $U_{k,n}$ is greater than the maximum of U on the boundary. We consider the set of points in $\bar{D}_h^-$ which lie on the positively sloping characteristics passing through the points (k, n) and $(k, n-1)$. Using (3.1) to write $L_h U_{k,n} \geq 0$ we have

$$(1 + \alpha_{k,n})(U_{k-1,n+1} - U_{k-1,n}) \geq (1 - A_{k,n})(U_{k,n} - U_{k,n-1}) \\ + (A_{k,n} - \alpha_{k,n})(U_{k,n} - U_{k-1,n+1}) - \gamma_{k,n}U_{k,n} \geq 0,$$

the second inequality following from the conditions (3.6), (3.8) and (3.9), together with the assumption that $U_{k,n}$ is the maximum and is non-negative. Next we assert that

$$(3.11) \quad (1 + \alpha_{k-j+1,n+j-1})(U_{k-j,n+j} - U_{k-j,n+j-1}) \\ \geq (A - \alpha)_{k-j+1,n+j-1}(U_{k,n} - U_{k-j,n+j}) \geq 0,$$

for $j=1, 2, \dots, k$. The second inequality of (3.11) follows from (3.9) and the assumption that $U_{k,n}$ is the maximum. The first inequality we have proved for $j=1$, hence we proceed by induction. We will show that if (3.11) holds for any j with $1 \leq j \leq k-1$, then it holds also for $j+1$. From $L_h U_{k-j,n+j} \geq 0$, we have

$$(1 + \alpha_{k-j,n+j})(U_{k-j-1,n+j+1} - U_{k-j-1,n+j}) \geq (1 - A_{k-j,n+j})(U_{k-j,n+j} - U_{k-j,n+j-1}) \\ + (A - \alpha - \gamma)_{k-j,n+j}U_{k-j,n+j} - (A - \alpha)_{k-j,n+j}U_{k-j-1,n+j+1}.$$

Multiplying (3.11) by the quantity

$$(A - \alpha - \gamma)_{k-j,n+j}/(A - \alpha)_{k-j+1,n+j-1}$$

and adding the resulting inequality to the above inequality, we obtain

$$(1 + \alpha_{k-j,n+j})(U_{k-j-1,n+j+1} - U_{k-j-1,n+j}) \\ \geq (A - \alpha)_{k-j,n+j}(U_{k,n} - U_{k-j-1,n+j+1}) - \gamma_{k-j,n+j}U_{k,n} \\ + \left\{ (1 - A_{k-j,n+j}) - (A - \alpha - \gamma)_{k-j,n+j} \left(\frac{1 + \alpha}{A - \alpha} \right)_{k-j+1,n+j-1} \right\} \\ \cdot (U_{k-j,n+j} - U_{k-j,n+j-1}).$$

The second term on the right side is non-negative by (3.6) and the assumption that $U_{k,n}$, the maximum, is non-negative. In the last term on the right side, the first factor is non-negative by (3.9) and (3.10), while the second factor is non-negative by the induction hypothesis (3.11). Thus (3.11) holds for $j+1$ and hence for $j=1, 2, \dots, k$.

For $j=k$, the first inequality of (3.11) can be written

$$U_{0,n+k} - U_{0,n+k-1} \geq \left(\frac{A - \alpha}{1 + \alpha} \right)_{1,n+k-1} (U_{k,n} - U_{0,n+k}).$$

Since it was assumed that the maximum does not occur on the boundary, the second factor on the right side is positive. From this we conclude that $U_{0,n+k} > U_{0,n+k-1}$ for some $k \geq 1$, $n \geq 1$, which contradicts the hypothesis (3.5). Hence the maximum must be attained on the boundary. The result for $\gamma_{k,n} \equiv 0$ follows by setting $\gamma_{k,n} \equiv 0$ in the proof.

Let us now consider the region D_h^+ , for which we have the following maximum principle.

THEOREM 3.2. *Let $L_h U \geq 0$ on D_h^+ . Assume that the conditions*

$$(3.12) \quad K(y) - ha(x, y) > 0,$$

$$(3.13) \quad 1 \pm \frac{h}{2} b(x, y) > 0,$$

$$(3.14) \quad c \leq 0,$$

are satisfied on D_h^+ . Then the maximum of U on $\bar{D}_h^+$, if non-negative, is attained on the boundary. If $c \equiv 0$, then the result holds without requiring that the maximum be non-negative.

Proof. Let $M \geq 0$ denote the maximum value of U on $\bar{D}_h^+$ and suppose that M is greater than the maximum of U on the boundary. Then there exists a point (x, y) in D_h^+ at which $U = M$, and for at least one of its four neighboring points $U < M$. Then, referring to the equation (2.4), it follows from the conditions (3.12), (3.13) and (3.14) that $L_h U(x, y) < 0$, contrary to hypothesis. Thus the maximum must be attained on the boundary.

COROLLARY. *Let $L_h U \geq 0$ on D_h^+ , $U \leq 0$ on Γ_h^+ and $U \leq M$ on γ_h with $M > 0$. Suppose the conditions (3.12), (3.13) and (3.14) are satisfied on D_h^+ . If $(x, 0)$ is a point of γ_h at which $U = M$, then $U(x, 0) > U(x, h)$.*

Proof. By Theorem 3.2, $U \leq M$ throughout D_h^+ . If (x, h) is a boundary point of $\bar{D}_h^+$, then $U(x, h) \leq 0$ by hypothesis, and the theorem is trivially satisfied. Therefore we can assume that (x, h) is in D_h^+ . Suppose that $U(x, h) = M$. Then if $c(x, h) < 0$ we find immediately from (2.4) that $L_h U(x, h) < 0$, contrary to hypothesis. On the other hand, if $c(x, h) = 0$, we must have $L_h U(x, h) = 0$ which implies that $U = M$ at each of the four neighbors of the point (x, h) . In particular, $U(x, 2h) = M$. Repeating this argument a finite number of times, we either contradict the assumption that $L_h U \geq 0$ on D_h^+ , or we obtain $U = M$ at some point of Γ_h^+ , contrary to hypothesis.

We may now state and prove the following maximum principle for the whole domain D_h .

THEOREM 3.3. *Let $L_h U \geq 0$ on D_h with U satisfying (3.5) on Γ_h^- . Assume*

that the conditions (3.6) through (3.10) are satisfied on D_h^- , and the conditions (3.12) through (3.14) are satisfied on D_h^+ . Furthermore, suppose that on γ_h ,

$$(3.15) \quad a(x, 0) \leq 0,$$

$$(3.16) \quad 1 + \frac{h}{2} b(x, 0) > 0,$$

$$(3.17) \quad 1 - \frac{y_2}{2} b(x, 0) > 0,$$

$$(3.18) \quad c(x, 0) \leq 0.$$

Then the maximum of U on $\bar{D}_h$, if non-negative, is attained on the boundary. If $c \equiv 0$, then the result holds without the requirement that the maximum be non-negative.

Proof. Let M be the maximum value of U on the boundary Γ_h . Then in order that the theorem be meaningful, we must require that $M \geq 0$. Let M_1 be the maximum of U on γ_h and suppose that $M_1 > M$. Then the function $V = U - M$ satisfies $L_h V \geq 0$ on D_h , $V \leq 0$ on Γ_h , and $V \leq M_1 - M$ on γ_h with $M_1 - M > 0$. Let $(x, 0)$ be a point of γ_h where $U = M_1$. Then by the corollary to Theorem 3.2, $V(x, h) < M_1 - M$, which means that $U(x, h) < M_1$. Moreover, since V also satisfies (3.5), we have $U \leq M_1$ on D_h^- by Theorem 3.1. By virtue of the conditions (3.15) through (3.18), the formula (2.5) gives us $L_h U(x, 0) < 0$. This however contradicts the assumption that $L_h U \geq 0$ on D_h . Hence we must have $M_1 \leq M$. Then by Theorems 3.1 and 3.2 we immediately obtain $U \leq M$ throughout D_h .

Since our difference equation was obtained from the differential equation (1.1), it is desirable that the conditions (3.6) through (3.10) and (3.12) through (3.18) be translated into conditions on the coefficients of the differential equation. It is clear that because the coefficients are assumed to be continuous on $\bar{D}$, the conditions (3.14) and (3.18) are satisfied if and only if $c \leq 0$ on $\bar{D}^+$, while (3.13), (3.16) and (3.17) are satisfied if h is sufficiently small. If we also assume that there is a $\delta > 0$, such that $a(x, y) \leq 0$ for $0 \leq y \leq \delta$, then the conditions (3.12) and (3.15) will be satisfied for h sufficiently small, since K is uniformly positive for $y \geq \delta$. The conditions in D_h^- are investigated in the following.

THEOREM 3.4. *If*

$$(3.19) \quad c \leq 0 \text{ on } \bar{D}^-,$$

and

$$(3.20) \quad ya(x, y)[-K(y)]^{-1/2} \rightarrow 0 \text{ as } y \rightarrow 0,$$

uniformly on $\bar{D}^-$, then for h sufficiently small, the conditions (3.6), (3.7) and (3.8) are satisfied.

Proof. (3.6) follows immediately from (3.19) and the continuity of c . Recalling the definition of $\alpha_{k,n}$, we have

$$|\alpha_{k,n}| = \left| \frac{\lambda_n \lambda_{n+1}}{h} a(x, -y_n) \right| \leq |y_n a(x, -y_n) [-K(-y_n)]^{-1/2}|$$

using (2.12) and (2.13). Therefore, by the condition (3.20), there exists a $\delta_1 > 0$ such that $1 + \alpha_{k,n} > 0$ for all y_n with $y_n \leq \delta_1$. For $y_n > \delta_1$, the quantity

$$\lambda_{n+1}/h \leq \frac{1}{2} [-K(-y_n)]^{-1/2}$$

is uniformly bounded. Thus since $\lambda_n \rightarrow 0$ as $h \rightarrow 0$ and a is uniformly bounded on $\bar{D}^-$, $\alpha_{k,n} \rightarrow 0$ as $h \rightarrow 0$ for each n such that $y_n > \delta_1$. This proves that (3.7) is satisfied on all of D_h^- provided only that h is small.

Referring to (3.2) we may write

$$1 - A_{k,n} = \frac{2\lambda_{n+1}}{\lambda_n + \lambda_{n+1}} \left[1 + \frac{1}{2} \lambda_n b(x, -y_n) \right].$$

Since $\lambda_n \rightarrow 0$ as $h \rightarrow 0$ and b is uniformly bounded on $\bar{D}^-$, (3.8) will be satisfied for h sufficiently small.

THEOREM 3.5. *Let (3.20) hold on D^- , and let*

$$(3.21) \quad \frac{d}{dy} [(-K)^{1/2}] + a + b(-K)^{1/2} < 0$$

for $y < 0$ on $\bar{D}^-$. Furthermore, assume that the function H defined by (2.11) has a derivative of the form

$$(3.22) \quad H'(x) = x^{-\alpha} \tilde{H}(x)$$

near $x=0$, where $0 < \alpha < 1$ and $\tilde{H}$ is a function having three continuous derivatives with $\tilde{H}(x) \geq m > 0$ for $x \geq 0$. Then (3.9) is satisfied on D_h^- for h sufficiently small.

Proof. From (3.20) we obtain for nh sufficiently small

$$y_n \geq \frac{m}{1-\alpha} \left(\frac{nh}{2} \right)^{1-\alpha}, \quad n \geq 1; \quad \lambda_n < \frac{Mh}{2} \left(\frac{nh-h}{2} \right)^{-\alpha}, \quad n \geq 2;$$

where $\tilde{H}(x) \leq M$. Furthermore, by writing

$$\lambda_n - \lambda_{n+1} = \int_{(n-1)h/2}^{nh/2} \left[H'(x) - H' \left(x + \frac{h}{2} \right) \right] dx,$$

and noting that $H''(x) \leq (m\alpha/2)x^{-\alpha-1}$ for x sufficiently small, we find that for $n \geq 2$,

$$\lambda_n - \lambda_{n+1} > \frac{m\alpha h^2}{8} \left(\frac{nh + h}{2} \right)^{-\alpha-1}.$$

Combining these estimates, we obtain for $n \geq 2$,

$$y_n(\lambda_n - \lambda_{n+1})/\lambda_n\lambda_{n+1} > C_0 > 0,$$

with C_0 a constant. On the other hand, we have

$$\lambda_1 \leq \frac{M}{1-\alpha} \left(\frac{h}{2} \right)^{1-\alpha}, \quad \lambda_2 \leq \frac{M}{1-\alpha} h^{1-\alpha}.$$

If $0 \leq x \leq h/2$, (3.20) yields the estimate

$$H'(x) - H'\left(x + \frac{h}{2}\right) \geq C_1 x^{-\alpha},$$

where C_1 is a positive constant. This in turn, implies that

$$\lambda_1 - \lambda_2 \geq \frac{C_1}{1-\alpha} \left(\frac{h}{2} \right)^{1-\alpha}.$$

Combining these estimates we obtain

$$y_1(\lambda_1 - \lambda_2)/\lambda_1\lambda_2 > C_2 > 0.$$

Therefore, for all $n \geq 1$, the quantity $y_n(\lambda_n - \lambda_{n+1})/\lambda_n\lambda_{n+1}$ is greater than a fixed positive number provided that h and y_n are sufficiently small. We also have

$$\left| y_n \frac{\lambda_n + \lambda_{n+1}}{h} a(x, -y_n) \right| \leq \left| \frac{y_n a(x, -y_n)}{2(-K(-y_n))^{1/2}} \right| \frac{M}{m} \frac{2^\alpha}{1-\alpha}.$$

Thus we may choose h and y_n so small, say $y_n \leq \delta$, $\delta > 0$, that (3.9) is satisfied.

For $y_n > \delta$, we choose h so small that $y_{n-2} \geq \delta/2$. Then for $x \geq (nh - 2h)/2$ the function H has three continuous and uniformly bounded derivatives. We may then apply Taylor's theorem to the expression $A_{k,n} - \alpha_{k,n}$ to obtain

$$\begin{aligned} A_{k,n} - \alpha_{k,n} &= -\frac{1}{\lambda_n + \lambda_{n+1}} \frac{h^2}{4} (H_n')^3 \\ &\quad \cdot \left\{ \frac{H_n''}{(H_n')^3} + a(x, -y_n) + \frac{1}{H_n'} b(x, -y_n) + O(h) \right\} \\ &= -\frac{1}{\lambda_n + \lambda_{n+1}} \frac{h^2}{4} \left\{ -K(-y_n) \right\}^{-3/2} \left\{ \frac{d}{dy} (-K(-y_n))^{1/2} \right. \\ &\quad \left. + a(x, -y_n) + b(x, -y_n)(-K(-y_n))^{1/2} + O(h) \right\}, \end{aligned}$$

where $H_n = H(nh/2)$. Because of (3.22), the expression in the brackets can be made negative by choosing h small if $y_n > \delta$. This proves that (3.9) is satisfied for $y_n > \delta$.

It is of interest to ask how restrictive the condition (3.22) is on the function H . A partial answer is given by considering the class of functions

$$K(y) = -(-y)^m; \quad y \leq 0,$$

with m a positive integer. The corresponding function H' is given by

$$H'(x) = \left(\frac{m+2}{2} x \right)^{-m/(m+2)}$$

and thus this function satisfies (3.22) with $\alpha = m/(m+2)$ and $\tilde{H}(x) = (1-\alpha)^\alpha$.

THEOREM 3.6. *If for $y < 0$ on $\bar{D}^-$ the condition*

$$(3.23) \quad \begin{aligned} & 2(-K)^{1/4} \frac{d^2}{dy^2} \{(-K)^{-1/4}\} - 2 \frac{d}{dy} \{(-K)^{-1/2}\} a - \frac{1}{2K} a^2 \\ & - a_x - (-K)^{-1/2} a_y - \frac{1}{2} b^2 - (-K)^{1/2} b_x - b_y + 2c > 0 \end{aligned}$$

is satisfied, then for each $\delta > 0$ and for h sufficiently small, the condition (3.10) holds at each point of D_h^- where $y_n \geq \delta$.

Proof. Fix $\delta > 0$ and choose h so small that if $y_n \geq \delta$, then $y_{n-2} \geq \delta/2$. Application of Taylor's theorem then yields for the left side of (3.10),

$$\begin{aligned} & [(\lambda_{n-1} + \lambda_n)(\lambda_n + \lambda_{n+1})]^{-1} \frac{h^4}{8} (H'_n)^4 \left[-2(H'_n)^{-3/2} \frac{d^2}{dx^2} (H'_n)^{-1/2} + 2(H'_n)^{-1} H''_n a \right. \\ & \left. + \frac{1}{2} (H'_n)^2 a^2 - a_x - H'_n a_y - \frac{1}{2} b^2 - (H'_n)^{-1} b_x - b_y + 2c + O(h) \right], \end{aligned}$$

where a, a_x, a_y, b, b_x, b_y and c denote values of these functions at $(x, -y_n)$. Using (2.12) to replace H by K in the above expression we obtain for the left side of (3.10),

$$\begin{aligned} & [(\lambda_{n-1} + \lambda_n)(\lambda_n + \lambda_{n+1})]^{-1} \frac{h^4}{8K^2} \left\{ 2(-K)^{1/4} \frac{d^2}{dy^2} [(-K)^{-1/4}] - 2 \frac{d}{dy} [(-K)^{-1/2}] a \right. \\ & \left. - \frac{1}{2K} a^2 - a_x - (-K)^{-1/2} a_y - \frac{1}{2} b^2 - (-K)^{1/2} b_x - b_y + 2c + O(h) \right\}. \end{aligned}$$

Thus if we choose h sufficiently small, the condition (3.23) will assure the validity of (3.10) for $y_n \geq \delta$.

It is of interest to observe that the conditions (3.19), (3.21) and (3.23) are essentially the conditions under which the solution of the differential

equation (1.1) in D^- satisfies a maximum principle, as found by Agmon, Nirenberg and Protter [4].

4. The existence of the solution to the difference equation. For convenience, let us denote by the *Conditions A* the set of conditions under which the maximum principle, Theorem 3.3, holds. That is, we denote by *Conditions A*:

- (a) (3.6), (3.7), (3.8), (3.9) and (3.10) hold on D_h^- ,
- (b) (3.12), (3.13) and (3.14) hold on D_h^+ ,
- (c) (3.15), (3.16), (3.17) and (3.18) hold on γ_h .

THEOREM 4.1. *Let the Conditions A be satisfied. Then there exists a unique solution U to the difference equation (2.6) on D_h which satisfies the boundary conditions (2.7), for any given values of f , ϕ_1 and ϕ_2 .*

Proof. Let V be a solution of the homogeneous system of equations,

$$(4.1) \quad \begin{aligned} L_h V &= 0 \text{ on } D_h, \\ V &= 0 \text{ on } \Gamma_h. \end{aligned}$$

Then by Theorem 3.3, $V \leq 0$ on D_h . But we may also apply Theorem 3.3 to the function $-V$ to obtain $V \geq 0$ on D_h . Thus V must vanish identically on the whole domain D_h ; i.e., the homogeneous system has only the trivial solution $V \equiv 0$. But this implies that the system (2.6) and (2.7) has a unique solution for arbitrary right sides, i.e., for arbitrary values of f , ϕ_1 and ϕ_2 .

We now show that the system of linear equations (2.6) and (2.7) can be solved by means of the Gauss-Seidel iteration procedure. For this we number the P points of D_h in the following order. First we take the points of D_h^+ with the largest ordinate and number them in any order. Then we number the points on the next row down in any order, and continue this process until we have numbered all the points of D_h^+ and γ_h . Next we number the points $(1, n)$ of D_h^- in order of increasing n , then the points $(2, n)$ in order of increasing n , and so on. If we denote the value of U at the point i ($i=1, 2, \dots, P$) by U_i , and solve each of the equations

$$L_h U_i = f_i$$

for U_i , we obtain a linear system of equations of the form

$$U_i = \sum_{j=1}^{i-1} r_{ij} U_j + \sum_{j=i+1}^P r_{ij} U_j + s_i; \quad i = 1, 2, \dots, P.$$

In the Gauss-Seidel procedure, an arbitrary zeroth order approximation $U_i^{(0)}$, $i=1, 2, \dots, P$, to the solution is chosen, and successive approximations are calculated by the formula

$$U_i^{(m+1)} = \sum_{j=1}^{i-1} r_{ij} U_j^{(m+1)} + \sum_{j=i+1}^P r_{ij} U_j^{(m)} + s_i,$$

where $U_i^{(m)}$ denotes the m th approximation to U_i .

THEOREM 4.2. *Let the Conditions A be satisfied. Then for h sufficiently small and for arbitrary values $U_i^{(0)}$, $i=1, 2, \dots, P$, $U_i^{(m)} \rightarrow U_i$ as $m \rightarrow \infty$.*

Proof. If we let $V_i^{(m)}$ be the error in the m th approximation,

$$V_i^{(m)} = U_i - U_i^{(m)},$$

then the error terms satisfy the homogeneous system

$$(4.2) \quad V_i^{(m+1)} = \sum_{j=1}^{i-1} r_{ij} V_j^{(m+1)} + \sum_{j=i+1}^P r_{ij} V_j^{(m)}.$$

That is, $V_i^{(m)}$ can be considered the m th approximation to the solution of the homogeneous system (4.1), starting from an arbitrary zeroth order approximation. Assume that

$$V_i^{(m)} \leq M; \quad i = 1, 2, \dots, P,$$

for some $M > 0$. Then referring to (2.4), we see that since the point (x, y) which corresponds to $i=1$ is on the highest row of D_h^+ , the value $V_1^{(m+1)}$ is related to the values of $V^{(m)}$ at $(x-h, y)$, $(x+h, y)$ and $(x, y-h)$ at most. Furthermore, the conditions (3.12) and (3.14) imply that the coefficient of $U(x, y)$ in (2.4) is negative. Hence we have

$$V_1^{(m+1)} \leq \left\{ 1 - \frac{1 + hb/2 - h^2c}{2K + 2 - ha - h^2c} \right\} M.$$

By choosing h sufficiently small, the fraction on the right side can be made greater than a fixed number ρ with $0 < \rho < 1$. We conclude that

$$V_1^{(m+1)} \leq (1 - \rho)M$$

for h sufficiently small. By induction, this implies that for any point on the highest row of D_h we have the estimate

$$V_i^{(m+1)} \leq (1 - \rho)M.$$

We now consider the first point in the next row down, say the k th point. $V_k^{(m+1)}$ is related to the value of $V^{(m+1)}$ at the point immediately above (if that point belongs to D_h) and to the values of $V^{(m)}$ at the other adjacent points. Therefore, (2.4) gives us

$$V_k^{(m+1)} \leq (1 - \rho^2)M.$$

By induction we may conclude that this relation holds on the entire second row. Continuing this process, we arrive at the estimate

$$(4.3) \quad V_i^{(m+1)} \leq (1 - \rho^r)M,$$

at each point of D_h^+ , where r is the number of rows in D_h^+ . On γ_h , we use (2.5) to obtain

$$(4.4) \quad V_i^{(m+1)} \leq (1 - \rho_0^{r+1})M,$$

where $0 < \rho_0 \leq \rho$. Hence, by virtue of (4.3), (4.4) also holds on D_h^+ .

Due to the manner in which the points of D_h^- were ordered, the second sum does not appear in the formula (4.2) for $V_i^{(m+1)}$. But this means that $V_i^{(m+1)}$ satisfies the difference equation

$$L_h V_i^{(m+1)} = 0$$

on D_h^- , with $V^{(m+1)} \equiv 0$ on Γ_h^- . Therefore, by Theorem 3.1, the maximum of $V^{(m+1)}$ is attained on the boundary of D_h^- . That is, the bound (4.4) holds throughout D_h^- , and therefore throughout D_h . Since we may carry out the same procedure for $-V_i^{(m+1)}$, we obtain finally,

$$|V_i^{(m+1)}| \leq (1 - \rho_0^{r+1})M.$$

Thus if we now choose an arbitrary zeroth order approximation $U_i^{(0)}$, then

$$|V_i^{(m)}| \leq (1 - \rho_0^{r+1})^m M_0$$

where

$$M_0 = \max_i |U_i - U_i^{(0)}|.$$

Hence this shows that $V_i^{(m)} \rightarrow 0$ as $m \rightarrow \infty$.

5. A priori bounds. The maximum principle may also be employed to establish a priori bounds for functions defined on $\overline{D}_h$.

THEOREM 5.1. *Let the Conditions A be satisfied. Let U be any function defined on $\overline{D}_h$ such that the quantities*

$$B_1 = \max_{\Gamma_h} |U|, \quad B_2 = \max_{\Gamma_h^-} |U_{\bar{\tau}}|, \quad B_3 = \max_{D_h} |L_h U|,$$

are finite, where

$$U_{\bar{\tau}_{0,n}} = (U_{0,n} - U_{0,n+1})/\lambda_{n+1}.$$

Then for h sufficiently small,

$$(5.1) \quad |U| \leq B_1 + C(B_2 + B_3)$$

on D_h , where C is a finite, non-negative constant depending only on the domain D and the functions b and c .

Proof. Let

$$Y = \sup_D |y|.$$

We now define a mesh function E on D_h by the equations

$$\begin{aligned} E(-y_n) &= (1 - \mu\lambda_n)E(-y_{n-1}) \text{ on } D_h^-, \\ E(y) &= (1 + 2\mu h)E(y - h) \text{ for } y = nh; \quad n = 1, 2, \dots, \\ E(0) &= e^{2\mu Y}, \end{aligned}$$

where μ is a positive constant to be chosen. Let us choose h so small that $\mu h \leq 1/2$ and $\mu\lambda_n \leq 1/2$ for all n . Then E is a positive, nondecreasing function of y . Therefore, since on D_h^+ we have

$$E(nh) = (1 + 2\mu h)^n e^{2\mu Y} \leq e^{4\mu Y},$$

E is uniformly bounded on D_h , independent of h . To find a lower bound for E , we first note that due to our choice of h small, the inequalities

$$1 - \mu\lambda_n \geq 1/(1 + 2\mu\lambda_n) > e^{-2\mu\lambda_n}$$

are satisfied. Then on D_h^- ,

$$E(-y_n) = e^{2\mu Y} \prod_{i=1}^n (1 - \mu\lambda_i) > 1,$$

which implies that $E > 1$ on D_h .

If in addition to requiring that $\mu h \leq 1/2$, $\mu\lambda_n \leq 1/2$, we also take h so small that

$$1/y_2 \geq b(x, 0)/2,$$

we may write as a lower bound for $L_h E$ on D_h ,

$$L_h E \geq \mu^2 - 3|b|\mu + 2c.$$

Finally, on Γ_h^- ,

$$E_{\tau_0, n}^- = \mu E(-y_n) \geq \mu.$$

Therefore we may choose μ so large that for h small we have $L_h E \geq 1$ on D_h and $E_{\tau} \geq 1$ on Γ_h^- .

We now define functions V and W on D_h by

$$\begin{aligned} V &= U + B_1 + (B_2 + B_3)E, \\ W &= -U + B_1 + (B_2 + B_3)E. \end{aligned}$$

We then see easily that $L_h V \geq 0$ and $L_h W \geq 0$ on D_h , and $V_{\tau} \geq 0$ and $W_{\tau} \geq 0$ on Γ_h^- . Since $V \geq 0$ and $W \geq 0$ on Γ_h , the maxima of V and W are non-negative,

and the maxima of both functions occur on the boundary, by Theorem 3.3. The bound (5.1) for U then follows with $C = e^{4\mu Y} - 1$.

6. Convergence theorems.

THEOREM 6.1. *Suppose that the differential equation (1.1) has a solution u in D which satisfies the boundary conditions (1.3), and assume that u is of class $C^2(\bar{D})$. Assume that the Conditions A are satisfied for h sufficiently small and let U be the solution to the difference equation (2.6) with boundary conditions (2.7). Then $U \rightarrow u$ uniformly on D_h as $h \rightarrow 0$.*

Proof. Theorem 2.1 asserts that given $\epsilon > 0$, there exists an $h_0 > 0$ such that on D_h

$$|Lu - L_h u| < \epsilon/2C$$

for $h \leq h_0$, C being the fixed constant of Theorem 5.1. But since $Lu = L_h U = f$ on D_h , this estimate may be written

$$|L_h(U - u)| < \epsilon/2C.$$

Furthermore, since $U = \phi_1$ on Γ_h^+ and $u = \phi_1$ on Γ^+ and u and ϕ_1 are continuous on $\bar{D}^+$, there exists an $h_1 > 0$ such that

$$|U - u| < \epsilon/2$$

on Γ_h^+ , for $h \leq h_1$. Finally, on Γ_h^- , $U - u \equiv 0$. Hence, the bound (5.1) applied to $U - u$ yields at all points of D_h ,

$$|U - u| < \epsilon$$

for $h \leq h_2 = \max(h_0, h_1)$.

For the case $K(y) = y$ and $a \equiv b \equiv c \equiv 0$, Filippov [2] has proved a convergence theorem with somewhat weakened conditions on the derivatives of u at the boundary, provided that the curve Γ^+ satisfies certain conditions. We will prove an analogous theorem for the equation (1.1), under similar conditions on Γ^+ , and with additional restrictions on the function K .

Let us first extend the definition of the function G , defined for $y \leq 0$ by equation (2.1), to positive values of y by

$$(6.1) \quad G(y) = \int_0^y [K(\eta)]^{1/2} d\eta; \quad y > 0.$$

We will say that the curve Γ^+ satisfies *Condition B* if there exists an x_0 with $x_A < x_0 < x_B$ such that for each real number t , the curve

$$(6.2) \quad x - x_0 = tG(y)$$

intersects the curve Γ^+ at only one point.

LEMMA 6.1. *Let Γ^+ satisfy Condition B. Then if $0 < \theta < 1$, the transformation*

$$(6.3) \quad \begin{aligned} \bar{x} &= \theta(x - x_0) + x_0, \\ G(\bar{y}) &= \theta G(y); \end{aligned} \quad \text{sgn } \bar{y} = \text{sgn } y,$$

maps each point (x, y) of $\bar{D}$ into a point $(\bar{x}, \bar{y})$ of the interior D .

Proof. We note that $\bar{y}$ is uniquely defined by (6.3) since G is a monotone function for $y > 0$ and $y < 0$.

Suppose (x, y) lies in $\bar{D}$ with $y > 0$. Then for some number t , the point (x, y) lies on the curve (6.2). Therefore, since $\bar{y} < y$, the Condition B implies that $(\bar{x}, \bar{y})$ is a point of D .

For each point (x, y) in $\bar{D}^-$ we have,

$$y \leq 0, \quad x_A + G(y) \leq x \leq x_B - G(y).$$

Then the point $(\bar{x}, \bar{y})$ given by (6.3) is also a point of D , since

$$\bar{y} \leq 0, \quad x_A + G(\bar{y}) < \bar{x} < x_B - G(\bar{y}).$$

Let us define

$$F_\theta(y) = [K(y)/K(\bar{y})]^{1/2}.$$

Then for each fixed θ with $0 < \theta < 1$, F_θ is continuously differentiable and $F_\theta(y) \rightarrow 1$ as $\theta \rightarrow 1$ if $y \neq 0$. In the following theorem, we require that F_θ have smooth properties at $y = 0$. We summarize the properties under the *Conditions C*.

Conditions C:

- (a) For each fixed θ with $0 < \theta < 1$, F_θ is continuously differentiable on $\bar{D}$,
- (b) $F_\theta(y) \rightarrow 1$ and $F'_\theta(y) \rightarrow 0$ as $\theta \rightarrow 1$, uniformly on $\bar{D}$,
- (c) For $y < 0$, $y/\bar{y}$ is uniformly bounded.

We also need somewhat stronger Conditions A which we will denote by *Conditions A**:

Conditions A hold and there is a $\delta > 0$ such that

- (a) (3.8) holds with $1 - \delta$ on the right side,
- (b) (3.9), (3.10) and (3.12) hold with δ on the right side.

THEOREM 6.2. Suppose that the differential equation (1.1) has a solution u in D which satisfies the boundary conditions (1.3), where u is a function of class $C(\bar{D})$ and $C^2(D)$, with $u_y - (-K)^{1/2}u_x$ continuous on $D^- \cup \Gamma_1$. Assume moreover that the Conditions A* are satisfied for all h sufficiently small, and that the Conditions B and C and (3.20) are satisfied. Then the solution of the difference equation (2.6) with boundary conditions (2.7) tends uniformly to u as $h \rightarrow 0$.

Proof. We define a new function u_θ by

$$(6.4) \quad u_\theta(x, y) = u(\bar{x}, \bar{y}),$$

where (x, y) and $(\bar{x}, \bar{y})$ are related by (6.3). By Lemma 6.1, Conditions C and the assumption that u is of class $C^2(D)$, we find that u_θ is of class $C^2(\bar{D})$

for each θ with $0 < \theta < 1$. Furthermore, since $(\bar{x}, \bar{y})$ tends uniformly to (x, y) as $\theta \rightarrow 1$ and u is of class $C(\bar{D})$, if $\epsilon > 0$ is given, then there is a $\theta_0 < 1$ such that

$$|u - u_\theta| < \epsilon/3$$

on $\bar{D}$ for each θ with $\theta_0 \leq \theta < 1$.

Since u is a solution of (1.1) on D , for each θ with $0 < \theta < 1$, the function u_θ satisfies the differential equation

$$(6.5) \quad L_\theta u_\theta = K(y)u_{\theta xx} + u_{\theta yy} + a_\theta(x, y)u_{\theta x} + b_\theta(x, y)u_{\theta y} + c_\theta(x, y)u_\theta = f_\theta(x, y)$$

on $\bar{D}$, where

$$(6.6) \quad \begin{aligned} a_\theta(x, y) &= \theta F_\theta^2(y)a(\bar{x}, \bar{y}), \\ b_\theta(x, y) &= \theta F_\theta(y)b(\bar{x}, \bar{y}) - F_\theta'(y)/F_\theta(y), \\ c_\theta(x, y) &= \theta^2 F_\theta^2(y)c(\bar{x}, \bar{y}), \\ f_\theta(x, y) &= \theta^2 F_\theta^2(y)f(\bar{x}, \bar{y}). \end{aligned}$$

The Conditions C then imply that

$$(6.7) \quad a_\theta \rightarrow a, \quad b_\theta \rightarrow b, \quad c_\theta \rightarrow c, \quad f_\theta \rightarrow f \quad \text{as } \theta \rightarrow 1,$$

uniformly on $\bar{D}$. We next assert that if we define quantities $A_{\theta k, n}$, $\alpha_{\theta k, n}$ and $\gamma_{\theta k, n}$ by replacing a , b and c in equations (3.2), (3.3) and (3.4) by a_θ , b_θ and c_θ , then for h sufficiently small,

$$(6.8) \quad A_{\theta k, n} \rightarrow A_{k, n}, \quad \alpha_{\theta k, n} \rightarrow \alpha_{k, n}, \quad \gamma_{\theta k, n} \rightarrow \gamma_{k, n} \quad \text{as } \theta \rightarrow 1,$$

uniformly on D_n^- . The first and third limits of (6.8) are immediate consequences of (6.7) since the quantities multiplying b and c in (3.2) and (3.4), respectively, are uniformly bounded for h sufficiently small. For the second limit, we know from (2.13) that

$$\lambda_n \lambda_{n+1}/h \leq \frac{1}{2} \lambda_n [-K(-y_n)]^{-1/2}.$$

Hence $\lambda_n \lambda_{n+1}/h$ is uniformly bounded for $y_n \geq \delta > 0$, and the second limit follows for $y_n \geq \delta$. For $y_n < \delta$, we have

$$|\alpha_{\theta k, n}| = \left| \frac{\lambda_n \lambda_{n+1}}{h} \theta F_\theta^2(y) a(\bar{x}, \bar{y}) \right| \leq \left| \frac{y_n}{2\bar{y}} \theta F_\theta(y) \frac{\bar{y} a(\bar{x}, \bar{y})}{[-K(\bar{y})]^{1/2}} \right|,$$

and the right side can be made arbitrarily small by making y_n small, by virtue of the Conditions C and (3.20). But from the proof of Theorem 3.4, we observe that the condition (3.20) also implies that $\alpha_{k, n} \rightarrow 0$ as $y_n \rightarrow 0$. Hence if we choose δ sufficiently small, the difference $\alpha_{k, n} - \alpha_{\theta k, n}$ can be made arbitrarily small for $y_n < \delta$.

From (6.7) and (6.8) it follows that the Conditions A are satisfied by the

coefficients of the difference approximation to L_θ provided that h is small and θ is close to 1, since we have assumed that the stronger Conditions A* are satisfied for all h sufficiently small. Hence the difference problem corresponding to the differential equation (6.5) with boundary conditions

$$u_\theta(x, y) = u(\bar{x}, \bar{y}) \quad \text{for } (x, y) \text{ on } \Gamma,$$

has a unique solution U_θ , if h is small and θ is near 1. Moreover, since u_θ is twice continuously differentiable on $\bar{D}$ for each θ with $0 < \theta < 1$, U_θ tends uniformly to u_θ as $h \rightarrow 0$, by Theorem 6.1. That is, there exist numbers $h_1 > 0$ and $\theta_1 < 1$ such that on D_h ,

$$|u_\theta - U_\theta| < \epsilon/3$$

for each h and θ with $0 < h \leq h_1$, $\theta_1 \leq \theta < 1$.

Let us now compare the solution U of the difference approximation to $Lu = f$ with the solution U_θ of the difference approximation to $L_\theta u_\theta = f_\theta$. We have

$$L_h(U - U_\theta) = (f - f_\theta) = (a - a_\theta)U_{\theta_x} - (b - b_\theta)U_{\theta_y} - (c - c_\theta)U_\theta,$$

where U_{θ_x} and U_{θ_y} are difference quotients of U_θ defined in (2.3), (2.4) and (2.5). Since the functions a_θ and b_θ are independent of h , for each h we may choose θ so close to 1 that $|a - a_\theta| < h^2$ and $|b - b_\theta| < h^2$. Hence, due to the uniform boundedness of U_θ on $\bar{D}_h$ for h sufficiently small, it follows that for h small

$$L_h(U - U_\theta) < \epsilon/12C,$$

where C is the constant of Theorem 5.1. Furthermore, because u is continuous on $\bar{D}$, it is clear that if h is small and θ is near 1,

$$|U - U_\theta| < \epsilon/6$$

on the boundary Γ_h . Likewise, by selecting h and θ appropriately, it is possible to make

$$|U_{\bar{x}} - U_{\theta_{\bar{x}}}| < \epsilon/12C,$$

since $U = u$ and $U_\theta = u_\theta$ on Γ_1 , and du_θ/dy , which is given by

$$\frac{du_\theta}{dy}(x, y) = \theta F_\theta(y) [u_{\bar{y}}(\bar{x}, \bar{y}) - (-K(\bar{y}))^{1/2} u_{\bar{x}}(\bar{x}, \bar{y})]$$

is continuous on Γ_1 and tends to du/dy as $\theta \rightarrow 1$. If we now apply Theorem 5.1 to the function $U - U_\theta$, we find that on D_h ,

$$|U - U_\theta| < \epsilon/3$$

for h and θ such that $0 < h \leq h_2$, $\theta_2 \leq \theta < 1$. Taken together, these estimates imply that given $\epsilon > 0$, there is an $h_0 > 0$ such that if $0 < h \leq h_0$, then

$$|u - U| < \epsilon.$$

In one particular case, namely that in which the function K is given by

$$K(y) = |y|^m \operatorname{sgn} y$$

for some positive integer m , the function F_θ takes on an especially simple form, with the result that the required conditions are easily verified. From (2.1) and (6.1) we obtain

$$G(y) = \frac{2}{m+2} |y|^{(m+2)/2}.$$

Recalling that $G(\bar{y}) = \theta G(y)$, we find that

$$\bar{y} = \theta^{2/(m+2)} y.$$

Then we can easily calculate

$$F_\theta(y) = \theta^{-m/(m+2)}.$$

That is, F_θ depends only on θ and not on y . The Conditions C are clearly satisfied in this case.

In conclusion, it should be noted that the results are still valid for problems of the type:

- (a) to solve $Lu = f$ in D^+ with u given on Γ^+ and $\overline{AB}$,
- (b) to solve $Lu = f$ in D^- with u given on Γ_1 and $\overline{AB}$.

The maximum principles, a priori bounds and convergence theorems are applicable with obvious modifications in the hypotheses. Theorem 6.2 applies in either case (a) or (b) if the solution u is assumed to be twice continuously differentiable on $D \cup \overline{AB}$.

BIBLIOGRAPHY

1. F. Tricomi, *Sulle equazioni lineari alle derivate parziali di 2° ordine di tipo misto*, Atti Accad. Naz. dei Lincei (5) vol. 14 (1923).
2. A. F. Filippov, *On difference methods for the solution of the Tricomi problem*, Izv. Akad. Nauk SSSR. Ser. Mat. vol. 21 (1957) pp. 73-88.
3. P. Germain and R. Bader, *Sur quelques problèmes à l'équation de type mixte de Tricomi*, Office Nationale d'Etudes et de Recherches Aéronautiques, Publication No. 54, 1952.
4. S. Agmon, L. Nirenberg and M. H. Protter, *A maximum principle for a class of hyperbolic equations and applications to equations of mixed elliptic-hyperbolic type*, Comm. Pure Appl. Math. vol. 6 (1953) pp. 445-470.

UNIVERSITY OF CALIFORNIA,
RIVERSIDE, CALIFORNIA