

# INVARIANTS OF EUCLIDEAN REFLECTION GROUPS

BY  
LOUIS SOLOMON<sup>(1)</sup>

**1. Introduction and statement of results.** Let  $R^n$  be  $n$ -dimensional Euclidean space and let  $G$  be a finite group of orthogonal transformations of  $R^n$  generated by reflections. Let  $V = C^n$  be the complexification of  $R^n$ . Then  $G$  acts naturally in  $V$  and we say that  $G$  is a Euclidean reflection group in  $V$ . Let  $S$  be the  $C$ -algebra of complex-valued polynomial functions on  $V$ , let  $I(S)$  be the subalgebra of polynomials invariant under  $G$  and let  $F$  be the ideal of  $S$  generated by the homogeneous elements of positive degree in  $I(S)$ . Chevalley [2] has proved that

(a)  $I(S)$  is generated over  $C$  by  $n$  algebraically independent homogeneous polynomials  $f_1, \dots, f_n$  and the unit element.

(b)  $S/F$  as  $G$ -module affords the regular representation of  $G$ .

In view of (b) every irreducible  $G$ -module  $M$  occurs in  $S/F$  with multiplicity equal to  $\dim M$ . Since  $F$  is a homogeneous ideal,  $S/F = \sum_q (S/F)_q$  is naturally graded. We prove two theorems concerning the graded  $G$ -module structure of  $S/F$ .

The symmetric group  $G$  of degree  $n$  acts naturally as a Euclidean reflection group in  $V$  by permuting the elements of a basis. The irreducible characters of  $G$  are in 1-1 correspondence with partition diagrams of  $n$  nodes [6]. In the set of partition diagrams there is a natural ordering. The evidence suggests that characters which occur early in this ordering occur early in the decomposition  $S/F = \sum_q (S/F)_q$  in the sense that they occur for small values of  $q$ . On the other hand, a formula of Frobenius [6, p. 534] indicates that if a character occurs early in the partition ordering, then a reflection (transposition) fixes a large part of the corresponding representation space. These observations led to the following.

**THEOREM 1.** *Let  $G$  be a Euclidean reflection group and let  $M$  be an irreducible  $G$ -module. Let  $\gamma \in G$  be a reflection, let  $M_\gamma^-$  be the subspace of all  $x \in M$  such that  $\gamma x = -x$  and suppose that  $\dim M_\gamma^- = \dim M^-$  is independent of  $\gamma$ . If  $M$  is an irreducible constituent of  $(S/F)_q$  for precisely the values  $q_1(M), \dots, q_l(M)$ ,  $l = \dim M$ , then the average of the  $q_i(M)$  is*

$$\frac{\dim M^-}{\dim M} r$$

where  $r$  is the number of reflections in  $G$ .

---

Received by the editors June 7, 1963.

(1) This research was supported by the National Science Foundation under grant G-21514

The assumption that  $\dim M_\gamma^-$  is independent of  $\gamma$  is satisfied for all  $M$  when the reflections in  $G$  form a single conjugate class, and is satisfied for the irreducible modules  $E_p$  of Theorem 2 when  $G$  is a Weyl group. The extreme cases in Theorem 1 are given by  $\dim M^- = 0$ , corresponding to the principal character, and  $\dim M^- = \dim M$ , corresponding to the alternating character of  $G$ .

**THEOREM 2.** *Let  $L$  be a complex simple Lie algebra and let  $V$  be a Cartan subalgebra of  $L$ . Let  $G$  be the Weyl group of  $L$  acting in  $V$ . Let  $E_p$  be the  $G$ -module of alternating multilinear  $p$ -forms over  $V$ . Then  $E_p$  is irreducible and occurs as a constituent of  $(S/F)_q$  for precisely the values  $q = m_{i_1} + \cdots + m_{i_n}$ ,  $i_1 < \cdots < i_p$ , where  $m_i + 1$  is the degree of  $f_i$ . A basis for the isotypic component of  $S/F$  of type  $E_p$  is given by the set of minors of order  $p$  of the Jacobian matrix  $J$  of  $f_1, \dots, f_n$  reduced mod  $F$ .*

The irreducibility of the modules  $E_p$  for Weyl groups of the exceptional Lie algebras  $\mathfrak{E}_6, \mathfrak{E}_7$  was noticed by Frame [5]. It seems likely that Theorem 2 is true for all the Euclidean reflection groups. The proof we give depends on a theorem of Burnside [1] on Weyl groups which allows us to compute certain invariants of  $G$ .<sup>(2)</sup> Theorem 2 and its proof have the following corollaries:

(2a) All minors of  $J$  are linearly independent over  $C$  and remain linearly independent after reduction mod  $F$ . In particular, none of them vanish.

(2b) The algebra of invariants  $I(S/F \otimes E)$  is an exterior algebra over  $C$  on  $n$  generators.

(2c) If  $E = \sum_p E_p$  is the Grassmann algebra of  $V$ , then the algebra of invariants  $I(E \otimes E)$  is a truncated polynomial algebra over  $C$ , generated by the unit and an element  $w$  such that  $w^{n+1} = 0$ . The generator  $w$  may be identified with the Killing form.

(2d) For each  $p = 0, \dots, n$  there exists a homogeneous isomorphism

$$I(S/F \otimes E_p) \simeq I(S/F \otimes E_{n-p})$$

of graded vector spaces. Existence of this isomorphism for  $p = 1$  is equivalent to so-called double duality in the exponents  $m_i$ , the fact that if the  $m_i$  are arranged in increasing order  $m_1 \leq \cdots \leq m_n$  then  $m_i + m_{n-i+1}$  is independent of  $i$ .

The double duality in the exponents  $m_i$  was a long standing mystery for Weyl groups, explained a few years ago by Coleman [3] and Kostant [7]. Even if one assumes the double duality as known, the argument in (2d) does not furnish an explicit isomorphism  $I(S/F \otimes E_1) \simeq I(S/F \otimes E_{n-1})$ . We prove the existence of the isomorphism by computing the Poincaré series of both spaces. It would thus be interesting to give a direct invariant-theoretic proof for the double duality by exhibiting an isomorphism which is in some sense a natural one. We have not been able to do this, but present a line of argument which seems to lead in the right

<sup>(2)</sup> *Added in proof.* R. Steinberg has kindly shown me a proof of irreducibility which is independent of Burnside's theorem. His argument is valid for all the Euclidean groups.

direction. The elements of  $S \otimes E_1$  may be viewed as differential 1-forms on  $V$ . We study the space of those differential 1-forms on  $V$  which are skew invariant under  $G$ , in the sense that they are invariant under the rotation subgroup  $H$  of  $G$  and change sign under the elements outside  $H$ . It is not hard to show that this space is a free module over  $I(S)$  of rank  $n$ . From the double duality and the fact that  $G$  has a unique invariant quadratic form one concludes that among the polynomials  $f_1, \dots, f_n$  there is a unique polynomial  $f_n$  of greatest degree. Then assuming the double duality we prove the following.

**THEOREM 3.** *Let  $G$  be an irreducible Euclidean reflection group in  $V$ . Choose coordinates in  $V$  and let  $u_1, \dots, u_n$  be the minors of order  $n-1$  of  $J$  obtained by deleting the partial derivatives of  $f_n$ . If  $u_1, \dots, u_n$  are algebraically independent, then there exists a homogeneous derivation  $\hat{d}: S \rightarrow S \otimes E_1$  of  $S$ -modules such that  $\hat{d}f_1, \dots, \hat{d}f_n$  are a basis for the module of skew invariant differential 1-forms over  $I(S)$ .*

Granted the existence of the map  $\hat{d}$ , we can construct an explicit isomorphism  $I(S/F \otimes E_1) \simeq I(S/F \otimes E_{n-1})$ . We have been able to verify the algebraic independence of  $u_1, \dots, u_n$  in special cases but have no general argument. If the  $u_i$  are algebraically independent, then the Jacobian (determinant) of the  $u_i$  must be a constant multiple of  $(\det J)^{n-2}$ .

We work over the complex field  $\mathbb{C}$  as a matter of convenience, and irreducibility of modules will mean irreducibility over  $\mathbb{C}$ . The complex field is probably an alien here because a likely conjecture of Kostant states that all the absolutely irreducible representations of a Euclidean reflection group may be written with coefficients in  $\mathbb{R}$ . In any case, a real linear group which contains a reflection and is irreducible over  $\mathbb{R}$  remains irreducible over  $\mathbb{C}$ .

**2. Notation.** In this section we introduce some notation and collect some elementary facts about invariants and characters. Let  $G$  be a finite group of order  $g$ . By a graded  $G$ -module we mean a  $G$ -module which is a graded vector space  $M = \sum_{q \geq 0} M_q$  over  $\mathbb{C}$ , in which each homogeneous component  $M_q$  is a  $G$ -module finite dimensional over  $\mathbb{C}$ . Let  $\mu_q$  be the character of  $G$  corresponding to the module  $M_q$ . To the graded  $G$ -module  $M$  we let correspond the series

$$M(t, \gamma) = \sum_{q \geq 0} \mu_q(\gamma) t^q, \quad \gamma \in G.$$

For  $\gamma = 1$  this becomes the Poincaré series

$$M(t) = \sum_{q \geq 0} (\dim M_q) t^q$$

of the graded vector space  $M$ . All the tensor products we consider are tensor products over  $\mathbb{C}$ . If  $M, N$  are graded  $G$ -modules, then  $M \otimes N$  has a natural structure of graded  $G$ -module with the grading and  $G$ -module structure defined by

$$(M \otimes N)_q = \sum_{a+b=q} M_a \otimes N_b,$$

$$\gamma(x \otimes y) = \gamma x \otimes \gamma y, \quad x \in M, y \in N, \gamma \in G.$$

From the fact that the character of a tensor product (direct sum) of two  $G$ -modules is the product (sum) of the characters it follows that

$$(M \otimes N)(t, \gamma) = M(t, \gamma) N(t, \gamma).$$

We let  $I(M)$  denote the submodule of invariants of  $M$ , elements  $x \in M$  such that  $\gamma x = x$  for all  $\gamma \in G$ . For a finite dimensional  $M$  with character  $\mu$  the connection between invariants and characters is the formula

$$\dim I(M) = \frac{1}{g} \sum_{\gamma \in G} \mu(\gamma).$$

Thus for a graded  $M$  we have

$$I(M)(t) = \frac{1}{g} \sum_{\gamma \in G} M(t, \gamma).$$

The dual  $M^*$  of  $M$  has a natural  $G$ -module structure defined by

$$(\gamma f)(x) = f(\gamma^{-1}x), \quad x \in M, f \in M^*, \gamma \in G,$$

and we may extend this action to the algebra of polynomial functions on  $M$  or the Grassmann algebra of  $M$ . If  $\mu$  is the character of  $M$  then  $\mu^*(\gamma) = \mu(\gamma^{-1})$  is the character of  $M^*$ . We have an isomorphism  $M \cong M^*$  of  $G$ -modules if and only if the character of  $M$  is real. The space  $\text{Hom}_{\mathbb{C}}(M, N)$  has a natural  $G$ -module structure defined by

$$(\gamma \phi)(x) = \gamma(\phi(\gamma^{-1}x)), \quad x \in M, \phi \in \text{Hom}_{\mathbb{C}}(M, N), \gamma \in G,$$

and the submodule  $I(\text{Hom}_{\mathbb{C}}(M, N))$  is just the space  $\text{Hom}_G(M, N)$  of  $G$ -module homomorphisms. The natural isomorphism of vector spaces  $\text{Hom}_{\mathbb{C}}(M, N) \simeq N \otimes M^*$  is an isomorphism of  $G$ -modules and induces an isomorphism  $\text{Hom}_G(M, N) \simeq I(N \otimes M^*)$ . In particular we see that if  $M$  is irreducible then  $\dim I(N \otimes M^*)$  is the multiplicity of  $M$  in  $N$ , and that  $M$  is irreducible if and only if  $\dim I(M \otimes M^*) = 1$ .

3. Let  $G$  be a Euclidean reflection group in  $V$  and let  $S$  be the algebra of complex valued polynomial functions on  $V$ . Then  $S = \sum_{q \geq 0} S_q$  has a natural structure of graded  $G$ -module. A formula of Molien, easy to verify by assuming  $\gamma$  in diagonal form, states that

$$(3.1) \quad S(t, \gamma^{-1}) = \frac{1}{(1 - \omega_1(\gamma)t) \cdots (1 - \omega_n(\gamma)t)}$$

where  $\omega_1(\gamma), \dots, \omega_n(\gamma)$  are the eigenvalues of  $\gamma$  as linear transformation of  $V$ .

If  $M$  is a finite dimensional  $G$ -module we give  $S \otimes M$  the grading defined by  $(S \otimes M)_q = S_q \otimes M$ . Then Molien's formula implies

$$(3.2) \quad I(S \otimes M)(t) = \frac{1}{g} \sum_{\gamma \in G} \frac{\mu(\gamma^{-1})}{(1 - \omega_1(\gamma)t) \cdots (1 - \omega_n(\gamma)t)}$$

where  $\mu$  is the character of  $M$ . From Theorem (a) of Chevalley we see that

$$(3.3) \quad I(S)(t) = \frac{1}{(1 - t^{m_1+1}) \cdots (1 - t^{m_n+1})}.$$

Chevalley has also shown [2] that if  $p_1, \dots, p_k \in S$  form a  $C$ -basis for  $S/F$  when reduced mod  $F$ , then  $p_1, \dots, p_k$  are a basis for  $S$  as free module over  $I(S)$ . From this fact we readily deduce the following two lemmas.

**LEMMA 1.** *Let  $\tau_q$  be the character of  $(S/F)_q$ . Then*

$$\sum_q \tau_q(\gamma^{-1})t^q = \frac{(1 - t^{m_1+1}) \cdots (1 - t^{m_n+1})}{(1 - \omega_1(\gamma)t) \cdots (1 - \omega_n(\gamma)t)}.$$

**Proof.** Let  $p_1, \dots, p_k \in S$  form a  $C$ -basis for  $S/F$  when reduced mod  $F$ . Then the map  $\sum_i s_i p_i \rightarrow \sum_i s_i \otimes (p_i + F)$ ,  $s_i \in I(S)$  defines an isomorphism  $S \simeq I(S) \otimes S/F$  of graded  $G$ -modules. Since  $G$  acts trivially on  $I(S)$  we have  $S(t, \gamma) = I(S)(t)(S/F)(t, \gamma)$ . Thus from (3.1) and (3.3)

$$(S/F)(t, \gamma^{-1}) = \frac{(1 - t^{m_1+1}) \cdots (1 - t^{m_n+1})}{(1 - \omega_1(\gamma)t) \cdots (1 - \omega_n(\gamma)t)}$$

which proves the lemma.

If we let  $t \rightarrow 1$  we find  $\sum_q \tau_q(\gamma) = 0$  if  $\gamma \neq 1$  and  $\sum_q \tau_q(1) = g$ , so that  $\sum_q \tau_q$  is the character of the regular representation of  $G$ . Thus  $S/F$  affords the regular representation of  $G$ . If  $M$  is an irreducible  $G$ -module we let  $a_q(M)$  be the multiplicity of  $M$  in  $(S/F)_q$ . Since  $S/F$  contains  $M$  with multiplicity equal to  $\dim M$  we have

$$\sum_q a_q(M) = \dim M.$$

We view  $S \otimes M$  naturally as an  $S$ -module and then  $I(S \otimes M)$  is an  $I(S)$ -module.

**LEMMA 2.** *Let  $M$  be an irreducible  $G$ -module. Then  $I(S \otimes M)$  is a free module over  $I(S)$ . It has a basis over  $I(S)$  consisting of homogeneous elements in which the number of elements of degree  $q$  is  $a_q(M^*)$ . The rank of  $I(S \otimes M)$  as  $I(S)$ -module is equal to  $\dim M$ .*

**Proof.** In the proof of Lemma 1 we have remarked that  $S \simeq I(S) \otimes (S/F)$  and hence  $S \otimes M \simeq I(S) \otimes (S/F) \otimes M$ . Since  $G$  acts trivially on  $I(S)$  we see by averaging over the group that  $I(S \otimes M) \simeq I(S) \otimes I(S/F \otimes M)$ . Thus  $I(S \otimes M)$  is free over  $I(S)$  and we may choose as basis a  $C$ -basis for  $I(S/F \otimes M)$ . This may be

chosen as a union of  $C$ -bases for the  $I((S/F)_q \otimes M)$ . But  $\dim I((S/F)_q \otimes M)$  is the multiplicity of the irreducible  $M^*$  in  $(S/F)_q$  so  $\dim I((S/F)_q \otimes M) = a_q(M^*)$ . The rank of  $I(S \otimes M)$  as  $I(S)$ -module is thus  $\sum_q a_q(M^*) = \dim M^* = \dim M$ . The argument shows that  $I(S \otimes M)$  is free over  $I(S)$  for any  $G$ -module  $M$ .

4. To prove Theorem 1 we simply compute the Poincaré series  $I(S \otimes M^*)(t)$  in two ways and compare the results for  $t = 1$ . Set  $a(t) = \sum_q a_q(M)t^q$ . From Lemma 2 with  $M$  replaced by  $M^*$  we have

$$I(S \otimes M^*)(t) = \frac{a(t)}{(1 - t^{m_1+1}) \cdots (1 - t^{m_n+1})}$$

and thus from (3.2) we see that

$$(4.1) \quad \frac{1}{g} \sum_{\gamma \in G} \frac{\mu(\gamma)}{(1 - \omega_1(\gamma)t) \cdots (1 - \omega_n(\gamma)t)} = \frac{a(t)}{(1 - t^{m_1+1}) \cdots (1 - t^{m_n+1})}$$

where  $\mu$  is the character of  $M$ . Let  $G_1$  be the set of elements of  $G$ , distinct from the identity, which fix an  $n - 1$  dimensional subspace of  $V$ . For  $\gamma \in G_1$  the eigenvalues  $\omega_i(\gamma)$  are  $1, 1, \dots, 1, \omega$  where  $\omega$  is a root of unity. Now the fact that  $G$  may be written as a real orthogonal group implies  $\omega = -1$ . The left-hand side of (4.1) becomes

$$\frac{1}{g} \left[ \frac{\mu(1)}{(1-t)^n} + \frac{1}{(1-t)^{n-1}(1+t)} \sum_{\gamma \in G_1} \mu(\gamma) + \cdots \right]$$

where  $\cdots$  denotes terms which have at most  $(1-t)^{n-2}$  in the denominator. Since  $a(1) = \sum_q a_q(M) = \dim M = \mu(1)$  we have

$$\frac{1}{(1-t)^{n-1}(1+t)} \sum_{\gamma \in G_1} \mu(\gamma) + \cdots = \frac{ga(t)}{(1-t^{m_1+1}) \cdots (1-t^{m_n+1})} - \frac{a(1)}{(1-t)^n}.$$

Now multiply both sides by  $(1-t)^{n-1}$  and let  $t \rightarrow 1$ . Using the known formula [4; 8]  $g = \prod_i (m_i + 1)$  for the group order we find

$$(4.2) \quad \frac{1}{2} \sum_{\gamma \in G_1} \mu(\gamma) = -a'(1) + \frac{1}{2} \left( \sum_i m_i \right) a(1)$$

where  $a'$  denotes the derivative with respect to  $t$ . For  $\gamma \in G_1$ ,  $\mu(\gamma) = \dim M_\gamma^+ - \dim M_\gamma^-$  is, by assumption, independent of  $\gamma$ . The number of elements in  $G_1$  is the number of reflections in  $G$  which is known [4; 8] to be  $\sum_i m_i$ . From the definition of  $a(t)$  we have  $a'(1) = \sum_q q a_q(M)$ . If we insert this information in (4.2) we find

$$\frac{1}{2} \left( \sum_i m_i \right) (\dim M^+ - \dim M^-) = - \sum_q q a_q(M) + \frac{1}{2} \left( \sum_i m_i \right) (\dim M^+ + \dim M^-)$$

so that

$$\sum_q q a_q(M) = \left( \sum_i m_i \right) \dim M^-.$$

The average of the  $q_i(M)$  is then

$$\frac{\sum_q q a_q(M)}{\sum_q a_q(M)} = \frac{\dim M^-}{\dim M} r$$

with  $r = \sum_i m_i$ . The significance of the integer  $r = \sum_i m_i$  in this formula becomes clear if one observes, with Lemma 1, that  $r$  is the largest integer  $q$  for which  $(S/F)_q \neq 0$ .

5. To prove Theorem 2 we shall need some results from an earlier article [9] together with a lemma concerning certain invariants of the symmetric group. Let  $E = \sum_p E_p$  be the Grassman algebra of  $V$ . The homogeneous component  $E_p$  of degree  $p$  is the space of all  $p$ -linear alternating functions on  $V$ . We identify  $E_0$  with  $\mathbb{C}$  and  $E_1$ , as vector space, with  $S_1$ . The group  $G$  acts naturally on  $E$  and on  $S \otimes E$ . Choose a coordinate system  $x_1, \dots, x_n$  in  $V$  and let  $d: S \otimes E \rightarrow S \otimes E$  be the  $\mathbb{C}$ -linear map defined by

$$d: s \otimes x_{i_1} \wedge \cdots \wedge x_{i_p} \rightarrow \sum_{j=1}^p \frac{\partial s}{\partial x_j} \otimes x_j \wedge x_{i_1} \wedge \cdots \wedge x_{i_p}, \quad s \in S.$$

If we identify  $S$  with  $S \otimes \mathbb{C}$  then  $dx_i = d(x_i \otimes 1) = 1 \otimes x_i$  so that the elements of  $S \otimes E_p$  may be written in the form

$$\sum_{i_1 < \cdots < i_p} s_{i_1 \dots i_p} dx_{i_1} \cdots dx_{i_p}, \quad s_{i_1 \dots i_p} \in S.$$

It is clear that  $S \otimes E$  is just the algebra of differential forms on  $V$  and that  $d$  is exterior differentiation. Since  $d$  commutes with the action of  $G$  on  $S \otimes E$  it follows that  $d$  maps  $I(S \otimes E)$  into  $I(S \otimes E)$ . In particular, the differentials  $df_i$  are invariants of  $S \otimes E$ . We have shown in [9] that the  $\mathbb{C}$ -algebra  $I(S \otimes E)$  of invariant differential forms is an exterior algebra on  $n$  generators over the  $\mathbb{C}$ -algebra  $I(S)$  of invariant polynomials, and is in fact generated over  $I(S)$  by the differentials  $df_1, \dots, df_n$  of the polynomial invariants  $f_1, \dots, f_n$  and the unit element. It follows that

$$(5.1) \quad I(S \otimes E_p)(t) = \frac{\sigma_p(t^{m_1}, \dots, t^{m_n})}{(1 - t^{m_1+1}) \cdots (1 - t^{m_n+1})}, \quad p = 1, \dots, n,$$

where  $\sigma_p(t_1, \dots, t_n)$  is the  $p$ th elementary symmetric function in the indeterminates  $t_1, \dots, t_n$ .

LEMMA 3. Let  $x_1, \dots, x_n$  be indeterminates, let  $G$  be the symmetric group on  $x_1, \dots, x_n$  and let  $E$  be the exterior algebra on  $x_1, \dots, x_n$  over  $\mathbb{C}$ . The group  $G$  acts naturally on the commutative algebra  $E \otimes E$ . Set

$$u = \sum_i x_i \otimes x_i, \quad v = \sum_{i < k} x_i \otimes x_k + x_k \otimes x_i.$$

Then

$$I(E_p \otimes E_p) = Cu^p \oplus Cu^{p-1}v, \quad p = 1, \dots, n-1.$$

**Proof.** Clearly both  $u^p$  and  $u^{p-1}v$  are in  $I(E_p \otimes E_p)$ . To hold the indices in check we let  $\Omega$  denote the set of increasing sequences  $i_1 < \dots < i_p$  of  $p$  integers chosen from  $1, \dots, n$ , we let  $(i)$  denote an element of  $\Omega$ , let  $\{i\}$  be the corresponding unordered set, and write  $x_{(i)} = x_{i_1} \wedge \dots \wedge x_{i_p}$ . Suppose  $y = \sum c_{(i),(k)} x_{(i)} \otimes x_{(k)} \in I(E_p \otimes E_p)$  where  $c_{(i),(k)} \in C$  and the sum is over all pairs  $(i), (k)$  of elements of  $\Omega$ . If for given sets  $\{i\}, \{k\}$  the intersection  $\{i\} \cap \{k\}$  contains fewer than  $p-1$  elements, then there exist two distinct indices, say  $k_\alpha, k_\beta$ , which are distinct from all elements of  $\{i\}$ . Now apply the transposition  $(k_\alpha k_\beta)$  of  $G$  to each term in the sum  $y$ . The invariance of  $y$  shows that  $c_{(i),(k)} = -c_{(i),(k)}$  and hence  $c_{(i),(k)} = 0$ . Thus we may write  $y = y_1 + y_2$  where  $y_1$  is a linear combination of elements  $x_{(i)} \otimes x_{(k)}$  such that  $\{i\} \cap \{k\}$  contains  $p$  elements, in other words  $\{i\} = \{k\}$ , and  $y_2$  is a linear combination of elements  $x_{(i)} \otimes x_{(k)}$  such that  $\{i\} \cap \{k\}$  contains  $p-1$  elements. Invariance of  $y$  implies the invariance of  $y_1$  and  $y_2$ . We thus have  $y_1 = \sum_{(i)} b_{(i)} x_{(i)} \otimes x_{(i)}$  with  $b_{(i)} \in C$  and invariance of  $y_1$  shows that all the  $b_{(i)}$  are equal, say  $b_{(i)} = b$ . Then  $y_1 = b \sum_{(i)} x_{(i)} \otimes x_{(i)}$  is a  $C$ -multiple of  $u^p = p! \sum_{(i)} x_{(i)} \otimes x_{(i)}$ . Similarly with slightly more effort one sees that  $y_2$  is a  $C$ -multiple of  $u^{p-1}v$ . Thus the elements  $u^p, u^{p-1}v$  span  $I(E^p \otimes E^p)$ . For  $p < n$  both  $u^p$  and  $u^{p-1}v$  are not zero and hence linearly independent over  $C$ . This proves the lemma. The argument breaks down for  $p = n$  only because  $u^{n-1}v = 0$  and in that case we have  $I(E_n \otimes E_n) = Cu^n$ . The elements  $u, v$  also satisfy the relations  $u^{n+1} = 0$  and  $v^2 = 0$ .

We are now in position to prove Theorem 2. Since  $L$  is simple,  $G$  acting in  $V$  is an irreducible group. A theorem of Burnside [1] states that there exists a coordinate system  $x_1, \dots, x_n$  in  $V$  such that  $G$  acting on  $V^* \simeq E_1$  includes the symmetric group  $H$  on  $x_1, \dots, x_n$ . In this coordinate system the Killing form must be

$$\frac{a}{2} \sum_i x_i^2 + b \sum_{i < k} x_i x_k \in I(S_2)$$

where  $a, b$  are real numbers. We cannot have  $a = 0$  because the form is positive definite. We let  $I(E \otimes E)$  denote the elements of  $E \otimes E$  invariant under  $G$  and  $I_H(E \otimes E)$  the elements invariant under  $H$ . Under the map  $f \rightarrow \sum_i (\partial f / \partial x_i) \otimes x_i$  of  $S_2 \rightarrow E_1 \otimes E_1$  the Killing form maps into  $au + bv \in I(E_1 \otimes E_1)$  where  $u, v \in I_H(E_1 \otimes E_1)$  are the invariants of Lemma 3. Since  $I(E \otimes E) \subseteq I_H(E \otimes E)$ , Lemma 3 shows that  $\dim I(E_p \otimes E_p) \leq 2$ . Suppose  $\dim I(E_p \otimes E_p) = 2$  for some  $p = 1, \dots, n-1$ . We prove that  $\dim I(E_{n-1} \otimes E_{n-1}) = 2$ . If  $p = n-1$  there is nothing to prove so assume  $p < n-1$ . From Lemma 3 we see that  $u^p \in I(E_p \otimes E_p)$  and  $u^{p-1}v \in I(E_p \otimes E_p)$ . Since  $v^2 = 0$  it follows that both  $au^p v = u^{p-1}v(au + bv)$  and  $au^{p+1} + bu^p v = u^p(au + bv)$  are in  $I(E_{p+1} \otimes E_{p+1})$ . Since  $p < n-1$  we



have  $u^p v \neq 0$  and since  $a \neq 0$  it follows that  $\dim I(E_{p+1} \otimes E_{p+1}) = 2$ . We conclude by induction that  $\dim I(E_{n-1} \otimes E_{n-1}) = 2$ . Let  $Z$  be the 1-dimensional  $G$ -module defined by the homomorphism  $\gamma \rightarrow \det \gamma$ ,  $\gamma \in G$ . Then  $E_{n-1} \simeq E_1 \otimes Z$  as  $G$ -modules. Since  $\det \gamma = \pm 1$ ,  $Z \otimes Z \simeq C$  is the trivial  $G$ -module. Then  $E_{n-1} \otimes E_{n-1} \simeq E_1 \otimes E_1$  as  $G$ -modules so that  $\dim I(E_1 \otimes E_1) = 2$  which contradicts the irreducibility of  $E_1$ . Thus  $\dim I(E_p \otimes E_p) = 1$  for all  $p$  and hence  $E_p$  is irreducible for all  $p$ .

Let  $\theta: S \rightarrow S/F$  be the natural map and extend  $\theta$  to a map, denoted again  $\theta$ , of  $S \otimes E \rightarrow S/F \otimes E$  by letting it be the identity on  $E$ . Then  $\theta$  is a homomorphism of  $G$ -modules and of  $C$ -algebras. Since  $\theta$  is a homomorphism of  $G$ -modules we certainly have  $\theta I(S \otimes E) \subseteq I(S/F \otimes E)$ . Actually we have  $\theta I(S \otimes E) = I(S/F \otimes E)$  because, using complete reducibility of the representations of  $G$ , we may choose a graded  $G$ -module  $T$  such that  $S = F \oplus T$  and then  $\theta: I(T \otimes E) \rightarrow I(S/F \otimes E)$  is an isomorphism. Set  $z_i = \theta(df_i)$ . Since  $I(S \otimes E)$  is generated over  $I(S)$  by the  $df_i$  and the unit element, and since every element  $s \in I(S)$  may be written as  $s = s_0 + s_1$  with  $s_0 \in C$  and  $s_1 \in F$ , it follows that the  $z_i$  together with the unit element generate  $I(S/F \otimes E)$  as algebra over  $C$ . Thus the  $\binom{n}{p}$  elements  $z_{i_1} \cdots z_{i_p}$ ,  $i_1 < \cdots < i_p$  generate  $I(S/F \otimes E_p)$  as vector space over  $C$ . In fact they form a basis for  $I(S/F \otimes E_p)$  because Lemma 2 shows that  $\dim I(S/F \otimes E_p) = \dim E_p = \binom{n}{p}$ . Thus the  $z_{i_1} \cdots z_{i_p}$  are linearly independent over  $C$ .

The Killing form induces a natural isomorphism  $E_p \simeq E_p^*$  of  $G$ -modules and hence a natural isomorphism  $S/F \otimes E_p \simeq \text{Hom}(E_p, S/F)$  of  $G$ -modules. Under this isomorphism the invariants  $I(S/F \otimes E_p)$  correspond to  $\text{Hom}_G(E_p, S/F)$ . For  $(i) \in \Omega$ , let  $\phi_{(i)}$  be the image in  $\text{Hom}_G(E_p, S/F)$  of  $z_{i_1} \cdots z_{i_p}$ . The linear independence of the  $z_{i_1} \cdots z_{i_p}$  implies linear independence of the  $\phi_{(i)}$ . Now if  $M$  is an irreducible  $G$ -module and  $\phi_1, \dots, \phi_s$  are linearly independent  $G$ -module homomorphisms of  $M$  into a  $G$ -module  $N$  then the sum  $\sum_i \phi_i(M)$  is direct. This follows at once from Schur's lemma by induction on the number of summands. In the case at hand this means that the sum  $\sum_{(i)} \phi_{(i)}(E_p)$  is direct. Since the number of summands is  $\binom{n}{p} = \dim E_p$ , the sum  $\sum_{(i)} \phi_{(i)}(E_p)$  is the isotypic component of  $S/F$  of type  $E_p$ . Thus  $E_p$  occurs as an irreducible constituent of  $(S/F)_q$  for precisely the values  $q = m_{i_1} + \cdots + m_{i_p}$ ,  $i_1 < \cdots < i_p$ . From the definition of  $\phi_{(i)}$  as the image of  $\theta(df_{i_1} \cdots df_{i_p})$  it follows that a basis for  $\phi_{(i)}(E_p)$  is given by the  $\binom{n}{p}$  minors of  $J$  which involve  $f_{i_1}, \dots, f_{i_p}$ , reduced mod  $F$ . This completes the proof of Theorem 2.

For the corollaries we argue as follows:

(2a) The linear independence over  $C$  of the minors of  $J$  after reduction mod  $F$  amounts to the linear independence of the elements  $\theta(df_{i_1} \cdots df_{i_p})$  over  $C$ . This we have shown.

(2b) The proof of the theorem shows that  $I(S/F \otimes E)$  is generated as algebra over  $C$  by the  $z_i = \theta(df_i)$  and the unit element. Since  $z_i z_j = -z_j z_i$ ,  $I(S/F \otimes E)$  is a homomorphic image of an exterior algebra on  $n$  generators. But

$\dim I(S/F \otimes E) = \dim E = 2^n$  so  $I(S/F \otimes E)$  is in fact an exterior algebra on the  $z_i = \theta(df_i)$ .

(2c) The proof of the theorem shows that  $I(E_p \otimes E_p) = \mathbb{C}w^p$  for all  $p = 1, \dots, n$  where  $w = au + bv$  may be identified with the Killing form. Suppose we have an isomorphism  $E_p \simeq E_q$  of  $G$ -modules. Then  $\binom{n}{p} = \dim E_p = \dim E_q = \binom{n}{q}$  so  $q = p$  or  $q = n - p$ . Suppose  $q = n - p$ . Let  $\chi_p$  be the character of  $E_p$  and let  $\gamma \in G$  be a reflection. Then  $\chi_p(\gamma)$  is the  $p$ th elementary symmetric function of the eigenvalues  $1, 1, \dots, 1, -1$  so that  $\chi_p(\gamma) = \binom{n-1}{p-1} - \binom{n-1}{p}$ . Now  $\chi_p(\gamma) = \chi_{n-p}(\gamma)$  shows  $n - p = p$ . Thus in any case  $q = p$ . It follows that  $I(E_p \otimes E_q) = 0$  for  $q \neq p$  and hence  $I(E \otimes E) = \sum_p I(E_p \otimes E_p)$  is generated over  $\mathbb{C}$  by the unit element and an element  $w = au + bv$  which satisfies  $w^{n+1} = 0$  and which may be identified with the Killing form.

(2d) From Theorem 2 or directly from (5.1) we conclude that  $I(S/F \otimes E_p)(t) = \sigma_p(t^{m_1}, \dots, t^{m_n})$ . Thus one has a homogeneous isomorphism

$$(5.2) \quad I(S/F \otimes E_1) \simeq I(S/F \otimes E_{n-1})$$

of graded vector spaces if and only if there exists an integer  $k$  such that  $t^k \sigma_1(t^{m_1}, \dots, t^{m_n}) = \sigma_{n-1}(t^{m_1}, \dots, t^{m_n})$ . Comparing coefficients on both sides shows that this condition is equivalent to the existence of an integer  $k$  such that  $k + m_i + m_{n-i+1} = m_1 + \dots + m_n$ . This is equivalent in turn to the statement that  $m_i + m_{n-i+1}$  is independent of  $i$ , the double duality. The same kind of coefficient comparison shows that double duality implies the isomorphisms

$$I(S/F \otimes E_p) \simeq I(S/F \otimes E_{n-p}).$$

For later use we remark that (5.2) is equivalent to the existence of a homogeneous isomorphism

$$(5.3) \quad I(S \otimes E_1) \simeq I(S \otimes E_{n-1})$$

of graded vector spaces.

6. Let  $M, N$  be  $G$ -modules. We say that  $M$  and  $N$  are skew isomorphic if there exists a 1-1  $\mathbb{C}$ -linear map  $\theta$  of  $M$  onto  $N$  such that  $\theta\gamma x = (\det \gamma)\gamma\theta x$  for all  $x \in M$  and all  $\gamma \in G$ . We call  $\theta$  a skew isomorphism between  $M$  and  $N$ . Since  $\det \gamma = \pm 1$  the relation of skew isomorphism is symmetric. Again we let  $Z$  denote the 1-dimensional  $G$ -module defined by the homomorphism  $\gamma \rightarrow \det \gamma$  and let  $z$  be a generator of  $Z$ . If we set  $\tilde{M} = M \otimes Z$  we see that  $x \rightarrow x \otimes z$ ,  $x \in M$ , defines a skew isomorphism between  $M$  and  $\tilde{M}$ . Since  $Z \otimes Z \simeq \mathbb{C}$ ,  $\tilde{\tilde{M}}$  and  $M$  are isomorphic as  $G$ -modules.

From Lemma 1 we conclude that  $t^r(S/F)(t^{-1}, \gamma) = \det \gamma(S/F)(t, \gamma)$  where  $r = \sum_i m_i$ . Thus  $\tau_{r-q}(\gamma) = (\det \gamma)\tau_q(\gamma)$  for all  $\gamma \in G$  and all  $q = 0, \dots, r$  so that  $(S/F)_q$  and  $(S/F)_{r-q}$  are skew isomorphic. If  $M$  is an irreducible  $G$ -module and  $N$

is any  $G$ -module, then  $\hat{M}$  is irreducible and the multiplicity of  $M$  in  $N$  is equal to the multiplicity of  $\hat{M}$  in  $\hat{N}$ . Hence the

**THEOREM.** *If  $M$  is an irreducible  $G$ -module, then the multiplicity of  $M$  in  $(S/F)_q$  is equal to the multiplicity of  $\hat{M}$  in  $(S/F)_{r-q}$  where  $r = \sum_i m_i$ . Thus in the notation of Theorem 1, with suitable ordering,  $q_i(\hat{M}) = r - q_i(M)$ .*

We say that  $x \in M$  is a skew invariant if  $\gamma x = (\det \gamma)x$  for all  $\gamma \in G$ . Let  $\hat{I}(M)$  denote the subspace of skew invariant elements of  $M$ . Then the map  $x \rightarrow x \otimes z$  defines a natural isomorphism  $\hat{I}(M) \simeq I(\hat{M})$  of vector spaces. It follows that we have an isomorphism

$$(6.1) \quad \hat{I}(S \otimes M) \simeq I(S \otimes \hat{M})$$

of graded vector spaces which is homogeneous of degree zero. From Lemma 2 we see that  $\hat{I}(S \otimes M)$  is free over  $I(S)$  of rank equal to  $\dim M$ .

Since  $E_{n-1} \simeq \hat{E}_1$  is an isomorphism of  $G$ -modules, the isomorphism (5.3) equivalent to the double duality becomes

$$(6.2) \quad \hat{I}(S \otimes E_1) \simeq I(S \otimes E_1).$$

Now  $I(S \otimes E_1)$  is generated freely over  $I(S)$  by the  $df_i$ . If we can construct the homogeneous derivation  $\hat{d}$  of Theorem 3 then  $\hat{I}(S \otimes E_1)$  is generated freely over  $I(S)$  by the  $\hat{d}f_i$  and the homogeneous isomorphism (6.2) is defined by  $df_i \rightarrow \hat{d}f_i$ . Thus in this formulation the double duality is equivalent to the existence of the map  $\hat{d}$ .

In connection with (6.2) it is worth noting that the homogeneous isomorphism  $\hat{I}(S \otimes E_0) \simeq I(S \otimes E_0)$  amounts to the familiar fact that every skew invariant polynomial may be written as an invariant polynomial multiplied by  $\det J$ .

7. Let  $q = m_1 + \cdots + m_{n-1}$ . We have a sequence of natural maps

$$S_q \otimes E_{n-1} \xrightarrow{\psi_1} S_q \otimes E_1^* \xrightarrow{\psi_2} \text{Hom}(E_1, S_q) \xrightarrow{\psi_3} \text{Hom}(S_1, S_q)$$

where  $\psi_1$  is a skew isomorphism of  $G$ -modules induced by the natural duality in the Grassman algebra, where  $\psi_2$  is an isomorphism of  $G$ -modules induced by the natural isomorphism of vector spaces, and  $\psi_3$  is the isomorphism of  $G$ -modules induced by the identification of  $E_1$  with  $S_1$ . The composite map

$$\psi: S_q \otimes E_{n-1} \rightarrow \text{Hom}(S_1, S_q)$$

is a skew isomorphism of  $G$ -modules. Let  $\eta = \psi(df_1 \cdots df_{n-1})$ . Since  $df_1 \cdots df_{n-1} \in I(S_q \otimes E_{n-1})$  we have  $\eta \in \hat{I}(\text{Hom}(S_1, S_q))$  so that  $\eta$  is a skew homomorphism of  $S_1$  into  $S_q$ . Since  $S_1$  is irreducible,  $\eta$  must be injective, and from the definition of the map  $\psi$  we see that  $\eta x_i = u_i$  where  $u_i$  is the minor of order  $n-1$  of  $J$  obtained by deleting the derivatives of  $f_n$  and the derivatives with respect to  $x_i$ .

Our map  $d: S \rightarrow S \otimes E_1$  is homogeneous of degree  $-1$  and commutes with the action of  $G$ . Let  $\hat{d} = d \circ \eta$ . Then  $\hat{d}$  is a skew isomorphism of  $S_1$  into  $S_{q-1} \otimes E_1$ . Since  $S$  is a polynomial ring we may extend  $\hat{d}$  to a derivation  $\hat{d}: S \rightarrow S \otimes E_1$  of  $S$ -modules. By induction on the degree of a homogeneous element one sees that  $\hat{d}\gamma s = (\det \gamma)\gamma \hat{d}s$  for all  $\gamma \in G$  and all  $s \in S$ . Thus  $\hat{d}$  maps  $I(S)$  into  $\hat{I}(S \otimes E_1)$ . Since  $\hat{d}x_i = du_i = \sum_k (\partial u_i / \partial x_k) dx_k$  we have

$$(7.1) \quad \hat{d}f = \sum_i \sum_k \frac{\partial f}{\partial x_i} \frac{\partial u_i}{\partial x_k} dx_k.$$

We claim that the elements  $\hat{d}f_1, \dots, \hat{d}f_n$  are linearly independent over  $S$ . If not, then we have a relation  $\sum_i s_i \hat{d}f_i = 0$  where  $s_i \in S$  and where  $s_1$ , say, is not zero. Then multiplication by  $\hat{d}f_2 \cdots \hat{d}f_n$  shows that  $\hat{d}f_1 \cdots \hat{d}f_n = 0$ . On the other hand, computing directly from (7.1) shows that

$$\hat{d}f_1 \cdots \hat{d}f_n = \det \left( \frac{\partial f_i}{\partial x_k} \right) \det \left( \frac{\partial u_i}{\partial x_k} \right) dx_1 \cdots dx_n$$

which is not zero in view of our assumption about the algebraic independence of the  $u_i$ . Thus the  $\hat{d}f_i$  are linearly independent over  $S$  and the sum  $P = \sum_i I(S) \hat{d}f_i$  is direct. Since  $G$  is a real group it has an invariant quadratic form  $f_1$  and hence  $m_1 = 1$ . The degree of the map  $\hat{d}$  is thus  $q-1 = m_1 + \cdots + m_{n-1} - 1 = m_2 + \cdots + m_{n-1}$  and the Poincaré series for the graded vector space  $P$  is thus

$$P(t) = \frac{t^{m_2 + \cdots + m_{n-1}} (t^{m_1} + \cdots + t^{m_n})}{(1 - t^{m_1+1}) \cdots (1 - t^{m_n+1})}.$$

On the other hand, using (5.1), (6.1) and the double duality we see that  $P(t) = I(S \otimes E_{n-1})(t) = \hat{I}(S \otimes E_1)(t)$ . Since  $P \subseteq \hat{I}(S \otimes E_1)$  is an inclusion of graded vector spaces we have  $P = \hat{I}(S \otimes E_1)$ . Thus  $\hat{I}(S \otimes E_1)$  is freely generated over  $I(S)$  by  $\hat{d}f_1, \dots, \hat{d}f_n$  and Theorem 3 is proved.

8. For the symmetric group on  $n$  letters  $x_1, \dots, x_n$  we can give the following construction for the skew invariant differential 1-forms. Let  $\sigma_1, \dots, \sigma_n$  be the elementary symmetric functions of  $x_1, \dots, x_n$  and let  $\Delta(x_1, \dots, x_{n-1}) = \prod_{1 \leq i < j \leq n-1} (x_i - x_j)$  be the fundamental skew invariant polynomial for the symmetric group on the letters  $x_1, \dots, x_{n-1}$ . Then a basis for the skew invariant differential 1-forms over the algebra of symmetric functions is given by the forms

$$\omega_k = \sum_i (-1)^{i+1} \Delta(x_1, \dots, \tilde{x}_i, \dots, x_n) \frac{\partial \sigma_k}{\partial x_i} dx_i, \quad k = 1, \dots, n,$$

where  $\tilde{x}_i$  means that the letter  $x_i$  is to be omitted.

## REFERENCES

1. W. Burnside, *The determination of all groups of rational linear substitutions of finite order which contain the symmetric group in the variables*, Proc. London Math. Soc. (2) **10** (1912), 284–308.
2. C. Chevalley, *Invariants of finite groups generated by reflections*, Amer. J. Math. **77** (1955), 778–782.
3. A. J. Coleman, *The Betti numbers of the simple Lie groups*, Canad. J. Math. **10** (1958), 349–356.
4. H. S. M. Coxeter, *The product of the generators of a finite group generated by reflections*, Duke Math. J. **18** (1951), 765–782.
5. J. S. Frame, *The classes and representations of the groups of 27 lines and 28 bitangents*, Ann. Mat. Pura Appl. **32** (1951), 83–119.
6. G. Frobenius, *Über die Charaktere der symmetrischen Gruppe*, S.-B. Preuss. Akad. Wiss. Berlin (1900), 516–534.
7. B. Kostant, *The principal three dimensional subgroup and the Betti numbers of a complex simple Lie group*, Amer. J. Math. **81** (1959), 973–1032.
8. G. C. Shephard and J. A. Todd, *Finite unitary reflection groups*, Canad. J. Math. **6** (1954), 274–304.
9. L. Solomon, *Invariants of finite reflection groups*, Nagoya Math. J. **22** (1963), 57–64.

INSTITUTE FOR ADVANCED STUDY,  
PRINCETON, NEW JERSEY  
HAVERFORD COLLEGE,  
HAVERFORD, PENNSYLVANIA