

GENERALIZED COMMUTING PROPERTIES OF MEASURE-PRESERVING TRANSFORMATIONS⁽¹⁾

BY
ROY L. ADLER

1. Introduction. Let (X, \mathfrak{B}, m) be a measure space where X is a set of elements, \mathfrak{B} a σ -field of measurable subsets of X , and m a countably additive measure defined on \mathfrak{B} . A measure-preserving transformation T of the measure space (X, \mathfrak{B}, m) is almost everywhere a one-to-one mapping of X into itself such that if $B \in \mathfrak{B}$ then TB and $T^{-1}B \in \mathfrak{B}$ with $m(TB) = m(T^{-1}B) = m(B)$. Let G be the group of all measure-preserving transformations of X onto itself with I denoting the identity transformation on X . Associated with a measure-preserving transformation T is a sequence $C_n(T)$, $n = 0, 1, 2, \dots$, of subfamilies of G defined inductively as follows:

$$\begin{aligned} C_0(T) &= \{S : S \in G, S = I \text{ a.e.}\}, \\ C_n(T) &= \{S : S \in G, STS^{-1}T^{-1} \in C_{n-1}(T)\}. \end{aligned}$$

In view of the observation that $C_1(T)$ is the set of measure-preserving transformations which commute with T almost everywhere $C_n(T)$ will be called the *n*th class of generalized T -commuting transformations. It is clear that

$$C_n(T) \subseteq C_{n+1}(T), \quad n = 0, 1, 2, \dots$$

If there exists an integer N such that $C_N(T) = C_{N+1}(T)$ then $C_n(T) = C_{n+1}(T)$ for all $n \geq N$; and in this case we define $N(T) = \min\{N : C_N(T) = C_{N+1}(T)\}$, otherwise we set $N(T) = \infty$. We call $N(T)$ the *generalized commuting order* of T . If $RT_1R^{-1} = T_2$ a.e., where $R, T_1, T_2 \in G$ then it is clear that $N(T_1) = N(T_2)$, i.e., $N(T)$ is a conjugacy invariant or spatial isomorphism invariant for measure-preserving transformations. It is of course not to be expected that the converse holds.

2. Pure point spectrum. In order to avoid pathologies which can arise with too liberal a choice of measure space we impose the mild restriction that (X, \mathfrak{B}, m) be a Lebesgue space [8] with $m(X) = 1$. A measure-preserving transformation T is said to be ergodic if $TB = B$, $B \in \mathfrak{B}$ implies

Received by the editors December 11, 1963.

⁽¹⁾ The material in this paper was included in the author's doctoral dissertation submitted to Yale University (1961) under the direction of Professor S. Kakutani.

$m(B) = 0$ or $m(X - B) = 0$. A measure-preserving transformation T induces a unitary operation U_T on $L^2(X, \mathfrak{B}, m)$ defined by $U_T: f(x) \rightarrow f(Tx)$. The proper values and the proper functions of this unitary operator are also called proper values and proper functions of the measure-preserving transformation; furthermore, it is clear that if a complex number c is a proper value of T then $|c| = 1$. The proper value theorem [3, p. 34] relates the concept of ergodicity of a measure-preserving transformation with its proper values. A measure-preserving transformation T on a space of finite measure is ergodic if and only if the number 1 is a simple proper value of T ; moreover, if a measure-preserving transformation T is ergodic the absolute value of every proper function of T is constant a.e., every proper value is simple, and the set of all proper values of T is a subgroup of the circle group. A measure-preserving transformation is said to have pure point spectrum if $L^2(X, \mathfrak{B}, m)$ can be spanned by its proper functions. In connection with generalized commuting properties of measure-preserving transformations the following theorem can be proved.

THEOREM. *If a measure-preserving transformation T and all of its powers are ergodic and T has pure point spectrum then its commuting order is two. Furthermore, $C_0(T)$, $C_1(T)$ and $C_2(T)$ are subgroups of G .*

We shall discuss this theorem and its proof in the light of the representation theorem for ergodic measure-preserving transformations with pure point spectrum on Lebesgue spaces. P. R. Halmos and J. von Neumann [4] showed such a transformation could be considered as measure theoretically the same as a rotation $T_a: x \rightarrow ax$ which is Haar measure-preserving on some compact separable Abelian group⁽²⁾. They also proved that a rotation $T_a: x \rightarrow ax$ on a compact separable Abelian group X is ergodic if and only if $\{a^n: n = 0, \pm 1, \pm 2, \dots\}$ is dense in X . Such groups admitting ergodic rotations are called monothetic and are discussed in [2] and [5]. The above theorem can now be stated in the following more detailed form.

THEOREM. *Let T_a and all its powers be ergodic on a compact separable Abelian group X ; then $N(T_a) = 2$. Furthermore, $C_0(T_a)$, $C_1(T_a)$ and $C_2(T_a)$ are subgroups of G which can be characterized as follows: (i) $C_0(T_a)$ is, as defined, the family of measure-preserving transformations almost everywhere equal to the identity; (ii) $C_1(T_a)$ is the family of measure-preserving transformations almost everywhere equal to rotations on the group; (iii) $C_2(T_a)$ is the family of measure-preserving transformations almost everywhere equal to rotations composed with continuous automorphisms of the group X .*

⁽²⁾ The standard facts concerning topological groups used in this work are contained in [6] and [7]. By separable we mean satisfying the second axiom of countability.

We shall establish the theorem in a sequence of propositions the first of which is to be found in [4] with only an indication of its proof.

PROPOSITION 1. *Let T_a be an ergodic rotation on a compact separable Abelian group X . If $S \in C_1(T_a)$ i.e. $ST_a = T_aS$ a.e., there exists $b \in X$ such that $S = T_b$ a.e.*

Proof. Let $x^* \in X^*$ the character group of X . Since S commutes with T_a a.e., we have

$$x^*(ST_a x) = x^*(T_a Sx) = x^*(a \cdot Sx) = x^*(a)x^*(Sx) \quad \text{a.e.}$$

Thus $x^*S(\cdot)$ is a proper function of T_a associated with the proper value $x^*(a)$. But $x^*(\cdot)$ also is a proper function T_a associated with the proper value $x^*(a)$. The proper value theorem asserts that all proper values of an ergodic measure-preserving transformation are simple. Hence for each $x^* \in X^*$

$$x^*(Sx) = \phi(x^*)x^*(x) \quad \text{a.e.,}$$

where $\phi(x^*)$ is a constant of absolute value 1 which depends only on the character x^* . This relation yields

$$x_1^* x_2^*(Sx) = \phi(x_1^* x_2^*) x_1^* x_2^*(x) \quad \text{a.e.,}$$

$$x_1^* x_2^*(Sx) = x_1^*(Sx) x_2^*(Sx) = \phi(x_1^*) \phi(x_2^*) x_1^*(x) x_2^*(x) \quad \text{a.e.}$$

for $x_1^*, x_2^* \in X^*$ which in turn imply

$$\phi(x_1^* x_2^*) = \phi(x_1^*) \phi(x_2^*),$$

$$\phi(x^{*-1}) = \overline{\phi(x^*)}$$

for $x_1^*, x_2^*, x^* \in X^*$. In other words ϕ is a homomorphism of the character group X^* into the multiplicative group of complex numbers of absolute value 1. The character group of a compact group is discrete so ϕ is automatically continuous. Thus ϕ is a character on X^* , i.e., $\phi \in X^{**}$. By Pontrjagin's duality theorem, there exists an element $b \in X$ such that $\phi(x^*) = x^*(b)$. Thus $x^*(Sx) = x^*(b)x^*(x) = x^*(bx)$ a.e. Let N_x be the zero measure set of points x where $x^*(Sx) \neq x^*(bx)$. Since X is separable X^* is countable so that $N = \bigcup_{x^* \in X^*} N_x$ is also of measure zero. Therefore for all x^* we have $x^*(Sx) = x^*(bx)$, $x \in X - N$. From the Peter-Weyl theorem there are sufficiently many characters to distinguish the elements of X ; consequently $Sx = bx$, $x \in X - N$.

The commutativity of X states that if $S = T_b$ a.e. for some $b \in X$, then $S \in C_1(T_a)$. This fact along with Proposition 1 completely characterizes the first class of generalized commuting transformations of T_a .

PROPOSITION 2. *Let T_a be an ergodic rotation on a compact separable Abelian group X . If $S \in C_2(T_a)$, then there exists an element $b \in X$ and a*

continuous automorphism K of X such that $S = T_b K$ a.e.

Proof. Essentially we are characterizing those measure-preserving transformations which supply spatial isomorphisms between ergodic rotations. By hypothesis $ST_a S^{-1} T_a^{-1} \in C_1(T_a)$. Proposition 1 assures the existence of an element $c \in X$ such that $ST_a S^{-1} T_a^{-1} = T_c$ a.e. or $ST_a S^{-1} = T_d$ a.e., where $d = ac$. For each $x^* \in X^*$ the function $x^* S(\cdot)$ is a proper function of T_a associated with the proper value $x^*(d)$ because $x^*(ST_a x) = x^*(T_a Sx) = x^*(d \cdot Sx) = x^*(d)x^*(Sx)$ a.e. Next we shall show that for each $x^* \in X^*$ there exists a character denoted by $K^* x^*$ such that $K^* x^*(a) = x^*(d)$, for assuming the contrary there would exist $y^* \in X^*$ such that $y^*(d) \neq x^*(d)$ for any $x^* \in X^*$. As is well known, proper functions associated with different proper values are orthogonal in $L^2(X)$; so $y^* S(\cdot)$ having $y^*(d)$ as a proper value would be orthogonal to every $x^* \in X^*$. Since X^* is a complete orthonormal system in $L^2(X)$ a contradiction would immediately arise. For each $x^* \in X^*$ the character $K^* x^*$ is uniquely defined for suppose $L^* x^*$ were another satisfying $L^* x^*(a) = x^*(d)$. We would have $L^* x^*(a) = K^* x^*(a)$; that is the two characters would agree on the generating element a which means that they must agree everywhere. The above relation between $K^* x^*$ and x^* then reveals that $K^* x^*(\cdot)$ and $x^* S(\cdot)$ are proper functions of T_a both having the same proper value $x^*(d)$. Since T_a is ergodic, the proper value theorem asserts that $x^*(d)$ is simple. Therefore for each $x^* \in X^*$ there exists a complex number $\phi(x^*)$ depending on x^* such that

$$(*) \quad \phi(x^*) K^* x^*(x) = x^*(Sx) \quad \text{a.e.}$$

In addition it is clear that $|\phi(x^*)| = 1$. The set

$$\{x^* S : x^* \in X^*\} = \{\phi(x^*) K^* x^* : x^* \in X^*\}$$

is the image of X^* under the unitary operator U_s ; consequently, since X^* is complete in $L^2(X)$, so is $\{K^* x^* : x^* \in X^*\}$. Thus we can conclude that the mapping $K^* : x^* \rightarrow K^* x^*$ maps X^* onto itself. From (*) we obtain

$$\begin{aligned} \phi(x_1^* x_2^*) K^* x_1^* x_2^*(x) &= x_1^* x_2^*(Sx) = x_1^*(Sx) x_2^*(Sx) \\ &= \phi(x_1^*) K^* x_1^*(x) \phi(x_2^*) K^* x_2^*(x) \quad \text{a.e.} \end{aligned}$$

for $x_1^*, x_2^* \in X^*$. Since characters are continuous we obtain

$$\phi(x_1^* x_2^*) K^* x_1^* x_2^*(x) = \phi(x_1^*) \phi(x_2^*) K^* x_1^*(x) K^* x_2^*(x)$$

everywhere. In particular for $x = e$, the unit element of the group, the equation yields

$$\phi(x_1^* x_2^*) = \phi(x_1^*) \phi(x_2^*)$$

and consequently

$$K^*x_1^*x_2^* = K^*x_1^*K^*x_2^*$$

for $x_1^*, x_2^* \in X^*$. In addition it follows that

$$\phi(x^{*-1}) = \overline{\phi(x^*)}$$

and

$$K^*x^{*-1} = (K^*x^*)^{-1}$$

for $x^* \in X^*$. Thus ϕ is a homomorphism of X^* into the complex numbers of absolute value 1 and K^* is an endomorphism of X^* . That K^* is one-to-one and thus an automorphism of X^* results also from (*); for suppose $K^*x^* = e^*$, the identity character. Then

$$x^*(Sx) = \phi(x^*)K^*x^*(x) = \phi(x^*)e^*(x) = \text{constant} \quad \text{a.e.}$$

Since S is a measure-preserving transformation of X onto itself, $x^*(x) = \text{constant}$ a.e., and by continuity $x^*(x) = \text{constant}$ everywhere on X . But the only constant character is e^* ; therefore $x^* = e^*$. The continuity of ϕ and K^* is automatically provided by X^* being discrete. Thus ϕ is a character on X^* and the duality theorem of Pontrjagin asserts the existence of an element $b \in X$ such that $\phi(x^*) = x^*(b)$ for $x^* \in X^*$. Furthermore, the continuous automorphism K^* induces a continuous automorphism K^{**} on the character group X^{**} of X^* defined by $K^{**}x^{**}(x^*) = x^{**}(K^*x^*)$. The duality theorem is invoked once again to guarantee the existence of a continuous automorphism K of X satisfying $K^*x^*(x) = x^*(Kx)$ for all $x^* \in X^*$. Then from (*) we have

$$x^*(Sx) = x^*(b)x^*(Kx) = x^*(b \cdot Kx) = x^*(T_bKx) \quad \text{a.e.}$$

for all $x^* \in X^*$. From the countability of X^* and the Peter-Weyl theorem we conclude as before that $S = T_bK$ a.e.

Observe that

$$(T_bK)T_a(T_bK)^{-1}T_a^{-1} = T_{a^{-1} \cdot K(a)}$$

for any $b \in X$ and any continuous automorphism K on X . Thus if $S = T_bK$ a.e. then $S \in C_2(T_a)$. This fact along with Proposition 2 completely characterizes $C_2(T_a)$.

LEMMA. *If T_a is an ergodic rotation on a compact separable Abelian group X which is spatially isomorphic to a rotation T_d composed with a continuous automorphism K , i.e., there exists a measure-preserving transformation S of X onto itself such that $ST_aS^{-1} = T_dK$ a.e., then all characters $x^* \in X^*$ have finite orbits under K .*

Proof. For fixed $x^* \in X^*$ consider the Fourier expansion

$$x^*(Sx) \sim \sum_{\gamma \in \Gamma} c_\gamma x_\gamma^*(x),$$

where the characters are indexed with a countable set Γ with the coefficients c_γ square summable. The automorphism K maps characters into characters, and consequently we shall denote by $x_{K^{-1}\gamma}^*$ the character defined by $x_{K^{-1}\gamma}^*(x) = x_\gamma^*(Kx)$. Since

$$x^*(T_a Sx) = x^*(ST_d Kx) \quad \text{a.e.},$$

$$x^*(a) \cdot x^*(Sx) = x^*(ST_d Kx) \quad \text{a.e.}$$

we obtain from the Fourier expansion

$$\sum_{\gamma \in \Gamma} x^*(a) c_\gamma x_\gamma^*(x) \sim \sum_{\gamma \in \Gamma} c_\gamma x_\gamma^*(d) x_\gamma^*(Kx) \sim \sum_{\gamma \in \Gamma} c_\gamma x_\gamma^*(d) x_{K^{-1}\gamma}^*(x).$$

Reindexing the series we have

$$\sum_{\gamma \in \Gamma} x^*(a) c_\gamma x_\gamma^*(x) \sim \sum_{\gamma \in \Gamma} c_{K\gamma} x_{K\gamma}^*(d) x_\gamma^*(x).$$

Therefore,

$$x^*(a) c_\gamma = c_{K\gamma} x_{K\gamma}^*(d),$$

whence,

$$|c_\gamma| = |c_{K\gamma}|.$$

Suppose $x_{\gamma_0}^*$ has an infinite orbit under K , i.e., $x_{\gamma_0}^*, x_{K\gamma_0}^*, \dots, x_{K^n\gamma_0}^*, \dots$ were all distinct. Since $|c_{\gamma_0}| = |c_{K\gamma_0}| = \dots = |c_{K^n\gamma_0}| = \dots$ and the coefficients $c_\gamma, \gamma \in \Gamma$, are square summable we would have $c_{\gamma_0} = 0$. Thus in the Fourier expansion of x^*S if the coefficient $c_\gamma \neq 0$ then there exists an integer n depending upon γ such that $x_{K^n\gamma}^* = x_\gamma^*$. Because $\{x^*S : x^* \in X^*\}$ is complete in $L^2(X)$ and every $x^*S, x^* \in X^*$ can be expanded in terms of characters whose orbits under K are finite, the set of characters whose orbits are finite under K is also complete in $L^2(X)$. This combined with the fact that characters are orthogonal to each other shows that all characters must have finite orbits under K .

PROPOSITION 3. *Let T_a and all of its powers $T_{a^n} = T_a^n, n = 0, 1, 2, \dots$, be ergodic rotations on a compact separable Abelian group X . Then $C_3(T_a) = C_2(T_a)$.*

Proof. Let $S \in C_3(T_a)$. From Proposition 2 we have $ST_a S^{-1} T_a^{-1} = T_b K$ a.e., where K is a continuous automorphism of X and T_b is some rotation on X . Thus $ST_a S^{-1} = T_d K$, where $d = K(a)b$. Consider the function $x^*S(\cdot)$, where x^* is some character of X . By the preceding lemma there exists an integer n depending on x^* such that $x^*(K^n x) = x^*(x)$; consequently,

$$x^*(ST_a^n x) = x^*((T_d K)^n Sx) = x^*(d_n \cdot K^n Sx) = x^*(d_n) x^*(Sx) \quad \text{a.e.},$$

where $d_n = d \cdot Kd \dots K^{n-1}d$. The function $x^*S(\cdot)$ is a proper function of

T_a^n which by hypothesis is ergodic. Invoking the proper value theorem we have that x^*S is almost everywhere equal to a constant $\phi(x^*)$ multiplied by some character, denoted by L^*x^* , i.e.,


$$x^*(Sx) = \phi(x^*)L^*x^*(x) \quad \text{a.e.}$$

for $x^* \in X^*$. But this is the previous relation (*). From this and the orthogonality of characters follows that L^*x^* is uniquely defined. Then proceeding in exactly the same manner as Proposition 2 we show that there exists $c \in X$ and a continuous automorphism L of X such that $S = T_cL$ a.e., in other words $S \in C_2(T_a)$.


COUNTEREXAMPLE. In order to demonstrate the essential role in Proposition 3 played by the hypothesis that all powers of T_a be ergodic consider the finite cyclic group $G_9 = \{0, 1, 2, 3, 4, 5, 6, 7, 8\}$ the integers with addition modulo 9. The rotation $T_a: x \rightarrow x + a$ (modulo 9) is ergodic if and only if a is a generating element of G_9 , i.e., if and only if a is relatively prime to 9. Furthermore, since 9 is not a prime number every ergodic rotation has a power which is not ergodic. We can enumerate all of the automorphisms of G_9 by K_k , where k is an integer less than and relatively prime to 9. The automorphism K_k is completely determined by the relation $K_k(1) = k$. The possible choices for k are $k = 1, 2, 4, 5, 7$ and 8. We denote by I the identity automorphism and by J the involution automorphism, $J: x \rightarrow -x$ (modulo 9). Since $K^n(1) = k^n$ (modulo 9), $2^6 \equiv 4^3 \equiv 5^6 \equiv 7^3 \equiv 8^2 \equiv 1$ (modulo 9) and $5^3 \equiv 4$ (modulo 9), we have the following relations:

$$K_1 = (K_2)^6 = (K_4)^3 = (K_5)^6 = (K_7)^3 = (K_8)^2 = I, \\ (K_5)^3 = K_4, K_3 = J.$$

Consider the rotation $T_1: x \rightarrow x + 1$ (modulo 9) and the permutation T_1K_4 on G_9 . Their behavior on G_9 is described in the diagrams below:



The transformations T_1 and T_1K_4 are spatially isomorphic; the relation $S^{-1}T_1S = T_1K_4$ is satisfied by a permutation S whose behavior on G_9 is determined by the following diagram:



Thus $S^3 = I$. Next we shall show that S is not of the form $T_a K$, where T_a is some rotation on G_9 and K is one of the automorphisms of G_9 listed above. Suppose $S = T_a K$. Then $(T_a K)^3 = I$. From this we obtain

$$T_b K^3 = I,$$

where $b \equiv K^2(a) + K(a) + a$ (modulo 9). Thus K^{-3} is an automorphism that is also a rotation: the only such automorphism is $K^{-3} = K^3 = I$. This reduces the possibilities for K to those automorphisms whose cube is the identity, i.e., $K = K_4$ or $K = K_7$. Suppose $S = T_a K_4$. We get

$$2 = S(1) = T_a K_4(1) = 4 + a \pmod{9}$$

consequently, $a = 7$. On the other hand

$$0 = S(2) = T_7 K_4(2) = 8 + 7 = 6 \pmod{9}$$

which is a contradiction. The remaining possibility is $S = T_a K_7$. Here, however, we get

$$2 = S(1) = T_a K_7(1) = a + 7 \pmod{9}$$

so that $a = 4$. On the other hand

$$0 = S(2) = T_4 K_7(2) = 7 + 14 = 3 \pmod{9}$$

which is again impossible. Therefore S is not in $C_3(T_1)$.

3. Anzai's skew product transformations. Let X, Y be unit intervals with Borel measurability and Lebesgue measure and $Z = X \times Y$ be the unit square with the usual direct product measurability and measure. We shall consider the following skew product measure-preserving transformation defined on Z ; let $T_{\gamma, \alpha}$ denote the measure-preserving transformation with α -function defined by $T_{\gamma, \alpha}: (x, y) \rightarrow (x + \gamma, y + \alpha(x))$ (additions modulo 1) where γ is an irrational number and $\alpha(\cdot)$ a real-valued measurable function defined on X . Conditions for ergodicity of $T_{\gamma, \alpha}$ along with proof that it is measure-preserving can be found in Anzai's paper [1]. Two other results from [1] upon which the subsequent work depends are the following: (Proper value criterion) The value $e^{2\pi i \lambda}$ is a proper value of $T_{\gamma, \alpha}$ if and only if there exists an integer p and a real-valued meas-

urable function $\theta(\cdot)$ on X such that $\lambda - p\alpha(x) = \theta(x + \gamma) - \theta(x)$ (modulo 1) a.e.; (Spatial isomorphism criterion) If S is a measure-preserving transformation such that $ST_{\gamma,\alpha}S^{-1} = T_{\gamma,\beta}$ a.e. where $T_{\gamma,\alpha}$ and $T_{\gamma,\beta}$ are ergodic skew product transformations with α -function and β -function, respectively, then S has either the form

$$(i) \quad S: (x, y) \rightarrow (x + u, y + \theta(x))$$

(additions modulo 1) where u is a real number and $\theta(\cdot)$ is a real-valued measurable function on X such that

$$\beta(x + u) - \alpha(x) = \theta(x + \gamma) - \theta(x) \pmod{1} \quad \text{a.e.}$$

or

$$(ii) \quad S: (x, y) \rightarrow (x + u, \theta(x) - y)$$

(additions modulo 1) where u and $\theta(\cdot)$ now satisfy

$$\beta(x + u) + \alpha(x) = \theta(x + \gamma) - \theta(x) \pmod{1} \quad \text{a.e.}$$

We shall restrict ourselves to the most tractable class of skew product transformation, namely, when $\alpha(x) = \nu x$ for some nonzero integer ν .

THEOREM. *The commuting order $N(T_{\gamma,\nu}) = 2$. Furthermore, $C_0(T_{\gamma,\nu})$, $C_1(T_{\gamma,\nu})$ and $C_2(T_{\gamma,\nu})$ are subgroups of (X) which can be characterized in the following manner: (i) $C_0(T_{\gamma,\nu})$ is as defined, the family of measure-preserving transformations almost everywhere equal to the identity; (ii) $C_1(T_{\gamma,\nu})$ is the family of measure-preserving transformations almost everywhere equal to some power of a ν th root of $T_{\gamma,\nu}$ composed with a transformation R of the form $R: (x, y) \rightarrow (x, y + c)$ where c is some constant; (iii) $C_2(T_{\gamma,\nu})$ is the family of measure-preserving transformations almost everywhere equal to one of the forms*

$$\begin{aligned} (x, y) &\rightarrow (x + u, y + Kx + c), \\ (x, y) &\rightarrow (-x + u, y + Kx + c), \\ (x, y) &\rightarrow (x + u, Kx - y + c), \\ (x, y) &\rightarrow (-x + u, Kx - y + c), \end{aligned}$$

where K is an arbitrary integer and u and c are arbitrary real numbers.

This theorem will be established in a sequence of propositions.

PROPOSITION 1. *If $S \in C_1(T_{\gamma,\nu})$, i.e., $ST_{\gamma,\nu} = T_{\gamma,\nu}S$ a.e., then S almost everywhere is of the form $S(x, y) \rightarrow (x + n\gamma/\nu, y + nx + c)$ where c is some real constant; in other words*

$$S = (T_{\gamma,\nu})^{n/\nu} R \quad \text{a.e.,}$$

where $(T_{\gamma,\nu})^{1/\nu}$ is a ν th root of T , and R is a measure-preserving transformation

defined by $R: (x, y) \rightarrow (x, y + c)$ (additions modulo 1) for some real constant c .

Proof. Applying Anzai's result to S we see that it must have the form:

$$(i) \quad S: (x, y) \rightarrow (x + u, y + \theta(x)),$$

where $\nu u = \theta(x + \gamma) - \theta(x)$ (modulo 1) a.e. for some real-valued measurable function $\theta(\cdot)$ or

$$(ii) \quad S: (x, y) \rightarrow (x + u, \theta(x) - y),$$

where $2\nu x + \nu u = \theta(x + \gamma) - \theta(x)$ (modulo 1) a.e.

Case (ii) can be immediately eliminated: for it follows that the function $e^{2\pi i \theta(\cdot)}$ is a "generalized proper value" (see [3, p. 57]) of the rotation $T_\gamma: x \rightarrow x + \gamma$ (modulo 1) on the circle, i.e.,

$$e^{2\pi i \theta(x+\gamma)} = e^{2\pi i \nu u} e^{2\pi i 2\nu x} e^{2\pi i \theta(x)} \quad \text{a.e.}$$

but for T_γ the only "generalized proper values" are proper functions themselves, i.e.,

$$e^{2\pi i \theta(x)} = e^{2\pi i n x + c}$$

for some integer n and real number c ; therefore

$$2\nu x + \nu u = n(x + \gamma) - nx = n\gamma \quad (\text{modulo } 1) \quad \text{a.e.}$$

which is impossible.

Suppose S satisfies case (i); then the function $e^{2\pi i \theta(\cdot)}$ is a proper function with $e^{2\pi i n \gamma}$ the associated proper value of the ergodic rotation T_γ whose only proper values are $e^{2\pi i n \gamma}$, each of which corresponds to a proper function which is almost everywhere equal to a constant times the function which takes the values $e^{2\pi i n x}$. Therefore $e^{2\pi i \nu u} = e^{2\pi i n \gamma}$ and $e^{2\pi i \theta(x)} = e^{2\pi i (n x + c)}$ a.e. for some n from which we see that S has the form

$$S: (x, y) \rightarrow (x + n\gamma/\nu, y + nx + c)$$

(additions modulo 1) where c is some real constant. Since they also commute with $T_{\gamma, \nu}$, the ν th roots of $T_{\gamma, \nu}$ must be of the above form with $n = 1$ from which we can conclude that

$$(T_{\gamma, \nu})^{1/\nu}: (x, y) \rightarrow (x + \gamma/\nu, y + x - (\nu - 1)\gamma/2 + p/\nu)$$

(additions modulo 1), where $p = 0, 1, 2, \dots, \nu - 1$. The constant c then can be adjusted so that

$$S = (T_{\gamma, \nu})^{n/\nu} R \quad \text{a.e.,}$$

where $R: (x, y) \rightarrow (x, y + c)$ (additions modulo 1).

It is easy to verify that transformations almost everywhere equal to one of the forms $(T_{\gamma, \nu})^{n/\nu} R$ commute with $T_{\gamma, \nu}$ almost everywhere. This fact along with Proposition 1 completely characterizes $C_1(T_{\gamma, \nu})$.

PROPOSITION 2. If $S \in C_2(T_{\gamma,\nu})$ then S almost everywhere satisfies one of the forms

- (i) $(x, y) \rightarrow (x + u, y + Kx + c),$
- (ii) $(x, y) \rightarrow (x + u, Kx - y + c),$
- (iii) $(x, y) \rightarrow (-x + u, Kx + y + c),$
- (iv) $(x, y) \rightarrow (-x + u, Kx - y + c),$

where K is an arbitrary integer and u and c are arbitrary real numbers.

Proof. By Proposition 1

$$ST_{\gamma,\nu}S^{-1}T_{\gamma,\nu}^{-1} = (T_{\gamma,\nu})^{n/\nu}R \quad \text{a.e.,}$$

where $R: (x, y) \rightarrow (x, y + c)$; whence

$$(*) \quad ST_{\gamma,\nu}S^{-1} = (T_{\gamma,\nu})^{(n+\nu)/\nu}R \quad \text{a.e.}$$

The transformation on the right is the ergodic skew product transformation $T_{(n+\nu)\gamma/\nu,\beta}$ with β -function which satisfies $\beta(x) = (n + \nu)x + c'$ in which all the constants involved are lumped together in c' ; whereupon we write (*) as $ST_{\gamma,\nu}S^{-1} = T_{(n+\nu)\gamma/\nu,\beta}$. Applying Anzai's criterion for proper values and the familiar argument of "generalized proper values" used before (Proposition 1) we see that the proper values of $T_{\gamma,\nu}$ are precisely the positive and negative integral powers of $e^{2\pi i\gamma}$ whereas the proper values of $T_{(n+\nu)\gamma/\nu,\beta}^\dagger$ are precisely the positive and negative integral powers of $e^{2\pi i(n+\nu)\gamma/\nu}$. Since $T_{\gamma,\nu}$ is isomorphic to $T_{(n+\nu)\gamma/\nu,\beta}$, we have that $e^{2\pi i\gamma}$ is some power of $e^{2\pi i(n+\nu)\gamma/\nu}$ and vice versa. This can only happen if $(n + \nu)/\nu = \pm 1$. For the case $(n + \nu)/\nu = 1$ we can apply Anzai's spatial isomorphism criterion which yields that S has either the form

$$S: (x, y) \rightarrow (x + u, y + \theta(x))$$

(additions modulo 1), where u is a real number and $\theta(\cdot)$ is a real-valued measurable function such that

$$(n + \nu)(x + u) + c' - \nu x = \theta(x + \gamma) - \theta(x) \quad (\text{modulo } 1) \quad \text{a.e.}$$

or

$$S: (x, y) \rightarrow (x + u, \theta(x) - y)$$

(additions modulo 1), where

$$(n + \nu)(x + u) + c' + \nu x = \theta(x + \gamma) - \theta(x) \quad (\text{modulo } 1) \quad \text{a.e.}$$

In either case the argument of "generalized proper values" demonstrates that $\theta(x) = Kx + c$ a.e. for some integer K and real number c ; consequently, S almost everywhere satisfies one of the forms (i) or (ii) in the statement of the proposition. For the case $(n + \nu)/\nu = -1$ we have that $n = -2\nu$ and $\beta(x) = -\nu x + c'$. If Q denotes the mapping $(x, y) \rightarrow (-x, -y)$ then

$$QT_{-\gamma,\beta}Q^{-1} = T_{\gamma,\beta'},$$

where $\beta'(x) = \nu x - c'$. From this and relation (*) we have

$$QST_{\gamma,\nu}S^{-1}Q^{-1} = T_{\gamma,\beta'} \quad \text{a.e.}$$

to which we apply the considerations of the previous case to obtain that QS satisfies almost everywhere one of the forms (i) and (ii) in the statement of the proposition. Hence S must satisfy almost everywhere one of the forms (iii) or (iv).

It is easy to check that all transformations almost everywhere equal to one of the four forms belong to $C_2(T_{\gamma,\nu})$. This fact along with Proposition 2 completely characterizes $C_2(T_{\gamma,\nu})$.

PROPOSITION 3. $C_3(T_{\gamma,\nu}) = C_2(T_{\gamma,\nu})$.

Proof. If $S_3 \in C_3(T_{\gamma,\nu})$ then $S_3 T_{\gamma,\nu} S_3^{-1} T_{\gamma,\nu}^{-1} = S_2$ a.e., where S_2 has one of the four forms described in the previous proposition. Thus we can write

$$S_3 T_{\gamma,\nu} S_3^{-1} = T \quad \text{a.e.,}$$

where T has one of the forms

- (i) $(x, y) \rightarrow (x + u, y + Kx + c),$
- (ii) $(x, y) \rightarrow (x + u, Kx - y + c),$
- (iii) $(x, y) \rightarrow (-x + u, y + Kx + c),$
- (iv) $(x, y) \rightarrow (-x + u, Kx - y + c),$

where K is an arbitrary integer and u and c are arbitrary real numbers.

For case (i) we show $u = \pm \gamma$ exactly as in the previous proposition; in addition $K \neq 0$ if T is to be isomorphic to $T_{\gamma,\nu}$. Furthermore we can apply the isomorphism criterion of Anzai in the same manner as in Proposition 2 to solve for S_3 ; whereupon it is clear that S_3 is one of the four forms of $C_2(T_{\gamma,\nu})$. Cases (ii), (iii) and (iv) can be reduced to case (i) by taking squares and considering $S_3 T_{\gamma,\nu}^2 S_3^{-1} = T^2$, and we solve for S_3 in the identical manner obtaining the same results.

4. Some unsolved problems. 1. What are $N(T_a)$ and $C_n(T_a)$ for ergodic rotations T_a whose powers are not necessarily ergodic?

2. What are $N(T)$ and $C_n(T)$, where T is the shift transformation on a two-sided infinite direct product of a measure space with itself consisting of a finite number of points?

3. Give examples for $N(T) = n$ for each integer n including $N(T) = \infty$.

4. In general is $C_n(T)$ a subgroup of G ?

REFERENCES

1. H. Anzai, *Ergodic skew product transformations on the torus*, Osaka Math. J. 3 (1951), 83-99.
2. H. Anzai and S. Kakutani, *Bohr compactifications of a locally compact abelian group*. I, II, Proc. Imp. Acad. Tokyo 19 (1943), 476-480; ibid. 19 (1943), 533-539.
3. P. R. Halmos, *Lectures on ergodic theory*, Publications of the Mathematical Society of Japan, no. 3, The Mathematical Society of Japan, Tokyo, 1956.

4. P. R. Halmos and J. von Neumann, *Operator methods in classical mechanics*. II, Ann. of Math. (2) 43 (1942), 332-350.
5. P. R. Halmos and H. Samelson, *On monothetic groups*, Proc. Nat. Acad. Sci. U.S.A. 28 (1942), 254-258.
6. L. H. Loomis, *An introduction to abstract harmonic analysis*, Van Nostrand, New York. 1953.
7. L. Pontrjagin, *Topological groups*, Princeton Univ. Press, Princeton, N. J., 1948.
8. V. A. Rohlin, *On fundamental ideas of measure theory*, Amer. Math. Soc. Transl. No. 71 (1952).

INTERNATIONAL BUSINESS MACHINES CORPORATION,
YORKTOWN HEIGHTS, NEW YORK