

SUBGROUPS OF THE MULTIPLICATIVE GROUP OF A DIVISION RING

BY

R. J. FAUDREE⁽¹⁾

Introduction. The general question of what groups can be embedded in the multiplicative group of a division ring is an unsolved problem, but some parts of the problem have been completely solved. It is well known that an abelian group can be embedded in a division ring if and only if its torsion subgroup is locally cyclic (see [4]). B. H. Neumann (see [10]) and A. I. Malcev (see [9]) showed that all ordered groups can be embedded in a division ring and Amitsur solved the problem for finite groups (see [2]). These results exhaust the major work that has been done on the problem to date.

Iwasawa showed that any torsion-free nilpotent group can be ordered (see [7]); therefore, by Neumann's and Malcev's result any torsion-free nilpotent group can be embedded in a division ring. This paper will extend this result by showing that certain torsion-free solvable groups can be embedded in a division ring. This paper will also extend the results on abelian, finite and ordered groups by giving an embedding theorem for a group G which has an abelian, finite or ordered group H as a normal subgroup and a certain normal structure between H and G .

An embedding theorem. Let Γ denote a fixed division ring, x an indeterminate over Γ and θ an automorphism of Γ . $\Gamma\langle x; \theta \rangle$ will denote the Hilbert division ring of all formal power series in x over Γ (see [9]) $\Gamma\langle x; \theta \rangle = \{ \sum_{i=0}^{\infty} a_i x^i \mid a_i \text{ in } \Gamma \text{ and } n \text{ any integer} \}$, where addition and equality are defined as in a commutative power series ring and multiplication is determined by the equation $x \cdot a = \theta(a) \cdot x$ for all a in Γ . Let $\Gamma[x; \theta]$ denote the ring of polynomials in $\Gamma\langle x; \theta \rangle$, that is the ring generated by Γ and x ; and let $\Gamma(x; \theta)$ denote the division ring generated by Γ and x . $\Gamma[x; \theta]$ is an Ore polynomial ring (see [11]) and thus has a quotient field which is uniquely determined up to isomorphism. Therefore $\Gamma(x; \theta) = \{ f(x)g^{-1}(x) \mid f(x), g(x) \neq 0 \text{ in } \Gamma[x; \theta] \} = \{ g^{-1}(x)f(x) \mid f(x), g(x) \neq 0 \text{ in } \Gamma[x; \theta] \}$. This structure of $\Gamma(x; \theta)$ is of importance.

Received by the editors April 4, 1965.

⁽¹⁾ This represents a portion of the author's Doctoral dissertation for the Ph.D. at Purdue University.

In the following discussion G will denote a group, H and H_i subgroups of G ; and D will denote a division ring and D_i subdivision rings of D . The multiplicative group of D will be denoted by D^* , and if G is a subgroup of D^* then $D_i = \{H_i\}$ will mean that D_i is the division ring generated by H_i .

An ascending normal series $\{H_i \mid 0 \leq i \leq \alpha, H_0 = H \text{ and } H_\alpha = G\}$ will be called a *completely infinite supercyclic series from H to G* if each factor group H_{i+1}/H_i is infinite cyclic. If such a series exists G is said to be *completely infinite supercyclic from H to G* . In the case where $H = \{e\}$, the identity subgroup, G is a *completely infinite supercyclic group*.

The class of completely infinite supercyclic groups is a large one. Let G be a solvable group, say $G^{(n)} = \{e\}$, where $G^{(i)}$ is the i th derived group. If $G^{(i)}/G^{(i+1)}$ is a free abelian group for each i , then there obviously is a completely infinite supercyclic series from $G^{(i+1)}$ to $G^{(i)}$; hence such a series exists from $\{e\} = G^{(n)}$ to G . The free solvable groups comprise an important subclass of the above class. Let F be a free group. By Schreiers Theorem $F^{(i)}$ is also a free group so $F^{(i)}/F^{(i+1)}$ is a free abelian group; hence $F/F^{(n)}$ is a completely infinite supercyclic group for any positive integer. J. F. Bowers discussed a class of groups which he called completely infinite polycyclic groups (see [3]) which are also completely infinite supercyclic groups. A finitely generated torsion-free nilpotent group has a completely infinite supercyclic series, in fact one of finite length (see [13, p. 241]).

Let G be a subgroup of D^* such that $D = \langle G \rangle$, let $\{H_i \mid 0 \leq i \leq \alpha\}$ be a completely infinite supercyclic series from H to G with $H_{i+1} \cdot H_i = (x_{i+1} \cdot H_i)$ for x_{i+1} in H_{i+1} , and let $D_i = \langle H_i \rangle$. If x_{i+1} is transcendental over D_i ($0 \leq i < \alpha$) then we call D an *Ore extension of D_0 determined by the completely infinite supercyclic series $\{H_i \mid 0 \leq i \leq \alpha\}$* .

We will now give some elementary properties of Ore extensions. The following proposition is an easy consequence of elementary Ore polynomial ring theory:

PROPOSITION 1. *Let Γ be a subdivision ring of a division ring D and x an element of D which is transcendental over Γ . If $x\Gamma x^{-1} = \Gamma$, then D_1 , the division ring generated by Γ and x , is isomorphic to $\Gamma(x; \theta)$ where θ is the automorphism of Γ induced by x .*

Thus if D is an Ore extension of D_0 determined by the completely infinite supercyclic series $\{H_i \mid 0 < i < \alpha\}$, then D_{i+1} is isomorphic to $D_i(x_{i+1}; \theta_{i+1})$ where θ_{i+1} is the automorphism of D_i induced by x_{i+1} .

PROPOSITION 2. *Let θ and τ be automorphisms of a division ring D , and x and y indeterminates over D . Then θ can be extended to be an automorphism of $D(x; \tau)$ onto $D(y; \theta\tau\theta^{-1})$ which takes x into y .*

Since $x \cdot d = \tau(d)x$ and $y \cdot \theta(d) = \theta\tau(d) \cdot y$ for all d in D , the map determined by $x \rightarrow y$ and $d \rightarrow \theta(d)$ for all d in D is an automorphism of $D(x; \tau)$ onto $D(y; \theta\tau\theta^{-1})$.

PROPOSITION 3. Let G be a group with a normal subgroup H and two completely infinite supercyclic series $\{H_i \mid 0 \leq i \leq \alpha_1\}$ and $\{H'_i \mid 0 \leq i \leq \alpha_2\}$ from H to G . If D is an Ore extension of D_0 determined by the series $\{H_i \mid 0 \leq i \leq \alpha_1\}$, then D is an Ore extension of D_0 determined by the series $\{H'_i \mid 0 \leq i \leq \alpha_2\}$.

Proposition 3 is obvious if $\alpha_1 = 1$ and the proof can be completed by a straightforward application of transfinite induction and Proposition 1.

THEOREM 1. Let $\{H_i \mid 0 \leq i \leq \alpha\}$ be a completely infinite supercyclic series from H to G and let D be an Ore extension of D_0 determined by the series $\{H_i \mid 0 \leq i \leq \alpha\}$. If θ is an automorphism of G such that $\theta|_H$ can be extended to an automorphism of D_0 , then θ can be extended to an automorphism of D .

If $\alpha = 1$, then Propositions 1 and 2 imply that the Theorem is true. With Proposition 3 the proof can be completed by means of transfinite induction.

THEOREM 2. Let G be a group with a normal subgroup H and a completely infinite supercyclic series $\{H_i \mid 0 \leq i \leq \alpha\}$ from H to G with $H_{i+1}/H_i = (x_{i+1} \cdot H_i)$ for x_{i+1} in H_{i+1} . If H can be embedded in a division ring D_0 with $D_0 = \{H\}$ such that the automorphism of H induced by x_{i+1} can be extended to an automorphism of D_0 , then G can be embedded in a division ring D which is an Ore extension of D_0 determined by the completely infinite supercyclic series $\{H_i \mid 0 \leq i \leq \alpha\}$.

Consider the division ring $D_1 = D_0(y_1; \theta_1)$ where θ_1 is the automorphism of D_0 induced by x_1 and y_1 an indeterminate over D_0 . Since $y_1 h y_1^{-1} = \theta_1(h) = x_1 h x_1^{-1}$ for all h in H , the map of H_1 into D_1^* determined by $x_1 \rightarrow y_1$ and $h \rightarrow h$ for all h in H is an isomorphism onto $Gp(H, y_1)$. This completes the first step of an induction proof. Using Theorem 1 to extend automorphisms the proof of Theorem 2 can be completed by transfinite induction.

For a detailed look at a construction of a division ring like the one used in Theorems 1 and 2 look at a paper by M. Ikeda (see [6]).

Before stating some consequences of Theorems 1 and 2 some definitions will be given. A group G is said to have *property E* if G can be embedded in a division ring. A group G has *property EE* if G can be embedded in some division ring D such that any automorphism of G can be extended to an automorphism of D ; and an ordered group G has *property EE** if G can be embedded in some division ring D such that any order automorphism of G can be extended to D .

It is important to note that the above definition just requires the existence of one such division ring with the extending property for every automorphism, not every division in which G can be embedded.

In view of these definitions we have:

COROLLARY 1. If a group G has a normal subgroup H with *property EE* and a

completely infinite supercyclic series from H to G , then G has property E . If in addition H is a characteristic subgroup, then G has property EE .

COROLLARY 2. *Let G be a group with a normal subgroup H with property EE^* and a completely infinite supercyclic series $\{H_i \mid 0 \leq i \leq \alpha\}$ from H to G with $H_{i+1}/H_i = (H_{i+1} \cdot H_i)$. If each x_{i+1} induces an order automorphism on H , then G has property E .*

Since the identity group has property EE , we have the following:

COROLLARY 3. *If G is a completely infinite supercyclic group, then G has property EE .*

COROLLARY 4. *Let G be a solvable group, say $G^{n+1} = 1$. If $G^{(i)}/G^{(i+1)}$ is a free abelian group ($0 \leq i \leq n$), then G has property EE . In particular if $F^{(n)}$ is a free group then $F/F^{(n)}$ has property EE .*

COROLLARY 5. *A finitely-generated torsion-free nilpotent group has property EE .*

The results of Neumann and Malcev (see [10] and [9]) prove that the groups of Corollaries 3, 4, and 5 have property E since they can be ordered. But it is not true that all completely infinite supercyclic groups can be ordered, for the group $G_p(x, y \mid yxy^{-1} = x^{-1})$ is such a group.

The construction of Theorem 2 is not sufficient to completely determine which torsion-free solvable groups have property E . $G^{(i)}/G^{(i+1)}$ may not be free abelian, even though G is a torsion-free solvable group; for example the additive rationals Q^+ , has this property. This is not really a difficulty for by a modification of the proofs, the assumption in Theorem 2 that H_i/H_{i+1} is infinite cyclic can be replaced by the condition that H_i/H_{i+1} is a subgroup of Q^+ , since any such group is a union of a countable tower of infinite cyclic groups. A difficulty still arises since there are finitely-generated torsion-free solvable groups which are not completely infinite supercyclic groups. $G = G_p(b_1, b_2, c \mid b_i = b_i^c = b_i^{-1}, [b_1, b_2] = C^{2^x}, i = 1 \text{ or } 2)$ is a supersolvable torsion-free group, but every supersolvable series of G has a factor of order 2 (see [3] and [5]).

The remainder of the paper will be devoted to determining which finite, abelian or ordered groups have property EE or EE^* .

Property EE for finite groups. In this section we will prove that all finite groups with property E have property EE using Amitsur's classification for such groups (see [2]). First some notation will be given.

Let m and r be relatively prime integers. Put $s = (r - 1, m)$, $t = m/s$ and $n = \text{minimal integer satisfying } r^n \equiv 1 \pmod{m}$. Denote by $G_{m,r}$ group generated by two elements A and B satisfying the relations $A^m = 1$, $B^n = A^t$ and $BAB^{-1} = A^r$. Let Q denote the rational field and e_m a fixed primitive m th root of unity. σ_r will represent the automorphism of $Q(e_m)$ determined by $e_m \rightarrow e_m^r$. $\mathfrak{A}_{m,r} = (Q(e_m), \sigma_r, e_s)$

will denote the cyclic algebra determined by the field $Q(e_m)$, automorphism σ_r , and element $e_s = e_m^t$, as defined by Albert (see [1, p. 74]).

If G is a group $A(G)$, $I(G)$, $Z(G)$ and $|G|$ will denote the automorphism group, inner-automorphism group, center of group and order of group respectively. \mathfrak{T} , \mathfrak{O} and \mathfrak{S} will denote the binary tetrahedral, binary octahedral and binary icosahedral group respectively. They are isomorphic to G_1 , G_2 and G_3 respectively where $G_i = G_p(A, C \mid A^{2(i+2)} = 1, C^3 = A^{(i+2)} \text{ and } CAC^{-1} = A^{-1}C)$. Also \mathfrak{T} , \mathfrak{O} and \mathfrak{S} have order 24, 48 and 120 respectively, have center of order 2 and have automorphism groups of order 24, 48 and 60 respectively. These groups are discussed in an article by Vincent (see [12]) and in Amitsur's paper (see [2]).

With these definitions we can state an important theorem proved by Amitsur (see [2]).

THEOREM 3. *A finite group G has property E if and only if G is one of the following types:*

- (1) *A group $G_{m,r}$ with property E ;*
- (2) *A group $\mathfrak{T} \times G_{m,r}$, where $G_{m,r}$ has property E , $(6, |G_{m,r}|) = 1$, and other conditions hold (see [2]);*
- (3) *The groups \mathfrak{O} and \mathfrak{S} .*


We will use this theorem to prove that a finite group with property E has property EE . First we will prove some lemmas.

LEMMA 1. *If $G_{m,r}$ has property E , then $G_{m,r}$ has property EE .*

Proof. If $G_{m,r}$ has property E then $A \rightarrow e_m$ and $B \rightarrow \sigma_r$ defines an embedding of $G_{m,r}$ into the division ring $\mathfrak{U}_{m,r} = (Q(e_m), \sigma_r, e_s)$, (see [2]). Using this identification we have $\mathfrak{U}_{m,r} = (Q(A), B^t, A)$. Let θ be an automorphism of G . Direct calculation verifies that $A \rightarrow \theta(A)$ and $B \rightarrow \theta(B)$ defines an automorphism of $\mathfrak{U}_{m,r}$ which extends the automorphism θ .

LEMMA 2. *\mathfrak{T} has property EE .*

Proof. \mathfrak{T} can be embedded in $\mathfrak{U}_{4,-1}$ and \mathfrak{O} can be embedded in $\mathfrak{U}_{8,-1}$. \mathfrak{O} has an isomorphic copy of \mathfrak{T} as a normal subgroup. Thus with appropriate identification we have the following diagram (see [2]).



Since \mathfrak{T} is normal in \mathfrak{O} there is an element x in \mathfrak{O} which induces an outer-automorphism θ on \mathfrak{T} . But $x\mathfrak{T}x^{-1} = \mathfrak{T}$ implies $x\mathfrak{U}_{4,-1}x^{-1} = \mathfrak{U}_{4,-1}$, so θ can be

extended to $\mathfrak{A}_{4,-1}$. Since $|A(\mathfrak{T})/I(\mathfrak{T})| = 2$ the proof is complete, for any inner-automorphism can be extended.

LEMMA 3. \mathfrak{D} has property E.

Proof. Since $|A(\mathfrak{D})/I(\mathfrak{D})| = 2$ it is sufficient to show that one outer-automorphism of \mathfrak{D} can be extended to $\mathfrak{A}_{8,-1} \cdot \mathfrak{A}_{8,-1} \cong Q(2^{1/2}) \otimes_Q \mathfrak{A}_{4,-1}$ and $\mathfrak{A}_{4,-1}$ is isomorphic to the rational quaternions. Identifying elements under the isomorphisms mentioned above we have

$$\mathfrak{D} = \{ \pm (1 \pm i \pm j \pm k)/2, \pm \{1, i, j, k\}, \\ \pm (1 \pm \{i, j, k\})/2^{1/2}, \pm (i \pm \{j, k\})/2^{1/2}, \pm (j \pm k)/2^{1/2} \}$$

is a subgroup of $\mathfrak{A}_{8,-1}$ which generates $\mathfrak{A}_{8,-1}$ (see [2]). The map $2^{1/2} \rightarrow -2^{1/2}$ obviously defines an automorphism θ of $\mathfrak{A}_{8,-1}$ and $\theta|_{\mathfrak{D}} = \mathfrak{D}$. θ is an outer-automorphism since $2^{1/2}$ is in the center of $\mathfrak{A}_{8,-1}$, thus an outer-automorphism of \mathfrak{D} can be extended to $\mathfrak{A}_{8,-1}$.

LEMMA 4. \mathfrak{Z} has property EE.

Proof. \mathfrak{Z} can be embedded in $\mathfrak{A}_{10,-1} \cong Q(5^{1/2}) \otimes_Q \mathfrak{A}_{4,-1}$ and with appropriate identification we have that \mathfrak{Z} is the subgroup of $\mathfrak{A}_{10,-1}$ generated by j, i_1 , and e where e is a primitive 5th root of unity and $i_1 = (e^2 - e^3 + (e - e^4)j)5^{1/2}$ (see [2]). \mathfrak{Z} also generates $\mathfrak{A}_{10,-1}$. Obviously the map $5^{1/2} \rightarrow -5^{1/2}$ defines an outer-automorphism θ of $\mathfrak{A}_{10,-1}$ since $5^{1/2}$ is in the center of $\mathfrak{A}_{10,-1}$. Direct calculation verifies that $\theta|_{\mathfrak{Z}} = \mathfrak{Z}$. Therefore θ is an extension of an outer-automorphism of \mathfrak{Z} , which completes the proof since $|A(\mathfrak{Z})/I(\mathfrak{Z})| = 2$.

With these lemmas we are prepared to prove

THEOREM 4. A finite group with property E also has property EE.

Proof. Groups of type (1) and (3) of Theorem 3 have property EE by Lemmas 1, 3 and 4. If $G = \mathfrak{T} \times G_{m,r}$ has property E, then G can be embedded in $\mathfrak{A}_{4,-1} \otimes_Q \mathfrak{A}_{m,r}$ in the natural way (see [2]). There is no loss of generality in assuming \mathfrak{T} is in $\mathfrak{A}_{4,-1}$ and $G_{m,r}$ is $\mathfrak{A}_{m,r}$. If θ is an automorphism of G then $\theta = \theta|_{\mathfrak{T}}$ and $\theta_2 = \theta|_{G_{m,r}}$ are automorphisms of \mathfrak{T} and $G_{m,r}$ respectively since $(6, |G_{m,r}|) = 1$. θ_1 and θ_2 can be extended to automorphism θ'_1 and θ'_2 of $\mathfrak{A}_{4,-1}$ and $\mathfrak{A}_{m,r}$ respectively by lemmas 1 and 2. Thus $\theta'_1 \otimes \theta'_2$ is an automorphism of $\mathfrak{A}_{4,-1} \otimes_Q \mathfrak{A}_{m,r}$ which extends θ . This completes proof of Theorem 4.

Property EE for abelian groups.

THEOREM 5. If G is an abelian group with property E, then G has property EE.

Proof. Let G be an abelian group generated by elements x_i ($i \in I$) which is embeddable in a field. The construction of Cohn (see [4]) yields a field K which is

generated by the x_i as a field, with the multiplicative relations holding between the x_i as defining relations. Thus every relation Φ holding in K between the x_i follows from the defining relations in G . Hence if $\theta: x_i \rightarrow x'_i$ is an automorphism of G , any relation Φ between the x_i also holds between the x'_i and conversely. Therefore $x_i \rightarrow x'_i$ defines an automorphism of K which extends θ .

Property EE^* for ordered groups. An ordered group G can be embedded in the formal power series ring D of G over any division ring K (see [10]). An element ψ of D is a map on G into K such that the subset of G on which ψ takes nonzero values is a well-ordered subset. If ψ and ϕ are in D and g in G , then addition and multiplication in D are defined componentwise as follows:

$$\begin{aligned}(\psi + \phi)(g) &= \psi(g) + \phi(g), \\ \psi \cdot \phi(g) &= \sum_{rs=g} \psi(r) \phi(s).\end{aligned}$$

THEOREM 6. *If G is an ordered group, then G has property EE^* .*

Proof. Let Γ be the formal power series ring of G over the rational field \mathcal{Q} . If ψ is in Γ , then ψ is a function of G into \mathcal{Q} such that $\psi(g) \neq 0$ if and only if g is in S_ψ , a well ordered subset of G . Let π_g be the function defined as follows:

$$\pi_g(h) = 1 \text{ if } g = h \text{ and zero otherwise.}$$

The map

$$\begin{aligned}\sigma: G &\rightarrow \Gamma \\ g &\rightarrow \pi_g \text{ for all } g \text{ in } G\end{aligned}$$

is the embedding of G into Γ (see [12]).

Under this identification G is a subset of Γ . Let D be the division ring in Γ generated by G , and let θ be an order automorphism of G . Define the following map:

$$\theta': D \rightarrow \Gamma$$

$$\theta'(\psi)(g) = \psi(\theta^{-1}(g)) \text{ for all } g \text{ in } G.$$

$\theta'(\psi)(g) = 0$ if and only if $\theta^{-1}(g)$ is in S_ψ , or equivalently if g is in $\theta(S_\psi)$. Since θ is an order automorphism, $\theta(S_\psi)$ is a well-ordered subset of G ; since θ' is well defined. Direct calculation verifies that if ψ and ζ are in D then

$$\theta'(\psi + \zeta)(g) = \theta'(\psi)(g) + \theta'(\zeta)(g)$$

and

$$\theta'(\psi \cdot \zeta)(g) = (\theta'(\psi) \cdot \theta'(\zeta))(g).$$

Therefore θ' is an isomorphism of D into Γ . Also obviously $\theta'(\pi_g) = \pi_{\theta(g)}$, hence $\theta' \mid G = \theta$. Since $\theta(G) = G$ and $D = \{G\}$, $\theta(D) = D$, thus θ is an automorphism of D extending θ .

In conclusion we summarize some of the results of Theorems 1, 2, 4, 5 and 6.

THEOREM 7. *Let G be a group with a normal subgroup H and a completely infinite supercyclic series from H to G .*

(i) *If H is finite or abelian, then G has property E if and only if H has property E .*

(ii) *If H is finite or abelian and characteristic in G , then G has property EE if and only if G has property E .*

(iii) *If H is an ordered subgroup of G such that every automorphism of H induced by an element of G is an order automorphism, then G has property E .*

BIBLIOGRAPHY

1. A. A. Albert, *Structure of algebras*, Amer. Math. Soc. Colloq. Publ. Vol. 24, Amer. Math. Soc., Providence, R. I., 1939.
2. S. A. Amitsur, *Finite subgroups of division rings*, Trans. Amer. Math. Soc. **80** (1955), 361–386.
3. J. F. Bowers, *On composition series of polycyclic groups*, J. London Math. Soc. **35** (1960), 433–444.
4. P. M. Cohen, *Eine Bemerkung über die multiplicative Gruppe eines Körpers*, Arch. Math. **13** (1962), 344–348.
5. K. A. Hirsch, *On infinite soluble groups*. IV, J. London Math. Soc. **27** (1952), pp. 81–85.
6. M. Ikeda, *Schiefkörper unendlichen Ranges über dem Zentrum*, Osaka Math. J. **14** (1962), 135–144.
7. K. Iwasawa, *On linearly ordered groups*, J. Math. Soc. Japan **1** (1948), 1–9.
8. N. Jacobson, *Structure of rings*, Amer. Math. Soc. Colloq. Publ. Vol. 37, Amer. Math. Soc., Providence, R. I., 1956.
9. A. I. Malcev, *On the embedding of group algebras in division algebras*, Dokl. Akad. Nauk SSSR **60** (1948), 1499–1501.
10. B. H. Neumann, *On ordered division rings*, Trans. Amer. Math. Soc. **66** (1949), 202–252.
11. O. Ore, *Theory of non-commutative polynomials*, Ann. of Math. **34** (1933), 480–508.
12. G. Vincent, *Les groupes linéaires finis sans points fixes*, Comment. Math. Helv. **20** (1947), 117–171.
13. H. Zassenhaus, *The theory of groups*, Macmillan, New York, 1958.

UNIVERSITY OF CALIFORNIA,
BERKELEY, CALIFORNIA