

DECOMPOSABLE MODULES

BY
EBEN MATLIS

1. Introduction. The concept of an h -local domain was introduced in [8, §8] as a simultaneous generalization of both a Dedekind domain and a local domain. It is a domain in which localizations with respect to maximal ideals behave properly. In this paper we give some further characterizations of an h -local domain and investigate its ideal structure. The tools found here, together with the theory of cotorsion modules of [8], are then used to settle some questions concerning decomposability of modules that have descended to us from the theory of Abelian groups. The point here is that the machinery of cotorsion modules and h -local domains is useful in solving problems that have a classical history, and were posed long before the development of this machinery.

In §5 we characterize those h -local domains that have the property that every finitely generated torsion module is a direct sum of cyclic modules. This generalizes and proves the converse of [3, Theorem 14]. In §6 we find the class of all integral domains having the property that every torsion-free module of finite rank is a direct sum of modules of rank one. This generalizes and proves the converse of [3, Theorem 12]. Finally, in §7 we find the class of all Noetherian domains having the property that every homomorphic image of the quotient field is indecomposable. This answers a question raised in [7, p. 573].

2. The ideal structure of an h -local domain. Throughout this paper R will denote an integral domain (not a field), Q its quotient field, and K the R -module Q/R . We will let H denote the ring $\text{Hom}_R(K, K)$. H is the completion of R in the R -topology [8, Proposition 6.4]. R is said to be an h -local domain if the following two conditions are satisfied:

- (1) Each nonzero ideal of R is contained in only finitely many maximal ideals of R ;
 - (2) Each nonzero prime ideal of R is contained in only one maximal ideal of R .
- R is said to be h -semilocal if it satisfies merely condition (1).

These types of rings can be characterized by finiteness conditions on the topology of Ω_p , the space of prime ideals of R ; or of Ω , the space of maximal ideals of R . A set in Ω_p is called closed if it consists of all of the prime ideals of R containing a given ideal of R . A similar statement is true for Ω . It is then easy to see that R is h -local if and only if every proper closed set in Ω_p is a finite union of isolated closed sets (a closed set is called *isolated*, if it contains only one closed point).

It is also obvious that R is h -semilocal if and only if Ω has the cofinite topology (i.e., the open sets of Ω are the complements of finite sets).

Examples of h -local domains are the following:

- (a) Local domains (domains with only one maximal ideal, and no chain conditions assumed).
- (b) Dedekind rings.
- (c) Noetherian domains of Krull dimension one.
- (d) Domains in which every nonzero prime ideal is maximal, and no maximal ideal is contained in the union of the remaining ones (see [8, Proposition 8.7]).

In [8, Theorem 8.5] we proved the following characterization of h -local domains.

THEOREM. *The following statements are equivalent:*

- (1) R is an h -local domain.
- (2) $S(M) \otimes_R R_M \cong Q$ for all $M \in \Omega$, where $S(M) = \bigcap R_N$ for all $N \in \Omega - \{M\}$.
- (3) $K \cong \sum \bigoplus K_M$, where M ranges over all maximal ideals in Ω .
- (4) $H \cong \prod H(M)$, where M ranges over all maximal ideals in Ω , and where $H(M) = \text{Hom}_{R_M}(K_M, K_M)$ is the completion of R_M .

DEFINITION. An ideal I of R will be called *colocal* if it is contained in only one maximal ideal M of R . I will then be said to belong to M . An R -module will be called *local* if it is isomorphic to R/I , where I is a colocal ideal of R . We have the following elementary properties of colocal ideals;

- (1) If I_1, \dots, I_n are colocal ideals belonging to M , then $I_1 \cap \dots \cap I_n$ is a colocal ideal belonging to M .
- (2) If I and J are colocal ideals belonging to different maximal ideals, then I and J are relatively prime; i.e., $I + J = R$.
- (3) Let $I = \bigcap_{j=1}^n I_j$, where each I_j is a colocal ideal belonging to a maximal ideal M_j , and the M_j 's are all distinct. Then $I = I_1 I_2 \cdots I_n$ and $R/I \cong R/I_1 \oplus \cdots \oplus R/I_n$.
- (4) If A is a local R -module, and if B and C are submodules of A such that $B + C = A$, then either $B = A$ or $C = A$.

LEMMA 2.1. *Let I be a colocal ideal of R belonging to a maximal ideal M . Then:*

- (1) $R_M \otimes_R R/I \cong R/I$, and $I_M \cap R = I$.
- (2) If N is a maximal ideal different from M , then $R_N \otimes_R R/I = 0$.

Proof. Let $J = \{r \in R \mid \text{there is an } s \in R - M \text{ such that } sr \in I\}$. Then $J = I_M \cap R$, and $I \subset J$. Hence $I_M = J_M$. Now $1 \notin J$, and so J is a colocal ideal belonging to M . Let L be any colocal ideal belonging to M , and let N be a maximal ideal different from M . Since L is not contained in N , it meets $R - N$, and thus $L_N = R_N$. Hence we have $I_N = R_N$ and $J_N = R_N$ for all maximal ideals N different from M . But $I_M = J_M$ also, and thus we have $I = \bigcap I_N = \bigcap J_N = J$, where the intersection ranges over all maximal ideals N of R . Therefore $I_M \cap R = I$.

Define $f: R/I \rightarrow R_M \otimes_R R/I$ by $f(\bar{x}) = 1 \otimes \bar{x}$ for all $\bar{x} \in R/I$. If $\bar{x} \in \text{Ker } f$, then there exists an $s \in R - M$ such that $s\bar{x} = 0$. Thus if $\bar{x} = x + I$, $x \in R$, we have $sx \in I$. Hence $x \in J = I$, and $\bar{x} = 0$. Therefore, f is 1 to 1. Let $\bar{y} \in R_M \otimes_R R/I$; then there exists $\bar{x} \in R/I$ and $s \in R - M$ such that $\bar{y} = (1/s) \otimes \bar{x}$. Now $Rs + I$ can be contained in no maximal ideal of R , and so $Rs + I = R$. Therefore, $s(R/I) = R/I$, and so there exists an element $\bar{z} \in R/I$ such that $\bar{x} = s\bar{z}$. We then have $\bar{y} = 1 \otimes \bar{z} = f(\bar{z})$. Thus f is onto, and so is an isomorphism.

If N is a maximal ideal different from M , then we have seen that $I_N = R_N$, and so $R_N \otimes_R R/I \cong R_N/I_N = 0$.

DEFINITION. Let $I = \bigcap_{j=1}^n I_j$ be a finite intersection of ideals, where I_j is a colocal ideal belonging to a maximal ideal M_j . We will say that this decomposition is *normal*, if $M_i \neq M_k$ for $i \neq k$. It is clear from elementary property (1) that every finite intersection of colocal ideals can be normalized.

LEMMA 2.2. Let $I = \bigcap_{j=1}^n I_j$ be a normal decomposition of an ideal I , where I_j is a colocal ideal belonging to a maximal ideal M_j . Then $I_j = I_{M_j} \cap R$, and so the I_j 's are unique.

Proof. By Lemma 2.1 we have

$$I_{M_k} \cap R = \left(\bigcap_{j=1}^n I_j \right)_{M_k} \cap R = \bigcap_{j=1}^n ((I_j)_{M_k} \cap R) = I_k \cap R = I_k.$$

THEOREM 2.3. Let R be an integral domain. Then R is an h -local ring if and only if every nonzero ideal of R is a finite intersection of colocal ideals.

Proof. Suppose that every nonzero ideal of R is a finite intersection of colocal ideals. Let I be a nonzero ideal of R . Then $I = \bigcap_{j=1}^n I_j$, where I_j is a colocal ideal belonging to a maximal ideal M_j . If M is any maximal ideal containing I , then M contains one of the I_j 's, and hence is equal to that M_j . Therefore, I is contained in only finitely many maximal ideals of R . If I is a prime ideal, then it can not be the intersection of two properly larger ideals, and so it is equal to one of the I_j 's. Thus I is contained in only one maximal ideal of R . Therefore, R is an h -local domain.

Conversely, suppose that R is an h -local domain. Let I be a nonzero ideal of R ; then I is contained in only a finite number of maximal ideals M_1, \dots, M_n . Let $I_j = I_{M_j} \cap R$ for $j = 1, \dots, n$. If N is a maximal ideal different from M_1, \dots, M_n , then I meets $R - N$ and so $I_N = R_N$. Thus, if we let M range over all maximal ideals of R , we have $\bigcap_{j=1}^n I_j = \bigcap_{j=1}^n I_j \cap (\bigcap_M R_M) = \bigcap_M I_M = I$.

It only remains to show that I_j is a colocal ideal belonging to M_j . Since $I \subset M_j$, we have $I_j \subset M_j$. Let N be any maximal ideal different from M_j . Then $(I_j)_N = (I_{M_j})_N \cap R_N$. But $R_{M_j} \otimes_R R_N \cong Q$, since R is h -local, and so

$$(I_{M_j})_N = (I \otimes_R R_{M_j}) \otimes_R R_N \cong Q.$$

Therefore, $(I_j)_N = Q \cap R_N = R_N$. It follows that I_j is not contained in N . Hence I_j is a colocal ideal belonging to M_j .

LEMMA 2.4. *Let R be an integral domain and I a nonzero projective ideal of R . Let M_1, \dots, M_n be maximal ideals of R . Then there exists an ideal of R isomorphic to I which is not contained in any of M_1, \dots, M_n ; (i.e., there exists an element $q \in I^{-1}$ such that $qI \not\subset M_1 \cup \dots \cup M_n$).*

Proof. The proof will be by induction on n . Suppose that $n=1$. Now $1 = \sum_{i=1}^i a_i q_i$, where $a_i \in I$ and $q_i \in I^{-1}$. Hence we can assume without loss of generality that $a_1 q_1 \notin M_1$, and so $q_1 I$ is not contained in M_1 . Suppose that $n > 1$, and that the lemma is true for $n-1$. We can assume that the M_i 's are all distinct. By the induction hypothesis we may assume that I is not contained in $M_2 \cup \dots \cup M_n$, but that I is contained in M_1 . Choose $p \in M_2 M_3 \dots M_n$ such that $p \notin M_1$. By the case $n=1$ there exists an element $q \in I^{-1}$ such that $qI \not\subset M_1$. Let $v = 1 - pq$; then we can not have $vI \subset M_i$ for any $i > 1$. For suppose that $vI \subset M_i$. Take $x_i \in I$ such that $x_i \notin M_i$; then $x_i - pqx_i = vx_i \in M_i$ and $pqx_i \in M_i$. Thus $x_i \in M_i$, which is a contradiction. We can not have $vI \subset M_1$, either. For take $x \in I$ such that $qx \notin M_1$; then $x \in M_1$, and if $x - pqx = vx \in M_1$, we would have $pqx \in M_1$. But $p \notin M_1$, and so $qx \in M_1$, which is a contradiction. Hence vI is not contained in any one of M_1, \dots, M_n .

COROLLARY 2.5. *Let R be an integral domain and I a projective ideal of R . If I is contained in only a finite number of maximal ideals of R , then I is generated by at most two elements.*

Proof. We can assume that I is not the zero ideal. By Lemma 2.4 there exists an element $q \in I^{-1}$ such that qI is not contained in any of the maximal ideals that contain I . Since qI is an ideal of R we then have that $I + qI = R$. Hence there exist elements a, b in I such that $1 = a + qb$. If $x \in I$, then $x = xa + (qx)b$, and so I is generated by a and b .

COROLLARY 2.6. *Let R be an integral domain, I a projective rank one R -module and J an ideal of R . If J is contained in only a finite number of maximal ideals of R , then we have $I/JI \cong R/J$.*

Proof. Since I is a finitely generated R -submodule of Q , there exists a nonzero element d in R such that dI is an ideal of R . Then $dI/JdI \cong I/JI$, and so we can assume that I is an ideal of R . By Lemma 2.4 there exists an element q in I^{-1} such that qI is not contained in any of the maximal ideals that contain J . Thus we have $qI + J = R$, and so $qI \cap J = qIJ$. Therefore, we have $R/J = (qI + J)/J \cong qI/(qI \cap J) = qI/qIJ \cong I/IJ$.

THEOREM 2.7. *Let R be an h -semilocal domain, and I a nonzero projective ideal of R . Then,*

- (1) *If J is a nonzero ideal of R , we have $I/JI \cong R/J$.*
- (2) *I is generated by two elements, one of which may be chosen arbitrarily.*
- (3) *We have $I^{-1}/R \cong R/I$.*

Proof.

- (1) This follows from Corollary 2.6.
- (2) Let a be a nonzero element of I , and let $J = aI^{-1}$. Then J is an ideal of R . Hence by Corollary 2.6, $I/JI \cong R/J$ is cyclic. Now $JI = aI^{-1}I = Ra$. Hence I is generated by a and one other element.
- (3) By part (2) I^{-1} is generated by 1 and one other element. Therefore, I^{-1}/R is cyclic. Since $(I^{-1})^{-1} = I$, the annihilator of I^{-1}/R is exactly I . Thus $I^{-1}/R \cong R/I$.

REMARKS. (1) Theorem 2.7 is of course a classical theorem for Dedekind rings. If R is actually semilocal (i.e., if R has only a finite number of maximal ideals), and if we take J to be the 0 ideal in Corollary 2.6, we obtain the theorem that every projective ideal of R is free on one generator. Thus there is an underlying unity in these two well-known theorems. They are merely special cases of Theorem 2.7 that postulates a weak type of finiteness condition that encompasses both. With the austerity of the assumptions goes a corresponding increased transparency in the nature of the proof.

(2) It is an open question whether Corollary 2.5 is the best result obtainable. In other words, if I is a projective ideal of an arbitrary integral domain, and if I is contained in only a finite number of maximal ideals, can one of the two generators of I be chosen from among any of the nonzero elements of I ? The affirmative answer for h -semilocal domains is provided by Theorem 2.7.

(3) Let R be an h -semilocal domain. If P is a projective R -module of finite rank n , then it follows from [1, Theorem 3] that P is a direct sum of a projective ideal I of R and a free R -module of rank $n-1$. By [3, Lemma 1] the only invariants of P are n and the ideal class of I . It follows that if I_1, \dots, I_n and J_1, \dots, J_n are projective ideals of R , then $I_1 \oplus \dots \oplus I_n$ and $J_1 \oplus \dots \oplus J_n$ are isomorphic if and only if $I_1 I_2 \dots I_n$ and $J_1 J_2 \dots J_n$ are isomorphic. By the proof of [3, Theorem 2] an R -module which is a direct sum of an infinite number of nonzero projective ideals is free.

THEOREM 2.8. *Let R be an h -local domain, and P a finitely generated R -module. Let H be the completion of R . Then the following statements are equivalent:*

- (1) *P is R -projective.*
- (2) *$H \otimes_R P$ is H -free.*

Proof. Assume that P is R -projective, and let F be a free R -module such that $\text{rank } F = \text{rank } P$. Let M be a maximal ideal of R . Then P_M is R_M -free, and so

$P_M \cong F_M$. Hence $K_M \otimes_R P \cong K_M \otimes_R P_M \cong K_M \otimes_R F_M \cong K_M \otimes_R F$. Since R is h -local, we have $K \cong \sum \oplus K_M$, where M ranges over all maximal ideals of R . Thus we have $K \otimes_R P \cong \sum \oplus (K_M \otimes_R P) \cong \sum \oplus (K_M \otimes_R F) \cong K \otimes_R F$. Therefore, $\text{Hom}_R(K, K \otimes_R P) \cong \text{Hom}_R(K, K \otimes_R F)$. Since P is finitely generated, we have by [8, Proposition 7.2] and [8, Corollary 3.3] that $H \otimes_R P \cong \text{Hom}_R(K, K \otimes_R P)$. Similarly we have $H \otimes_R F \cong \text{Hom}_R(K, K \otimes_R F)$. Thus $H \otimes_R P \cong H \otimes_R F$, and so $H \otimes_R P$ is H -free.

Conversely, if $H \otimes_R P$ is H -free, then by [8, Proposition 9.1] we have that P is R -projective.

Theorem 2.8 raises the question of which finitely generated projective H -modules are H -free? Theorem 3.2 will provide the answer for h -local domains. Since Theorem 2.8 will be a special case of Theorem 3.2, this will give a second proof of Theorem 2.8.

3. Modules over h -local domains. An R -module C is said to be *cotorsion*, if $\text{Hom}_R(Q, C) = 0$ and $\text{Ext}_R^1(Q, C) = 0$. If M is a maximal ideal of R and A is an R -module, we denote $\text{Hom}_R(R_M, A)$ by the symbol A^M . In [8] we showed that R is h -local if and only if $T \cong \sum \oplus T_M$ for every torsion R -module T , where M ranges over all maximal ideals of R . We then showed that if R is h -local and C is a cotorsion R -module, then $C = \prod C^M$. The next theorem repairs the gap in [8] by showing that these three conditions are all equivalent. It reveals the real reason why h -local domains are so useful.

THEOREM 3.1. *The following statements are equivalent:*

- (1) R is an h -local domain.
- (2) $T \cong \sum \oplus T_M$ for all torsion R -modules T , where M ranges over all maximal ideals of R .
- (3) $C \cong \prod C^M$ for all cotorsion R -modules C , where M ranges over all maximal ideals of R .

Proof. We have already proved ([8, Theorem 8.5] and [8, Corollary 8.6]) that (1) \Rightarrow (2) and (2) \Rightarrow (3). It remains to prove that (3) \Rightarrow (1). Hence assume that (3) holds.

Let P be a nonzero prime ideal of R . We will show first that P is contained in only one maximal ideal of R . Now R/P is torsion of bounded order, and hence is cotorsion. Thus by assumption we have $R/P \cong \prod (R/P)^M$. If there are at least two nonzero components in this product, then there exist ideals A and B of R , each properly containing P , such that $R/P = A/P \oplus B/P$. Thus we have $A \cap B = P$. Since P is a prime ideal, this is a contradiction. Hence there exists a maximal ideal M of R such that $R/P \cong (R/P)^M$, and $(R/P)^N = 0$ for every other maximal ideal N of R . Thus we have $R_M/P_M \cong (R/P)_M \cong [(R/P)^M]_M \cong (R/P)^M \cong R/P$. Hence $P_M \neq R_M$, and so $P \subset M$.

Let N be a maximal ideal of R different from M . We must show that P is not contained in N . Now R_N/P_N is torsion of bounded order as an R -module, and so is a cotorsion R -module. Thus we have $R_N/P_N \cong \prod (R_N/P_N)^L$, where L ranges over all maximal ideals of R . But $(R_N/P_N)^L$ is an R_N -module; and R_N/P_N is a local R_N -module, and hence indecomposable. Therefore, $(R_N/P_N)^L = 0$ for every $L \neq N$. In particular, $(R_N/P_N)^M = 0$.

Suppose that we have $P \subset N$. We define a map $\varphi: [(R/P)^M]_N \rightarrow (R_N/P_N)^M$ as follows. We have $[(R/P)^M]_N = \text{Hom}_R(R_M, R/P)_N$, and

$$(R_N/P_N)^M \cong \text{Hom}_{R_N}(R_M \otimes_R R_N, R_N/P_N).$$

Let $f \in \text{Hom}_R(R_M, R/P)$, $x \in R_M$, s and $t \in R - N$; then set

$$[\varphi(f \otimes (1/s))](x \otimes (1/t)) = f(x) \otimes (1/st).$$

Suppose that $f \otimes (1/s) \in \text{Ker } \varphi$; then for each $x \in R_M$ there exists $u_x \in R - N$ such that $u_x f(x) = 0$. But $f(x) \in R/P$, and $u_x \notin P$. Therefore, $f(x) = 0$ for all $x \in R_M$, and so $f = 0$. Thus we have $\text{Ker } \varphi = 0$. But we have seen above that $(R_N/P_N)^M = 0$. Hence we have $[(R/P)^M]_N = 0$, also. Therefore, $R_N/P_N \cong (R/P)_N \cong [(R/P)^M]_N = 0$. This contradicts $P \subset N$, and so we have that M is the only maximal ideal of R containing P .

Let M and N be two distinct maximal ideals of R . Then by what we have just shown, $M \cap N$ contains no nonzero prime ideal of R . Hence by [8, Corollary 8.2] we have $R_M \otimes_R R_N = Q$.

Let I be a nonzero ideal of R , and let $T = R/I$. We must show that I is contained in only a finite number of maximal ideals of R . Let \mathcal{M} be the collection of maximal ideals M of R such that $T^M \neq 0$. We will show first that \mathcal{M} is a finite set. Since T is a cotorsion R -module, we have $T \cong \prod T^M$, $M \in \mathcal{M}$. Let $S = \sum \oplus T^M$, $M \in \mathcal{M}$. Then $S \subset T$, and so S is torsion of bounded order. Therefore, S is a cotorsion R -module, and we have $S \cong \prod S^N$, where N ranges over all maximal ideals of R . If N is a maximal ideal such that $N \notin \mathcal{M}$, then $S^N \subset T^N = 0$. Therefore, we have $S \cong \prod S^M$, $M \in \mathcal{M}$.

Let N and M be two distinct maximal ideals of R . Then

$$(T^M)^N \cong \text{Hom}_R(R_N \otimes_R R_M, T) \cong \text{Hom}_R(Q, T) = 0.$$

Thus if $N \in \mathcal{M}$, we have $S^N = \text{Hom}_R(R_N, \sum_{M \in \mathcal{M}} \oplus T^M) \cong (T^N)^N \cong T^N$. Therefore, $S \cong \prod_{M \in \mathcal{M}} S^M \cong \prod_{M \in \mathcal{M}} T^M \cong T$. We also have $(T^M)_N \cong ([T^M]_M)_N \cong T^M \otimes_R Q = 0$. Therefore, if $N \notin \mathcal{M}$, we have $S_N \cong \sum_{M \in \mathcal{M}} \oplus (T^M)_N = 0$. Since $S \cong T$, we also have $T_N = 0$, and hence $(T/S)_N = 0$. Now let $M \in \mathcal{M}$. Then $T = T^M \oplus A$, $S = T^M \oplus B$, where $B \subset A$ are submodules of T . Hence $T_M \cong (T^M)_M \oplus A_M \cong T^M \oplus A_M$. But $T_M \cong R_M/I_M$ is a local R_M -module and hence is indecomposable. Therefore, we have $A_M = 0$, and it

follows that $(A/B)_M = 0$ also. Now $T/S \cong A/B$, and so $(T/S)_M = 0$. We have thus shown that $(T/S)_N = 0$ for every maximal ideal N of R . Hence we have $S = T$. But this means that the set \mathcal{M} is finite.

If $N \notin \mathcal{M}$, then we have seen above that $R_N/I_N = T_N = 0$. This says that I is not contained in N , and therefore, I is contained in only the maximal ideals belonging to \mathcal{M} , which is a finite set. Hence I is contained in only a finite number of maximal ideals of R . Thus R is an h -local ring, and the proof of the theorem is complete.

DEFINITION. Let R be an h -local ring, H its completion, and M a maximal ideal of R . Then HM is a maximal ideal of H and $H/HM \cong R/M$ (see [8, Proposition 5.10]). Let P be a projective H -module, and define $\text{rank}_M P$ to be the dimension of P/MP over R/M (or over H/HM). If P is a finitely generated H -module, then $\text{rank}_M P$ is finite.

THEOREM 3.2. *Let R be an h -local domain, and H its completion. Let P be a finitely generated, projective H -module. Then P is isomorphic to a finite direct sum of principal ideals of H . P is a free H -module if and only if $\text{rank}_M P$ is constant for all maximal ideals M of R .*

Proof. Since P is a direct summand of a finitely generated free H -module, P is a cotorsion R -module. Thus by Theorem 3.1 we have $P \cong \prod P^M$, where M ranges over all maximal ideals of R . By [8, Theorem 8.5] we have $H = \prod H(M)$, where M ranges over all maximal ideals of R and $H(M) = \text{Hom}_{R_M}(K_M, K_M)$. Since $H(M)/R_M$ is torsion-free and divisible, we have $P^M \cong \text{Hom}_R(H(M), P)$, and so P^M is an $H(M)$ -module. If N and M are different maximal ideals of R , we have by [8, Lemma 8.3] that $\text{Hom}_R(H(M), P^N) = 0$. This shows that $H = \prod H(M)$ acts componentwise on $P = \prod P^M$, and that P^M is the $H(M)$ -component of P . Therefore, P^M is a finitely generated, projective $H(M)$ -module.

Since P is a direct summand of a finitely generated, free H -module, $\text{rank}_M P$ is bounded. Let $n = \max(\text{rank}_M P)$, where M ranges over all maximal ideals of R . Let $A = \prod_{N \neq M} H(N)$, so that $H = H(M) \oplus A$. Then $R/M \cong H/MH \cong H(M)/MH(M) \oplus A/MA$, and so we have $A = MA$. Hence if $B = \prod_{N \neq M} P^N$, we have $MB = B$. Thus $P/MP \cong P^M/MP^M$, and so $\text{rank}_M P^M = \text{rank}_M P \leq n$.

It will follow from Proposition 4.1 of the next section, that since R_M is a local ring, $H(M)$ is a local ring also. Thus P^M , being a projective $H(M)$ -module, is a free $H(M)$ -module of rank $\leq n$. Therefore, $P^M = \sum_{i=1}^n \oplus A_{i,M}$, where $A_{i,M} = 0$ or $A_{i,M} \cong H(M)$. Thus we have $P \cong \prod P^M \cong \prod_M (\sum_{i=1}^n \oplus A_{i,M}) \cong \sum_{i=1}^n \oplus (\prod_M A_{i,M})$. Now $\prod_M A_{i,M}$ is isomorphic to H if $A_{i,M} \neq 0$ for every M , and it is isomorphic to a direct summand of H , otherwise. Hence P is isomorphic to a finite direct sum of principal ideals of H . It is also clear that P is a free H -module if and only if $\text{rank}_M P = n$ for every maximal ideal M of R .

THEOREM 3.3. *Let R be an h -local domain, and C an R -module. Then $\text{inj. dim.}_R C = \sup_M \text{inj. dim.}_{R_M} C_M$, where M ranges over all maximal ideals of R . Thus $\text{gl. dim. } R = \sup_M \text{gl. dim. } R_M$.*

Proof. We will prove first that $\text{inj. dim.}_R C \leq \sup_M \text{inj. dim.}_{R_M} C_M$. Hence we can assume that $\sup_M \text{inj. dim.}_{R_M} C_M = n < \infty$.

Case I. $n=0$. In this case C_M is an injective R_M -module for all maximal ideals M of R . Hence by [2, Chapter VI, Proposition 4.1.3], C_M is an injective R -module for all M . Let T be the torsion submodule of C , and let $S = C/T$. Then T_M is the torsion submodule of C_M , and so $C_M \cong T_M \oplus S_M$. Hence S_M is torsion-free and divisible. Since $S \subset S_M \subset Q \otimes_R S$, it follows that $S_M = Q \otimes_R S$. Thus $S = \bigcap S_M = Q \otimes_R S$, and so S is an injective R -module. It suffices then in this case to prove that T is an injective R -module, for then $C \cong T \oplus S$ will be an injective R -module.

Let I be a nonzero ideal of R and N a maximal ideal of R . Consider an exact sequence of the form:

$$(*) \quad 0 \rightarrow T \rightarrow A \rightarrow R_N/I_N \rightarrow 0.$$

Then A is a torsion R -module, and we have $A = \sum \oplus A_M$ and $T = \sum \oplus T_M$. From exact sequence $(*)$ we see that we have $A_M/T_M \cong (R_N/I_N)_M$. If $M \neq N$, we have $(R_N/I_N)_M \cong R/I \otimes_R Q = 0$ by [8, Corollary 8.2]. Thus $T_M = A_M$ for all $M \neq N$. Hence exact sequence $(*)$ splits if and only if the following exact sequence splits:

$$0 \rightarrow T_N \rightarrow A_N \rightarrow R_N/I_N \rightarrow 0.$$

But this sequence splits, because T_N is an injective R_N -module. Therefore $(*)$ splits, and so we have $\text{Ext}_R^1(R_N/I_N, T) = 0$. By Theorem 3.1 we have $R/I \cong \sum_N \oplus R_N/I_N$, and so we have $\text{Ext}_R^1(R/I, T) \cong \prod_N \text{Ext}_R^1(R_N/I_N, T) = 0$. Therefore, T is an injective R -module.

Case II: $n > 0$. Take an exact sequence:

$$(**) \quad 0 \rightarrow C \rightarrow E_0 \rightarrow E_1 \rightarrow \cdots \rightarrow E_{n-1} \rightarrow D \rightarrow 0,$$

where each E_i is an injective R -module. Then for each maximal ideal M of R we have an exact sequence:

$$(***) \quad 0 \rightarrow C_M \rightarrow (E_0)_M \rightarrow (E_1)_M \rightarrow \cdots \rightarrow (E_{n-1})_M \rightarrow D_M \rightarrow 0.$$

Now E_i is a direct sum of a torsion injective R -module and a torsion-free injective R -module. Hence using Theorem 3.1 we see that $(E_i)_M$ is a direct summand of E_i , and so is R -injective. Thus by [2, Chapter VI, Proposition 4.1.3], $(E_i)_M$ is an injective R_M -module. Since $\text{inj. dim.}_{R_M} C_M \leq n$, it follows that D_M is an injective R_M -module for all M . Therefore, by Case I, D is an injective R -module. Thus we see from exact sequence $(**)$ that $\text{inj. dim.}_R C \leq n$.

We next prove that $\sup_M \text{inj. dim.}_{R_M} C_M \leq \text{inj. dim.}_R C$; and for this we may assume that $\text{inj. dim.}_R C = n < \infty$. Then we have an exact sequence of the form (***) with $D = E_n$, and E_i an injective R -module for all $i \geq 0$. As above it follows that $(E_i)_M$ is an injective R_M -module for all $i \geq 0$. We then have an exact sequence of the form (***) for each maximal ideal M , with $D_M = (E_n)_M$. Therefore,

$$\sup_M \text{inj. dim.}_{R_M} C_M \leq n,$$

and the theorem is proved.

4. Local domains.

PROPOSITION 4.1. *Let R be a local domain with maximal ideal M , and let H be the completion of R . Then H is a commutative local ring with maximal ideal HM , and $H/HM \cong R/M$.*

Proof. By [8, Proposition 5.10], HM is a maximal ideal of R and $H/HM \cong R/M$. H is a commutative ring by [8, Proposition 5.1]. Suppose that J is an ideal of H that is not contained in HM . Then $H = HM + J$, and so there exist $g \in HM$ and $p \in J$ such that $1 = g + p$. Now we have $g = \sum_{i=1}^n h_i m_i$, where $h_i \in H$ and $m_i \in M$ for $i = 1, \dots, n$. Let s be a nonzero element of R , and set $y = [(1/s) + R] \in K$. Then $h_i(y)$ is an element of K that is annihilated by s , and so $h_i(y) \in Ry$. It follows that if x is any element of K , then there exists an element $r_{i,x} \in R$ such that $h_i(x) = r_{i,x}x$. Hence $g(x) = \sum_{i=1}^n h_i(m_i x) = \sum_{i=1}^n r_{i,x} m_i x = m_x x$, where $m_x \in M$. Therefore,

$$p(x) = x - g(x) = x - m_x x = (1 - m_x)x.$$

Let $u_x = 1 - m_x$; then u_x is a unit in R and we have $p(x) = u_x x$. It follows that p is an automorphism of K , and so $p^{-1} \in H$. Thus $J = H$, which shows that H is a local ring with maximal ideal HM .

COROLLARY 4.2. *Let R be a local domain. Then K is an indecomposable R -module.*

Proof. Since H is a local ring by Proposition 4.1, it can have no idempotents other than the identity. Therefore K is indecomposable.

REMARKS. It is not true in general that if R is a local domain and I is an ideal of R , then $K \otimes_R I$ is indecomposable, as we shall see in Theorem 7.4.

COROLLARY 4.3. *Let R be an h -local domain, and H its completion. Then H is a direct product of commutative, local rings.*

Proof. By [8, Theorem 8.5] we have $H \cong \prod H(M)$, where

$$H(M) = \text{Hom}_{R_M}(K_M, K_M)$$

for all maximal ideals M of R . By Proposition 4.1, $H(M)$ is a commutative local ring.

LEMMA 4.4. *Let R be an integral domain and H its completion. Let J be an ideal of R such that $J \cap R = 0$. Let P be an ideal of H that is maximal with respect to the properties that $J \subset P$ and $P \cap R = 0$. Then P is a prime ideal of H ; and if P is not a cotorsion R -module, then P is a maximal ideal of R such that H/P is torsion-free and divisible over R .*

Proof. It is clear from an application of Zorn's Lemma that there exist ideals P of H that are maximal with respect to the properties that $J \subset P$ and $P \cap R = 0$. Let P be such an ideal. Let a, b be elements of H such that $ab \in P$. Suppose that $a \notin P$ and $b \notin P$. Then we have $r = p + ha$ and $s = q + kb$, where r, s are nonzero elements of R , $p, q \in P$, and $h, k \in H$. Then $rs \in P \cap R = 0$, and so $r = 0$ or $s = 0$, which is a contradiction. Thus either $a \in P$ or $b \in P$; and so P is a prime ideal of H .

Suppose that P is not a cotorsion R -module. Assume that P is not a maximal ideal of H . Then there is an ideal A of H that is properly between P and H . But then $A \cap R \neq 0$, and so by [8, Proposition 7.8], A is a cotorsion R -module. Let S be the intersection of all cotorsion submodules of H that contain P . Now H/P is a torsion-free R -module, since P is a prime ideal of H and $P \cap R = 0$. Therefore [8, Proposition 2.8] applies, and we have that S/P is torsion-free and divisible, and that S is a cotorsion submodule of H . Thus S is an ideal of H , and $S \subset A$. Therefore, $S \neq P$ and $S \neq H$. Hence $S \cap R \neq 0$, and so there exists a nonzero element $r \in S \cap R$. By [8, Proposition 7.8], $P + Hr$ is a cotorsion R -module, and so $S \subset P + Hr$. But of course $P + Hr \subset S$, and so we have $S = P + Hr$. Therefore,

$$S/P = (P + Hr)/P \cong Hr/(P \cap Hr) = Hr/Pr \cong H/P.$$

Thus H/P is torsion-free and divisible. But by [8, Proposition 2.8] this implies that $H = S$. This contradiction shows that P is a maximal ideal of H . It then follows from [8, Proposition 7.8] that H/P is torsion-free and divisible.

THEOREM 4.5. *Let R be a local domain. Then the following statements are equivalent:*

- (1) *Every proper R -submodule of K is h -reduced.*
- (2) *If J is a nonzero ideal of H , then $J \cap R \neq 0$.*
- (3) *If J is any ideal of H and $I = J \cap R$, then $J = HI$.*
- (4) *Every nonzero element of H is an epimorphism of K .*

If any of these conditions hold, then H is an integral domain.

Proof. (1) \Rightarrow (2). Suppose that J is a nonzero ideal of H , and that $J \cap R = 0$. By Proposition 4.1, H is a local ring with maximal ideal HM , where M is the maximal ideal of R , and $H/HM \cong R/M$. Let P be an ideal of H maximal with respect to the properties that $J \subset P$ and $P \cap R = 0$. Since $P \neq HM$, we have by Lemma 4.4 that P is a cotorsion R -module and H/P is torsion-free. Thus we have an exact sequence:

$$0 \longrightarrow K \otimes_R P \xrightarrow{f} K \otimes_R H \longrightarrow K \otimes_R H/P \longrightarrow 0.$$

Since $K \otimes_R H \cong K$ by [8, Proposition 5.5], we have by assumption that either f is an epimorphism, or $f = 0$. If $f = 0$, then $K \otimes_R P = 0$; and so P is torsion-free and divisible, which contradicts the fact that P is cotorsion. Hence f is an epimorphism. But this implies that $K \otimes_R H/P = 0$, and so H/P is torsion-free and divisible. But this also contradicts the fact that P is cotorsion. Thus we must have that $J \cap R \neq 0$.

(2) \Leftrightarrow (3). This equivalence has been proved in [8, Proposition 7.8].

(2) \Rightarrow (4). Let g be a nonzero element of H . Then by assumption there exists a nonzero element r in $Hg \cap R$. Thus $r = kg$, where $k \in H$. Now $K = rK = g(k(K)) \subset g(K) \subset K$. Hence $K = g(K)$, and g is an epimorphism.

(4) \Rightarrow (1). Let L be a nonzero, h -divisible submodule of K . Then by [8, Lemma 3.1] we have that $L \cong K \otimes_R \text{Hom}_R(K, L)$. Therefore, $\text{Hom}_R(K, L) \neq 0$. Now $\text{Hom}_R(K, L) \subset H$, and every element of H is an epimorphism on K . Thus $L = K$, and so K has no proper, h -divisible submodules.

Suppose that (4) holds and that h, k are elements of H such that $hk = 0$. Since h and k are epimorphisms or zero, we must have that either $h = 0$ or $k = 0$. Thus H is an integral domain.

DEFINITION. We will say that a local domain is *closed*, if it satisfies the conditions of Theorem 4.5.

Examples of closed, local domains.

(1) A valuation ring is closed, since the submodules of K are linearly ordered.

(2) Let R be an analytically irreducible, Noetherian, local domain of Krull dimension one. Then the completion of R in the maximal ideal topology is equal to H (see [8, p. 49]). By definition, H is then an integral domain. Since the Krull dimension of H is equal to the Krull dimension of R , which is one, the only prime ideals of H are 0 and HM (where M is the maximal ideal of R). Hence by Lemma 4.4 we have $J \cap R \neq 0$ for every nonzero ideal J of H . Thus R is a closed, local domain.

THEOREM 4.6. *Let R be a closed, local domain. Then we have the following:*

- (1) *If \bar{Q} is the quotient field of H , then $\bar{Q} \cong H \otimes_R Q$ and $\bar{Q}/H \cong K$.*
- (2) *If I is a nonzero ideal of R , then $\bar{Q}/HI \cong Q/I$ and $H/HI \cong R/I$.*
- (3) *H is complete in the H -topology.*
- (4) *The categories of torsion R -modules and torsion H -modules are identical.*

- (5) *The categories of cotorsion R -modules and cotorsion H -modules are identical.*
 (6) *If S is any H -module, then $\text{inj. dim.}_R S = \text{inj. dim.}_H S$, and $\text{w. dim.}_R S = \text{w. dim.}_H S$.*
 (7) *$\text{gl. dim. } R = \text{gl. dim. } H$ and $\text{w. gl. dim. } R = \text{w. gl. dim. } H$.*

Proof. Let J be a nonzero ideal of H and $I = J \cap R$. By Theorem 4.5 we have $J = HI$; and by [8, Proposition 5.10] we have $H/J \cong R/I$. By [8, Proposition 5.5] we have $\text{Ext}_H^n(H/J, H \otimes_R Q) \cong \text{Ext}_R^n(R/I, H \otimes_R Q) = 0$ for every $n \geq 0$. Thus $H \otimes_R Q$ is an injective H -module. Consider the exact sequence:

$$0 \rightarrow R \rightarrow Q \rightarrow K \rightarrow 0.$$

If we tensor this sequence with H we obtain the exact sequence:

$$0 \rightarrow H \rightarrow H \otimes_R Q \rightarrow K \rightarrow 0$$

by [8, Proposition 5.5]. Since K is torsion as an R -module, it is a fortiori torsion as an H -module. Thus $H \otimes_R Q$ is the injective envelope of H , and so $H \otimes_R Q \cong \tilde{Q}$, the quotient field of H . From this last exact sequence we clearly have $\tilde{Q}/H \cong K$.

Now $J \cong H \otimes_R I$ by [8, Proposition 5.10] and [8, Proposition 7.8]. Thus we have $\tilde{Q}/HI \cong K \otimes_H HI \cong K \otimes_H (H \otimes_R I) \cong K \otimes_R I \cong Q/I$.

It is clear from Theorem 4.5 that the R -topology and the H -topology on an H -module are the same. In particular, H is complete in the H -topology.

It follows from [8, Proposition 5.5] that the categories of torsion R -modules and torsion H -modules are the same. And it follows from [8, Proposition 5.7] that the categories of cotorsion R -modules and cotorsion H -modules are the same.

If S is any H -module, then the injective dimension of S over R or over H is determined by the vanishing of $\text{Ext}_R^n(T, S)$ or $\text{Ext}_H^n(T, S)$, respectively, where T ranges over all torsion R -modules (which are the same as all torsion H -modules). These two Ext modules are isomorphic by [8, Proposition 5.5]. Thus we have $\text{inj. dim.}_R S = \text{inj. dim.}_H S$. A similar argument shows that $\text{w. dim.}_R S = \text{w. dim.}_H S$.

The weak global dimension of an integral domain is completely determined by its torsion modules. Thus by [2, Chapter VI, Proposition 4.1.2] we have $\text{w. gl. dim. } R = \text{w. gl. dim. } H$. By [8, Proposition 3.5] we have that $\text{gl. dim. } R = \sup \text{inj. dim.}_R A$, where A ranges over all torsion-free, cotorsion R -modules A . A similar statement is true for H . Since these two categories are identical, and since the injective dimensions of the modules in this category are the same whether considered over R or over H , we have that $\text{gl. dim. } R = \text{gl. dim. } H$.

DEFINITION. Almost maximal valuation rings and maximal valuation rings have been defined in terms of the existence of solutions of systems of simultaneous congruences. However, in view of [6, Theorem 4] and [6, Theorem 9] we may give the following alternative definitions. A valuation ring R is called *almost maximal* if K

is injective; it is called *maximal* if it is both almost maximal and complete in the R -topology.

PROPOSITION 4.7. *Let R be an integral domain and H its completion. Then R is an almost maximal valuation ring if and only if H is a maximal valuation ring.*

Proof. Suppose that H is a valuation ring. Then by [8, Proposition 5.10] the ideals of R are linearly ordered. Hence R is a valuation ring. Thus we may assume that R is a valuation ring. Then by [8, Proposition 8.8], H is a valuation ring. If \bar{Q} is the quotient field of H , then by Theorem 4.6, $\bar{Q}/H \cong K$ and $\text{inj. dim.}_R K = \text{inj. dim.}_H K$. The proposition now follows from the definitions.

PROPOSITION 4.8. *Let R be a valuation ring and H its completion. Then R is almost maximal if and only if there exists an injective R -module C such that $H \cong \text{Hom}_R(C, C)$.*

Proof. If R is almost maximal, then K is injective and $H \cong \text{Hom}_R(K, K)$. Conversely, suppose that there exists an injective R -module C such that $H \cong \text{Hom}_R(C, C)$. Since H is h -reduced, C is a torsion R -module. Thus by Theorem 4.6, C is an injective H -module and $H \cong \text{Hom}_H(C, C)$. Let A be any H -module. Since H is a valuation ring, we have $\text{w. gl. dim. } H = 1$, and so $\text{Tor}_2^H(A, C) = 0$. Thus using [2, Chapter VI, Proposition 5.1] we have $\text{Ext}_H^2(A, H) \cong \text{Ext}_H^2(A, \text{Hom}_H(C, C)) \cong \text{Hom}_H(\text{Tor}_2^H(A, C), C) = 0$. Therefore, $\text{inj. dim.}_H H = 1$, and so K is H -injective. Hence by Theorem 4.6, K is R -injective, and R is an almost maximal valuation ring.

5. Decomposable torsion modules. The fundamental theorem of Abelian groups shows that every finitely generated torsion module over a Dedekind ring is a direct sum of cyclic modules. I. Kaplansky has extended this result to include almost maximal valuation rings ([3, Theorem 14]). The present author has proved that if a Noetherian integral domain has this property, then it must be a Dedekind ring ([6, Proposition 15]). The main purpose of this section is to generalize the above result of Kaplansky and prove its converse (see Theorem 5.7). The story does not end here, for it would be desirable to remove the assumption in Theorem 5.7 that R is an h -local domain.

PROPOSITION 5.1. *Let R be an h -local Prufer domain, and let A be a direct sum of cyclic R -modules. Then any torsion and finitely generated R -submodule of A is again a direct sum of cyclic R -modules.*

Proof. Since R is a Prufer domain, the torsion submodule of A is a direct summand of A . Thus we may assume that A is a direct sum of a finite number of cyclic torsion R -modules. By Theorem 3.1 there are a finite number of maximal ideals

M_1, \dots, M_n of R such that $A \cong A_{M_1} \oplus \dots \oplus A_{M_n}$. Each A_{M_i} is a finite direct sum of cyclic torsion R_{M_i} -modules. Let B be an R -submodule of A . Then $B = B_{M_1} \oplus \dots \oplus B_{M_n}$, and $B_{M_i} \subset A_{M_i}$. By Lemma 2.1 the R -structure and the R_{M_i} -structure of A_{M_i} and B_{M_i} are the same. Thus we may assume without loss of generality that R is a local domain. But then R is a valuation ring, and the proposition is an immediate consequence of [6, Proposition 14].

PROPOSITION 5.2. *Let R be an h -local Prufer domain, and let B be a finitely generated R -module. Let $E(B)$ be the injective envelope of B . Then $E(B)$ is a finite direct sum of indecomposable injective R -modules.*

Proof. Since R is a Prufer domain, the torsion submodule of B is a direct summand of B . Hence $E(B)$ is the direct sum of the injective envelope of the torsion submodule of B and of a finite number of copies of Q . Thus we may assume that B is a torsion R -module. By Theorem 3.1 there are a finite number of maximal ideals M_1, \dots, M_n of R such that $B \cong B_{M_1} \oplus \dots \oplus B_{M_n}$. If M is a maximal ideal of R and C an injective R_M -module, then by [2, Chapter VI, Proposition 4.1.3] C is also an injective R -module. Hence we may assume that R is a local ring. But then R is a valuation ring, and so the proposition is an immediate consequence of [6, Proposition 1].

DEFINITION. Let R be an integral domain and A a nonempty subset of an R -module. Then we will define $O(A)$ to be the annihilator of A ; i.e., $O(A) = \{r \in R \mid rA = 0\}$.

LEMMA 5.3. *Let R be a valuation ring and B an R -module. Let y be a nonzero element of B and r an element of R such that $ry \neq 0$. Then $rO(ry) = O(y)$.*

Proof. Since $r \notin O(y)$, we have $O(y) \subset Rr$. Thus $O(y) = Jr$, where J is an ideal of R . Let $a \in O(ry)$; then $0 = ary$, and so $ar \in O(y)$. Therefore, $a \in J$ and we have $O(ry) \subset J$. Conversely, let $a \in J$. Then $ar \in O(y)$, and so $a \in O(ry)$. Thus $J \subset O(ry)$, and we have $J = O(ry)$. Hence $O(y) = Jr = rO(ry)$.

LEMMA 5.4. *Let R be a valuation ring, and let $E(K)$ be the injective envelope of K . Then $E(K)$ is an indecomposable injective R -module. Furthermore, if $y \in E(K)$, then $O(y)$ is a principal ideal of R .*

Proof. Let S and T be nonzero submodules of $E(K)$. Then $S \cap K \neq 0$ and $T \cap K \neq 0$, since $E(K)$ is an essential extension of K . Now the submodules of K are linearly ordered, and so we have that $(S \cap K) \cap (T \cap K) \neq 0$. Therefore, $S \cap T \neq 0$, and so $E(K)$ is an indecomposable, injective R -module.

Let y be a nonzero element of $E(K)$. Take a nonzero element a in R which is not a unit in R , and let $v = (1/a) + R$, a nonzero element of K . Then $Ry \cap Rv \neq 0$, and

so there exist elements r, s in R such that $ry = sv \neq 0$. Thus by Lemma 5.3 we have $O(y) = rO(ry) = rO(sv) = (r/s)O(v)$. Now $O(v) = Ra$, and so $O(y) = R(ra/s)$ is a principal ideal of R .

The following lemma has been proved by I. Kaplansky in [3, Theorem 14]⁽¹⁾. We give an independent proof that is based on the properties of indecomposable, injective modules.

LEMMA 5.5. *Let R be a valuation ring. Then R is an almost maximal valuation ring if and only if every finitely generated R -module is a direct sum of cyclic R -modules.*

Proof. Assume that R is an almost maximal valuation ring, and let B be a finitely generated R -module. By Proposition 5.2 the injective envelope $E(B)$ is a finite direct sum of indecomposable, injective R -modules. By [6, Theorem 4] every indecomposable, injective R -module is of the form Q/I , where I is an ideal of R . Now Q/I has the property that every finitely generated R -submodule is cyclic. Thus B is a finitely generated R -submodule of a direct sum of cyclic R -modules. Since B is a direct sum of its torsion submodule and a free R -module, we can assume without loss of generality that B is a torsion R -module. It then follows from Proposition 5.1 that B itself is a direct sum of cyclic R -modules.

Conversely assume that every finitely generated R -module is a direct sum of cyclic R -modules. By our definitions, to prove that R is an almost maximal valuation ring it is sufficient to prove that $E(K) = K$, where $E(K)$ is the injective envelope of K . Suppose that $E(K) \neq K$, and take $y \in E(K) - K$. Then by Lemma 5.4, $O(y)$ is a principal ideal of R , say $O(y) = Rb$, where b is a nonzero element of R and b is not a unit of R . Let $x = (1/b) + R$; then x is an element of $K \subset E(K)$ and $O(x) = Rb = O(y)$.

By hypothesis, $Rx + Ry$ is a direct sum of cyclic R -modules. Since $E(K)$ is an indecomposable, injective R -module by Lemma 5.4, $Rx + Ry$ must actually be a cyclic R -module. Hence there exists an element $z \in E(K)$ such that $Rx + Ry = Rz$. Thus there exist nonzero elements r, s in R such that $x = rz$ and $y = sz$. If r divides s , then there exists an element $d \in R$ such that $s = rd$. Then we have $y = sz = drz = dx$, and so y is an element of K , which is a contradiction. Therefore, s divides r , and so there exists an element $a \in R$ such that $r = as$. Hence we have $x = rz = asz = ay$. By Lemma 3 we then have $O(y) = aO(ay) = aO(x)$. Since $O(y) = Rb = O(x)$, it follows that $Rb = Rab$. Hence $Ra = R$, and so a is a unit in R . Therefore, $y = a^{-1}ay = a^{-1}x$ is an element of K . This contradiction shows that $E(K) = K$, and so R is an almost maximal valuation ring.

LEMMA 5.6. *Let R be a local domain. If every finitely generated, torsion R -module is a direct sum of cyclic R -modules, then R is an almost maximal valuation ring.*

⁽¹⁾ Also see Theorem 11.1 of Kaplansky's paper *Elementary divisors and modules*, Trans. Amer. Math. Soc. 66 (1949), 464–491.

Proof. Let a, b be nonzero elements of R . Suppose that $a \notin Rb$ and $b \notin Ra$. We will show that this yields a contradiction, which will therefore prove that R is a valuation ring. The lemma will then follow from Lemma 5.5.

Let $x = (1/a) + R$ and $y = (1/b) + R$, so that x and y are nonzero elements of K . Let $A = Rx + Ry$; then A is a finitely generated torsion R -module. Hence by assumption we have $A = A_1 \oplus \cdots \oplus A_n$, where each A_i is a cyclic torsion R -module. For each $i = 1, \dots, n$ let $p_i: A \rightarrow A$ be defined to be the projection on A_i with kernel $A_1 \oplus \cdots \oplus A_{i-1} \oplus A_{i+1} \oplus \cdots \oplus A_n$.

Now $A_i = p_i(A) = p_i(Rx) + p_i(Ry)$. But A_i is a local R -module, and so we must have either $A_i = p_i(Rx)$ or $A_i = p_i(Ry)$. We can assume that $A_i = p_i(Rx)$ for $i \leq k$ and that $A_i = p_i(Ry)$ for $i \geq k+1$. Let $B_1 = A_1 \oplus \cdots \oplus A_k$, and $B_2 = A_{k+1} \oplus \cdots \oplus A_n$, so that $A = B_1 \oplus B_2$. Since $ap_i(x) = p_i(ax) = p_i(0) = 0$, we have $p_i(x) \subset Rx$ for all $i \leq n$. Therefore $B_1 \subset Rx$. Similarly we have $B_2 \subset Ry$.

Since $A = B_1 \oplus B_2$, there exist $z_1 \in B_1$ and $z_2 \in B_2$ such that $x = z_1 + z_2$. Since $B_1 \subset Rx$ and $B_2 \subset Ry$, there exist elements r, s in R such that $z_1 = rx$ and $z_2 = sy$. Hence we have $(1/a) + R = [(r/a) + R] + [(s/b) + R] = [(r/a) + (s/b)] + R$. Thus there exists an element c in R such that $1/a = (rb + sa)/ab + c$. Hence $b = rb + sa + cab$, and so $(1-r)b = (s+cb)a$. If r is not a unit in R , then $1-r$ is a unit in R , and we would have $b \in Ra$ by the preceding equation. This is a contradiction to our assumptions, and so r must be a unit in R . Therefore, $x = r^{-1}rx = r^{-1}z_1 \in B_1$. It follows that $B_1 = Rx$. Similarly we have $B_2 = Ry$.

Thus we have $A = B_1 \oplus B_2 = Rx \oplus Ry$, and so $Rx \cap Ry = 0$. Let $u = 1/a$ and $v = 1/b$; then $Rx = Ru/R$ and $Ry = Rv/R$. Hence $0 = Ru/R \cap Rv/R = (Ru \cap Rv)/R$, and so $Ru \cap Rv = R$. Now we have $(Ra + Rb)^{-1} = Ru \cap Rv$. Thus we have shown that if a, b are elements of R such that $a \notin Rb$ and $b \notin Ra$, then $(Ra + Rb)^{-1} = R$.

Let t be a nonzero element of R and let $d = bt$ and $e = at$. If $d \in Re$, then $b \in Ra$, and so we have $d \notin Re$. Similarly we have $e \notin Rd$. Hence by what we have shown above we have $(Rd + Re)^{-1} = R$. Let $D = Ru + Rv$, and $s = abt$. Then $sD = Rd + Re$, and so $(sD)^{-1} = R$. But $(sD)^{-1} = (1/s)D^{-1}$, and so we have $D^{-1} = sR$.

If we take $t = 1$, then $s = ab$, and we have $D^{-1} = abR$. If we take $t = a$, then $s = a^2b$, and we have $D^{-1} = a^2bR$. Thus we have $abR = a^2bR$, which implies that $R = Ra$. Therefore, a is a unit in R , and so $b \in Ra$. This contradiction shows that if a and b are nonzero elements of R , then either $a \in Rb$, or $b \in Ra$. This shows that R is a valuation ring, and hence by Lemma 5.5 an almost maximal valuation ring. This completes the proof of the lemma.

The following theorem generalizes and proves the converse of [3, Theorem 14].

THEOREM 5.7. *Let R be an h -local domain. Then the following statements are equivalent:*

- (1) R is a Prufer domain and K is injective.
- (2) R_M is an almost maximal valuation ring for every maximal ideal M of R .
- (3) Every finitely generated torsion R -module is a direct sum of cyclic R -modules.

Proof. (1) \Rightarrow (2). Since R is a Prufer ring, R_M is a valuation ring for every maximal ideal M of R . By Theorem 3.3 we have $0 = \text{inj. dim.}_R K = \sup_M \text{inj. dim.}_{R_M} K_M$. Thus K_M is R_M -injective, and so R_M is a maximal valuation ring for every maximal ideal M of R .

(2) \Rightarrow (3). Let T be a finitely generated, torsion R -module. Then by Theorem 3.1 there exists a finite number of maximal ideals M_1, \dots, M_n of R such that $T \cong T_{M_1} \oplus \dots \oplus T_{M_n}$. By Lemma 2.1 the R -structure and the R_{M_i} -structure of T_{M_i} are the same, and so without loss of generality we may assume that R is a local ring. Then by assumption R is an almost maximal valuation ring. But then by Lemma 5.5, T is a direct sum of cyclic R -modules.

(3) \Rightarrow (1). Let M be a maximal ideal of R , and let T be a finitely generated, torsion R_M -module. Then by Lemma 2.1, T is a finitely generated, torsion R -module. Hence by assumption T is a direct sum of cyclic R -modules. But by Lemma 2.1 again, T is a direct sum of cyclic R_M -modules. Thus Lemma 5.6 applies to R_M , and R_M is an almost maximal valuation ring. Since this is true for all maximal ideals M of R , R is a Prufer domain; and by Theorem 3.3 we have

$$\text{inj. dim.}_R K = \sup_M \text{inj. dim.}_{R_M} K_M = 0.$$

6. Decomposable torsion-free modules. There is very little that is known about the theory of torsion-free modules over an integral domain. A beginning step in the program is to attempt to find out when a torsion-free module of finite rank is indecomposable. I. Kaplansky has proved that over a maximal valuation ring every torsion-free module of countable rank is a direct sum of modules of rank one ([3, Theorem 12]). This section is concerned with determining the class of all integral domains that have the property that every torsion-free module of finite rank is a direct sum of modules of rank one. In the course of this investigation we exhibit a number of different ways of constructing indecomposable torsion-free modules of rank two for various kinds of integral domains. This considerably sharpens the results in [7, Proposition 7 and Corollary].

Most of this section is devoted to proving the following theorem:

THEOREM 6.1. *Let R be an integral domain. Then the following statements are equivalent:*

- (1) *Every torsion-free R -module of finite rank is a direct sum of R -modules of rank one.*
- (2) *R is an h -local domain with at most two maximal ideals M_1 , M_2 and R_{M_1} , R_{M_2} are maximal valuation rings.*

DEFINITION. We will say that an integral domain has *property D*, if it satisfies condition (1) of Theorem 6.1.

LEMMA 6.2. *Let R be an integral domain with property D , and let S be a subring of Q containing R . Then S also has property D .*

Proof. Let A be a torsion-free S -module. Suppose that we have $A = A_1 \oplus A_2$, where A_1, A_2 are R -submodules of A . Let s be a nonzero element of S , and x a nonzero element of A_1 . Then $sx = y_1 + y_2$, where $y_i \in A_i$ for $i=1, 2$. Now there exist elements a and b in R such that $b \neq 0$ and $s = a/b$. Then $ax = bsx = by_1 + by_2$. But $ax \in A_1$, and so $by_2 = 0$. Hence $y_2 = 0$, and so $sx = y_1 \in A_1$. Therefore, A_1 is an S -module. Similarly A_2 is an S -module. Thus if R has property D , so has S .

LEMMA 6.3. *Let R be an integral domain, complete in the R -topology; and let J be a nonzero ideal of R . Then $K \otimes_R J$ has no proper, h -divisible R -submodules.*

Proof. We have $K \otimes_R J \cong Q/J$. Let B be a proper R -submodule of Q containing J . Since R is complete in the R -topology, R is cotorsion as an R -module. Therefore J is also a cotorsion R -module, and we have $\text{Ext}_R^1(Q, J) = 0$. Since B is a proper R -submodule of Q , we have $\text{Hom}_R(Q, B) = 0$. Thus we have an exact sequence:

$$0 = \text{Hom}_R(Q, B) \rightarrow \text{Hom}_R(Q, B/J) \rightarrow \text{Ext}_R^1(Q, J) = 0,$$

where the end R -modules are zero. Thus $\text{Hom}_R(Q, B/J) = 0$, and so B/J is h -reduced. Therefore, $K \otimes_R J$ has no proper, h -divisible R -submodules.

LEMMA 6.4. *Let R be an integral domain, complete in the R -topology; and let I and J be nonzero ideals of R . Consider an exact sequence:*

$$0 \longrightarrow I \xrightarrow{i} A \xrightarrow{p} J \longrightarrow 0,$$

where A is an R -module. Then if A is decomposable, the sequence splits.

Proof. Assume that A has nonzero R -submodules A_1 and A_2 such that $A = A_1 \oplus A_2$. Suppose the sequence does not split. Let p_i be the restriction of p to A_i for $i=1, 2$. Now p_1 is a monomorphism. For suppose that $\text{Ker } p_1 \neq 0$. Since A_1 and J are torsion-free R -modules of rank one, this means that $p_1 = 0$. Hence we have: $\text{Ker } p + A_2 = A$. If $A_2 \cap \text{Ker } p \neq 0$, then since A_2 and J are torsion-free R -modules of rank one, we have $A_2 \subset \text{Ker } p$. This is clearly a contradiction, and so we have $A_2 \cap \text{Ker } p = 0$. Therefore, $A = \text{Ker } p \oplus A_2$; but this means the exact sequence splits, which contradicts our assumption. Thus we have $\text{Ker } p_1 = 0$, and so p_1 is a monomorphism. Similarly p_2 is a monomorphism.

Let $J_1 = p_1(A_1)$ and $J_2 = p_2(A_2)$, so that $J = J_1 + J_2$. Define $f: A \rightarrow J_1 \oplus J_2$ by $f(x_1 + x_2) = (p_1(x_1), p_2(x_2))$, where $x_i \in A_i$ for $i = 1, 2$. Since p_i is an isomorphism of A_i onto J_i for $i = 1, 2$, it follows that f is an isomorphism. Define

$$\Delta: J_1 \cap J_2 \rightarrow J_1 \oplus J_2$$

by $\Delta(y) = (y, -y)$ for $y \in J_1 \cap J_2$; and define $\gamma: J_1 \oplus J_2 \rightarrow J$ by $\gamma((y_1, y_2)) = y_1 + y_2$, where $y_i \in J_i$ for $i = 1, 2$. We have that Δ is a monomorphism, γ is an epimorphism and $\text{Ker } \gamma = \text{Im } \Delta$. Since $\gamma f = p$, f maps I into $\text{Im } \Delta$. Thus we can define a map $g: I \rightarrow J_1 \cap J_2$ by $g = \Delta^{-1} f$. Then $\Delta g = f$, and g is an isomorphism. Then we have a commutative diagram with exact rows:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & I & \xrightarrow{f} & A & \xrightarrow{p} & J & \longrightarrow & 0 \\ & & \downarrow g & & \downarrow f & & \downarrow 1 & & \\ 0 & \longrightarrow & J_1 \cap J_2 & \xrightarrow{\Delta} & J_1 \oplus J_2 & \xrightarrow{\gamma} & J & \longrightarrow & 0 \end{array}$$

and the vertical maps are isomorphisms. Since the top row does not split, the bottom row does not split either.

Now we have an isomorphism $\mu: K \otimes_R (J_1 \oplus J_2) \rightarrow (K \otimes_R J_1) \oplus (K \otimes_R J_2)$ defined by $\mu(k \otimes (x_1, x_2)) = (k \otimes x_1, k \otimes x_2)$, where $k \in K$ and $x_i \in J_i$ for $i = 1, 2$. Define $\xi: K \otimes_R (J_1 \cap J_2) \rightarrow (K \otimes_R J_1) \oplus (K \otimes_R J_2)$ by $\xi = \mu(1 \otimes \Delta)$ and define

$$\eta: (K \otimes_R J_1) \oplus (K \otimes_R J_2) \rightarrow K \otimes_R J$$

by $\eta = (1 \otimes \gamma) \mu^{-1}$. Then we have a commutative diagram with exact rows:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & K \otimes_R (J_1 \cap J_2) & \xrightarrow{1 \otimes \Delta} & K \otimes_R (J_1 \oplus J_2) & \xrightarrow{1 \otimes \gamma} & K \otimes_R J & \longrightarrow & 0 \\ & & \downarrow 1 & & \downarrow \mu & & \downarrow 1 & & \\ 0 & \longrightarrow & K \otimes_R (J_1 \cap J_2) & \xrightarrow{\xi} & (K \otimes_R J_1) \oplus (K \otimes_R J_2) & \xrightarrow{\eta} & K \otimes_R J & \longrightarrow & 0. \end{array}$$

Hence $K \otimes_R J = \eta(K \otimes_R J_1) + \eta(K \otimes_R J_2)$. By Lemma 6.3, $K \otimes_R J$ has no proper h -divisible R -submodules, and thus we can assume that $\eta(K \otimes_R J_1) = K \otimes_R J$. Therefore, $(K \otimes_R J_1) \oplus (K \otimes_R J_2) = \text{Im } \xi + (K \otimes_R J_1)$. Now every element of

$$K \otimes_R (J_1 \cap J_2)$$

is of the form $k \otimes x$, where $k \in K$ and $x \in J_1 \cap J_2$. We have $\xi(k \otimes x) = (k \otimes x, -k \otimes x)$, and thus if $\xi(k \otimes x) \in K \otimes_R J_1$, then there exist elements $k_1 \in K$ and $x_1 \in J_1$ such that $(k \otimes x, -k \otimes x) = (k_1 \otimes x_1, 0)$. Hence $k \otimes x = 0$, and so we have $\text{Im } \xi \cap (K \otimes_R J_1) = 0$. Thus the bottom row of the above commutative diagram splits. It follows that the

top row of the above diagram splits. Therefore the following sequence is exact and splits:

$$0 \longrightarrow \operatorname{Hom}_R(K, K \otimes (J_1 \cap J_2)) \xrightarrow{(1 \otimes \Delta)_*} \operatorname{Hom}_R(K, K \otimes (J_1 \oplus J_2)) \\ \xrightarrow{(1 \otimes \gamma)_*} \operatorname{Hom}_R(K, K \otimes J) \longrightarrow 0.$$

Now by [8, Corollary 3.3] we have a commutative diagram:

$$\begin{array}{ccccccc} 0 \rightarrow & J_1 \cap J_2 & \xrightarrow{\Delta} & J_1 \oplus J_2 & \xrightarrow{\gamma} & J & \rightarrow 0 \\ & \downarrow & & \downarrow & & \downarrow & \\ 0 \rightarrow & \operatorname{Hom}_R(K, K \otimes (J_1 \cap J_2)) & \xrightarrow{(1 \otimes \Delta)_*} & \operatorname{Hom}_R(K, K \otimes (J_1 \oplus J_2)) & & & \\ & & & & \xrightarrow{(1 \otimes \gamma)_*} & \operatorname{Hom}_R(K, K \otimes J) & \rightarrow 0, \end{array}$$

where the bottom row is exact and splits. Since $J_1 \cap J_2$, $J_1 \oplus J_2$, and J are all torsion-free, cotorsion R -modules, we have by [8, Proposition 3.2] that the vertical maps of this last diagram are isomorphisms. Thus the top row of this last diagram splits, which contradicts the fact that we have proved that it does not split. The only way out of this dilemma is that the exact sequence provided by the lemma must have been split to begin with. This concludes the proof of Lemma 6.4.

PROPOSITION 6.5. *Let R be an integral domain, complete in the R -topology. If R has property D , then R is a maximal valuation ring.*

Proof. Let I and J be nonzero ideals of R and consider an exact sequence of the form:

$$0 \rightarrow I \rightarrow A \rightarrow J \rightarrow 0,$$

where A is an R -module. By property D , A is a direct sum of two R -modules of rank one. Then by Lemma 6.4, this exact sequence splits. This means that $\operatorname{Ext}_R^1(J, I) = 0$. Since we have $\operatorname{Ext}_R^2(R/J, I) \cong \operatorname{Ext}_R^1(J, I)$, it follows that $\operatorname{Ext}_R^2(R/J, I) = 0$. Therefore, $\operatorname{inj. dim.}_R I = 1$. From the exact sequence:

$$0 \rightarrow I \rightarrow Q \rightarrow K \otimes_R I \rightarrow 0$$

we can then conclude that $K \otimes_R I$ is an injective R -module. By Lemma 6.3, $K \otimes_R I \cong Q/I$ is an indecomposable injective R -module.

Let a and b be nonzero elements of R . Let $I = Ra \cap Rb$, $L_1 = Ra$, and $L_2 = Rb$. Then $L_i/I \subset Q/I$ for $i = 1, 2$; and we have $(L_1/I) \cap (L_2/I) = 0$. Since Q/I is an indecomposable, injective R -module, either $L_1/I = 0$ or $L_2/I = 0$. Hence either $a \in Rb$ or $b \in Ra$. Therefore, R is a valuation ring. Since we have shown that $K \cong Q/R$

is an injective R -module, R is an almost maximal valuation ring. But R is complete in the R -topology, and so R is a maximal valuation ring.

LEMMA 6.6. *Let R be a local domain and consider an exact sequence of the form:*

$$0 \longrightarrow R \xrightarrow{i} A \xrightarrow{p} Q \longrightarrow 0.$$

Then if A is a decomposable R -module, the exact sequence splits.

Proof. Suppose that $A = A_1 \oplus A_2$, where A_1, A_2 are nonzero R -submodules of A . If we tensor the exact sequence with K we obtain the isomorphisms: $K \cong K \otimes_R R \cong K \otimes_R A \cong (K \otimes_R A_1) \oplus (K \otimes_R A_2)$. But by Corollary 4.2, K is an indecomposable R -module. Therefore we can assume that $K \otimes_R A_2 = 0$. Since A_2 is torsion-free of rank one, we must then have that $A_2 \cong Q$.

If $A_2 \cap \text{Ker } p \neq 0$, then we must have $A_2 \subset \text{Ker } p = \text{Im } i \cong R$. But since R is h -reduced, this can not happen, and so $A_2 \cap \text{Ker } p = 0$. Since Q has no proper divisible R -submodules, we have $p(A_2) = Q$. Therefore, $A = A_2 + \text{Ker } p$, and so $A = A_2 \oplus \text{Ker } p$. Thus the exact sequence splits.

COROLLARY 6.7. *Let R be a local domain with property D . Then R is a maximal valuation ring.*

Proof. Consider an exact sequence of the form:

$$0 \rightarrow R \rightarrow A \rightarrow Q \rightarrow 0.$$

By property D , A is decomposable, and so by Lemma 6.6 the exact sequence splits. Therefore, we have $\text{Ext}_R^1(Q, R) = 0$; and so by [8, Proposition 5.2], R is complete in the R -topology. Hence by Proposition 6.5, R is a maximal valuation ring.

LEMMA 6.8. *Let R be an integral domain with exactly two maximal ideals. Then if R has property D , it is an h -local domain.*

Proof. Let M_1, M_2 be the maximal ideals of R and let $S = R_{M_1} + R_{M_2} \subset Q$. Since $R_{M_1} \cap R_{M_2} = R$, we have an exact sequence:

$$0 \rightarrow R \rightarrow R_{M_1} \oplus R_{M_2} \rightarrow S \rightarrow 0.$$

By Lemma 6.2, R_{M_1} has property D . Since R_{M_1} is a local ring, we can conclude from Corollary 6.7 that R_{M_1} is a maximal valuation ring, and hence is a cotorsion R_{M_1} -module. Since we have $\text{Ext}_R^1(Q, R_{M_1}) \cong \text{Ext}_{R_{M_1}}^1(Q, R_{M_1})$ by [2, Chapter VI, Proposition 4.1.3], it follows that R_{M_1} is a cotorsion R -module. Similarly R_{M_2} is a

cotorsion R -module, and thus so is $R_{M_1} \oplus R_{M_2}$. Therefore, since S is torsion-free, we have an imbedding: $R \subset H \subset R_{M_1} \oplus R_{M_2}$ by [8, Proposition 2.8], where H is the completion of R in the R -topology. If R were complete, then it would be a local domain by Proposition 6.5. Hence R is not complete, and so $R \neq H$. Thus H/R is not zero; by [8, Proposition 5.2] it is torsion-free and divisible. Since $H/R \subset S$, and since S is torsion-free of rank one, we have $H/R = S$. Therefore, $H = R_{M_1} \oplus R_{M_2}$. Since $R_{M_i} = H(M_i)$ for $i = 1, 2$, it follows from [8, Theorem 8.5] that R is an h -local domain.

PROPOSITION 6.9. *Let R be an h -local domain with more than two maximal ideals. Then R has an indecomposable, torsion-free R -module of rank two.*

Proof. Let M_1, M_2, M_3 be three distinct maximal ideals of R . Let V be a two dimensional vector space over Q with basis u and v . Choose an element $b \in M_3$ such that $b \notin M_1$ and $b \notin M_2$. Let $z = (1/b)u + v \in Q$. We define three R -submodules C_1, C_2 , and C_3 of V as follows. Let $C_1 = Qu \oplus R_{M_1}v$; $C_2 = R_{M_2}u \oplus Qv$; and $C_3 = R_{M_3}u \oplus R_{M_3}z$. We let $C = C_1 \cap C_2 \cap C_3$; we will show that C is an indecomposable, torsion-free R -module of rank two.

(1) We have $u, z \in C$; and $(R_{M_2} \cap R_{M_3})u \subset C$. Clearly $(R_{M_2} \cap R_{M_3})u \subset C$, and so $u \in C$. Certainly $z \in C_1$ and $z \in C_3$. Since $b \notin M_2$, we have $1/b \in R_{M_2}$; and so $z \in C_2$. Therefore z is in the intersection of the C_i 's, which is C .

(2) We have $C_1 = C_{M_1}$. Since $(R_{M_2} \cap R_{M_3})u \subset C$, we have $R_{M_1}(R_{M_2} \cap R_{M_3})u \subset C_{M_1}$. But $R_{M_1}(R_{M_2} \cap R_{M_3}) = Q$, since R is an h -local domain. Therefore, $Qu \subset C_{M_1}$. In particular, $(1/b)u \in C_{M_1}$. Now $z \in C \subset C_{M_1}$, and so $v = z - (1/b)u \in C_{M_1}$. Thus $C_1 = Qu \oplus R_{M_1}v \subset C_{M_1}$. But $C \subset C_1$, and so $C_{M_1} \subset (C_1)_{M_1} = C_1$. Thus we have $C_1 = C_{M_1}$.

(3) We have $C_2 = C_{M_2}$. Since $u \in C$, we have $R_{M_2}u \subset C_{M_2}$. Hence $(1/b)u \in C_{M_2}$. Thus $v = z - (1/b)u \in C_{M_2}$. Now $(R_{M_3} \cap R_{M_1})bv \subset C$. For let $a \in R_{M_3}$; then $abv = -au + abz \in C_3$, and so $R_{M_3}bv \subset C_3$. Since $Qv \subset C_2$, and $R_{M_1}v \subset C_1$, it follows that $(R_{M_3} \cap R_{M_1})bv \subset C$. Therefore, $R_{M_2}(R_{M_3} \cap R_{M_1})bv \subset C_{M_2}$. But we have

$$R_{M_2}(R_{M_3} \cap R_{M_1})b = Qb = Q,$$

since R is h -local. Hence $Qv \subset C_{M_2}$, and so $C_2 = R_{M_2}u \oplus Qv \subset C_{M_2}$. Since $C \subset C_2$, we have $C_{M_2} \subset (C_2)_{M_2} = C_2$. Thus $C_2 = C_{M_2}$.

(4) We have $C_3 = C_{M_3}$. Since u, z are in C , we have $C_3 = R_{M_3}u \oplus R_{M_3}z \subset C_{M_3}$. On the other hand, since $C \subset C_3$, we have $C_{M_3} \subset (C_3)_{M_3} = C_3$. Thus $C_3 = C_{M_3}$.

Now C is a torsion-free R -module of rank two, since it contains u and z . Suppose that C is decomposable. We then have $C = D_1 \oplus D_2$, where D_1, D_2 are R -submodules of C of rank one.

(5) We can assume that $(D_1)_{M_1} = Qu$ and $(D_2)_{M_2} = Qv$. For we have $C_1 = C_{M_1} = (D_1)_{M_1} \oplus (D_2)_{M_1}$. If both $(D_1)_{M_1}$ and $(D_2)_{M_1}$ are reduced, then C_1 is reduced. But

$Qu \subset C_1$, and so C_1 is not reduced. Thus we can assume that $(D_1)_{M_1}$ is not reduced. Therefore, $(D_1)_{M_1}$ is isomorphic to Q . Since Qu is the divisible submodule of C_1 , we have $(D_1)_{M_1} \subset Qu$. But then we must have $(D_1)_{M_1} = Qu$, since both modules have rank one. A similar argument for C_2 shows that we must have either $(D_1)_{M_2} = Qv$, or $(D_2)_{M_2} = Qv$. Since $(D_1)_{M_1} = Qu$, and D_1 has rank one, we can not have $(D_1)_{M_2} = Qv$. Thus $(D_2)_{M_2} = Qv$.

It follows from (5) that we have $D_1 \subset Qu$ and $D_2 \subset Qv$. Let $E_1 = (D_1)_{M_3}$ and $E_2 = (D_2)_{M_3}$. Then $E_1 \subset Qu$, $E_2 \subset Qv$, and $C_3 = C_{M_3} = E_1 \oplus E_2$. Now C_3 is a free R_{M_3} -module of rank two, and so E_1 is a free R_{M_3} -module with generator x_1 , and E_2 is a free R_{M_3} -module with generator x_2 .

There exists elements q_1, q_2 in Q such that $x_1 = q_1u$ and $x_2 = q_2v$. Since $z \in C_3$, there exist elements d_1, d_2 in R_{M_3} such that $(1/b)u + v = z = d_1x_1 + d_2x_2 = d_1q_1u + d_2q_2v$. Thus we have $1 = d_2q_2$. Since $q_2v = x_2 \in C_3$, there exist elements a_1, a_2 in R_{M_3} such that $q_2v = a_1u + a_2z = (a_1 + a_2/b)u + a_2v$. Thus we have $a_1 + a_2/b = 0$ and $a_2 = q_2$. Combining these results we have $0 = bd_2(a_1 + a_2/b) = bd_2a_1 + d_2a_2 = bd_2a_1 + 1$. Hence $1/b = -d_2a_1 \in R_{M_3}$. But $b \in M_3$, and so $1 = (1/b)b \in R_{M_3}M_3$. But this is contrary to the fact that $R_{M_3}M_3$ is a proper ideal of R_{M_3} . This contradiction proves that C is an indecomposable, torsion-free R -module of rank two.

LEMMA 6.10. *Let R be an h -local domain with exactly two maximal ideals M_1 and M_2 . If R_{M_1} and R_{M_2} are maximal valuation rings, then R has property D .*

Proof. Assume that R_{M_1}, R_{M_2} are maximal valuation rings. Let A be a torsion-free R -module of rank one. Then we have $\text{inj. dim.}_{R_{M_i}} A_{M_i} \leq 1$ for $i=1, 2$ by [6, Theorem 4]. Hence by Theorem 3.3 we have that $\text{inj. dim.}_R A \leq 1$. There are only three possibilities for A ; either $A \cong Q$, or A is isomorphic to an ideal of R , or A is a cotorsion R -module. For suppose that A is not isomorphic to Q , or to an ideal of R . We may clearly assume that $A \subset Q$. If $A_{M_1} = Q$ and $A_{M_2} = Q$, then $A = A_{M_1} \cap A_{M_2} = Q$, contrary to our assumption. Thus we can assume that $A_{M_2} \neq Q$. Then there exists a nonzero element $r \in R$ such that $rA_{M_2} \subset R_{M_2}$, and so A_{M_2} is isomorphic to an ideal of R_{M_2} . If we have $A_{M_1} \neq Q$ also, then there exists a nonzero element $s \in R$ such that $sA_{M_1} \subset R_{M_1}$. But then we would have $rsA \subset rA_{M_2} \cap sA_{M_1} \subset R_{M_2} \cap R_{M_1} = R$, and so A would be isomorphic to an ideal of R . This contradiction shows that $A_{M_1} = Q$. Hence we have $A = A_{M_1} \cap A_{M_2} = Q \cap A_{M_2} = A_{M_2}$. Therefore A is isomorphic to an ideal of R_{M_2} , and so is a cotorsion R_{M_2} -module. But then by [2, Chapter VI, Proposition 4.1.3] A is also a cotorsion R -module.

In the course of the above proof we have seen that A is a cotorsion R -module if and only if $A \neq Q$ and either $A = A_{M_1}$ or $A = A_{M_2}$.

Let C be a torsion-free R -module of rank $n < \infty$. We will prove that C is a direct sum of n R -modules of rank one by induction on n . The case $n=1$ is trivial, and so

we assume that the assertion is true for torsion-free R -modules of rank $< n$, and that $n > 1$. We can then clearly assume that C is reduced.

Suppose that C contains a nonzero, pure, cotorsion R -submodule D of rank $< n$. Then C/D is a torsion-free R -module of rank $< n$, and we have an exact sequence:

$$\text{Ext}_R^1(Q \otimes_R C/D, D) \rightarrow \text{Ext}_R^1(C/D, D) \rightarrow \text{Ext}_R^2(K \otimes_R C/D, D).$$

The first term of this sequence is zero, since D is a cotorsion R -module; and the last term is zero, since $\text{inj. dim.}_R D = 1$. Therefore, $\text{Ext}_R^1(C/D, D) = 0$; and so $C \cong D \oplus C/D$. By the induction hypothesis both D and C/D are direct sums of torsion-free R -modules of rank one. Hence C is also such a direct sum. Thus we can assume that C does not contain any nonzero, pure, cotorsion R -submodules of rank $< n$.

Let B be a pure R -submodule of C of rank $n-1$, and let $A = C/B$. Then A is a torsion-free R -module of rank one, and we have an exact sequence:

$$(*) \quad 0 \rightarrow B \rightarrow C \rightarrow A \rightarrow 0.$$

Case I. $A \cong Q$.

Let $V = Q \otimes_R C$; then V is the injective envelope of C . We can assume that C_{M_1} and C_{M_2} are contained in V . Since R_{M_1} is a maximal valuation ring we have $\text{Ext}_{R_{M_1}}^1(Q, B_{M_1}) = 0$. Thus if we tensor exact sequence $(*)$ with R_{M_1} it splits, and we obtain $C_{M_1} \cong B_{M_1} \oplus Q_1$, where Q_1 is an R -submodule of V that is isomorphic to Q .

Let $E = Q_1 \cap C$. Since V is the injective envelope of C , we have that E is a nonzero R -submodule of C , and thus E has rank one. Now E is a pure submodule of C . For if x is an element of C and r an element of R such that rx is a nonzero element of E , then $rx \in Q_1$, and so $x \in Q_1 \cap C = E$. Thus C/E is a torsion-free R -module. Hence $C_{M_1}/E_{M_1} \cong (C/E)_{M_1}$ is a torsion-free R_{M_1} -module. Since $E_{M_1} \subset Q_1$, we have $C_{M_1}/E_{M_1} \cong B_{M_1} \oplus Q_1/E_{M_1}$. If $E_{M_1} \neq Q_1$, then Q_1/E_{M_1} is a nonzero torsion R_{M_1} -module, contrary to the fact that C_{M_1}/E_{M_1} is torsion-free. Thus we have $E_{M_1} = Q_1$. Now we also have $E_{M_2} \subset Q_1$, and so $E = E_{M_1} \cap E_{M_2} = Q_1 \cap E_{M_2} = E_{M_2}$. We can not have $E = Q_1$, since $E \subset C$ and C is reduced. Thus by our earlier remarks, E is a pure cotorsion R -submodule of C of rank one. This contradicts our assumption that C has no such submodules, and so in the presence of this condition Case I can not arise.

Case II. A is isomorphic to an ideal I of R .

We have $\text{Ext}_R^1(I, B) \cong \text{Ext}_R^2(R/I, B)$. But $\text{inj. dim.}_R B = 1$, and so $\text{Ext}_R^2(R/I, B) = 0$. Therefore, $\text{Ext}_R^1(I, B) = 0$, and so exact sequence $(*)$ splits. Thus we have $C \cong B \oplus I$, and by the induction hypothesis C is a direct sum of R -modules of rank one.

If $\text{Hom}_R(C, R) \neq 0$, then there exists an exact sequence of the form $(*)$ with A isomorphic to a nonzero ideal I of R . Thus by Case II we can assume without loss of generality that $\text{Hom}_R(C, R) = 0$.

Case III. A is a cotorsion R -module.

By our earlier remarks we can assume that $A = A_{M_2}$ and $A \not\cong Q$. Now by the induction hypothesis we have $B = B_1 \oplus \cdots \oplus B_{n-1}$, where each B_i is an R -submodule of B of rank one. Since B_1 is pure in B and B is pure in C , it follows that B_1 is pure in C . By our assumptions B_1 is not a cotorsion R -module, and $B_1 \not\cong Q$. Therefore, B_1 is isomorphic to an ideal of R . Hence $\text{Hom}_R(B_1, R) \neq 0$. Since $\text{Hom}_R(B_1, R) \subset \text{Hom}_R(B, R)$, we have that $\text{Hom}_R(B, R) \neq 0$. Thus $\text{Hom}_R(B, R)$ is a nonzero, torsion-free R -module. Now we have an exact sequence:

$$0 = \text{Hom}_R(C, R) \rightarrow \text{Hom}_R(B, R) \rightarrow \text{Ext}_R^1(A, R).$$

Therefore, $\text{Ext}_R^1(A, R)$ is not a torsion R -module.

Since R is an h -local domain, we have $R_{M_1} + R_{M_2} = Q$. Of course $R = R_{M_1} \cap R_{M_2}$, and thus we have an exact sequence:

$$0 \rightarrow R \rightarrow R_{M_1} \oplus R_{M_2} \rightarrow Q \rightarrow 0.$$

Now

$$\text{Hom}_R(A, R_{M_1}) = \text{Hom}_R(A_{M_2}, R_{M_1}) = 0$$

by [8, Lemma 8.3]. Since $\text{Hom}_R(A, R) \subset \text{Hom}_R(A, R_{M_1})$, we have $\text{Hom}_R(A, R) = 0$ also. Thus from the last exact sequence we obtain an exact sequence:

$$0 \rightarrow \text{Hom}_R(A, R_{M_2}) \rightarrow \text{Hom}_R(A, Q) \rightarrow \text{Ext}_R^1(A, R) \rightarrow \text{Ext}_R^1(A, R_{M_1} \oplus R_{M_2}).$$

Since $\text{rank } A = 1$, we have $\text{Hom}_R(A, Q) \cong Q$. Now $R_{M_1} \oplus R_{M_2}$ is a cotorsion R -module and has injective dimension one over R . Thus $\text{Ext}_R^1(A, R_{M_1} \oplus R_{M_2}) = 0$. Hence the last exact sequence reduces to the following exact sequence:

$$0 \rightarrow \text{Hom}_R(A, R_{M_2}) \rightarrow Q \rightarrow \text{Ext}_R^1(A, R) \rightarrow 0.$$

Since $A \cong A_{M_2}$ is isomorphic to an ideal of R_{M_2} , we have that $\text{Hom}_R(A, R_{M_2})$ is not zero. Therefore, the above exact sequence shows that $\text{Ext}_R^1(A, R)$ is a torsion R -module. But we have shown that $\text{Ext}_R^1(A, R)$ contains a nonzero, torsion-free R -module. This contradiction shows that Case III can not arise.

Having now considered all possible cases, we have completed the proof of the lemma.

Proof of Theorem 6.1. Assume that R has property D . If R has only one maximal ideal, then R is a maximal valuation ring by Corollary 6.7. Hence we may assume that R has more than one maximal ideal. Let M_1, M_2 be two different maximal ideals of R , and let S be the complement in R of $M_1 \cup M_2$. Then R_S has only two maximal ideals; and by Lemma 6.2 R_S has property D . Thus by Lemma 6.8, R_S is an h -local domain. This proves that every nonzero prime ideal of R is contained in only one maximal ideal of R . Suppose that R has still a third maximal

ideal M_3 . Let T be the complement of $M_1 \cup M_2 \cup M_3$ in R . Then by what we have just proved R_T is an h -local domain with only three maximal ideals. R_T has property D by Lemma 6.2. But this contradicts Proposition 6.9. Thus R is an h -local domain with only two maximal ideals M_1, M_2 and R_{M_1}, R_{M_2} are maximal valuation rings.

Conversely, if R is a maximal valuation ring then Kaplansky has proved a stronger result than property D (see [3, Theorem 12] or [6, Theorem 9]). If R is an h -local domain with exactly two maximal ideals M_1, M_2 and R_{M_1} and R_{M_2} are maximal valuation rings, then R has property D by Lemma 6.10.

This completes the proof of Theorem 6.1.

REMARKS. It is an open question whether or not there can exist an integral domain with two maximal ideals M_1, M_2 such that R_{M_1}, R_{M_2} are maximal valuation rings. Kaplansky has proved in [3, Theorem 12] that if R is a maximal valuation ring, then every torsion-free R -module of countable rank is a direct sum of R -modules of rank one. It is a reasonable conjecture that the converse of this is true; namely, if an integral domain has this property, then it must be a maximal valuation ring.

It is of some interest to find as many kinds of integral domains as possible for which the possession of property D will imply that the domain is a maximal valuation ring. The next proposition and its corollary will provide a partial answer to this question.

An immediate corollary of a result that Kaplansky has proved in [4] is that if R is a Noetherian domain with property D , then R is a complete, discrete, valuation ring. A trivial modification of his proof will prove the next proposition. We will give the full proof of the proposition for, a priori, it is not clear which portion of Kaplansky's proof we will be using.

DEFINITION. We say that an integral domain is *powerful*, if Q is a countably generated R -module. R is *locally powerful*, if R_M is powerful for every maximal ideal M of R .

PROPOSITION 6.11. *Let R be a locally powerful integral domain. Then R has property D if and only if it is a maximal valuation ring.*

Proof. Assume that R has property D . Suppose that R is not a maximal valuation ring. Then by Theorem 6.1, R is an h -local domain with exactly two maximal ideals M_1, M_2 and R_{M_1}, R_{M_2} are maximal valuation rings. By assumption K_{M_1} is a countably generated R_{M_1} -module. Since K_{M_1} is a torsion R_{M_1} -module, it follows from Lemma 2.1 that a set of R_{M_1} generators of K_{M_1} is also a set of R generators of K_{M_1} . Thus K_{M_1} is a countably generated R -module. Similarly K_{M_2} is a countably generated R -module. Since $K = K_{M_1} \oplus K_{M_2}$, K is also a countably generated R -module. Therefore, Q is a countably generated R -module. Thus there exist sequences $\{a_n\}$ and $\{b_n\}$ of elements of R such that $\{1/a_n\}$ generates Q over R and $a_{n+1} = b_{n+1}a_n$.

Now R has infinitely many units. For if we take p to be an element in M_1 that is not in M_2 , and q to be an element in M_2 that is not in M_1 , then $p+q^i$ is a unit for each positive integer i , and these units are all distinct. Thus there exists a unit $u \in R$ such that $u \neq \pm 1$.

Since R is an h -local domain and R_{M_1} , R_{M_2} are complete in the R -topology, we have $H = R_{M_1} \oplus R_{M_2}$, where H is the completion of R . Thus $h = (u, u^{-1})$ is an element of H . Now the R -topology on R is given by the submodules Ra_n . Therefore, we can find a Cauchy sequence $\{d_n\}$ of elements of R such that d_n converges to h in H , and such that $d_n - d_{n+1} \in Ra_n$.

Let V be a two dimensional vector space over Q with basis x and y , and define a sequence of elements $\{w_n\}$ of V by setting $w_n = x + d_n y$. Let N be the R -submodule of V generated by the elements $y, w_0, (1/a_1)w_1, \dots, (1/a_n)w_n, \dots$. Then N is a torsion-free R -module of rank two. Since R has property D , N is decomposable; and thus there exists an idempotent endomorphism φ of N such that $\varphi(N)$ is of rank one. Modifying φ by a scalar, we may assume that $\varphi(x)$ and $\varphi(y)$ are linear combinations of x and y with coefficients in R ; say $\varphi(x) = r_1 x + r_2 y$ and $\varphi(y) = r_3 x + r_4 y$, with the elements r_i in R . The matrix of φ has determinant 0 and trace $\neq 0$; i.e.,

$$(1) \quad r_1 r_4 = r_2 r_3, \quad \text{and} \quad r_1 + r_4 \neq 0.$$

Since $w_n = x + d_n y$ is divisible by a_n in N , so is

$$(2) \quad \varphi(x + d_n y) - (r_1 + d_n r_3)(x + d_n y) = [(r_2 + d_n r_4) - d_n(r_1 + d_n r_3)]y.$$

A computation will show (exactly as in [4]) that $Qy \cap N = Ry$; and thus we have $(r_2 + d_n r_4) - d_n(r_1 + d_n r_3) \equiv 0 \pmod{Ra_n}$. Passing to the limit in H we obtain:

$$(3) \quad (r_2 + hr_4) - h(r_1 + hr_3) = 0.$$

Since $h = (u, u^{-1})$, equation (3) holds with h replaced by either u or u^{-1} .

If $r_1 = 0$, then $r_4 \neq 0$ and either $r_2 = 0$ or $r_3 = 0$. From this we deduce that $u = u^{-1}$, so that $u^2 = 1$, which is a contradiction. Hence $r_1 \neq 0$. We then have by (1):

$$(4) \quad r_2 + hr_4 = (r_2/r_1)(r_1 + hr_3).$$

Comparing (3) and (4) we find that both u and u^{-1} are equal to r_2/r_1 or $-r_3/r_1$. Since $u \neq u^{-1}$, it must be the case that one is r_2/r_1 and the other $-r_3/r_1$. But then $r_2 r_3 = -r_1^2$, whence $r_1 + r_4 = 0$, contradicting (1).

Thus we have that R has only a single maximal ideal, and so is a maximal valuation ring. Of course if R is a maximal valuation ring, then R has property D by Theorem 6.1.

COROLLARY 6.12. *Let R be an integral domain with the descending chain condition on prime ideals. Then R has property D if and only if R is a maximal valuation ring.*

Proof. Assume that R has property D . Let M be a maximal ideal of R . Then by Lemma 6.2, R_M also has property D , and is a local ring. Therefore by Theorem 6.1, R_M is a maximal valuation ring. Hence the prime ideals of R_M are linearly ordered. Since the prime ideals of R_M are in one to one, order preserving, correspondence with the prime ideals of R that are contained in M , R_M also has the descending chain condition on prime ideals. Thus R_M has only finitely many prime ideals. Let P be the minimal, nonzero prime ideal of R , and let s be a nonzero element of P . Let S be the multiplicatively closed subset of R_M consisting of the powers of s . Then every prime ideal of R_M meets S , and hence $(R_M)_S$ has no nonzero prime ideals. Therefore, $(R_M)_S$ is a field, and so $(R_M)_S = Q$, the quotient field of R_M . Thus the quotient field of R_M is a countably generated R_M -module. We have thus proved that R is a locally powerful domain. It then follows from Proposition 6.11 that R is a maximal valuation ring. The converse follows from Theorem 6.1.

COROLLARY 6.13. *Let R be a Noetherian domain. Then R can not have two distinct rank one prime ideals P_1, P_2 such that R_{P_1}, R_{P_2} are complete, discrete, valuation rings.*

Proof. Suppose that R has two such prime ideals P_1 and P_2 . Let S be the complement in R of $P_1 \cup P_2$. Then R_S has the same property as R . Hence we may assume that R has Krull dimension one, and two maximal ideals P_1 and P_2 such that R_{P_1}, R_{P_2} are complete discrete valuation rings. Then R is an h -local domain, and so by Theorem 6.1, R has property D . But R is locally powerful, and thus we have a contradiction by Proposition 6.11. This proves the corollary.

REMARKS. It is an open question whether a Noetherian domain can have even one rank one prime ideal P such that R_P is a complete, discrete, valuation ring without already being a complete, discrete, valuation ring.

7. Indecomposable h -divisible modules. Throughout this section, unless otherwise specified, R will be a local, Noetherian domain of Krull dimension one. We will let M denote the maximal ideal of R , and E the injective envelope of R/M . If A is an R -module, we will denote the dual $\text{Hom}_R(A, E)$ of A by A^* . We have a canonical imbedding of A into A^{**} , and thus $A^* = 0$ if and only if $A = 0$.

Let \hat{R} be the completion of R in the M -adic topology on R . Then $\hat{R} = H$, the completion of R in the R -topology. If I is a nonzero ideal of R and \hat{I} its completion in \hat{R} , then $\hat{I} = \hat{R} \otimes_R I$ and we have $\hat{R}/\hat{I} \cong R/I$. The \hat{R} -injective envelope of \hat{R}/\hat{M} is again E ; and if A is a torsion R -module, then A is an \hat{R} -module and $\text{Hom}_{\hat{R}}(A, E) \cong \text{Hom}_R(A, E) = A^*$. If D is an R -module with D.C.C. (descending chain condition), we have $D \cong \text{Hom}_{\hat{R}}(D^*, E)$ by [5, Corollary 4.3].

If A is an R -module, we will let $t(A)$ denote the torsion submodule of A .

LEMMA 7.1. *Let S be a torsion-free, reduced R -module of rank one (i.e., S is a proper R -submodule of Q). Then the following statements are equivalent:*

- (1) $K \otimes_R S$ is an indecomposable R -module.
- (2) $(K \otimes_R S)^*$ is an indecomposable R -module.
- (3) $t(S^*)$ is an indecomposable R -module.

Proof. (1) \Leftrightarrow (2). By [7, Theorem 1], $K \otimes_R S$ has D.C.C. Hence we have $K \otimes_R S \cong \text{Hom}_{\hat{R}}((K \otimes_R S)^*, E)$. Since $(K \otimes_R S)^*$ is a cotorsion R -module by [8, Proposition 1.3], its R and \hat{R} direct summands coincide by [8, Proposition 5.6]. Thus $K \otimes_R S$ is an indecomposable R -module if and only if $(K \otimes_R S)^*$ is an indecomposable R -module.

(2) \Leftrightarrow (3). By [8, Proposition 1.3], $(K \otimes_R S)^*$ is a torsion-free, cotorsion R -module. Thus by [8, Corollary 3.3], $(K \otimes_R S)^*$ is an indecomposable R -module if and only if $K \otimes_R (K \otimes_R S)^*$ is an indecomposable R -module. It only remains to show that $K \otimes_R (K \otimes_R S)^* \cong t(S^*)$.

Now we have an exact sequence:

$$0 \rightarrow S \rightarrow Q \rightarrow K \otimes_R S \rightarrow 0,$$

and hence an exact sequence:

$$0 \rightarrow (K \otimes_R S)^* \rightarrow Q^* \rightarrow S^* \rightarrow 0.$$

Tensoring this last sequence with K , we obtain the isomorphisms: $\text{Tor}_1^R(K, S^*) \cong K \otimes_R (K \otimes_R S)^*$. By [2, Chapter VII, Proposition 2.2] we have that $t(S^*) \cong \hat{\text{Tor}}_1^R(K, S^*)$. Thus $t(S^*) \cong K \otimes_R (K \otimes_R S)^*$, and the proof of the lemma is complete.

COROLLARY 7.2. *Let I be an ideal of R . Then the following statements are equivalent:*

- (1) $K \otimes_R I$ is an indecomposable R -module.
- (2) \hat{I} is an indecomposable R -module.
- (3) I^* is an indecomposable R -module.

Proof. (1) \Leftrightarrow (2). Since $\hat{I} \cong \hat{R} \otimes_R I$, we have by [8, Proposition 7.2] that $\hat{I} = \text{Ext}_R^1(K, I)$. Thus the equivalence of (1) and (2) follows from [8, Corollary 3.3].

(1) \Leftrightarrow (3). From the exact sequence: $0 \rightarrow I \rightarrow R$, we obtain the exact sequence: $E = R^* \rightarrow I^* \rightarrow 0$. Thus $t(I^*) = I^*$, and so the equivalence of (1) and (3) follows from Lemma 7.1.

LEMMA 7.3. *Assume that \hat{R} has only two prime ideals. Let B be any finitely generated R -submodule of E . Then any divisible R -submodule of E/B is an indecomposable R -module.*

Proof. If X is any R -submodule of E , define X' to be the annihilator of X in \hat{R} . Let $J = B'$; then by [5, Theorem 4.2] we have that $\hat{R}/J \cong B^*$. Since B^* has D.C.C. by [5, Corollary 4.3], it follows that J is an \hat{M} -primary ideal of \hat{R} .

Let D/B be a divisible R -submodule of E/B , where D is an R -submodule of E containing B . Suppose that we have $D/B = D_1/B \oplus D_2/B$, where D_1, D_2 are R -submodules of D containing B . Then $D_1 + D_2 = D$ and $D_1 \cap D_2 = B$. By [5, Theorem 4.2] we then have that $(D_1 \cap D_2)' = D_1' + D_2'$. Therefore, $J = D_1' + D_2'$, and D_1', D_2' are ideals of \hat{R} .

Let P be the other prime ideal of \hat{R} . Suppose that neither D_1' nor D_2' are \hat{M} -primary ideals of \hat{R} . Then $D_1' \subset P$ and $D_2' \subset P$. Hence $J = D_1' + D_2' \subset P$. Since J is an \hat{M} -primary ideal, this is a contradiction. Thus we can assume that D_1' is an \hat{M} -primary ideal of \hat{R} . Therefore, \hat{R}/D_1' has D.C.C.

Now by [5, Theorem 4.2] we have $D_1 \cong (\hat{R}/D_1')^*$, and by [5, Corollary 4.3] we have that $(\hat{R}/D_1')^*$ is a finitely generated \hat{R} -module. Since the \hat{R} -structure and the R -structure of D_1 are the same, D_1 is a finitely generated R -module. Therefore, D_1/B is a finitely generated R -module. But D_1/B is a direct summand of D/B , and hence is a divisible R -module. Thus we have $D_1/B = 0$, and so D/B is an indecomposable R -module.

THEOREM 7.4. *The following statements are equivalent:*

- (1) \hat{R} has exactly two prime ideals.
- (2) $K \otimes_R S$ is an indecomposable R -module for every torsion-free R -module S of rank one.
- (3) $K \otimes_R I$ is an indecomposable R -module for every ideal I of R .

Proof. (1) \Rightarrow (2). Let S be a reduced, torsion-free R -module of rank one (i.e., S is a nonzero proper R -submodule of Q). By [7, Theorem 1], S/MS has D.C.C. Since S/MS is annihilated by M , it has finite length, and thus is finitely generated. Thus there exists a nonzero, finitely generated R -submodule I of S such that $MS + I = S$. Hence we have $M(S/I) = S/I$. If r is any nonzero, nonunit of R , there exists an integer $n > 0$ such that $M^n \subset Rr$. Hence $S/I = r(S/I)$, and so S/I is divisible. We can assume without loss of generality that I is a nonzero ideal of R .

We have an exact sequence:

$$0 \rightarrow (S/I)^* \rightarrow S^* \rightarrow I^* \rightarrow 0.$$

Since S/I is divisible, $(S/I)^*$ is torsion-free. Thus we have $t(S^*) \subset t(I^*)$. Now as we have seen in the proof of Lemma 7.1, we have $t(S^*) \cong K \otimes_R (K \otimes_R S)^*$. Therefore $t(S^*)$ is a divisible submodule of $t(I^*)$.

Let $B = (R/I)^*$; then we have an exact sequence:

$$0 \rightarrow B \rightarrow E \rightarrow I^* \rightarrow 0,$$

and so $I^* \cong E/B$. Thus $t(I^*) = I^*$, and so $t(S^*)$ is isomorphic to a divisible submodule of E/B . Since R/I has D.C.C., B is a finitely generated R -submodule of E by [5, Theorem 4.2]. Therefore, Lemma 7.3 applies and $t(S^*)$ is indecomposable. It follows from Lemma 7.1 that $K \otimes_R S$ is indecomposable.

(2) \Rightarrow (3). Trivial.

(3) \Rightarrow (1). Suppose that \hat{R} has more than two prime ideals. Then we have $n > 1$ distinct prime ideals P_1, \dots, P_n of rank 0 in \hat{R} . Thus in \hat{R} we have $0 = J \cap L$, an irredundant decomposition of 0, where J is a P_1 -primary ideal; and P_2, \dots, P_n belong to the ideal L . Now $J+L$ is an \hat{M} -primary ideal of \hat{R} . For if $P_j \supset J+L$ for some j , $1 \leq j \leq n$, then $P_j \supset J$; and so $P_j = P_1$. But $P_1 \not\supset L$, which is a contradiction. Thus $J+L$ is an \hat{M} -primary ideal of \hat{R} . Let $I = (J+L) \cap R$. Since $J+L$ is \hat{M} -primary, we have $\hat{I} = \hat{R}I = J+L$.

If Y is any ideal of \hat{R} , let $\text{Ann}(Y, E) = \{x \in E \mid Yx = 0\}$. We set $B = \text{Ann}(J, E)$ and $C = \text{Ann}(L, E)$. By [5, Theorem 4.2] we have $B+C = \text{Ann}(J \cap L, E)$ and $B \cap C = \text{Ann}(J+L, E)$. Since $J \cap L = 0$ and $J+L = \hat{I}$, we have $B+C = E$ and $B \cap C = \text{Ann}(\hat{I}, E)$.

Let $A = \text{Ann}(\hat{I}, E) = B \cap C$. Then we have $E/A = (B+C)/A \cong B/A \oplus C/A$. By [5, Theorem 4.2], $E/A \cong \text{Hom}_{\hat{R}}(\hat{I}, E)$; and so we have $(E/A)^* \cong \hat{I}$. Since \hat{I} is an indecomposable R -module by Corollary 7.2, we must have $(B/A)^* = 0$ or $(C/A)^* = 0$. Thus $B = A$, or $C = A$. If $B = A$, then $J = \hat{I}$, and so $J \supset L$. Thus $0 = J \cap L = L$, which contradicts the irredundancy of the decomposition of 0. Similarly we can not have $C = A$. This contradiction shows that \hat{R} has exactly two prime ideals.

REMARKS. It is a matter of classical ideal theory that the following conditions are equivalent (see [9]).

- (1) \hat{R} has exactly two prime ideals.
- (2) Every zero divisor of \hat{R} is nilpotent.
- (3) The zero ideal of \hat{R} is a primary ideal.
- (4) The integral closure of R is a valuation ring.
- (5) Every ring extension of R in Q is a local ring.

Theorem 7.4 provides the solution for a problem raised in [7, p. 573]; namely we have the following corollary.

COROLLARY 7.5. *Let R be a Noetherian integral domain. Then every R -homomorphic image of Q is indecomposable if and only if R is a local domain of Krull dimension one and \hat{R} has exactly two prime ideals.*

Proof. Suppose that every R -homomorphic image of Q is indecomposable. Then by [7, Theorem 2], R is a local domain of Krull dimension one. The corollary now follows directly from Theorem 7.4.

The following theorem can be derived without any difficulty from classical ideal theory (see [9]). We introduce it here mainly because it can now be proved by the

methods of this section, and because it is intimately related to the theorems of this section. It is of interest to compare this theorem with Theorem 4.5.

THEOREM 7.6. *Let R be a Noetherian domain of Krull dimension one. Then \hat{R} is an integral domain if and only if every proper R -submodule of Q is finitely generated.*

Proof. Assume that \hat{R} is an integral domain. It follows that R is then a local domain. Let S be an R -submodule of Q which is not finitely generated. We can assume that $R \subset S$. Since S/R has D.C.C. by [7, Theorem 1], we can further assume that every proper R -submodule of S/R is finitely generated. Now S/R is divisible. For, if not, there exists a nonzero element $r \in R$ such that $r(S/R) \neq S/R$. Thus $rS + R$ is finitely generated, from which it follows that S is finitely generated. This contradiction shows that S/R is divisible.

Let E be the injective envelope of R/M , where M is the maximal ideal of R . Since $(S/R)^* \neq 0$, and since E has no proper divisible submodule by [7, Proposition 2], it follows that there exists an epimorphism of S/R onto E . Let A/R be the kernel of this epimorphism, where A is an R -submodule of E . Then we have an exact sequence:

$$0 \rightarrow A \rightarrow S \rightarrow E \rightarrow 0.$$

Tensoring this exact sequence with K we obtain an exact sequence:

$$0 \rightarrow \text{Tor}_1^R(K, E) \rightarrow K \otimes_R A \rightarrow K \otimes_R S \rightarrow 0.$$

Since $\text{Tor}_1^R(K, E) \cong E$, this sequence yields the isomorphism: $K \otimes_R A \cong E \oplus K \otimes_R S$. But $K \otimes_R A$ is indecomposable by Theorem 7.4, and so $K \otimes_R S = 0$. This means that $S = Q$. Therefore we have proved that every proper R -submodule of Q is finitely generated.

Conversely, if every proper R -submodule of Q is finitely generated, then \hat{R} is an integral domain by Theorem 4.5.

BIBLIOGRAPHY

1. H. Bass, *Projective modules over algebras*, Ann. of Math. **73** (1961), 532–542.
2. H. Cartan and S. Eilenberg, *Homological algebra*, Princeton Univ. Press, Princeton, N. J., 1956.
3. I. Kaplansky, *Modules over Dedekind rings and valuation rings*, Trans. Amer. Math. Soc. **72** (1952), 327–340.
4. ———, *Decomposability of modules*, Proc. Amer. Math. Soc. **13** (1962), 532–535.
5. E. Matlis, *Injective modules over Noetherian rings*, Pacific J. Math. **8** (1958), 511–528.
6. ———, *Injective modules over Prufer rings*, Nagoya Math. J. **15** (1959), 57–69.
7. ———, *Some properties of Noetherian domains of dimension 1*, Canad. J. Math. **13** (1961), 569–586.
8. ———, *Cotorsion modules*, Mem. Amer. Math. Soc. No. 49, (1964).
9. D. G. Northcott, *General theory of one dimensional local rings*, Proc. Glasgow Math. Assoc. **2** (1956), 159–169.

NORTHWESTERN UNIVERSITY,
EVANSTON, ILLINOIS