

## REPRESENTATIONS FOR TRANSFORMATIONS CONTINUOUS IN THE $BV$ NORM

BY

J. R. EDWARDS AND S. G. WAYMENT

**Abstract.** Riemann and Lebesgue-type integrations can be employed to represent operators on normed function spaces whose norms are not stronger than sup-norm by  $T(f) = \int f d\mu$  where  $\mu$  is determined by the action of  $T$  on the simple functions. The real-valued absolutely continuous functions on  $[0, 1]$  are not in the closure of the simple functions in the  $BV$  norm, and hence such an integral representation of an operator is not obtainable. In this paper the authors develop a  $v$ -integral whose structure depends on fundamental functions different than simple functions. This integral is as computable as the Riemann integral. By using these fundamental functions, the authors are able to obtain a direct, analytic representation of the linear functionals on  $AC$  which are continuous in the  $BV$  norm in terms of the  $v$ -integral. Further, the  $v$ -integral gives a characterization of the dual of  $AC$  in terms of the space of fundamentally bounded set functions which are convex with respect to length. This space is isometrically isomorphically identified with the space of Lipschitz functions anchored at zero with the norm given by the Lipschitz constant, which in turn is isometrically isomorphic to  $L^\infty$ . Hence a natural identification exists between the classical representation and the one given in this paper. The results are extended to the vector setting.

**1. Introduction.** Riemann and Lebesgue integration are intimately related to the sup-norm topology on the function space and to function approximation in the sup-norm topology by step functions and simple functions respectively, where a simple function is a finite sum of scalars times characteristic functions of Lebesgue measurable sets. This close relationship is perpetuated in various abstractions of the integration process, including those given in [2], [3], [4], [11], [12], and [13]. In [3] the authors demonstrated that if a space of functions is contained in the sup-norm closure of a collection of simple functions, then any transformation  $T(f)$  which is continuous in a norm weaker than sup-norm has an integral representation. In 1909 F. Riesz [10] showed that the dual of  $C[0, 1]$  is  $BV_0^1$  (the functions of bounded variation with  $\|f\|_{BV} = V_0^1 f$ ) via his celebrated Stieltjes integral representation theorem. Obtaining a representation for the functionals on  $BV_0^1$  has been a problem

---

Presented to the Society, January 24, 1970 under the title *A representation for functionals continuous in the  $BV$  norm*; received by the editors November 28, 1969 and, in revised form, March 25, 1970.

*AMS 1969 subject classifications.* Primary 2645, 2649, 2825, 2850; Secondary 2646.

*Key words and phrases.* Generalized Stieltjes-type integral,  $v$ -integral, integral representation, convex with respect to length, fundamentally bounded, Lipschitz function, dual space, vector-valued functions, absolute continuity, bounded variation, semiabsolute continuity, semibounded variation.

Copyright © 1971, American Mathematical Society

of interest since that time. T. H. Hildebrandt [6] has given a representation for the continuous functionals on the class  $BV_0^1$  in terms of a Stieltjes-type integral involving functions defined on a space whose elements are collections of non-overlapping intervals. Using the notation introduced in [6] one can obtain a characterization for the continuous functionals on the absolutely continuous functions as follows: For  $I_1$  and  $I_2$  in  $\mathfrak{I}$ , define the distance between  $I_1$  and  $I_2$  to be  $d(I_1, I_2) = \mu(I_1 \Delta I_2)$ , the Lebesgue measure of the symmetric difference. Then  $\beta$  is continuous on the metric space  $\mathfrak{I}$  if and only if  $\alpha$  is absolutely continuous. Since any metric space is normal, it follows from Theorem 2 in [1, p. 262] that the continuous functionals are characterized by integrals with respect to regular bounded additive measures on the field generated by the closed sets in  $\mathfrak{I}$ . We remark that one can obtain similar representations for vector-valued functions using the results in [4]. Dunford-Schwartz indicate [1, p. 374] that such characterizations are not entirely satisfactory. Hildebrandt also has given [5] a representation with respect to a more conventional integral for the functionals on  $BV_0^1$  which are continuous in the weak topology, and indicates that by his method a representation can be obtained for the functionals on  $BV_0^1$  continuous in the sup-norm topology. Such a representation is also obtainable from the main theorem in [3]. Furthermore, the work in [3] seems to indicate that the historical type of integral representation theorem cannot be obtained for the functionals on a function space with norm stronger than sup-norm, or at least in a function space with a norm in which the simple functions are not dense. The space  $BV_0^1$  with the variational norm is such a space since for any continuous  $BV_0^1$  function  $f$  and any simple function  $s$  it follows that  $\|f - s\|_{BV} \geq \max \{V_0^1 f, V_0^1 s\}$ .

The technique for obtaining integral type representation theorems has been to consider the action of a functional  $T$  on a dense subset of functions (historically the characteristic functions of a certain class of sets) from which a set function is obtained. Integration of a function  $f$  with respect to this set function then gives  $T(f)$ . It is well known that the step functions are dense in the set  $S$  of saltus functions [8, p. 219] or break functions [6] of bounded variation in the  $BV_0^1$  norm and we shall show that the closure of the polygonal functions in the  $BV_0^1$  norm is precisely the set  $AC$  of absolutely continuous functions. Identification of an appropriate decomposition of a polygonal function into elementary functions leads to the notion of a generalized Stieltjes-type integral of a set function  $K$  with respect to a point function  $f$ . For brevity we shall denote the integral by  $v \int K \cdot df$ , and shall refer to it as the  $v$ -integral. The characterization of  $(S \oplus AC)^*$  is then made in terms of the  $v$ -integral and a class of set functions.

Dunford-Schwartz [1, p. 343] give a characterization of  $AC^*$  in terms of a Lebesgue integral of the product of the derivative  $f'$  of  $f$  and a function  $g$  from  $L^\infty$ . The  $v$ -integral representation given in this paper enjoys a computability not shared by the above mentioned representation and requires no knowledge of Lebesgue integration. The space of Lipschitz functions on  $[0, 1]$  are shown to be isometric

and isomorphic to the collection of set functions which characterize  $AC^*$  and then the identification of the  $g$  in the Dunford-Schwartz representation with the  $K$  in the representation given herein is made. In this process we give a representation of  $AC^*$  in terms of an integral involving a Lipschitz function (which is a generalization of the Stieltjes integral and also is a generalization of the integral given by Lane [7]).

Consider the functional  $T(f)=f(1)-f(0)$  on  $BV_0^1$ . If  $T$  is restricted to  $AC$ , then the representation given in [1] is  $T(f)=L \int_0^1 f'g \, d\mu$  (we suppose  $f(0)=0$  for simplicity). The  $g \in L^\infty$  associated with  $T$  is thus in the equivalence class generated by  $g \equiv 1$ . If  $C$  represents the Cantor function, then  $C \in BV_0^1$  but  $C \notin AC$ . Since  $f'g$  is integrable (Lebesgue) for  $f \in BV_0^1$  and  $g \in L^\infty$ , we can consider the action of the integral representation on  $C$  and find  $T(C)=0$ . In fact, for each  $T$  on  $AC$ , the integral representation in [1] gives  $L \int_0^1 C'g = 0$ . However, the representation given herein for  $T$  which maps  $f \in AC$  to  $f(1)-f(0)$  extends immediately to all of  $BV_0^1$  and gives for each  $f \in BV_0^1$  the value  $f(1)-f(0)$ . We give in the last section of this paper a more complete discussion of this issue and a sufficient condition on the set function  $K$  (or on the associated Lipschitz function  $L$ ) so that  $v \int K \cdot df$  generates a continuous map on  $BV_0^1$ .

**2. A characterization of  $AC$ .** In this section we identify the desired dense set of functions and establish that an appropriate limit with respect to the net of partitions exists. Since  $f-g=A$  for some constant  $A$  implies  $\|f-g\|_{BV}=0$ , we shall always choose the representative element  $f$  from each class in  $BV_0^1$  so that  $f(0)=0$ .

**2.1. THEOREM.** *The space  $AC$  of absolutely continuous functions with the  $BV_0^1$  norm is complete and is the closure of the set  $P$  of polygonal functions in the  $BV_0^1$  norm.*

**Proof.** If  $f$  is absolutely continuous, then  $f'$  is an  $L^1$  function. Hence there exists a sequence  $s_i$  of step functions converging to  $f'$  in the  $L^1$  norm. Let  $p_i(x) = \int_0^x s_i \, d\mu$ . Then  $p_i$  is polygonal and

$$\|f-p_i\|_{BV} = L \int |(f-p_i)'| \, d\mu = \|f'-s_i\|_{L^1}.$$

Consequently  $\langle p_i \rangle$  converges to  $f$  in the  $BV_0^1$  norm.

Conversely if  $\langle p_i \rangle$  is a Cauchy sequence of polygonal functions which is Cauchy in the  $BV_0^1$  norm, then  $\langle p_i' \rangle = \langle s_i \rangle$  is a sequence of  $L^1$  functions Cauchy in the  $L^1$  norm. Thus there exists an  $L^1$  function  $f'$  such that  $\langle s_i \rangle \xrightarrow{L^1} f'$ , or equivalently  $\langle p_i \rangle \xrightarrow{BV} f$  where  $f$  is the indefinite integral  $\int_0^x f' \, d\mu$ .

It is possible to show that Cauchy sequences  $\langle p_i \rangle$  of polygonal functions converge to an  $AC$  function with a calculus type argument which does not require any knowledge of  $L^1$ . Since such a proof gives insight and shows that the sequence  $\langle p_i \rangle$  is ultimately equi-absolutely continuous we next outline such a proof.

Since  $\langle p_i \rangle$  is Cauchy in the  $BV_0^1$  norm, it is Cauchy in the weaker sup-norm and the pointwise limit  $f$  is continuous. If  $\sigma$  is any partition of  $[0, 1]$ , then

$$(1) \quad \sum_{\sigma} |f(x_{i+1}) - f(x_i)| \leq \sum_{\sigma} [|f(x_{i+1}) - p_n(x_{i+1})| + |p_n(x_{i+1}) - p_n(x_i)| + |p_n(x_i) - f(x_i)|].$$

The sum of the first and third terms on the right of (1) can be made small by sufficiently large choice of  $n$ . The sum of the middle terms on the right of (1) is less than  $\|p_n\|_{BV}$ . But  $\sup_n \|p_n\|_{BV}$  is finite and hence the left side of (1) is bounded independent of  $\sigma$ , and is an increasing function of  $\sigma$ . Thus  $f \in BV$ .

Choose  $M$  so that  $n, m > M$  implies  $\|p_n - p_m\|_{BV} < \varepsilon$  and let  $\Delta_i f = f(x_{i+1}) - f(x_i)$ . Then

$$(2) \quad \sum_{\sigma} |\Delta_i(f - p_m)| \leq \sum_{\sigma} |\Delta_i(f - p_n)| + \sum_{\sigma} |\Delta_i(p_n - p_m)|,$$

and hence

$$\begin{aligned} \sum_{\sigma} |\Delta_i(f - p_m)| &\leq \limsup_n \sum_{\sigma} |\Delta_i(f - p_n)| + \limsup_n \sum_{\sigma} |\Delta_i(p_n - p_m)| \\ &\leq 0 + \limsup_n \|p_n - p_m\|_{BV} \leq \varepsilon, \end{aligned}$$

and hence  $\langle p_n \rangle \xrightarrow{BV} f$ .

Finally we show that  $f$  is absolutely continuous. Given  $\varepsilon > 0$ , choose  $M$  so that  $n, m \geq M$  implies  $\|p_n - p_m\|_{BV} < \varepsilon/3$ . For  $p_M$  there is a  $\delta$  such that if  $\sum |y_i - x_i| < \delta$  then  $\sum |p_M(y_i) - p_M(x_i)| < \varepsilon/3$ . Hence for any finite collection  $\{[x_i, y_i]\}$  of intervals such that  $\sum |y_i - x_i| < \delta$  we have

$$\begin{aligned} \sum |f(y_i) - f(x_i)| &= \sum |f(y_i) - p_n(y_i) + p_n(y_i) - p_M(y_i) + p_M(y_i) \\ &\quad - p_M(x_i) + p_M(x_i) - p_n(x_i) + p_n(x_i) - f(x_i)| \\ (3) \quad &\leq \sum [|f(y_i) - p_n(y_i)| + |p_n(y_i) - p_M(y_i) + p_M(x_i) - p_n(x_i)| \\ &\quad + |p_M(y_i) - p_M(x_i)| + |p_n(x_i) - f(x_i)|]. \end{aligned}$$

Except for the peculiar grouping of the second term on the right of (3), this is a straightforward decomposition. The sum of all first and fourth terms can be made less than  $\varepsilon/3$  for sufficiently large  $n$ . The sum of the second terms on the right of (3) is dominated by  $\|p_n - p_M\|_{BV} < \varepsilon/3$  and the sum of the third terms on the right of (3) is less than  $\varepsilon/3$  because of the absolute continuity of  $p_M$ . This shows that  $f \in AC$  and it is also clear that if  $\langle p_n \rangle$  is a Cauchy sequence of  $AC$  functions in the  $BV_0^1$  norm, then for  $\varepsilon > 0$  there exists  $\delta$  and  $N$  such that  $\sum |y_i - x_i| < \delta$  and  $n > N$  implies  $\sum |p_n(y_i) - p_n(x_i)| < \varepsilon$ .

**2.2. THEOREM.** *Let  $pf_{\sigma}$  represent the polygonal function with corners at precisely the points  $(x_i, f(x_i))$  for  $x_i \in \sigma$ . Then  $f \in AC$  implies  $\lim_{\sigma} pf_{\sigma} = f$  in the  $BV_0^1$  norm.*

**Proof.** Let  $q_n$  be a sequence of polygonal functions converging to  $f$  in the  $BV_0^1$  norm. Let  $\sigma$  represent the values of  $x$  at which  $q_n$  has corners. Then  $\|pf_{\sigma} - q_n\|_{BV}$

$\leq \|f - q_n\|$  since  $\sum_{\sigma} |\Delta_i(pf_{\sigma} - q_n)|$  is precisely the variation of  $pf_{\sigma} - q_n$  and is also an approximating sum to the value of  $\|f - q_n\|_{BV}$  which is obtained as the limit with respect to subdivisions of a nondecreasing function of subdivisions. Thus if  $\|f - q_n\|_{BV} < \varepsilon$  we conclude that for  $\sigma'$  beyond  $\sigma$ ,

$$\|f - pf_{\sigma'}\|_{BV} \leq \|f - q_n\|_{BV} + \|q_n - pf_{\sigma'}\|_{BV} \leq \|f - q_n\|_{BV} + \|q_n - f\|_{BV} < 2\varepsilon.$$

**2.3. THEOREM.** Let  $|\sigma|$  represent  $\max \{|x_{i+1} - x_i|\}$  for  $x_{i+1}, x_i \in \sigma$ . Then  $f \in AC$  implies  $\lim_{|\sigma| \rightarrow 0} pf_{\sigma} = f$  in  $BV_0^1$  norm.

**Proof.** Let  $\sigma_1$  be a partition such that  $\sigma_2$  finer than  $\sigma_1$  implies  $\|pf_{\sigma_2} - pf_{\sigma_1}\|_{BV} < \varepsilon/3$ . Let  $\delta_1$  be a number such that  $\sum |y_i - x_i| < \delta_1$  implies  $\sum |f(y_i) - f(x_i)| < \varepsilon/3$ . Let  $N$  be the number of points in  $\sigma_1$  and choose  $\sigma'$  to be any partition so that  $|\sigma'| = \delta < \delta_1/2N$ . Let  $\sigma = \sigma' * \sigma_1$ , that is,  $\sigma$  is the union of all points  $x$  such that  $x \in \sigma'$  or  $x \in \sigma_1$ . Since  $\sigma$  is finer than  $\sigma_1$ , we have  $\|pf_{\sigma} - f\| < \varepsilon/3$ . Let  $A = \{[x_i, x_{i+1}] \mid x_i, x_{i+1} \in \sigma \text{ and either } x_i \in \sigma_1 \text{ or } x_{i+1} \in \sigma_1\}$ . Then if  $[z_i, z_{i+1}] \in A$  we conclude that  $|z_{i+1} - z_i| < \delta < \delta_1/2N$ , and hence  $\sum_A |z_{i+1} - z_i| < \delta_1$ . Since  $\|pf_{\sigma} - pf_{\sigma'}\|_{BV} = \sum_A \|pf_{\sigma} - pf_{\sigma'}\|_{BV} \leq \sum_A \|pf_{\sigma}\|_{BV} + \sum_A \|pf_{\sigma'}\|_{BV} < \varepsilon/3 + \varepsilon/3$ , we have

$$\|f - pf_{\sigma'}\|_{BV} \leq \|f - pf_{\sigma}\|_{BV} + \|pf_{\sigma} - pf_{\sigma'}\|_{BV} < \varepsilon$$

for any  $\sigma'$  with  $|\sigma'| < \delta$ .

**3. The integral.** In this section we define and discuss the concepts relevant to the integral involved in the representation theorems.

**3.1. DEFINITION.** A half open interval  $(a, b] \subset (0, 1]$  will be called a fundamental set.

**3.2. DEFINITION.** Let  $K$  be a set function defined on the fundamental sets. The generalized Stieltjes-type integral of the set function  $K$  with respect to  $f$  (or briefly, the  $v$ -integral), denoted by  $v \int K \cdot df$  is defined to be

$$v \int K \cdot df = \lim_{\sigma} \sum_{\sigma} (f(x_{i+1}) - f(x_i)) K((x_i, x_{i+1})).$$

**3.3. THEOREM.** If  $f$  is a continuous function satisfying  $f(0) = 0$ , then the Riemann integral  $R \int f dx = v \int K \cdot df$  where  $K((a, b]) = 1 - b$ .

**Proof.** We have  $v \int K \cdot df = \lim_{\sigma} \sum_{\sigma} (f(x_{i+1}) - f(x_i))(1 - x_{i+1}) = s \int (1 - x) df$ , where the last integral is the Stieltjes integral.

**3.4. DEFINITION.** The set function  $K$  is said to be convex relative to length provided that if the fundamental set  $H$  is the union of a finite number of fundamental sets  $\{H_i\}_{i=1}^n$ , then  $K(H) = \sum_{i=1}^n \lambda_i K(H_i)$  where  $\lambda_i$  is the ratio of the length of  $H_i$  to the length of  $H$ .

**3.5. DEFINITION.** The set function  $K$  is said to be fundamentally bounded if  $|K(H)| \leq B$  for some  $B$  and all fundamental sets  $H$ . If  $WK$  is the least bound, then we shall say  $WK$  is the fundamental bound for  $K$ .

3.6. EXAMPLE. Define  $K((a, b)) = 0$  if  $b \leq \frac{1}{2}$  or if  $a > \frac{1}{2}$  and define  $K((\frac{1}{2}, b)) = 1/(b - \frac{1}{2})$ . Finally for  $a < \frac{1}{2} < b$  define  $K((a, b)) = [(b - \frac{1}{2})/(b - a)]K((\frac{1}{2}, b)) = 1/(b - a)$ . Then  $K$  is a convex set function which is not fundamentally bounded and for each function whose derivative exists from the right at  $\{\frac{1}{2}\}$  we have  $v \int K \cdot df = f'(\frac{1}{2}^+)$ . Thus  $v \int K \cdot df$  generates a continuous linear functional on  $C_1[0, 1]$ , in the  $C_1$  norm. We remark that Theorem 3.3 contains an example of a fundamentally bounded, but not convex, set function for which  $v \int K \cdot df$  exists for a certain set of functions.

3.7. DEFINITION. Let  $H = (a, b]$  be a fundamental set. We define the fundamental function

$$\begin{aligned}\Psi_H(t) &= 0 && \text{if } t \leq a, \\ &= (t - a)/(b - a) && \text{if } a < t < b, \\ &= 1 && \text{if } t \geq b.\end{aligned}$$

3.8. REMARK. The fundamental functions form a basis for the linear space of polygonal functions which are zero at zero.

#### 4. The integral representation for $AC^*$ .

4.1. LEMMA. Let  $K$  denote a convex set function. Suppose  $p$  is a polygonal function and that  $\sigma = \{x_i\}$  is the partition determined by the corners of  $p$ . If  $\sigma' = \{x_{ij}\}$ ,  $x_i < x_{ij} < x_{i+1}$  is a refinement of  $\sigma$ , then

$$\sum_{\sigma} [p(x_{i+1}) - p(x_i)]K(H_i) = \sum_{\sigma'} [p(x_{ij+1}) - p(x_{ij})]K(H_{ij}).$$

Hence, it follows that  $v \int K \cdot dp = \sum_{\sigma} \Delta_i p K(H_i)$ .

**Proof.** For each  $i$  we show that

$$[p(x_{i+1}) - p(x_i)]K(H_i) = \sum_j [p(x_{ij+1}) - p(x_{ij})]K(H_{ij}),$$

from which the lemma follows. First observe that  $p$  has constant slope on the interval  $[x_i, x_{i+1}]$ . Hence, for each  $j$  we have

$$[p(x_{ij+1}) - p(x_{ij})]/[p(x_{i+1}) - p(x_i)] = [x_{ij+1} - x_{ij}]/[x_{i+1} - x_i].$$

Since  $K$  is convex, then

$$\begin{aligned}K(H_i) &= \sum_j ([x_{ij+1} - x_{ij}]/[x_{i+1} - x_i])K(H_{ij}) \\ &= \sum_j ([p(x_{ij+1}) - p(x_{ij})]/[p(x_{i+1}) - p(x_i)])K(H_{ij}),\end{aligned}$$

from which the above desired equality follows.

4.2. THEOREM.  $T$  is an element of  $AC^*$  (the dual of  $AC$  with  $BV$  norm) if and only if there exists a unique fundamentally bounded set function  $K$  which is convex with respect to length such that for  $f \in AC$ ,  $T(f) = v \int K \cdot df$ . Furthermore if  $WK$  is the fundamental bound, then  $\|T\| = WK$ .

**Proof.** For each fundamental set  $H \subset (0, 1]$  let  $\Psi_H$  denote the corresponding elementary function and define the set function  $K$  by  $K(H) = T(\Psi_H)$ . If  $f \in AC$ , then from 2.2 we have that  $pf_\sigma$  converges to  $f$  in the  $BV$  norm. Hence  $T(f) = \lim_\sigma T(pf_\sigma) = \lim_\sigma T(\sum_\sigma \Delta_i pf_\sigma \Psi_{H_i}) = \lim_\sigma \sum_\sigma \Delta_i f K(H_i) = v \int K \cdot df$ . Further,

$$\|T\| = \sup_{\|f\|_{BV}=1} |T(f)| \geq \sup_{\Psi_H} |T(\Psi_H)| = \sup |K(H)| = WK,$$

and

$$\begin{aligned} \|T\| &= \sup_{\|f\|_{BV}=1} |T(f)| = \sup_{\|f\|_{BV}=1} \left| v \int K \cdot df \right| = \sup_{\|f\|_{BV}=1} \left| \lim_\sigma \sum_\sigma \Delta_i f K(H_i) \right| \\ &\leq \sup_{\|f\|_{BV}=1} \lim_\sigma \sum_\sigma |\Delta_i f| |K(H_i)| \leq \sup_{\|f\|_{BV}=1} \lim_\sigma \sum_\sigma |\Delta_i f| WK \\ &\leq \sup_{\|f\|_{BV}=1} \|f\|_{BV} WK = WK. \end{aligned}$$

The uniqueness of  $K$  follows from 4.1.

Conversely, if  $K$  is a convex fundamentally bounded set function then from the definition of the integral we must show  $\lim_\sigma \sum_\sigma \Delta_i f K(H_i)$  exists. From 2.1 we see  $\sum_\sigma \Delta_i f K(H_i) = \sum_\sigma \Delta_i pf_\sigma K(H_i) = v \int K \cdot dpf_\sigma$ . Since

$$\left| v \int K \cdot dpf_\sigma - v \int K \cdot dpf_{\sigma'} \right| = \left| \sum_{\sigma, \sigma'} \Delta_i (pf_\sigma - pf_{\sigma'}) K(H_i) \right| \leq \|pf_\sigma - pf_{\sigma'}\|_{BV} WK$$

we conclude that  $v \int K \cdot dpf_\sigma$  is Cauchy and  $v \int K \cdot df$  exists. Furthermore, it is immediate that  $|v \int K \cdot df| \leq \|f\|_{BV} WK$  and hence that if  $\langle f_\alpha \rangle$  converges to  $f$  in the  $BV$  norm, then  $v \int K \cdot df_\alpha$  converges to  $v \int K \cdot df$ . Hence  $T(f) = v \int K \cdot df$  is a continuous linear operator (the linearity of the integral follows from considering approximations to the integral).

**4.3. REMARK.** We note that one does not need convexity of  $K$  to generate a linear functional from  $v \int K \cdot df$ . This is somewhat surprising until one realizes that in case of the Lebesgue integral one does not require additivity of a measure  $\mu$  to generate a linear functional  $L \int f d\mu$ . The point is, once one has generated a linear functional from a bounded set function, then there exists an additive (in the case of Lebesgue) or a convex (in Theorem 4.2) set function which generates the same transformation. Thus in Theorem 3.3 in which the nonconvex set function  $K_1((a, b)) = 1 - b$  generated the transformation  $R \int f dx$ , we see that the (unique) convex set function  $K_2((a, b)) = 1 - (a + b)/2$  generates the same transformation via  $v \int K_2 \cdot df$ .

**4.4. REMARK.** From 2.3 we conclude that Definition 3.2, Theorem 3.3, Lemma 4.1 and Theorem 4.2 can be restated in terms of  $\lim |\sigma| \rightarrow 0$  rather than in terms of  $\lim_\sigma$  which is a limit with respect to the net of partitions. Hence the  $v$ -integral is as computable as the Riemann integral.

**5. The integral representation for  $S^*$ .** If  $s \in S$ , that is, if  $s$  is a saltus function [8, p. 205] or break function [6], then it is easy to see that there exists a sequence

$\langle p_n \rangle$  of step functions converging to  $s$  in the  $BV$  norm. Hence the representation possible for  $S^*$  is in terms of more conventional integrals which resemble Stieltjes integrals and are usually called left Cauchy and right Cauchy integrals. We shall give only a brief sketch of the development for comparison with the  $v$ -integral of §§3 and 4, calling attention to [6] for further development. We again assume  $s(0)=0$  for each  $s \in S$ .

5.1. LEMMA. *If  $s \in S$  then  $s = s_r + s_l$  where  $s_r$  is continuous from the right and  $s_l$  is continuous from the left at each point of  $[0, 1]$ .*

5.2. LEMMA. *If  $s_r$  and  $s_l$  are as in Lemma 5.1, then there exists sequences  $\langle p_n^r \rangle$  and  $\langle p_n^l \rangle$  of step functions continuous from the right and left respectively which converge in the  $BV$  norm to  $s_r$  and  $s_l$  respectively.*

5.3. DEFINITION. If  $\Phi_x^r(t)=0$  for  $t < x$  and  $\Phi_x^r(t)=1$  for  $t \geq x$  then  $\Phi_x^r$  is said to be a right elementary function. A function  $\Phi_x^l(t)$  is defined to be a left elementary function if  $\Phi_x^l(t)=0$  for  $t \leq x$  and  $\Phi_x^l(t)=1$  for  $t > x$ .

5.4. LEMMA. *The right and left elementary functions form a basis for the linear spaces of right and left continuous step functions respectively.*

5.5. DEFINITION. A set of points consisting of a single point will be defined to be an elementary set.

5.6. DEFINITION. The right Cauchy integral of a function  $f$  with respect to the elementary function (point function)  $K$  is defined to be

$$rv \int K \cdot df = \lim_{\sigma} \sum_{\sigma} [f(x_{i+1}) - f(x_i)] K(x_i)$$

provided the limit with respect to the net of partitions exists. The left Cauchy integral  $lv \int K \cdot df$  is defined to be  $\lim_{\sigma} \sum_{\sigma} [f(x_{i+1}) - f(x_i)] K(x_{i+1})$ .

5.7. THEOREM.  *$T$  is an element of  $S^*$  (the dual of  $S$  with  $BV$  norm) if and only if there exist two unique bounded point functions  $K_l$  and  $K_r$  such that for  $s \in S$ ,  $T(s) = rv \int K_r ds_r + lv \int K_l ds_l$ . Furthermore if  $WK = \max \{ \sup K_l(t), \sup K_r(t) \}$ , then  $\|T\| = WK$ .*

**Proof.** Since  $s = s_r + s_l$  and there exist sequences  $\langle p_n^r \rangle$  and  $\langle p_n^l \rangle$  converging to  $s_r$  and  $s_l$  respectively, we have

$$\begin{aligned} T(s) &= \lim_n T(p_n^r) + \lim_n T(p_n^l) \\ &= \lim_n \sum_{\sigma_n^r} [p_n^r(x_{i+1}) - p_n^r(x_i)] T(\Phi_{x_{i+1}}^r) + \lim_n \sum_{\sigma_n^l} [p_n^l(x_{i+1}) - p_n^l(x_i)] T(\Phi_{x_i}^l). \end{aligned}$$

If we define  $K_l(t) = T(\Phi_t^l)$  and  $K_r(t) = T(\Phi_t^r)$ , then  $T(s) = \lim_n T(p_n^r) + \lim_n T(p_n^l) = lv \int K_l ds_l + rv \int K_r ds_r$ . The remainder of the proof is straightforward.



5.8. REMARK. A more revealing representation for  $T(s)$  can be obtained by representing  $s(t)$  as follows:

$$\begin{aligned} s(t) &= \sum_x ((s(x+0) - s(x-0))\phi_x^1(t) + (s(x) - s(x-0))\delta(x, t)) \\ &= \sum_x a(x)\phi_x^1(t) + \sum_x b(x)\delta(x, t) \end{aligned}$$

where  $a(x)$  and  $b(x)$  are denumerably nonzero and  $\delta(x, t)$  is the Kronecker  $\delta$  function. T. H. Hildebrandt [5] has used such a representation to represent functionals continuous in the weak topology and observes (and we use his notation) that if  $g \in BV$ , then  $\lim_\sigma g_\sigma = g$  in the sup-norm topology. We remark further that if  $s \in BV$  is a saltus or break function, then  $\lim_\sigma s_\sigma = s$  in the  $BV$  norm, and consequently we refer the interested reader to the development in [5] for a more revealing representation theorem than that given in this paper for  $S^*$ .

If we denote by  $B$  the set of all bounded point functions and if we denote by  $\Omega$  the set of all fundamentally bounded convex set functions with norms given by the fundamental bound, then  $(S \oplus AC)^* = S^* \oplus AC^* = (B \oplus B) \oplus \Omega$ . If one desires to consider functions which are not zero at zero then one must form the direct sum of the preceding characterizations with  $R$ , where  $R$  denoted the real numbers.

We now return to a calculus for  $v \int K \cdot df$  defined on  $f \in AC$  and make several identifications.

6. **Some consequences of the calculus of the  $v$ -integral.** In this section we let  $\mathcal{L}$  denote the normed space of Lipschitz functions on  $[0, 1]$  which are zero at zero and such that  $\|f\|_{\mathcal{L}}$  is given by the Lipschitz constant.

6.1. DEFINITION. Suppose  $K \in \Omega$ . Then for a  $K$ - $v$ -integrable function  $f$  define  $v \int_a^b K \cdot df = v \int K \cdot df_{ab}$  where  $f_{ab}$  is defined by  $f_{ab}(x) = f(a)$  for  $x \leq a$ ,  $f_{ab}(x) = f(x)$  for  $a \leq x \leq b$  and  $f_{ab}(x) = f(b)$  for  $x \geq b$ .

6.2. THEOREM. Suppose  $f \in AC$  and  $K \in \Omega$ . Then  $v \int_a^b K \cdot df$  exists for  $a, b \in [0, 1]$ . Furthermore, for  $a < b < c$ ,  $v \int_a^c K \cdot df = v \int_a^b K \cdot df + v \int_b^c K \cdot df$ .

**Proof.** Since  $f \in AC$  then  $f_{ab} \in AC$ . Hence the first result follows from 4.2. Suppose  $\sigma$  is finer than the partition  $\{0, a, b, c, 1\}$ . Then

$$\sum_{\sigma} \Delta_i f K(x_i, x_{i+1}] = \sum_a^b \Delta_i f K(x_i, x_{i+1}] + \sum_b^c \Delta_i f K(x_i, x_{i+1}].$$

Therefore, the desired result follows by taking the limit over  $\sigma$ .

6.3. THEOREM. Suppose  $f \in AC$  and  $K \in \Omega$ . Define  $F(x) = v \int_0^x K \cdot df$ . Then  $F(x)$  is  $AC$ .

**Proof.** Suppose  $\varepsilon > 0$ . Since  $f \in AC$  then there is a  $\delta > 0$  such that if  $\{(x_i, x_{i+1})\}$  is a collection of intervals such that  $\sum \Delta x_i < \delta$ , then  $\sum |\Delta_i f| < \varepsilon/WK$ . Then

$$\begin{aligned} \sum |\Delta_i f| &= \sum \left| \int_0^{x_{i+1}} K \cdot df - \int_0^{x_i} K \cdot df \right| = \sum \left| \int_{x_i}^{x_{i+1}} K \cdot df \right| \\ &\leq \sum WK \|f_{x_i x_{i+1}}\|_{BV} = WK \sum \|f_{x_i x_{i+1}}\|_{BV} \leq WK\varepsilon/WK = \varepsilon. \end{aligned}$$

Hence  $F \in AC$ .

In the next three theorems we establish the isomorphism between  $\Omega$  and  $\mathcal{L}$ .

6.4. LEMMA. *If  $K \in \Omega$ , then there is a unique Lipschitz function  $F_K$  anchored at zero given by  $F_K(x) = xK((0, x])$ . Furthermore, the Lipschitz constant is  $WK$ .*

**Proof.** Since  $K$  is convex with respect to length,  $F_K(b) - F_K(a) = (b-a)K((a, b])$ . Since  $K$  is fundamentally bounded it follows that  $F_K$  is Lipschitz, and since  $WK$  is the fundamental bound for  $K$  it follows that  $WK$  is the Lipschitz constant for  $F_K$ .

6.5. LEMMA. *If  $F$  is a Lipschitz function such that  $F(0) = 0$  with Lipschitz constant  $L$ , then the set function  $K_F$  given by  $K_F((a, b]) = (F(b) - F(a))/(b-a)$  where  $a, b \in [0, 1]$  is convex with respect to length and is fundamentally bounded with fundamental bound  $WK_F = L$ .*

**Proof.** Since  $F$  is Lipschitz it follows that  $K_F$  is fundamentally bounded and that  $WK_F = L$ . We now show that  $K$  is convex with respect to length. Suppose  $(a, b] \subset [0, 1]$  and that  $\{x_i\}$  partition  $(a, b]$ . Then

$$\begin{aligned} \sum [\Delta x_i / (b-a)] K_F((x_i, x_{i+1}]) &= \sum [\Delta x_i / (b-a)] [\Delta_i F / \Delta x_i] \\ &= (b-a)^{-1} \sum \Delta_i F \\ &= (F(b) - F(a)) / (b-a) = K_F((a, b]). \end{aligned}$$

6.6. REMARK. In view of 6.5, an alternate form for the representation theorem 4.2 is  $T(f) = \lim_{\sigma} \sum_{\sigma} \Delta_i f \Delta_i F / \Delta_i x = \int df dF/dx$ , the last integral being the Hellinger integral<sup>(1)</sup>.

We summarize in the following theorem.

6.7. THEOREM. *The space  $\Omega$  is isometrically isomorphic to  $\mathcal{L}$  with the isomorphism given by  $K \leftrightarrow F_K$ . Hence, it follows that  $AC^*$  is isometrically isomorphic to  $\mathcal{L}$ .*

We now relate 4.2 to the classical representation theorem given in [1] and show that, for at least some elements  $T \in AC^*$ , Theorem 4.2 gives a more delineating representation of the norm preserving extension of  $T$  on  $BV$  than that of [1]. First we observe the following lemma without proof.

6.8. LEMMA. *The closure of  $C_1$ , the continuously differentiable functions on  $[0, 1]$  anchored at zero, in the  $BV$  norm is  $AC$ .*

<sup>(1)</sup> The authors were unaware of the Hellinger integral when the original manuscript was prepared and wish to acknowledge the referee for this contribution.

6.9. THEOREM. Suppose that  $g_n \in L^\infty$ ,  $n=0, 1, 2, \dots$ , and that  $g_n$  converges to  $g_0$  in the  $L^1$  norm. Further, suppose  $G_n(x) = L \int_0^x g_n$  (Lebesgue integral) for each  $n$ . Then for  $f \in \mathcal{L}$ ,  $v \int K_{G_n} \cdot df$  converges to  $v \int K_{G_0} \cdot df$ .

**Proof.** Observe that

$$\begin{aligned} \left| v \int K_{G_n} \cdot df - v \int K_{G_0} \cdot df \right| &= \left| \lim_{\sigma} \sum \Delta_i f [K_{G_n}((x_i, x_{i+1})) - K_{G_0}((x_i, x_{i+1}))] \right| \\ &= \lim_{\sigma} \left| \sum \Delta_i f \left[ \left( \int_{x_i}^{x_{i+1}} g_n - g_0 \right) / \Delta x_i \right] \right| \\ &\leq \lim_{\sigma} \sup \sum |\Delta_i f / \Delta x_i| \int_{x_i}^{x_{i+1}} |g_n - g_0| \\ &\leq L \int_0^1 |g_n - g_0| \end{aligned}$$

where  $L$  is the Lipschitz constant for  $f$ . Since  $g_n$  converges to  $g$  in the  $L^1$  norm, it follows that  $v \int K_{G_n} \cdot df$  converges to  $v \int K_{G_0} \cdot df$ .

6.10. THEOREM. Suppose  $f \in AC$  and  $g \in L^\infty$  and suppose  $G \in \mathcal{L}$  is given by  $G(x) = L \int_0^x g$ . Then  $v \int K_G \cdot df = L \int f'g$ .

This theorem essentially says that the most natural of all possible relations exists between our representation and that in [1], i.e., the isomorphism between the characterization of  $AC^*$  as  $L^\infty$  and the characterization of  $AC^*$  as  $\Omega$  isomorphic to  $\mathcal{L}$  is given by  $g \leftrightarrow T \leftrightarrow K_G \leftrightarrow G = L \int g$ .

**Proof of 6.10.** We first establish 6.10 for the case  $f \in C_1$  and  $g \in C$ . Then  $v \int K \cdot df$  is given by

$$\begin{aligned} \lim_{\sigma} \sum \Delta_i f K_G((x_i, x_{i+1})) &= \lim_{\sigma} \sum (\Delta_i f)(\Delta_i G / \Delta x_i) \\ &= \lim_{\sigma} \sum (\Delta_i f / \Delta x_i)(\Delta G_i / \Delta x_i) \Delta x_i \\ &= \lim_{\sigma} \sum f'(\xi_i) g(\eta_i) \Delta x_i \end{aligned}$$

by the Mean Value Theorem, where  $x_i < \xi_i$ ,  $\eta_i < x_{i+1}$ . But by Bliss' Theorem [9, p. 224] the above limit is the Riemann integral  $R \int f'(x)g(x) dx$ , which establishes the result in this special case.

Next the result is established for  $f \in C_1$  and  $g \in L^\infty$ . There is a bounded sequence  $\{g_n\}$  in  $C$  which converges to  $g$  in the  $L^1$  norm. Then Theorem 6.9 implies  $v \int K_{G_n} \cdot df$  converges to  $v \int K_G \cdot df$  where  $G_n(x) = L \int_0^x g_n$  for each  $n$ .

For each  $n$ , we know  $v \int K_{G_n} \cdot df = L \int f'g_n$  and by the Bounded Convergence Theorem  $L \int f'g_n$  converges to  $L \int f'g$ . Hence,  $v \int K_G \cdot df = L \int f'g$ .

Finally we establish the full result. We suppose  $f \in AC$  and  $g \in L^\infty$ . Lemma 6.8 implies that there is a sequence  $\{f_n\}$  in  $C_1$  which converges to  $f$  in the  $BV$  norm. Hence, it follows that  $\lim_n v \int K_G \cdot df_n = v \int K_G \cdot df$ . For each  $n$  we have shown that

$v \int K_G \cdot f_n = \int f'_n g$ . Since  $f_n$  converges to  $f$  in the  $BV$  norm, it follows that  $f'_n$  converges to  $f'$  in the  $L_1$  norm. Since for  $f$  in  $L^1$  and  $g$  in  $L^\infty$  we have  $|\int fg| < \|g\|_\infty \int |f|$ , it follows that  $\lim_n L \int f'_n g = L \int f' g$ , from which we have  $v \int K_G \cdot df = L \int f' g$ .

6.11. COROLLARY. Suppose  $f \in AC$  and  $K \in \Omega$ , and suppose  $F(x) = v \int_0^x K \cdot df$ . Then, the relation between  $F$  and  $f$  is given by  $F' = f' G'_K$  a.e.

We close this section with the following development which gives a sufficient condition on a convex set function  $K$  such that the  $v$ -integral with respect to  $K$  of every  $BV$  function will exist (Corollary 6.14).

6.12. THEOREM<sup>(2)</sup>. Suppose  $\{K_n\}$  is a sequence of fundamentally bounded set functions which converge to the set function  $K$  in the fundamental bound sense. Suppose further that  $f \in BV$  which is  $v$ -integrable with respect to each  $K_n$ . Then  $K$  is fundamentally bounded,  $f$  is  $v$ -integrable with respect to  $K$ , and  $\lim_n v \int K_n \cdot df = v \int K \cdot df$ .

**Proof.** Observe that

$$\left\| v \int K_n \cdot df - v \int K_m \cdot df \right\| = \left\| v \int (K_n - K_m) \cdot df \right\| \leq W(K_n - K_m) \|f\|_{BV}.$$

Hence,  $\lim_n v \int K_n \cdot df$  exists. Call the limit  $V$ . For  $\varepsilon > 0$ , there is an  $N$  such that  $n \geq N$  implies  $W(K_n - K) < (\varepsilon/3) \|f\|_{BV}$ , and such that  $\|V - v \int K_n \cdot df\| < \varepsilon/3$ . Choose  $\sigma'$  such that  $\sigma$  finer than  $\sigma'$  implies  $\|v \int K_n \cdot df - \sum_\sigma (\Delta_i f) K_n((x_i, x_{i+1}))\| < \varepsilon/3$ . Then for  $\sigma$  finer than  $\sigma'$ ,

$$\begin{aligned} \left\| V - \sum_\sigma (\Delta_i f) K((x_i, x_{i+1})) \right\| &\leq \left\| V - v \int K_n \cdot df \right\| + \left\| v \int K_n \cdot df - \sum_\sigma (\Delta_i f) K_n((x_i, x_{i+1})) \right\| \\ &\quad + \left\| \sum_\sigma (\Delta_i f) K_n((x_i, x_{i+1})) - \sum_\sigma (\Delta_i f) K((x_i, x_{i+1})) \right\| \\ &< \varepsilon/3 + \varepsilon/3 + W(K_n - K) \|f\|_{BV} < \varepsilon. \end{aligned}$$

Hence, the theorem is established.

6.13. THEOREM. If  $G$  is polygonal, the  $v \int K_G \cdot df$  exists for every  $f \in BV$ .

**Proof.** Suppose  $f \in BV$ . Let  $\sigma_G = \{a_i\}$  denote the partition of  $[0, 1]$  determined by  $G$ . Then for  $\sigma > \sigma_G$ ,  $\sum_\sigma \Delta_i f K_G((x_i, x_{i+1})) = \sum_\sigma (\Delta_i f) (\Delta_i G / \Delta x_i) = \sum_\sigma (\Delta_i f) \alpha_j$  where  $\alpha_j$  is the slope of  $G$  in the interval  $(a_j, a_{j+1})$ , where  $[x_i, x_{i+1}] \subset [a_j, a_{j+1}]$ . Therefore,

$$\sum_\sigma (\Delta_i f) K_G((x_i, x_{i+1})) = \sum_j (f(a_{j+1}) - f(a_j)) \alpha_j = v \int K_G \cdot df.$$

Observe that if  $K \in \Omega$  and  $f \in BV$ , then  $|\lim_\sigma \sum \Delta_i f K((x_i, x_{i+1}))| \leq WK \|f\|_{BV}$ . Therefore, if  $K \in \Omega$  integrates all  $BV$  functions in the  $v$ -sense, then it defines a linear

<sup>(2)</sup> The original Theorem 6.12 and proof require that the  $K_n$ 's be convex. The authors gratefully acknowledge the referee for this improved version.

functional on  $BV$ . This observation along with Theorems 6.12 and 6.13 yield the following.

6.14. COROLLARY. *If  $g$  is in the sup-norm closure of the step functions, and if  $G(x) = L \int_0^x g$ , then  $T(f) = v \int K_G \cdot df$  exists for every  $f \in BV$  and hence  $T \in BV^*$ .*

7. **The  $v$ -integral in the vector-valued setting.** In this section we suppose  $X$  is a Banach space and that  $Y$  is an  $LNS$ , and  $B[X, Y]$  is the  $LNS$  of bounded operators from  $X$  into  $Y$ . The variation of an  $X$ -valued function  $f$  is defined to be  $V_0^1 f = \sup_\sigma \sum \|\Delta_i f\|$  and  $BV(X)$  denotes the  $LNS$  of  $X$ -valued functions of bounded variation which are  $\theta_X$  (the additive identity of  $X$ ) at 0, with  $\|f\|_{BV(X)} = V_0^1 f$ . Define the space of  $X$ -valued absolutely continuous functions,  $AC(X)$ , in the natural, analogous fashion. We mean by an  $X$ -valued polygonal function a function of the form  $p = \sum \Psi_{H_i} \cdot x_i$ , where  $x_i \in X$  for each  $i$ . Let  $\Gamma$  denote the closure of  $P$ , the  $X$ -valued polygonal functions, in the  $BV$  norm. It is not known if  $\Gamma = AC$ . However, we show  $\Gamma \subset AC$  in the following theorem.

7.1. THEOREM. *If  $\{p_n\}$  is a sequence of polygonal functions which is Cauchy in the  $BV$  norm, then there is a function  $f \in AC$  such that  $\{p_n\}$  converges to  $f$  in the  $BV$  norm.*

The proof follows as in the second proof of 2.1.

The following theorem generalizes two theorems from earlier sections. The proof follows as earlier and is therefore omitted.

7.2. THEOREM. (i) *If  $f \in \Gamma$ , then  $\lim_\sigma pf_\sigma = f$  where the convergence is in the  $BV$  norm.* (ii) *If  $K$  is a  $B[X, Y]$ -valued set function which is convex with respect to length,  $p \in P$ , and  $\sigma$  is the partition of  $[0, 1]$  determined by  $p$ , then for  $\sigma' > \sigma$ ,*

$$\sum_{\sigma'} [K((x_i, x_{i+1}))](\Delta_i p) = \sum_{\sigma} [K((x_i, x_{i+1}))](\Delta_i p).$$

Hence,  $p$  is  $v$ -integrable with respect to  $K$  and  $v \int K \cdot df = \sum_{\sigma} [K((x_i, x_{i+1}))](\Delta_i p)$ .

7.3. THEOREM. *Suppose  $K$  is a fundamentally bounded set function which is convex with respect to length and takes its values in  $B[X, Y]$ . Then the map  $T(f) = v \int K \cdot df$  is a bounded linear operator from  $\Gamma$  into  $\bar{Y}$ , the completion of  $Y$ . Furthermore,  $\|T\| = WK$ .*

The proof follows as in 4.2.

7.4. THEOREM. *Suppose  $T$  is a bounded linear operator from  $\Gamma$  into  $Y$ . Then, there is a unique, fundamentally bounded set function  $K$  with values in  $B[X, Y]$  which is convex with respect to length such that  $T(f) = v \int K \cdot df$  for each  $f \in \Gamma$ . Furthermore,  $\|T\| = WK$ .*

**Proof.** Define the set function  $K$  on the set of intervals with values in  $B[X, Y]$  by  $K(H)x = T(\Psi(H)x)$ . The remainder of the proof now follows as in 4.2.

In the event  $Y$  is complete, then the above two theorems yield the following corollary.

7.5. COROLLARY. Suppose  $Y$  is complete. Then, a linear operator  $T$  from  $\Gamma$  into  $Y$  is bounded if and only if there is a unique, fundamentally bounded set function  $K$  with values in  $B[X, Y]$  which is convex with respect to length, such that  $T(f) = v \int K \cdot df$  for each  $f \in \Gamma$ . Furthermore,  $\|T\| = WK$ .

An alternative way to define the variation of an  $X$ -valued function is  $SV_0^1 f = \sup_{\sigma} \{ \sup \{ \| \sum_{\sigma} \alpha_i \Delta_i f \| : |\alpha_i| \leq 1 \text{ for each } i \} \}$ , which is called the semivariation of  $f$ . The space of functions of bounded semivariation which evaluate to  $\theta_x$  at 0, and with norm equal to the semivariation is denoted by  $BSV$ . The notion of semi-absolute continuity is defined in a fashion analogous to that of bounded semivariation and  $SAC$  denotes the corresponding functions space. Let  $S\Gamma$  denote the closure of  $P$  in the  $BSV$  norm. By a proof similar to that of 7.1 it follows that  $S\Gamma \subset SAC$ .

7.6. LEMMA. Suppose  $p, p' \in P$  and  $\sigma$  and  $\sigma'$  are the partitions of  $[0, 1]$  determined by  $p$  and  $p'$  respectively. Then  $\|p - p'\|_{BSV} = \sup \{ \| \sum_{\sigma * \sigma'} \alpha_i \Delta_i(p - p') \| : |\alpha_i| \leq 1 \}$ .

**Proof.** Let  $\{x_i\} = \sigma * \sigma'$ . Then

$$\begin{aligned} \|p - p'\|_{BSV} &= \sup \left\{ \sup \left\{ \left\| \sum_{\pi} B_j \Delta_j(p - p') \right\| : |B_j| \leq 1 \right\} : \pi \text{ is a partition of } [0, 1] \right\} \\ &= \sup \left\{ \sup \left\{ \left\| \sum_{\pi} B_j [(x_{ij+1} - x_{ij}) / (x_{i+1} - x_i)] \Delta_i(p - p') \right\| : |B_j| \leq 1 \right\} : \right. \\ &\quad \left. \pi > \sigma * \sigma' \right\} \\ &= \sup \left\{ \left\| \sum_{\sigma * \sigma'} \alpha_i \Delta_i(p - p') \right\| : |\alpha_i| \leq 1 \right\}. \end{aligned}$$

The last equality follows because  $|\sum_j B_j [(x_{ij+1} - x_{ij}) / (x_{i+1} - x_i)]| \leq 1$  for  $|B_j| \leq 1$ .

Lemma 7.6 allows us to observe that the analogies of 7.2, 7.3, 7.4 can be established in the semivariational setting by essentially the same proofs. The analogous theorem to Corollary 7.5 is stated below for comparison. Observe that for the special case  $Y = R$ , 7.5 and 7.7 imply that  $\Gamma^* = (S\Gamma)^*$ .

7.7. THEOREM. Suppose  $Y$  is complete. Then, a linear operator  $T$  from  $S\Gamma$  into  $Y$  is bounded if and only if there is a unique, fundamentally bounded set function  $K$  with values in  $B[X, Y]$  which is convex with respect to length, such that  $T(f) = v \int K \cdot df$  for each  $f \in S\Gamma$ . Furthermore,  $\|T\| = WK$ .

Let  $\Omega(B[X, Y])$  denote the space of set functions  $K$  which are fundamentally bounded and which are convex with respect to length and with norm  $\|K\| = WK$ . Let  $\mathcal{L}(B[X, Y])$  denote the space of  $B[X, Y]$ -valued Lipschitz functions  $f$  on  $[0, 1]$  (i.e., the difference quotients are uniformly bounded) such that  $f(0) = \theta_{B[X, Y]}$  with norm  $\|f\|_{BV} = L$ , the Lipschitz constant of  $f$ .

The results obtained in 6.1–6.7 and 6.12–6.14 all carry over to the vector setting and we summarize them in the following theorems.

7.8. THEOREM. *The space  $\Omega(B[X, Y])$  is isometrically isomorphic to the space  $\mathcal{L}(B[X, Y])$  and the isometry is given by  $G \leftrightarrow K_G$ . Hence,  $\Gamma^*$  and  $(S\Gamma)^*$  are each isometrically isomorphic to  $\mathcal{L}(B[X, Y])$ .*

7.9. THEOREM. *If  $f \in BV$ , and  $\{K_n\}$  is a sequence of fundamentally bounded set functions which converge to  $K$  in the fundamental bound sense, and if  $f$  is  $v$ -integrable with respect to  $K_n$  for each  $n$ , then  $f$  is  $v$ -integrable with respect to  $K$  and  $v \int K \cdot df = \lim_n v \int K_n \cdot df$ .*

7.10. THEOREM. *If  $G$  is polygonal, then  $v \int K_G \cdot df$  exists for every  $f \in BSV$ .*

7.11. COROLLARY. *A sufficient condition that  $T(f) = v \int K_G \cdot df$  be a bounded linear operator from  $BV$  into  $\bar{Y}$  is that  $G$  be in the closure of the polygonal functions in  $\mathcal{L}(B[X, Y])$ .*

#### REFERENCES

1. N. Dunford and J. Schwartz, *Linear operators. I: General theory*, Pure and Appl. Math., vol. 7, Interscience, New York, 1958. MR 22 #8302.
2. R. J. Easton and D. H. Tucker, *A generalized Lebesgue-type integral*, Math. Ann. **181** (1969), 311–324.
3. J. R. Edwards and S. G. Wayment, *A unifying integral representation theorem*, Math. Ann. **187** (1970), 317–328.
4. ———, *Integral representations for continuous linear operators in the setting of convex topological vector spaces*, Trans. Amer. Math. Soc. (to appear).
5. T. H. Hildebrandt, *Linear continuous functionals on the space  $(BV)$  with weak topologies*, Proc. Amer. Math. Soc. **17** (1966), 658–664. MR 33 #1710.
6. ———, *Linear operations on functions of bounded variation*, Bull. Amer. Math. Soc. **44** (1938), 75.
7. R. Lane, *The integral of a function with respect to a function*, Proc. Amer. Math. Soc. **5** (1954), 59–66. MR 15, 514.
8. I. P. Natanson, *Theory of functions of a real variable*, GITTL, Moscow, 1957; English transl., Ungar, New York, 1961. MR 26 #6309.
9. E. J. Purcell, *Calculus with analytic geometry*, Appleton-Century-Crofts, New York, 1965.
10. F. Riesz, *Sur les opérations fonctionnelles linéaires*, C. R. Acad. Sci. Paris **149** (1909), 974–977.
11. D. H. Tucker, *A note on the Riesz representation theorem*, Proc. Amer. Math. Soc. **14** (1963), 354–358. MR 26 #2865.
12. D. H. Tucker and S. G. Wayment, *Absolute continuity and the Radon-Nikodym theorem*, J. Reine Angew. Math. (to appear).
13. D. H. Uherka, *Generalized Stieltjes integrals and a strong representation theorem for continuous maps on a function space*, Math. Ann. **182** (1969), 60–66.

UTAH STATE UNIVERSITY,  
LOGAN, UTAH 84321