

NOETHERIAN INTERSECTIONS OF INTEGRAL DOMAINS

BY

WILLIAM HEINZER AND JACK OHM⁽¹⁾

Abstract. Let $D < R$ be integral domains having the same quotient field K and suppose that there exists a family $\{V_i\}_{i \in I}$ of 1-dim quasi-local domains having quotient field K such that $D = R \cap \{V_i \mid i \in I\}$. The goal of this paper is to find conditions on R and the V_i in order for D to be noetherian and, conversely, conditions on D in order for R and the V_i to be noetherian. An important motivating case is when the set $\{V_i\}$ consists of a single element V and V is a valuation ring. It is shown, for example, in this case that (i) if V is centered on a finitely generated ideal of D , then V is noetherian and (ii) if V is centered on a maximal ideal of D , then D is noetherian if and only if R and V are noetherian.

Our terminology is that of Zariski-Samuel [15] and Bourbaki [2]. All rings considered are commutative rings with identity, and except for several propositions concerning flat ring extensions, we deal only with integral domains. If S and T are sets, we use $S \setminus T$ to denote $\{s \in S \mid s \notin T\}$; and if V is a valuation ring, we use $V(-)$ to denote the value of $-$ with respect to an associated valuation. If D is a subring of a ring R and P is a prime ideal of D , then we write R_P for the quotient ring of R with respect to the multiplicative system $D \setminus P$. We use $\bar{R}$ to denote the integral closure of an integral domain R ; and finally, $\subset$ denotes containment and $<$ strict containment.

1. R -Krull domains. In proving the main theorems of this paper, we found that we were frequently using techniques that resembled those encountered in dealing with Krull rings, and toward the end of our work it became clear to us that a natural setting for our study is one that generalizes the language and ideas of classical Krull ring theory. Thus, the concept of an R -Krull domain will play a central role in our development, and we shall carry over to these rings a surprisingly large number of results, suitably generalized, from the theory of Krull rings.

We begin by fixing some notation and terminology which will remain in effect throughout the paper. R and D will always denote integral domains with the same

Presented to the Society, January 24, 1969 under the title *Noetherian and non-Noetherian commutative rings*; received by the editors October 12, 1968 and, in revised form, February 2, 1971.

AMS 1970 subject classifications. Primary 13E05, 13F05, 13B20.

Key words and phrases. 1-dim quasi-local integral domain, flat ring extension, Krull domain, essential valuation ring.

⁽¹⁾ The authors received partial support from the National Science Foundation during the writing of this paper. During completion of the paper the first named author was supported as an NRC-ONR Research Associate.

Copyright © 1972, American Mathematical Society

quotient field K . A set of domains $\{D_i\}_{i \in I}$ with the same quotient field K will be said to have *FC* (*finiteness condition*) if $0 \neq \xi \in K$ implies ξ is a unit of D_i for all but finitely many i , which is equivalent to the assertion that $0 \neq d \in D$ implies d is a unit of D_i for all but finitely many i . A nonempty set $\{V_i\}_{i \in I}$ of 1-dim quasi-local domains with quotient field K will be called a *set of R -representatives for D* if $\{V_i\}_{i \in I}$ has FC, $R \not\subseteq \bigcap_{i \in I} V_i$, and $D = R \cap (\bigcap_{i \in I} V_i)$. Note that when such a set exists, we must have $D < R$. If $\{V_i\}_{i \in I}$ is a set of R -representatives for D , we use P_i to denote the center of V_i on D , i.e. P_i is the intersection of the maximal ideal of V_i with D . V_i is said to be *irredundant* (in the intersection $D = R \cap (\bigcap_{i \in I} V_i)$) if $R \cap (\bigcap_{j \neq i} V_j) \not\subseteq V_i$.

A valuation ring is called *rational* if its associated value group is order isomorphic to an additive subgroup of the rational numbers. A set $\{V_i\}_{i \in I}$ of R -representatives for D will be called a *set of rational R -representatives for D* if every V_i , $i \in I$, is a rational valuation ring; and D will be called an *R -Krull domain* if there exists a set of rational R -representatives for D . Thus, an R -Krull domain is a domain D having the same quotient field K as R and such that there exists a set $\{V_i\}_{i \in I}$ of rational valuation rings with quotient field K such that $\{V_i\}_{i \in I}$ has FC, $R \not\subseteq \bigcap_{i \in I} V_i$, and $D = R \cap (\bigcap_{i \in I} V_i)$.

The concept of an R -Krull domain is a continuation of the line of thought begun in [12, §5]. When $R = K$ and each V_i is noetherian, then the notion of R -Krull domain coincides with that of Krull ring. Our goal in this section is to find conditions on R and the V_i in order for D to be noetherian and conversely conditions on D in order for R and the V_i to be noetherian. We begin by generalizing some well-known results on Krull rings.

1.1. LEMMA. *Let $\{D_\alpha\}$ be a set of overrings contained in K of the integral domain D , and let S be a multiplicative system of D . If $\{D_\alpha\}$ has FC and $D = \bigcap_\alpha D_\alpha$, then $D_S = \bigcap_\alpha (D_\alpha)_S$. In particular, if $\{V_i\}$ is a set of R -representatives for D , then $D_S = R_S \cap (\bigcap_{j \in J} V_j)$, where $\{V_j\}_{j \in J}$ is the set of V_i such that $P_i \cap S = \emptyset$.*

Proof. Let $\xi \in \bigcap_\alpha (D_\alpha)_S$. By the FC, $\xi \notin D_\alpha$ for at most finitely many α , say $\xi \notin D_1, \dots, D_n$. There exists $s_i \in S$ such that $s_i \xi \in D_i$, $i = 1, \dots, n$; so if $s = s_1 \cdots s_n$, then $s\xi \in D$ and $\xi \in D_S$. For the second assertion, observe that $(V_i)_S = K$ if $P_i \cap S \neq \emptyset$.

1.2. COROLLARY. *Let $\{V_i\}_{i \in I}$ be a set of R -representatives for D . If P is a prime ideal of D such that $R \not\subseteq D_P$, then $P_i \subset P$ for some $i \in I$.*

Proof. Take $S = D \setminus P$ in 1.1.

The next lemma plays an important role in what follows. Its proof is basically that of [12, Theorem 5.1, p. 330].

1.3. LEMMA. *If D, R are domains with quotient field K and V is a rational valuation ring with quotient field K such that $R \not\subseteq V$ and $D = R \cap V$, then $V = D_P$, where P is the center of V on D .*

Proof. $D_P \subset V$ is immediate, so we shall concern ourselves with the reverse inclusion. If $\xi \in V$, we must show that there exists $d \in D \setminus P$ such that $d\xi \in R$. For this it is sufficient to exhibit an element $d \in R$ such that $V(d) = 0$ and $d\xi \in R$. Since V and D have the same quotient field, $\xi = a/b$, $a, b \in D$; and since $R \not\subset V$, there exists $r \in R$ such that $V(r) < 0$. If $b \notin P$, then b is the required element. On the other hand, if $V(b) > 0$, then since V is a rational valuation ring, there exist integers $m, n \geq 1$ such that $V(r^n b^m) = 0$; thus, $d = r^n b^m$ is the required element. Q.E.D.

The example of [12, Example 5.3, p. 330] shows that 1.3 is no longer valid if one merely assumes that V is a rank 1 valuation ring rather than a rational valuation ring.

1.4. PROPOSITION. *Let D be an R -Krull domain and let $\{V_i\}_{i \in I}$ be a set of rational R -representatives for D . Then $\mathcal{E} = \{D_P \mid R \not\subset D_P \text{ and } D_P = V_i \text{ for some } i\}$ is also a set of rational R -representatives for D .*

Proof. Since any $D_P \in \mathcal{E}$ equals some V_i , certainly $\mathcal{E}$ has FC and each $D_P \in \mathcal{E}$ is a rational valuation ring. It therefore remains to see that $D = R \cap \{D_P \mid D_P \in \mathcal{E}\}$. $\subset$ is clear, so suppose x is in the right-hand side of the equality but is not in D . Then there exists at least one V_j such that $x \notin V_j$, and by the FC there exist at most finitely many such V_j ; call these $V_1, \dots, V_n$. Then $D = R' \cap V_1 \cap \dots \cap V_n$, where $R' = R \cap (\bigcap_{i \in I \setminus \{1, \dots, n\}} V_i)$. $D \neq R'$ since $x \in R' \setminus D$, so we may assume that V_1 is irredundant in this intersection. It follows by 1.3 that V_1 is a quotient ring of D . Therefore $V_1 \in \mathcal{E}$, and hence $x \in V_1$, a contradiction.

1.5. THEOREM. *Let D be an R -Krull domain, let $\{V_i\}_{i \in I}$ be a set of rational R -representatives for D , and let P_i denote the center of V_i on D . If P is a prime ideal of D such that $R \not\subset D_P$, then the following are equivalent:*

- (i) $P_i \subset P$ implies $P_i = P$, for all $i \in I$.
- (ii) There exists $j \in I$ such that $V_j = D_P$.
- (iii) There exists $j \in I$ such that $P_j = P$ and V_j is irredundant.
- (iv) $P = P_j$ for some $j \in I$, P is minimal, and no prime ideal of R lies over P .
- (v) P is minimal.

Proof. (i) $\Rightarrow$ (ii). By 1.1, $D_P = R_P \cap (\bigcap_{j \in J} V_j)$, where $\{V_j\}_{j \in J}$ are those V_i having center $P_i \subset P$. $D_P \neq R_P$, since otherwise $R \subset D_P$; so $J \neq \emptyset$. J is then finite by the FC, since $P_i \subset P$ implies $P_i = P$ by (i). Therefore we can delete finitely many elements from $\{V_j\}_{j \in J}$ to get an irredundant intersection $D_P = R_P \cap V_1 \cap \dots \cap V_t$. By 1.3, $V_1 = D_P$.

(ii) $\Rightarrow$ (iii). $V_j = D_P$ implies P is minimal and V_j is the only V_i centered on P . If V_j is redundant, then $D = R \cap (\bigcap_{i \neq j} V_i)$. But then (i) $\Rightarrow$ (ii) applied to the set of R -representatives $\{V_i\}_{i \neq j}$ implies that there exists V_i , $i \neq j$, such that V_i is centered on P , a contradiction.

(iii) $\Rightarrow$ (iv). V_j is irredundant implies $V_j = D_{P_j}$ by 1.3; and since V_j is 1-dim, it follows that P_j is minimal. If now Q is a prime ideal of R such that $Q \cap D = P$, then

$V_j = D_P \subset R_Q$. But V_j is a rank 1 valuation ring, so then $V_j = R_Q$; and hence $R \subset R_Q = V_j = D_P$, a contradiction to our hypothesis.

(iv) $\Rightarrow$ (v) $\Rightarrow$ (i). Trivial.

1.6. *Application.* Let D be an R -Krull domain and let $\{V_i\}_{i \in I}$ be a set of rational R -representatives for D . (ii) $\Leftrightarrow$ (iii) $\Leftrightarrow$ (v) of 1.5 shows that $\{\text{irredundant } V_i\text{'s}\} = \{V_i \mid R \not\subset V_i \text{ and } V_i = D_{P_i}\} = \{D_P \mid P \text{ is minimal and } R \not\subset D_P\}$. We shall denote this set by $\mathcal{E}$. Note that the characterization $\mathcal{E} = \{D_P \mid P \text{ is minimal and } R \not\subset D_P\}$ shows that $\mathcal{E}$ is independent of the choice of the set of rational R -representatives $\{V_i\}$. Also, we have already seen in 1.4 that $\mathcal{E}$ is itself a set of rational R -representatives for D . We shall call $\mathcal{E}$ the set of *essential R -representatives* for D , and we shall call the centers on D of the elements of $\mathcal{E}$ the *essential primes* of D . Thus, $\mathcal{E}$ is a set of irredundant rational R -representatives for D ; and given any set of rational R -representatives for D , the subset of irredundant elements is exactly $\mathcal{E}$. (If one merely assumes that the V_i are rank 1 valuation rings rather than rational valuation rings, the example of [12, Example 5.3, p. 330] shows that not every irredundant V_i need be in $\mathcal{E}$.)

We shall frequently use in what follows a couple of facts involving flatness, one of which is the following criterion [14, Theorem 2, p. 795]. Let $D \subset R$ be domains with the same quotient field. Then R is flat over D if and only if for any prime ideal P of D , either $R \subset D_P$ or $PR = R$. The second fact is that if R is flat over D and contained in the quotient field of D , then D is noetherian implies that R is noetherian [14, Corollary, p. 796].

We are now ready to examine the noetherian properties of R -Krull domains. Theorem 1.10 constitutes part of the main result of this section. Its proof requires the corollary to the next proposition.

1.7. **PROPOSITION.** *Let $\mathcal{R}$ be a ring and let $\{\mathcal{R}_\alpha\}$ be a set of flat $\mathcal{R}$ -algebras (or $\mathcal{R}$ -modules). Let A be an ideal of $\mathcal{R}$ such that for each maximal ideal $M \supset A$ of $\mathcal{R}$ there exists an $\mathcal{R}_\alpha$ such that $M\mathcal{R}_\alpha \neq \mathcal{R}_\alpha$. If A' is an ideal of $\mathcal{R}$ such that $A'\mathcal{R}_\alpha \subset A\mathcal{R}_\alpha$ for each α , then $A' \subset A$. Moreover, if for each α , $A\mathcal{R}_\alpha$ is finitely generated as an ideal of $\mathcal{R}_\alpha$ (or finitely generated as an $\mathcal{R}$ -module) and if there exists a finitely generated ideal $B \subset A$ such that $B\mathcal{R}_\alpha \neq A\mathcal{R}_\alpha$ for at most a finite number of α , then A is finitely generated.*

Proof. If $b \in A'$, then $(A:b)\mathcal{R}_\alpha = A\mathcal{R}_\alpha : b$ (where $A\mathcal{R}_\alpha : b = \{y \in \mathcal{R}_\alpha \mid by \in A\mathcal{R}_\alpha\}$), [2a, Exercise 22, p. 65] or [11, 18.10, p. 58]. Since $b\mathcal{R}_\alpha \subset A'\mathcal{R}_\alpha \subset A\mathcal{R}_\alpha$ we have $(A:b)\mathcal{R}_\alpha = \mathcal{R}_\alpha$ for each α . Thus $A:b$ is contained in no maximal ideal of $\mathcal{R}$, so $A:b = \mathcal{R}$, i.e. $b \in A$ and $A' \subset A$. Assume now that $A\mathcal{R}_\alpha$ is finitely generated for each α and that $B \subset A$, B finitely generated, is such that $B\mathcal{R}_\alpha = A\mathcal{R}_\alpha$ for all but a finite number of α , say $\alpha = 1, 2, \dots, n$. We can choose $A_i \subset A$, A_i finitely generated, such that $A_i\mathcal{R}_i = A\mathcal{R}_i$. Then $B' = B + A_1 + \dots + A_n$ is such that $A\mathcal{R}_\alpha \subset B'\mathcal{R}_\alpha$ for each α . Thus $A = B'$ and A is finitely generated. Q.E.D.

The following corollary includes the well-known statement that a quasi-semilocal domain which is locally noetherian is noetherian [15, II, Lemma 2, p. 327].

1.8. COROLLARY. *Let $\mathcal{R}$ be a ring and let $\{\mathcal{R}_\alpha\}$ be a set of flat $\mathcal{R}$ -algebras (or $\mathcal{R}$ -modules) such that for each proper ideal A of $\mathcal{R}$, $A\mathcal{R}_\alpha \neq \mathcal{R}_\alpha$ for at least one and at most a finite number of $\mathcal{R}_\alpha$. If each $\mathcal{R}_\alpha$ is a noetherian ring (or a noetherian $\mathcal{R}$ -module), then $\mathcal{R}$ is noetherian.*

For an incidental application of 1.8, let us consider the following question: Let $\mathcal{R}$ be a noetherian ring and $\mathcal{R}_1, \dots, \mathcal{R}_n$ be a set of flat overrings of $\mathcal{R}$ which are contained in the total quotient ring of $\mathcal{R}$. Is $\mathcal{R}' = \bigcap_{i=1}^n \mathcal{R}_i$ necessarily noetherian? The answer is "no" in general. In fact [7, Example 2.10] gives an example of a noetherian domain D having quotient rings D_1, D_2 such that $D_1 \cap D_2$ is not noetherian. However, the following is true.

1.9. PROPOSITION. *If $\mathcal{R}$ is a noetherian ring and $\mathcal{R}_1, \dots, \mathcal{R}_n$ are quasi-semilocal flat overrings of $\mathcal{R}$ which are contained in the total quotient ring of $\mathcal{R}$, then $\mathcal{R}' = \bigcap \mathcal{R}_i$ is noetherian.*

Proof. The $\mathcal{R}_i$ are flat over $\mathcal{R}$ and hence also are noetherian (see, for example, [1a, Corollary 1, p. 804]). Since every nonunit of $\mathcal{R}'$ is also a nonunit in some $\mathcal{R}_i$, it follows that the set of nonunits of $\mathcal{R}'$ is exactly the union of the finite set of contracted maximal ideals of the $\mathcal{R}_i$. Therefore every maximal ideal of $\mathcal{R}'$ is the contraction of a maximal ideal of some $\mathcal{R}_i$. Finally, since the $\mathcal{R}_i$ are contained in the total quotient ring of $\mathcal{R}$ and are flat over $\mathcal{R}$, they are also flat over $\mathcal{R}'$ [1b, Corollary 1, p. 803]. Thus, by 1.8, $\mathcal{R}'$ is noetherian. Q.E.D.

The next theorem is central to this paper.

1.10. THEOREM. *Let D be an R -Krull domain, let $\{V_i\}_{i \in I}$ be a set of rational R -representatives for D , and suppose that the center on D of every irredundant V_i is maximal. Then R is flat over D , and D is noetherian if and only if R and the irredundant V_i 's are noetherian.*

Proof. By 1.6 the irredundant V_i are precisely the elements of $\mathcal{E}$, i.e. the essential R -representatives for D . Let P be any prime ideal of D such that $R \not\subseteq D_P$. Since the center on D of any element of $\mathcal{E}$ is assumed maximal, 1.5(i) is satisfied by P ; and hence by 1.5(iv), P is itself the center of an element of $\mathcal{E}$ and therefore maximal, and no prime ideal of R lies over P . This implies that $PR = R$. Thus, $R \not\subseteq D_P$ implies $PR = R$, which proves that R is flat over D . It follows that D is noetherian implies R is noetherian.

We use 1.8 for the converse. As we have seen, for any prime ideal P of D , either $R \subseteq D_P$, in which case $PR \neq R$, or $R \not\subseteq D_P$, in which case P is the center of an element V of $\mathcal{E}$ and $PV \neq V$. Thus, since $\mathcal{E}$ has FC and each element of $\mathcal{E}$ is a quotient ring of D , 1.8 applies. Q.E.D.

Note that another way of stating 1.10 is the following: Let D be an R -Krull domain whose essential primes are maximal. Then R is flat over D , and D is noetherian if and only if R is noetherian and the essential primes of D are finitely generated. Moreover, when the essential primes of D are maximal, R is completely

determined by these primes, for it follows from the flatness of R [14, p. 795] that then $R = \bigcap \{D_Q \mid Q \text{ a nonessential prime of } D\}$.

We shall next examine a few examples to determine the bounds for possible generalizations of 1.10. A simple example in [3, p. 282] shows that there exist 1-dim local domains R, V with $D = R \cap V$ being 1-dim quasi-local but not noetherian; so the hypothesis that the V_i are rational valuation rings in 1.10 cannot be replaced by the hypothesis that the V_i are 1-dim local. (The D in this example is also integrally closed.)

The following modification of [12, Example 5.3, p. 330] shows that the assumption that the V_i are *rational* valuation rings is also crucial for the flatness assertion of 1.10.

1.11. *Example of $D = R \cap V$ such that V is a rank 1 valuation ring centered on a maximal ideal of D and yet R is not flat over D .* Let k be the rational numbers, let x be an indeterminate, and let V be the extension of the p -adic valuation v of k to $k(x)$ obtained by defining $V(a_0 + a_1x + \cdots + a_nx^n) = \inf \{v(a_i) + ip\}$, $i = 0, \dots, n$. Let $D = k[x] \cap V$, let M denote the center of V on D , and let P denote the center of the x -adic valuation ring of $k(x)$ on D . Then $P < M$, so there exists a rank 2 valuation ring W of $k(x)$ such that $W \supset D$ and W has prime ideals lying over M and P . If now we let $R = k[x] \cap W$, then $D = R \cap V$.

It is shown in [12, p. 331] that M is the only nonminimal prime ideal of D and that any other prime ideal of D is the contraction of a prime ideal of $k[x]$. Since M is the center of W on D , it follows that every prime ideal of D is the contraction of a prime ideal of R . Thus, it will follow that R is not flat over D if we show that $D \neq R$. For sufficiently large n , $x/p^n \notin V$ and hence $\notin D$; but $W(x)$ is infinitely larger than $W(p)$, so $x/p^n \in W$ and hence $\in R = k[x] \cap W$. Q.E.D.

The following can happen if one omits the condition that the center of each irredundant V_i be maximal in Theorem 1.10:

(i) D may be noetherian, but R may not be noetherian,

(ii) $R, V_i, i \in I$, may all be noetherian but D may not be noetherian. (We show in 1.17 that when D is noetherian the irredundant V_i are always noetherian.) An example which shows (i) is constructed in [4, §2], where R is an A -transform of a 3-dim normal affine ring D and $\{V_i\}$ is a finite set of discrete rank 1 valuation rings. To see (ii), it suffices to have a 2-dim, quasi-local, nonnoetherian Krull domain D having a minimal prime P such that $P = \sqrt{(x)}$ for some $x \in D$. For then take $R = D[1/x]$ and $V = D_P$. R is 1-dim Krull and hence noetherian, and $D = R \cap V$ since V is the only essential valuation ring of D which does not contain $1/x$. The existence of such a D is proved in [9, p. 28].

The following corollary, whose proof is immediate, includes [6, Proposition 1] as a special case.

1.12. COROLLARY. *Let D be a Krull domain with the set of essential valuation rings $\mathcal{E}$, and let $\mathcal{M}$ be any subset of $\{V \in \mathcal{E} \mid \text{the center of } V \text{ on } D \text{ is maximal}\}$. If $R = \bigcap \{V \mid V \in \mathcal{E} \setminus \mathcal{M}\}$, then R is noetherian if and only if D is noetherian.*

We next prove a generalization of 1.12. The generalization requires the following lemma which is taken from [13].

1.13. LEMMA. *Let R be an integral domain and let $\mathcal{P}$ be a set of minimal primes of R . If $\{R_P \mid P \in \mathcal{P}\}$ has FC and $R = \bigcap \{R_P \mid P \in \mathcal{P}\}$, then $\mathcal{P}$ is the set of all minimal primes of R .*

Proof. Let Q be a minimal prime of R and suppose that $Q \notin \mathcal{P}$. Choose $y \neq 0 \in Q$, and let $P_1, \dots, P_t$ be the primes in $\mathcal{P}$ such that $y \in P_i$ ($t \geq 1$ since otherwise $1/y \in \bigcap R_P = R$). Choose $x \in (P_1 \cap \dots \cap P_t) \setminus Q$.

Claim. $x^n/y \in \bigcap R_P = R$, for some n . For any n , $x^n/y \in R_P$, $P \neq P_1, \dots, P_t$, since $1/y \in R_P$, $P \neq P_1, \dots, P_t$. Moreover, R_{P_i} is 1-dim quasi-local implies $x^{n_i} \in yR_{P_i}$ for some n_i , $i = 1, \dots, t$. Choose $n = \max \{n_i\}$, and then $x^n \in yR \subset Q$, a contradiction to the choice of x .

1.14. THEOREM. *Let D be a domain, and let $\mathcal{M} = \{\text{minimal prime ideals of } D\}$. Assume $D = \bigcap \{D_P \mid P \in \mathcal{M}\}$, and that $\{D_P \mid P \in \mathcal{M}\}$ has FC. Let $\mathcal{N}$ be a subset of $\mathcal{M}$ such that every $P \in \mathcal{N}$ is maximal. Then $R = \bigcap \{D_Q \mid Q \in \mathcal{M} \setminus \mathcal{N}\}$ is flat over D .*

Proof. By 1.13, the minimal prime ideals of R are exactly $\{QD_Q \cap R \mid Q \in \mathcal{M} \setminus \mathcal{N}\}$. Therefore $PR = R$ for any $P \in \mathcal{N}$, for otherwise some minimal prime of R would necessarily contract to P by the maximality of P . If now Q is any prime of D such that $R \not\subset D_Q$, then $P \subset Q$ for some $P \in \mathcal{N}$, by 1.2. Therefore $P = Q$ and hence $QR = R$, which proves that R is flat over D .

1.15. COROLLARY. *With the notation and hypotheses of 1.14, R and $\{D_P \mid P \in \mathcal{N}\}$ are noetherian if and only if D is noetherian.*

Proof. R is flat over D , so D is noetherian implies R is noetherian. For the converse, observe that for any prime ideal M of D , if $MR = R$, then $M \in \mathcal{N}$ by 1.2; and then apply 1.8. Q.E.D.

Our next theorem considerably sharpens 1.10 in the direction “ D is noetherian implies the irredundant V_i are noetherian”. It and 1.10 should be considered the principal results of this section.

1.16. LEMMA. *Let A be a nonzero finitely generated ideal of an integral domain D with quotient field K . Then $A:A = \{\xi \in K \mid \xi A \subset A\}$ is a ring integral over D .*

Proof. It is immediate that $A:A$ is a ring. To see the integrality, we use a standard argument (see [15a, p. 216]). Let $A = (a_1, \dots, a_n)$ and let $d \in A:A$. Then $da_i = \sum_{j=1}^n y_{ij}a_j$, with $y_{ij} \in D$. Therefore $\det(dZ - Y) = 0$, where Z is the identity matrix and $Y = (y_{ij})$; and this is the desired integral equation for d .

1.17. THEOREM. *Let D, R, V be domains with quotient field K , and suppose that $D = R \cap V = R \cap \bar{D}$, that $D \neq R$, and that V is 1-dim quasi-local with center P on D . If P is finitely generated, then V is a noetherian valuation ring. (Note that if V is integrally closed, then $\bar{D} \subset V$, so $D = R \cap V = R \cap \bar{D}$.)*

Proof. By replacing D, R by their quotient rings with respect to the multiplicative system $D \setminus P$, we are reduced (by 1.1) to considering the case where D is quasi-local with maximal ideal P . Then either $PP^{-1} = D$ or $PP^{-1} = P$.

If $PP^{-1} = D$, then P is invertible. Since D is quasi-local, this implies that P is principal [2a, Theorem 4, p. 148]. Also, since P is contained in the height 1 maximal ideal of V , $\bigcap_{n=1}^{\infty} P^n = 0$ [12, Corollary 1.4, p. 323]. These two properties imply that D is a noetherian valuation ring ([2c, Proposition 9, p. 109]); so $D = V$, and we are done.

If, on the other hand, $PP^{-1} = P$, then $P^{-1} = P:P$. But $P:P$ is integral over D by 1.16, and hence $P^{-1} \subset \bar{D}$. If now $Q \neq P$ is any other prime ideal of D , then $R \subset D_Q$ by 1.2. Therefore if $r \in R$ and if $\partial(r) = \{d \in D \mid dr \in D\}$, then $\partial(r) \not\subset Q$. Thus, $\sqrt{\partial(r)} = P$; so P is finitely generated implies that $P^n \subset \partial(r)$ for some $n \geq 1$. Hence $rP^n \subset D$ and $r \in P^{-n}$ (where by definition $P^{-n} = (P^n)^{-1}$). Choose $r \in R \setminus D$. Then there exists $n \geq 1$ such that $r \in P^{-n}$, $r \notin P^{-n+1}$. Then $rP^{n-1} \subset R$ and $rP^{n-1} \subset P^{-n}P^{n-1} \subset P^{-1} \subset \bar{D}$; so $rP^{n-1} \subset R \cap \bar{D} = D$, and hence $r \in P^{-n+1}$, a contradiction to the choice of n . Q.E.D.

1.18. COROLLARY. Suppose $D = R \cap \{V_i \mid i \in I\}$, where $\{V_i\}_{i \in I}$ is a set of 1-dim quasi-local integrally closed domains having FC. If each V_i is centered on a finitely generated maximal ideal of D , then D is noetherian if and only if R is noetherian.

Proof. Apply 1.17 and 1.10.

The condition $R \cap \bar{D} = D$ of 1.17 cannot be weakened to $R \cap \bar{V}$ is integral over D , as the following example shows.

1.19. Example of domains R, V, D such that $D = R \cap V$, $R \not\subset V$, V is 1-dim quasi-local and the center of V on D is maximal, yet R, D are noetherian and V is not. The R and D will, in fact, be 2-dim local domains with R integral over D . Let $k_0 < k$ be fields with k a finite algebraic extension of k_0 , and let x, y be indeterminates over k ; let $R = k[x, y]_{(x, y)}$ ($= k + M$, where M is the maximal ideal of R); let $\bar{V}$ be the rank 1 valuation ring of $k(x, y)$, trivial on k , defined (using the procedure of [2c, p. 160]) by setting $\bar{V}(x) = 1$, $\bar{V}(y) = \pi$; and let $V = k_0 + N$, where N is the maximal ideal of $\bar{V}$. Note that $\bar{V} = k + N$ and that $R \cap N = M$. Let $D = R \cap V$. Then $D = k_0 + M$, so R is a finite integral extension of D , and hence D is noetherian by Eakin's theorem [3] (see [10] for a simplified proof of this theorem). On the other hand V is not noetherian since $\bar{V}$ is not.

It is possible to construct a more complicated example than the above which has the additional property that $R \not\subset \bar{D}$.

As another application of 1.17 and 1.10, we mention the following corollary, which includes a situation treated in [15, II, p. 328].

1.20. COROLLARY. Let V be a rank 1 valuation ring and R be a quasi-semilocal ring, let $D = R \cap V$, and suppose D, R, V all have the same quotient field. If there exists an element x which is a unit of R and a nonunit of V , then D is noetherian if and only if V, R are noetherian.

Proof. The existence of x implies that V is irredundant. Since every nonunit of D is either a nonunit of R or a nonunit of V , the set of nonunits of D is precisely the union of the finite number of contracted maximal ideals from R and V . Thus, D is quasi-semilocal, and the condition on x then implies that the center of V on D is maximal; so 1.17 and 1.10 apply. Q.E.D.

The converse to 1.17 is false; for it can be seen that the example of [12, Example 5.3, p. 330] gives a domain D which is an irredundant intersection of rank 1 valuation rings, all but one of which are noetherian and such that an infinite number of them have nonfinitely generated center. The existence of Krull rings having minimal primes which are not finitely generated (see for example, [4]) gives another source of counterexamples to the converse of 1.17. However, if one imposes the further condition that the center of V be maximal, then the converse is easily proved. Thus,

1.21. PROPOSITION. *Let R, V, D, P be as in 1.17. If P is maximal and V is a noetherian valuation ring, then P is finitely generated.*

Proof. If $M \neq P$ is a maximal ideal of D , then $D_M \supset R$ by 1.2. Also, $PR = R$ and $V = D_P$ by 1.5. Therefore there exists a finitely generated ideal P_0 of D such that $P_0R = R = PR$ and $P_0V = PV$. Then $P_0D_M = PD_M$ for every maximal ideal of D , so $P_0 = P$. Q.E.D.

It is perhaps worth noting that this proposition yields immediately (and hence provides a simple proof of) the main result, Theorem 3, p. 338, of [5], which asserts that if a domain D has the property that every D_P , P maximal, is a rank 1 discrete valuation ring and $D = \bigcap D_P$ is an irredundant intersection, then D is noetherian.

Let D be an integral domain with quotient field K . We remind the reader that an element $\xi \in K$ is called *almost integral* (or *quasi-integral*) over D if there exists a nonzero $d \in D$ such that $d\xi^i \in D$ for all $i = 1, 2, \dots$. An element which is integral over D is almost integral over D , and the concepts coincide when D is noetherian. (See for example, [2c, p. 20].)

1.22. COROLLARY (TO 1.17). *Let $D \neq R$ be domains with quotient field K . If W is a valuation ring of K such that $D = R \cap W$ and such that W is maximal with respect to this property (i.e. there does not exist a valuation ring U of K such that $U > W$ and $R \cap U = D$), then D is noetherian implies W is noetherian.*

Proof. By 1.17, it suffices to see that W has rank 1. If not, there exists a nontrivial valuation ring $W' > W$. If M' is the maximal ideal of W' , then $M' \subset W$; also $M' \cap D \neq 0$ since D and W' have the same quotient field. Therefore, if $x \in W' \cap R$ and $d \neq 0 \in M' \cap D$, then $dx^n \in M' \cap R \subset D$, $n = 1, 2, \dots$. Thus, x is almost integral over D ; and since D is noetherian, this implies x is integral over D . Therefore $x \in W$, so $W' \cap R = D$, a contradiction to the choice of W . Q.E.D.

The 1-dim assumption in 1.17 is essential, as is seen by the following example. Take $R = k[x, y, z, 1/x]$, where x, y, z are indeterminates over the field k . Let V_1

be the x -adic valuation ring of $k(x, y, z)$, and let V be the composite of V_1 with some nondiscrete valuation ring containing $k[y, z]$ of the residue field ($\cong k(y, z)$) of V_1 . Then $R \cap V = k[x, y, z] = D$ since $R \cap V_1 = k[x, y, z]$, but the maximal ideal of V is not finitely generated since V has a nondiscrete homomorphic image.

We conclude this section with the following intrinsic characterization of R -Krull domains. It is modeled after the corresponding statement for Krull rings in [11, p. 115].

1.23. THEOREM. *Let $D < R$ be integral domains having the same quotient field K . Then D is an R -Krull domain if and only if*

(i) *if P is a minimal prime of D such that $R \not\subseteq D_P$, then D_P is a rational valuation ring, and*

(ii) *if $x \in D$, $x \neq 0$, then $xD = (xR \cap D) \cap \{Q_j \mid j \in J\}$, where $\{Q_j\}_{j \in J}$ is a finite (possibly empty) family of primary ideals for minimal primes P_j of D .*

Proof. ($\Rightarrow$) From 1.6, $\{D_P \mid P \text{ is minimal and } R \not\subseteq D_P\}$ is precisely the set $\mathcal{E}$ of essential R -representatives for D . Thus each such D_P is a rational valuation ring and $D = R \cap \{D_P \mid D_P \in \mathcal{E}\}$. Hence if $x \in D$, $x \neq 0$, then $xD = xR \cap \{xD_P \mid D_P \in \mathcal{E}\}$. Since $\mathcal{E}$ has FC, $xD_P \neq D_P$ for only a finite number of the $D_P \in \mathcal{E}$, say $\{D_{P_j} \mid j \in J\}$. If $Q_j = xD_{P_j} \cap D$, then Q_j is P_j -primary and $xD = (xR \cap D) \cap \{Q_j \mid j \in J\}$.

($\Leftarrow$) Let $\{P_i \mid i \in I\}$ be the set of minimal primes of D such that $R \not\subseteq D_{P_i}$. By assumption each D_{P_i} is a rational valuation ring. Also we have $D_{P_i} < R_{P_i}$. Hence $R_{P_i} = K$, i.e. each nonzero ideal of R meets $D \setminus P_i$. If $x \in D$, $x \neq 0$, and $xD = (xR \cap D) \cap \{Q_j \mid j \in J\} \subset P_i$, then we claim that $P_i \in \{P_j \mid j \in J\}$. For $xR \cap D \not\subseteq P_i$ implies that some $Q_j \subset P_i$. Thus the set $\{D_{P_i} \mid i \in I\}$ has FC. It only remains to show that $D = R \cap \{D_{P_i} \mid i \in I\}$. We note first that if a minimal prime P_j of D is such that $R \subset D_{P_j}$, then $xR \cap D \subset xD_{P_j} \cap D$. Hence we may assume that the minimal primes P_j in (ii) are such that $R \not\subseteq D_{P_j}$. Thus for nonzero $x \in D$, we have from (ii) that $xD = (xR \cap D) \cap \{xD_{P_i} \mid i \in I\}$. If $\xi \in R \cap \{D_{P_i} \mid i \in I\}$ write $\xi = y/x$, $y, x \in D$. Then $y \in xR \cap D$, and $y \in xD_{P_i}$ for each i . Hence $y \in (xR \cap D) \cap \{xD_{P_i} \mid i \in I\} = xD$ and $\xi \in D$. Q.E.D.

2. Generalizations. We shall now generalize some of the results of §1 to the case that the V_i are no longer assumed to be rational valuation rings but instead are merely assumed to be 1-dim quasi-local. Again we begin by sufficiently generalizing the terminology of Krull rings to fit our need. As before, D and R will denote domains with the same quotient field K . If $\mathcal{V} = \{V_i\}_{i \in I}$ is a set of R -representatives for D , $D'(\mathcal{V})$ will denote the domain $R \cap (\bigcap_{i \in I} \overline{V_i})$, which will now play an important role.

2.1. LEMMA. *If $\mathcal{V} = \{V_i\}_{i \in I}$ is a set of R -representatives for D , then $D'(\mathcal{V})$ is almost integral over D (and hence $D'(\mathcal{V})$ is integral over D if D is noetherian).*

Proof. If $\xi \in D'(\mathcal{V})$, then by the FC, $\xi \notin V_i$ for at most finitely many i , say $\xi \notin V_1, \dots, V_n$. Since $\xi \in \overline{V_i}$, ξ is almost integral over V_i ; therefore we can choose

$a_i \neq 0 \in V_i$, and hence in D , such that $a_i \xi^j \in V_i$ for all $j=1, 2, \dots$. Let $d=a_1 \cdots a_n$, and it follows that $d \xi^j \in D$ for all $j=1, 2, \dots$.

2.2. LEMMA. *If $\mathcal{V}$ is a set of R -representatives for D , then $D'(\mathcal{V})$ is R -Krull and integral over D if and only if $\bar{D} \cap R$ is R -Krull. When this occurs, then $D'(\mathcal{V}) = \bar{D} \cap R$.*

Proof. It suffices to show that $D'(\mathcal{V})$ is integral over D implies $D'(\mathcal{V}) = \bar{D} \cap R$ and that $\bar{D} \cap R$ is R -Krull implies $D'(\mathcal{V}) = \bar{D} \cap R$. We always have $D'(\mathcal{V}) \supset \bar{D} \cap R$; and if $D'(\mathcal{V})$ is integral over D , the reverse inclusion is immediate. Suppose then that $\bar{D} \cap R$ is R -Krull. Every element of $D'(\mathcal{V})$ is then an element of R which is almost integral over D by 2.1 and hence which is a fortiori almost integral over $\bar{D} \cap R$. But the rational R -representatives of an R -Krull domain are rational valuation rings and therefore are completely integrally closed, so it follows that any element of R which is almost integral over $\bar{D} \cap R$ is in $\bar{D} \cap R$. Thus, $D'(\mathcal{V}) \subset \bar{D} \cap R$. Q.E.D.

We call D a *generalized R -Krull domain* provided there exists a set $\mathcal{V}$ of R -representatives for D such that $D'(\mathcal{V})$ is an R -Krull domain and is integral over D . Thus, if R and D are domains with quotient field K , then D is a generalized R -Krull domain if there exists a set $\{V_i\}_{i \in I}$ of 1-dim quasi-local domains with quotient field K such that $\{V_i\}_{i \in I}$ has FC, $R \not\subset \bigcap_{i \in I} V_i$, and $D = R \cap (\bigcap_{i \in I} V_i)$ and such that $D'(\mathcal{V}) = R \cap (\bigcap_{i \in I} \bar{V}_i)$ is R -Krull and integral over D . 2.2 shows that when D is a generalized R -Krull domain, the domain $D'(\mathcal{V})$, even though it is defined in terms of a set $\mathcal{V}$ of R -representatives for D , is equal to $\bar{D} \cap R$ and hence does not depend on the choice of the set $\mathcal{V}$. Therefore, in discussing a generalized R -Krull domain, we need only write D' rather than $D'(\mathcal{V})$ for this domain. We remind the reader that R can equal K in these considerations; thus, for example, noetherian domains whose principal ideals have no imbedded components are included in our generalized R -Krull domains.

The difficulty in dealing with a generalized R -Krull domain D is that one no longer has available a set of R -representatives with the good properties of the essential R -representatives of an R -Krull domain. (We illustrate these difficulties later in Examples 2.14 and 2.15.) However, we can single out a set of primes of D which will successfully generalize the notion of essential primes of an R -Krull domain. Thus, we shall say that a prime ideal P of the generalized R -Krull domain D is an *essential prime* of D if every prime of D' lying over P is an essential prime of the R -Krull domain D' . We next give some important characterizations of essential primes of a generalized R -Krull domain, similar to those characterizations of essential primes of an R -Krull domain given in 1.5. 2.4 deals with characterizations which are intrinsic in the sense that they do not involve a set of R -representatives, while 2.6 gives some characterizations which do involve such a set.

We need the following lemma, which will be used in localizing.

2.3. LEMMA. Let $\mathcal{V} = \{V_i\}_{i \in I}$ be a set of R -representatives for the generalized R -Krull domain D , and let $\{V_j\}_{j \in J}$ be those V_i whose center on D does not intersect S . Then $D_S = R_S \cap (\bigcap_{j \in J} V_j)$ and $(D')_S = R_S \cap (\bigcap_{j \in J} \bar{V}_j)$; and hence if $D_S \neq R_S$, then D_S is generalized R_S -Krull and $(D')_S = (D_S)'$.

Proof. The first assertion follows from 1.1 and the second follows from the first and the definitions.

The following intrinsic characterizations of essential primes of a generalized R -Krull domain follow readily from the definitions and the properties of essential primes of an R -Krull domain given in §1.

2.4. PROPOSITION. Let D be a generalized R -Krull domain and let P be a prime ideal of D . The following statements are equivalent:

- (i) Each prime ideal P' of D' lying over P is an essential prime for the R -Krull domain D' .
- (ii) P is a minimal prime of D and $R \not\subseteq D'_P$ for each prime P' of D' lying over P .
- (iii) P is a minimal prime of D and R and D_P have no common overring $< K$.
- (iv) P is a minimal prime of D and no prime of R lies over P .
- (v) P is a minimal prime of D and $R_P = K$.

We now wish to relate the essential primes of the generalized R -Krull domain D to a set $\{V_i\}$ of R -representatives for D . Let P_i denote the center of V_i on D . If P is an essential prime of D , then from 2.4 and 1.2 we have $P \in \{P_i\}$. Hence each essential prime of D is a minimal element of the set $\{P_i\}$. To classify more specifically the essential primes of D in terms of the set $\{P_i\}$, we will make use of the following lemma.

2.5. LEMMA. Let D be a domain having a set of R -representatives $\{V_i\}_{i \in I}$. Let P_i denote the center of V_i on D , let P be a minimal element (with respect to inclusion) of the set $\{P_i\}$, and let $\{V_j\}_{j \in J}$ be those V_i having center P on D . Let $\{P'_\beta\}$ be the set of contractions to $D' = R \cap \{\bar{V}_i \mid i \in I\}$ of the nonzero prime ideals of the $\bar{V}_j$, $j \in J$. Then any prime ideal P' of D' lying over P contains $\bigcap P'_\beta$.

Proof. Let $x \in \bigcap P'_\beta$. Then, for any $j \in J$, $V_j[x]$ is a finite V_j -module, and x is in the Jacobson radical of $V_j[x]$. Therefore, since V_j is 1-dim quasi-local, some power of x is in the nonzero conductor of V_j in $V_j[x]$, and thus is in V_j . Since J is finite by the finiteness condition on R -representatives, there exists then a positive integer m such that $x^m \in V_j$ for all $j \in J$. Therefore $x^m \in R \cap \{V_j \mid j \in J\} = D_P$, the equality following from 1.1. Thus, there exists $s \in D \setminus P$ such that $sx^m \in D \cap P'_\beta = P$. If then P' is any prime of D' lying over P , it follows that $x \in P'$.

2.6. THEOREM. Let D be a generalized R -Krull domain, and let $\{V_i\}$ be a set of R -representatives for D . Let P_i denote the center of V_i on D , and let P be a minimal element (with respect to inclusion) of the set $\{P_i\}$. The following statements are equivalent:

- (1) P is an essential prime of D .
 (2) $R \not\subseteq D'_{P'}$ for each prime ideal P' of D' lying over P .
 (3) Each V_i centered on P is such that R and V_i have no common overring $< K$.
 (4) Each V_i centered on P is such that $\bar{V}_i$ is an intersection of finitely many essential R -representatives for D' .

Moreover, if P is an essential prime of D , then $D_P = \bigcap \{V_i \mid V_i \text{ is centered on } P\}$.

Proof. (1) $\Rightarrow$ (2) and (1) $\Rightarrow$ (3) are immediate from 2.4.

(1) $\Rightarrow$ (4). If P is essential and V_i is centered on P , then each maximal ideal M of $\bar{V}_i$ lies over P in D and hence lies over an essential prime $M \cap D'$ of D' . Therefore $D'_{M \cap D'}$ is an essential R -representative for D' ; and since it is also then a rational valuation ring, $D'_{M \cap D'} = (\bar{V}_i)_M$. But $\bar{V}_i$ is the intersection of its localizations at maximal ideals, so $\bar{V}_i$ is an intersection of essential R -representatives for D' . That there are only finitely many such R -representatives follows from the finiteness condition on R -representatives, since each of these representatives is centered on P in D .

(4) $\Rightarrow$ (3). If R and V_i have a common overring $< K$ for some V_i centered on P , then they have a common valuation overring $W < K$. But $W \supset \bar{V}_i$; so if $\bar{V}_i$ is a finite intersection of rational valuation rings, then W must be one of them [11, p. 38]. But then W would be an essential R -representative for D' , a contradiction to $R \subset W$.

We need the following *observation* for the remainder of the proof: If P' is a prime ideal of D' lying over P such that $R \not\subseteq D'_{P'}$, then P' is an essential prime of D' . For, by 1.2, P' contains an essential prime Q' of D' , and then $Q = Q' \cap D \subset P$. If $Q < P$, the assumption that P is minimal in the set of centers $\{P_i\}$ implies $(V_i)_Q = K$ for all i . But then $R \subset R_Q = D_Q \subset D'_{Q'}$, the equality following from 2.3; and this contradicts the fact that Q' is an essential prime of D' . Thus, $Q = P$; and therefore $Q' = P'$ since D' is integral over D .

(2) $\Rightarrow$ (1). In view of 2.4, it suffices to show that P is a minimal prime of D . Since D' is integral over D , we need only show that each prime P' of D' lying over P is minimal. By the above observation, any such P' is an essential prime of D' and hence is minimal.

(3) $\Rightarrow$ (1). Let $\{V_j\}_{j \in J}$ be those V_i having center P on D , and let $\{P'_\beta\}$ denote the set of prime ideals of D' that are contractions of nonzero primes of the $\bar{V}_j$, $j \in J$. (3) assures that $R \not\subseteq D'_{P'_\beta}$ for each P'_β ; so by the above observation, each P'_β must be an essential prime of the R -Krull domain D' . Since D' has only finitely many essential primes lying over P , $\{P'_\beta\}$ is finite. By 2.5, each prime of D' lying over P must then be in the set $\{P'_\beta\}$, i.e. P is an essential prime of D .

For the final assertion of the theorem, observe that, by 2.4, P is minimal and $R_P = K$ for any essential prime P of D . Hence, by 2.3,

$$D_P = \bigcap \{V_i \mid V_i \text{ has center } P \text{ on } D\}. \quad \text{Q.E.D.}$$

We are now ready to examine noetherian properties of D , R and the V_i .

2.7. THEOREM. *Let D be a noetherian domain having a set $\{V_i\}_{i \in I}$ of R -representatives such that $R \not\subseteq \bigcap_{i \in I} \bar{V}_i$. Then D is a generalized R -Krull domain. Moreover, if $V \in \{V_i\}$ is centered on an essential prime of D , then V is noetherian. Thus, for example, if for each i , R and V_i have no common overring $< K$ and there are no proper containment relations among the centers P_i of the V_i on D , then all the V_i are noetherian.*

Proof. If D is noetherian, then the integral closure $\bar{D}$ of D is a Krull domain [11, Theorem 33.10, p. 118]. Hence by 2.2, D is a generalized R -Krull domain. If V is centered on an essential prime P of D , then $D_P \subset V \subset K$ and by 2.4, P is a minimal prime of D . Thus V lies between a 1-dim noetherian domain and its quotient field. By the Krull-Akizuki theorem [11, Theorem 33.2, p. 115], V is noetherian. The last assertion follows from 2.6.

2.8. THEOREM. *Let D be a generalized R -Krull domain and let $\{V_i\}$ be a set of R -representatives for D . If each V_i is centered on an essential maximal ideal of D , then R is a flat D -module. In this case, D is noetherian if and only if R and every finite intersection of the V_i are noetherian.*

Proof. Let P be a prime ideal of D . If $R \not\subseteq D_P$, then by 1.2, P contains and hence must equal the center of some V_i on D . Hence by 2.4, no prime of R lies over P . Since P is maximal, we have $PR = R$. Thus R is a flat D -module [14, p. 795]. Moreover, 2.6 implies that

$$D_P = \bigcap \{V_i \mid V_i \text{ has center } P \text{ on } D\}.$$

Hence if R and every finite intersection of the V_i are noetherian then, by 1.8, D is noetherian. Conversely, D noetherian implies R is noetherian since R is a flat D -module contained in the quotient field of D . To show that D noetherian implies that every finite intersection of the V_i is noetherian, it will suffice by the Krull-Akizuki theorem to show that if $P_1, \dots, P_n$ are essential primes of D , then $\bigcap_{i=1}^n D_{P_i}$ is 1-dim and noetherian. By 1.9, $\bigcap_{i=1}^n D_{P_i}$ is noetherian and since each nonunit of $\bigcap_{i=1}^n D_{P_i}$ is contained in $\bigcup_{i=1}^n P_i D_{P_i}$ we see that the contractions of the $P_i D_{P_i}$ are the only nonzero primes of $\bigcap_{i=1}^n D_{P_i}$. Hence $\bigcap_{i=1}^n D_{P_i}$ is 1-dim. Q.E.D.

In order to apply 2.8 in showing that R and the V_i noetherian imply D is noetherian, it is useful to have sufficient conditions that every finite intersection of the V_i be noetherian. We note the following consequence of the approximation theorem for independent valuation rings.

2.9. PROPOSITION. *Let $\{W_i\}_{i=1}^n$ be a family of 1-dim quasi-local domains with quotient field K such that each $\bar{W}_i$ is a finite intersection of valuation rings. Assume that for $i \neq j$, W_i and W_j are not dominated by a common valuation ring. Then $T = \bigcap_{i=1}^n W_i$ is 1-dim, has quotient field K , and if Q_i is the center of W_i on T , then the Q_i are the only nonzero primes of T and $T_{Q_i} = W_i$ for each i .*

Proof. The nonunits of T are contained in $\bigcup_{i=1}^n Q_i$ and hence every maximal ideal is some Q_i . Thus, all assertions will follow if we show $T_{Q_i} = W_i$ for each i . By the approximation theorem [15b, p. 45], there exists $y \in K$ having positive value in each valuation dominating W_i and negative value in each valuation dominating W_j for $j \neq i$. Thus $W_i[y]$ is a finite W_i -module with y in the Jacobson radical of $W_i[y]$. Hence for some positive integer m , y^m is in the conductor of W_i in $W_i[y]$ and therefore in the maximal ideal of W_i . Thus $1 + y^m$ is a unit of W_i having negative value in each valuation dominating W_j for $j \neq i$. Let $x = 1/(1 + y^m)$. Then $W_j[x]$ is a finite W_j -module, and for $j \neq i$, x is contained in the Jacobson radical of $W_j[x]$. We conclude that for some positive integer t , $x^t \in (\bigcap_{j \neq i} Q_j) \setminus Q_i$. Hence $(W_j)_{Q_i} = K$ for $j \neq i$. Thus by 1.1, we have $T_{Q_i} = \bigcap_{j=1}^n (W_j)_{Q_i} = W_i$. Q.E.D.

2.10. COROLLARY. Let $\{W_i\}_{i=1}^n$ be a family of 1-dim (noetherian) local domains with quotient field K such that for $i \neq j$, W_i and W_j are not dominated by a common valuation ring. Then $T = \bigcap_{i=1}^n W_i$ is a 1-dim (noetherian) semilocal domain.

2.11. COROLLARY. Let D be a generalized R -Krull domain. Assume that there exists a family $\{V_i\}$ of noetherian R -representatives for D such that each V_i is centered on a maximal ideal of D . If for each i , R and V_i , and V_j and V_i , for $j \neq i$, have no common overring $< K$, then R is noetherian implies D is noetherian.

Proof. By 2.10 every finite intersection of the V_i is noetherian. By 2.6 each V_i is centered on an essential prime of D . Hence by 2.8, R is noetherian implies D is noetherian.

If we restrict ourselves to the question of when D is noetherian implies R is noetherian, we can replace the assumption in 2.8 that each V_i is centered on an essential prime of D by an assumption only involving the V_i . We use $(V_i : \bar{V}_i)$ to denote the conductor of V_i in $\bar{V}_i$, i.e. $(V_i : \bar{V}_i) = \{a \in V_i \mid a\bar{V}_i \subset V_i\}$. Note that if $\bar{V}_i$ is a finite V_i -module, then $(V_i : \bar{V}_i) \neq 0$.

2.12. LEMMA. Let D be a domain with a set of R -representatives $\{V_i\}_{i \in I}$. Assume that I is finite, say $I = \{1, \dots, n\}$, that each V_i is centered on a maximal ideal of D and that $C_i = (V_i : \bar{V}_i) \neq 0$, $i = 1, \dots, n$. Then D is noetherian implies R is noetherian.

Proof. Let $D' = R \cap \{\bar{V}_i \mid i \in I\}$. $C = \bigcap_{i=1}^n (C_i \cap D)$ is a nonzero ideal of D and $CD' \subset D$; thus D' is a finite D -module and hence noetherian. In particular, R is noetherian if $R \subset \bigcap_{i=1}^n \bar{V}_i$; so we may assume $R \not\subset \bigcap_{i=1}^n \bar{V}_i$. Then since D is noetherian, D' is an R -Krull domain by 2.7. Let $\mathcal{E}'$ denote the essential R -representatives of D' and let P_i denote the center of V_i on D . For any $D'_{P'} \in \mathcal{E}'$, $P_i \subset P' \cap D$ for some i , by 1.2. It follows from our hypothesis that $P' \cap D$ is maximal; and hence P' is maximal, since D' is integral over D . Thus, D' is noetherian implies that R is noetherian by 1.10.

2.13. THEOREM. Let D be a domain having a set of R -representatives $\{V_i\}_{i \in I}$. Assume that each V_i is centered on a maximal ideal P_i of D and that $(V_i : \bar{V}_i) \neq 0$ for all $i \in I$. Then D is noetherian implies that R is noetherian.

Proof. By Cohen's theorem, it suffices to see that any prime ideal Q_0 of R is finitely generated. Let $Q = Q_0 \cap D$, and let A be a finite set of generators for Q . If $Q \subset P_i$ for some i , then Q is contained in only finitely many of the P_i , say $P_1, \dots, P_n$; and in this case, if $S = D \setminus \bigcup_{i=1}^n P_i$, then R_S is noetherian by 1.1 and 2.12. Thus, if $Q \subset P_i$ for some i , then we can find a finite subset B of Q_0 such that $BR_S = Q_0R_S$. Let $C = A$ if $Q \not\subset P_i$ for all $i \in I$, and otherwise let $C = A \cup B$. *Claim:* $Q_0 = CR$. It suffices to show that $Q_0R_{M_0} \subset CR_{M_0}$ for every maximal ideal M_0 of R . Let $M = M_0 \cap D$. If $R \subset D_M$, then $R_{M_0} = D_M$ and hence $Q_0R_{M_0} = QD_M = AD_M \subset CR_{M_0}$. If $R \not\subset D_M$, then $P_i \subset M$ for some $i \in I$ by 1.2; therefore $P_i = M$ since P_i is assumed maximal. If $Q \not\subset M$, then $Q_0R_{M_0} = R_{M_0} = AR_{M_0} = CR_{M_0}$. On the other hand, if $Q \subset M = P_i$, then $R_S \subset R_{M_0}$; and hence $Q_0R_S = BR_S$, which implies $Q_0R_{M_0} = BR_{M_0} \subset CR_{M_0}$. Q.E.D.

We conclude with two examples which illustrate the difficulties that can arise in considering generalized R -Krull domains.

2.14. *Example of a generalized R -Krull domain $D = R \cap V_1 \cap V_2$ such that V_2 is irredundant but its center on D is not minimal.* Let $k_0 < k$ be fields with k algebraic over k_0 , let x be an indeterminate over k and y be an element of $k[[x]]$ of strictly positive order such that x, y are algebraically independent over k ; let $R = k[x, y]_{(x, y)}[1/x]$; let V_1 be the x -adic valuation ring of $k(x, y)$; let $\bar{V}_2 = k[[x]] \cap k(x, y)$ ($= k + M_2$, where M_2 is the maximal ideal of the valuation ring $\bar{V}_2$); let $V_2 = k_0 + M_2$; and let $D = R \cap V_1 \cap V_2$. Note that $k[x, y]_{(x, y)} = k + M$, where M is the maximal ideal of $k[x, y]_{(x, y)}$. Moreover, $k + M \subset \bar{V}_2$ and $M_2 \cap k[x, y]_{(x, y)} = M$. Also, $R \cap V_1 = k[x, y]_{(x, y)}$. Therefore $D = k[x, y]_{(x, y)} \cap V_2 = k_0 + M$, and $D' = k[x, y]_{(x, y)} \cap \bar{V}_2 = k + M$. Thus, D' is integral over D and $R \not\subset D'_M = D'$. Therefore D is generalized R -Krull. However, M is not a minimal prime ideal of D since M is not a minimal prime of D' . Note also that no two of the rings R, V_1, V_2 have a common overring $< K$.

The following example is taken from [13]. It justifies the hypothesis that D' is integral over D in our results on " R and the V_i noetherian imply D is noetherian".

2.15. *Example of a domain $D = R \cap V$ with R and V noetherian, V centered on a maximal ideal of D , R and V having no common overring $< K$, and yet such that $D' = R \cap \bar{V}$ is not integral over D . Hence, by 2.1, D is not noetherian.* Let k be a field, and let x, y be indeterminates over k ; let $R = k[x, y, x/y]$ and let V_y be the y -adic valuation ring of $k(x, y)$. Note that $V_y = k(x) + M$, where M is the maximal ideal of V_y . Let $V = k(\xi) + M$, where ξ is chosen by means of the following lemma.

LEMMA. *There exists $\xi \in k(x) \setminus k$ such that $k(\xi) \cap k[x] = k$.*

We shall delay proving the lemma and proceed with the example.

Let $D = R \cap V$, and let $D' = R \cap V_y$. (Note that $V_y = \bar{V}$.) *Claim:* $D' = k[x, y]$. For, the essential valuations of $k[x, y]$ are exactly the p -adics, where p is an irreducible polynomial in $k[x, y]$. Therefore $k[x, y] = \bigcap V_p$, where V_p denotes the p -adic valuation ring. But the only V_p which does not contain R is V_y . Therefore $R \cap V_y \subset \bigcap V_p = k[x, y]$. This proves the claim.

Now let P denote the ideal $y \cdot k[x, y]$. Then $D = R \cap V = k[x, y] \cap V = k[x, y] \cap (k(\xi) + M) = (k[x] + P) \cap (k(\xi) + M) = (k[x] \cap k(\xi)) + P = k + P$, and D' is certainly not integral over D since both $y \cdot D'$ and $(x, y) \cdot D'$ lie over P ; hence, by 2.1, D is not noetherian.

We note that P is not minimal since, for example, $\{f \cdot y \cdot (x + y) \mid f \in k[x, y]\}$ is easily seen to be a prime ideal of D which is $< P$. Also, R is not flat over D' , since $(x, y) \cdot R \neq R$ and $R \not\subseteq D'_{(x, y)}$, and hence R is a fortiori not flat over D .

Proof of the above lemma. Let $\xi = (x^2 + 1)/x = 1/x + x$. Then x is a root of $Z^2 - \xi Z + 1$, so $[k(x) : k(\xi)] = 2$. Let W' denote the $1/x$ -adic valuation ring of $k(x)$ and W the x -adic valuation ring of $k(x)$. W and W' are then conjugates over $k(\xi)$ and hence $W \cap k(\xi) = W' \cap k(\xi)$. But

$$k[x] = \bigcap \{W_\alpha \mid W_\alpha \text{ is a valuation ring of } k(x) \text{ over } k \text{ distinct from } W'\}.$$

Hence each valuation ring of $k(\xi)$ over k is dominated by some W_α . Thus $k[x] \cap k(\xi) = (\bigcap_\alpha W_\alpha) \cap k(\xi) = k$.

REFERENCES

1. T. Akiba, a) *Remarks on generalized rings of quotients*, Proc. Japan Acad. **40** (1964), 801–806. MR **31** #4807.
- b) *Remarks on generalized rings of quotients*. II, J. Math. Kyoto Univ. **5** (1965), 39–44. MR **34** #185.
2. N. Bourbaki, a) *Algèbre commutative*. Chap. 1: *Modules plats*. Chap. 2: *Localisation*, Actualités Sci. Indust., no. 1290, Hermann, Paris, 1961. MR **36** #146.
- b) *Algèbre commutative*. Chap. 3: *Graduations, filtrations et topologies*. Chap. 4: *Idéaux premiers associés et décomposition primaire*, Actualités Sci. Indust., no. 1293, Hermann, Paris, 1961. MR **30** #2027.
- c) *Algèbre commutative*. Chap. 5: *Entièrs*. Chap. 6: *Valuations*, Actualités Sci. Indust., no. 1308, Hermann, Paris, 1964. MR **33** #2660.
3. P. M. Eakin, Jr., *The converse to a well known theorem on Noetherian rings*, Math. Ann. **177** (1968), 278–282. MR **37** #7360.
4. P. Eakin and W. Heinzer, *Non-finiteness in finite dimensional Krull domains*, J. Algebra **14** (1970), 333–340.
5. R. W. Gilmer, Jr., *OVERRINGS OF PRÜFER DOMAINS*, J. Algebra **4** (1966), 331–340. MR **34** #2609.
6. W. Heinzer, *On Krull overrings of an affine ring*, Pacific J. Math. **29** (1969), 145–149. MR **39** #5537.
7. W. Heinzer, J. Ohm and R. L. Pendleton, *On integral domains of the form $\bigcap D_P$, P minimal*, J. Reine Angew. Math. **241** (1970), 147–159. MR **41** #8393.
8. D. Lazard, a) *Thesis*, Paris, 1968.
- b) *Autour de la platitude*, Bull. Soc. Math. France **97** (1969), 81–128. MR **40** #7310.
9. K. R. Nagarajan, *Groups acting on Noetherian rings*, Nieuw Arch. Wisk. (3) **16** (1968), 25–29. MR **37** #5202.
10. M. Nagata, *A type of subrings of a noetherian ring*, J. Math. Kyoto Univ. **8** (1968), 465–467. MR **38** #4460.
11. ———, *Local rings*, Interscience Tracts in Pure and Appl. Math., no. 13, Interscience, New York, 1962. MR **27** #5790.
12. J. Ohm, *Some counter examples related to integral closure in $D[[x]]$* , Trans. Amer. Math. Soc. **122** (1966), 321–333. MR **34** #2613.

13. J. Ohm and R. Pendleton, *Weak Krull rings* (unpublished).
14. F. Richman, *Generalized quotient rings*, Proc. Amer. Math. Soc. **16** (1965), 794–799. MR **31** #5880.
15. O. Zariski and P. Samuel, *Commutative algebra*. Vols. I, II, University Series in Higher Math., Van Nostrand, Princeton, N. J., 1957, 1960. MR **19**, 833; MR **22** #11006.

DEPARTMENT OF MATHEMATICS, PURDUE UNIVERSITY, LAFAYETTE, INDIANA 47907*

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA 70803

* Current address of both authors.