

# UNIQUENESS OF HAAR SERIES WHICH ARE $(C, 1)$ SUMMABLE TO DENJOY INTEGRABLE FUNCTIONS

BY

WILLIAM R. WADE

ABSTRACT. A Haar series  $\sum \alpha_k \chi_k$  satisfies Condition H if  $\alpha_k \chi_k / k \rightarrow 0$  uniformly as  $k \rightarrow \infty$ . We show that if such a series is  $(C, 1)$  summable to a Denjoy integrable function  $f$ , except perhaps on a countable subset of  $[0, 1]$ , then that series must be the Denjoy-Haar Fourier series of  $f$ .

1. **Introduction.** The Haar functions  $\chi_0, \chi_1, \dots$  are a complete orthonormal system in the Hilbert space  $L^2[0, 1]$ . For the purposes of this paper we need only recall that  $\chi_0$  is identically 1 and that, given any positive integer  $n = 2^m + k$ , where  $0 \leq k < 2^m$ , the corresponding Haar function  $\chi_n$  takes on the value  $+\sqrt{2^m}$  on the open interval

$$(1) \quad \Delta(1, n) \equiv (2k/2^{m+1}, [2k+1]/2^{m+1}),$$

and takes on the value  $-\sqrt{2^m}$  on the open interval

$$(2) \quad \Delta(2, n) \equiv ([2k+1]/2^{m+1}, [2k+2]/2^{m+1}).$$

Furthermore, the support of that  $n$ th Haar function is precisely the closure of the union of intervals (1) and (2):

$$\text{Supp}[\chi_n] = [k/2^m, [k+1]/2^m].$$

The Denjoy integral (see [5, pp. 84–85]),  $(D) \int_a^b$ , is more general than either Lebesgue's integral or the improper Riemann integral.

The  $D$ -Haar Fourier series of a Denjoy integrable function  $f$  is a Haar series  $S(x) = \sum_{k=0}^{\infty} \alpha_k \chi_k(x)$  which is related to  $f$  by the following formula:

$$(3) \quad \alpha_k = (D) \int_0^1 f(x) \chi_k(x) dx.$$

If (3) holds and  $f$  is also Lebesgue integrable, then  $S$  is simply the Haar Fourier series of  $f$ .

---

Presented to the Society, January 18, 1972; received by the editors February 22, 1972.  
AMS (MOS) subject classifications (1969). Primary 4220, 4215, 4210; Secondary 4032, 4042.

Key words and phrases. Harr functions, Denjoy integral,  $(C, 1)$  summable.

Copyright © 1973, American Mathematical Society

The harmonic analysis of Haar series is greatly simplified by the following property of Haar Fourier series:

The Haar Fourier series of any Lebesgue integrable function  $f$  converges to  $f$  almost everywhere on  $[0, 1]$ .

The standard proof of this fact [1, pp. 47–50] uses only one consequence of the Lebesgue integrability of  $f$ : the derivative of an indefinite Lebesgue integral is equal to the integrand almost everywhere. Since this property is also shared by Denjoy's integral, that same proof will establish that

(4) *The D-Haar Fourier series of  $f$  converges to  $f$  a.e.*

2. **The uniqueness theorem.** In this section we state the uniqueness theorem proved in §4 and relate it to the research presented in [2] and [4].

A Haar series  $\sum_{k=0}^{\infty} \alpha_k \chi_k(x)$  satisfies *Condition G* if, given any  $t_0 \in [0, 1]$ ,

$$(5) \quad \lim_{j \rightarrow \infty} \alpha_{K_j} / \chi_{K_j}(t_0) = 0,$$

where  $K_1, K_2, \dots$  are all those indices  $p$  for which  $\chi_p(t_0) \neq 0$ . Lemma 3 of [2] shows that a D-Haar Fourier series always satisfies *Condition G*.

A Haar series  $\sum_{k=0}^{\infty} \alpha_k \chi_k(x)$  satisfies *Condition H* if

$$(6) \quad \lim_{n \rightarrow \infty} \alpha_n \chi_n(x) / n = 0 \quad \text{uniformly for } x \in [0, 1].$$

*Condition G* is equivalent to supposing that the limit in (6) exists pointwise, since if  $\chi_p(t_0) \neq 0$  then  $p \geq |\chi_p^2(t_0)| \geq p/8$ . Thus *Condition H* can be viewed as the uniform analogue of *Condition G*.

F. G. Arutjunjan [2] has shown that if a Haar series  $S$  satisfying *Condition G* converges, except perhaps on a countable subset of  $[0, 1]$ , to a Denjoy integrable function  $f$ , then  $S$  must be the D-Haar Fourier series of  $f$ .

We shall denote the  $n$ th partial  $(C, 1)$  sum of a Haar series  $S(x) = \sum_{k=0}^{\infty} \alpha_k \chi_k(x)$  by

$$(7) \quad \sigma_n(S; x) \equiv \frac{S_1(x) + \dots + S_{n+1}(x)}{n+1} \equiv \sum_{k=0}^n \left(1 - \frac{k}{n+1}\right) \alpha_k \chi_k(x)$$

where  $S_{n+1}(x) = \sum_{k=0}^n \alpha_k \chi_k(x)$ . If the sequence displayed in (7) converges at  $x$  then  $S$  is said to be  $(C, 1)$  summable at  $x$ . If  $S$  converges at  $x$  it is automatically  $(C, 1)$  summable at  $x$ ; the converse of this statement is false.

The question motivating this research was: Does Arutjunjan's result still hold if  $S$  is only  $(C, 1)$  summable off that countable set?

If  $S$  also satisfies *Condition H* (loosely, if  $S$  satisfies *Condition G* uniformly) the answer to this question is yes and is obtained as a corollary to the following result, which is proved in §4.

**Theorem.** *Let*

$$(8) \quad S(x) = \sum_{k=0}^{\infty} \alpha_k X_k(x)$$

*be a Haar series satisfying Condition H such that*

$$(9) \quad \limsup_{n \rightarrow \infty} |\sigma_n(S; x)| < \infty$$

*for all but countably many  $x$ 's in  $[0, 1]$ . Suppose further that  $f$  is a Denjoy integrable function such that*

$$(10) \quad \lim_{n \rightarrow \infty} \sigma_n(S; x) = f(x) \quad \text{a.e. in } [0, 1].$$

*Then  $S$  is the D-Haar Fourier series of  $f$ .*

This result is already known in the case that  $f$  is Lebesgue integrable (see [4]), but the techniques used here to establish Lemma 4 in the next section are radically different and substantially more complex due to the fact that a function can be conditionally Denjoy integrable.

### 3. Fundamental lemmas.

**Lemma 1.** *Let  $S$  be a Haar series satisfying Condition G and  $\Delta(i_0, k_0)$  be an interval of the form (1) or (2) such that*

$$(12) \quad S_{k_0+1}(x) \neq 0 \quad \text{for } x \in \Delta(i_0, k_0).$$

*Then given any  $t_0 \in [0, 1]$  there is an interval  $\Delta(i'_0, k'_0)$  of the form (1) or (2) such that  $t_0$  does not lie in the closure of  $\Delta(i'_0, k'_0)$  and such that*

$$(13) \quad S_{k'_0+1}(x) \neq 0 \quad \text{for } x \in \Delta(i'_0, k'_0).$$

This lemma was proved on pp. 225–226 of [4].

**Lemma 2.** *Let  $f$  be a Denjoy integrable function and  $\mathfrak{D}$  be any collection of nonoverlapping subintervals of a fixed interval  $\Delta(i_0, k_0)$  of the form (1) or (2). Define the function  $f^*$  by*

$$(14) \quad \begin{aligned} f^*(x) &= f(x) \quad \text{if } x \in \Delta(i_0, k_0) \sim \bigcup \mathfrak{D}, \\ &= \frac{1}{|\Delta|} (D) \int_{\Delta} f(t) dt \quad \text{if } x \in \Delta \in \mathfrak{D}. \end{aligned}$$

*Then the  $(k_0 + 1)$ st partial sum of the D-Haar Fourier series  $T$  of  $f$  can be written as*

$$T_{k_0+1}(x) = \frac{1}{|\Delta(i_0, k_0)|} (D) \int_{\Delta(i_0, k_0)} f^*(t) dt,$$

*for any  $x \in \Delta(i_0, k_0)$ .*

This lemma was proved on pp. 338–339 of [2].

**Lemma 3.** Let  $f$  be a Denjoy integrable function,  $T$  be its  $D$ -Haar Fourier series and  $\mathfrak{D}$  be any collection of nonoverlapping subintervals of a fixed interval  $\Delta(i_0, k_0)$  of the form (1) or (2). Suppose further that  $S^*$  is any Haar series whose partial  $(C, 1)$  sums satisfy

$$(15) \quad \lim_{n \rightarrow \infty} \int_{\Delta(i_0, k_0)} \sigma_n(S^*; t) dt = (D) \int_{\Delta(i_0, k_0)} f^*(t) dt,$$

where  $f^*$  is defined by (14). Then  $S_{k_0+1}^*(x) \equiv T_{k_0+1}(x)$  for  $x \in \Delta(i_0, k_0)$ .

**Proof.** The integral of a Haar function is zero, and the support of each Haar function  $\chi_{k_0+1}, \chi_{k_0+2}, \dots$  is either a subset of  $\Delta(i_0, k_0)$  or disjoint from it. Consequently, if  $n > k_0$ ,

$$(16) \quad \begin{aligned} & \int_{\Delta(i_0, k_0)} \sigma_n(S^*; t) dt \\ & \equiv \frac{k_0 + 1}{n + 1} \int_{\Delta(i_0, k_0)} \sigma_{k_0}(S^*; t) dt + \frac{n - k_0 + 1}{n + 1} \int_{\Delta(i_0, k_0)} S_{k_0+1}^*(t) dt. \end{aligned}$$

But  $\chi_0, \chi_1, \dots, \chi_{k_0}$  are all constant on  $\Delta(i_0, k_0)$  so the last summand of (16) is simply

$$\frac{n - k_0 + 1}{n + 1} |\Delta(i_0, k_0)| S_{k_0+1}^*(x)$$

for any  $x \in \Delta(i_0, k_0)$ . Solving (16) for  $S_{k_0+1}^*(x)$  we then obtain

$$\begin{aligned} S_{k_0+1}^*(x) &= \frac{n + 1}{n - k_0 + 1} \cdot \frac{1}{|\Delta(i_0, k_0)|} \int_{\Delta(i_0, k_0)} \sigma_n(S^*; t) dt \\ &\quad - \frac{k_0 + 1}{n - k_0 + 1} \cdot \frac{1}{|\Delta(i_0, k_0)|} \int_{\Delta(i_0, k_0)} \sigma_{k_0}(S^*; t) dt. \end{aligned}$$

Taking the limit of both sides of this equation as  $n \rightarrow \infty$  and applying (15) results in

$$S_{k_0+1}^*(x) = \frac{1}{|\Delta(i_0, k_0)|} (D) \int_{\Delta(i_0, k_0)} f^*(t) dt,$$

for any  $x \in \Delta(i_0, k_0)$ . According to Lemma 2, this is the value of  $T_{k_0+1}$  over  $\Delta(i_0, k_0)$ .

**Lemma 4.** Let  $f$  be Denjoy integrable,  $T$  be its Haar Fourier series and suppose that  $S(x) = \sum \alpha_k \chi_k(x)$  is a Haar series satisfying Condition H which is  $(C, 1)$  summable to  $f$  almost everywhere. Suppose further that on some fixed interval  $\Delta(i_0, k_0)$  of the form (1) or (2) that

$$(17) \quad S_{k_0+1}(x) \neq T_{k_0+1}(x) \quad \text{for } x \in \Delta(i_0, k_0).$$

Then given any  $M > 0$  there is an interval  $\Delta(i'_k, k'_0)$  contained in  $\Delta(i_0, k_0)$  such that  $|\sigma_{k'_0}(S; x)| > M$  for  $x \in \Delta(i'_0, k'_0)$  and such that  $T_{k'_0+1}$  and  $S_{k'_0+1}$  satisfy (17) on  $\Delta(i'_0, k'_0)$ .

**Proof.** We shall prove this lemma in six steps:

I. We begin by supposing the lemma is false; i.e., that there is an  $M_0$  such that if  $\Delta(j, n)$  is any subinterval of  $\Delta(i_0, k_0)$  of the form (1) or (2) then

$$(18) \quad |\sigma_n(S; x)| > M_0 \quad \text{for } x \in \Delta(j, n) \\ \text{implies } S_{n+1}(x) \equiv T_{n+1}(x) \quad \text{for } x \in \Delta(j, n).$$

II. We next shall use Lemma 1 to show that it is no loss of generality to suppose that  $|\alpha_n \chi_n(x)/n| < M_0$  for  $n \geq k_0$ .

III. We shall then construct a "maximal" class of intervals

$$\mathcal{D} \equiv \{\Delta(i_1, \rho_1), \Delta(i_2, \rho_2), \dots\}$$

on which  $S_{\rho_k+1}$  and  $T_{\rho_k+1}$  are identically equal.

IV. Next we shall construct a subseries  $S^*$  of  $S$  such that

(a)  $S^*$  is  $(C, 1)$  summable to the function  $f^*$  almost everywhere, where  $f^*$  is defined with respect to the class  $\mathcal{D}$  chosen in III by (14);

(b)  $S^*$  and  $S$  are identical in  $\Delta(i_0, k_0) \sim \mathcal{D}$ ;

(c)  $S_{k_0+1}^*(x) \equiv S_{k_0+1}(x)$  for  $x \in \Delta(i_0, k_0)$ .

V. We shall then show that the partial  $(C, 1)$  sums of this series  $S^*$  are bounded by  $13M_0$  by showing that if they are not, we are lead to a contradiction of the construction of  $S^*$  in step IV.

VI. Finally, we shall use IV and V to lead to the ultimate contradiction. Indeed, since any  $(C, 1)$  partial sum of a Haar series is Lebesgue integrable, we can use IV(a) and V to conclude that  $f^*$  is Lebesgue integrable and that (15) holds. Consequently, by Lemma 3,  $S_{k_0+1}^* \equiv T_{k_0+1}$  on  $\Delta(i_0, k_0)$ . But, by IV(c), this identity also implies  $S_{k_0+1} \equiv T_{k_0+1}$  on  $\Delta(i_0, k_0)$ . Since this and hypothesis (17) of this lemma are incompatible, assumption I was false; i.e., the proof of the lemma is complete by contradiction.

What remains, then, is to execute steps II through V:

II. By (6) we choose an integer  $2^Q$  so large that

$$(19) \quad |\alpha_n \chi_n(x)/n| < M_0 \quad \text{whenever } n \geq 2^Q$$

and for any  $x \in [0, 1]$ .

Using Lemma 1 successively on the points  $t_0 = 1/2^Q$ ,  $t_0 = 2/2^Q, \dots, t_0 = (2^Q - 1)/2^Q$  and the series  $S - T$ , we may suppose with no loss of generality that  $|\Delta(i_0, k_0)| < 1/2^Q$ . This fact together with (19) will assure us that

$$(20) \quad |\alpha_n \chi_n(x)/n| < M_0 \quad \text{for } n \geq k_0$$

and for any  $x \in [0, 1]$ .

III. If there is no subinterval  $\Delta(i_1, \rho_1)$  of  $\Delta(i_0, k_0)$  such that  $S_{\rho_1+1} \equiv T_{\rho_1+1}$  on  $\Delta(i_1, \rho_1)$  then set  $\mathcal{D} = \emptyset$ . Otherwise let  $\Delta(i_1, \rho_1)$  be the first such interval of the form (1) or (2).

Suppose that we have either terminated this process or have managed to choose  $N-1$  intervals  $\Delta(i_1, \rho_1), \dots, \Delta(i_{N-1}, \rho_{N-1})$ . If there is no subinterval  $\Delta(i_n, \rho_n)$  of  $\Delta(i_0, k_0)$  disjoint from  $\bigcup_{l=1}^{N-1} \Delta(i_l, \rho_l)$  such that  $S_{\rho_n+1} \equiv T_{\rho_n+1}$  then set

$$\mathcal{D} = \{\Delta(i_l, \rho_l): l = 1, 2, \dots, N-1\}.$$

Otherwise let  $\Delta(i_N, \rho_N)$  be the first such interval of the form (1) or (2).

If this process can be continued indefinitely, set

$$\mathcal{D} = \{\Delta(i_l, \rho_l): l = 1, 2, \dots\}.$$

Clearly  $\mathcal{D}$  is a collection of nonoverlapping intervals of the form (1) or (2). Define the function  $f^*$  by (14). Notice that if  $\mathcal{D}$  is empty then  $f^*$  is identically equal to  $f$ .

IV. We shall construct the series  $S^*$  by choosing a particular sequence of integers  $n_1 < n_2 < \dots$  which are indices of Haar functions whose support lies in the closure of  $\Delta(i_0, k_0)$ .

Let  $n_1$  be that integer such that  $\Delta(1, n_1) \cup \Delta(2, n_1)$  is the interval  $\Delta(i_0, k_0)$  without its midpoint. For instance, if  $\Delta(i_0, k_0) = (1/4, 1/2)$  then  $n_1 = 5$ . By (17),  $n_1$  is the first integer such that  $S_{k_0+1} \not\equiv T_{k_0+1}$  on  $\Delta(i_0, k_0)$ .

Let  $n_2$  be the very next integer such that the support of  $\chi_{n_2}$  lies in the closure of  $\Delta(i_0, k_0)$  and such that, if  $i_1$  and  $k_1$  are chosen so that  $\Delta(1, n_2) \cup \Delta(2, n_2)$  is the interval  $\Delta(i_1, k_1)$  without its midpoint, then  $S_{k_1+1} \not\equiv T_{k_1+1}$  on  $\Delta(i_1, k_1)$ . Throughout the following pages we shall denote the interval  $\Delta(j, n)$  without its midpoint as  $\Delta^*(j, n)$ .

We continue this process as long as possible, thereby generating subintervals  $\Delta(i_j, k_j)$  of  $\Delta(i_0, k_0)$  and integers  $n_j$  ( $j = 1, 2, \dots$ ) such that

$$(21) \quad \Delta(1, n_{j+1}) \cup \Delta(2, n_{j+1}) = \Delta^*(i_j, k_j)$$

and

$$(22) \quad S_{k_j+1} \not\equiv T_{k_j+1} \text{ on } \Delta(i_j, k_j) \text{ for } j = 0, 1, 2, \dots$$

Finally, using the sequence  $n_1, n_2, \dots$  just generated we set

$$(23) \quad S^*(x) \equiv S_{k_0+1}(x) + \sum_{j=1}^{\infty} \alpha_{n_j} \chi_{n_j}(x).$$

In case the process for selecting the  $n_j$ 's terminates after a finite number of steps,  $S^*$  is just a finite series. Note also that IV(c) is trivially satisfied.

We shall now show IV(a) and (b) are also satisfied by this  $S^*$ .

Indeed, by the disjointness of the collection  $\mathcal{D}$  and by the choice of the sequence  $\{n_j\}$ , if the support of  $\chi_n$  is contained in  $\Delta(i_0, k_0)$  but  $\chi_n$  does not appear in the sum (23), then it must be the case that  $\text{Supp}(\chi_n) \subseteq \Delta(i_l, \rho_l)$  for some  $l$ . In particular, the series  $S$  and  $S^*$  are the same series in the set  $\Delta(i_0, k_0) \sim \bigcup \mathcal{D}$ . By the hypotheses of this lemma, then

$$(24) \quad \lim_{n \rightarrow \infty} \sigma_n(S^*; x) \equiv f(x) = f^*(x) \quad \text{for almost every } x \text{ in } \Delta(i_0, k_0) \sim \bigcup \mathcal{D}.$$

On the other hand, if  $x_0 \in \Delta(i_l, \rho_l)$  for some  $l$  used to define  $\mathcal{D}$ , then the disjointness of  $\mathcal{D}$  means that the series (23) must be truncated at  $\rho_l$ . Since  $S$  and  $S^*$  were identical up to that point,  $S^*(x_0) \equiv S_{\rho_l+1}(x_0)$ . But by the choice of  $\rho_l$ ,  $S_{\rho_l+1} \equiv T_{\rho_l+1}$  on  $\Delta(i_0, \rho_0)$ . Lemma 2 and the fact  $f^*$  is constant on  $\Delta(i_l, \rho_l)$  now imply that  $T_{\rho_l+1} \equiv f^*$  on  $\Delta(i_l, \rho_l)$ . Combining these three facts we conclude that  $S^*(x_0) = f^*(x_0)$ . Since  $x_0$  was any point in any interval of  $\mathcal{D}$ , we can now conclude that

$$(25) \quad \lim_{n \rightarrow \infty} \sigma_n(S^*; x) = f^*(x) \quad \text{on each } \Delta \in \mathcal{D}.$$

Combining (24) and (25) we have IV(a).

V. Suppose that the partial  $(C, 1)$  sums of  $S^*$  are not bounded by  $13M_0$  on  $\Delta(i_0, k_0)$  and let  $L$  be the smallest index greater than or equal to  $n_1$  such that  $|\sigma_L(S^*; t_0)| > 13M_0$  for some  $t_0 \in \Delta(i_0, k_0)$ .

If we let  $n_p$  be the largest number in the sequence  $n_1, n_2, \dots$  which is less than or equal to  $L$ , then for some choice of  $j_p = 1$  or  $2$ ,

$$(26) \quad |\sigma_L(S^*; x)| > 13M_0 \quad \text{for } x \in \Delta(j_p, n_p).$$

Indeed, if  $L = n_1$  then (26) is trivial by (21). Otherwise we use the least property of  $L$ .

We first begin by noting that

$$(27) \quad S_{n_p+1} \not\equiv T_{n_p+1} \quad \text{on } \Delta^*(i_{p-1}, k_{p-1}).$$

Indeed, if (27) were false, then  $S_{n_p+1} = S_{k_{p-1}+1} + \alpha_{n_p} \chi_{n_p}$ , and a corresponding equation involving  $T$  and its  $n_p$ th coefficient, say  $\beta_{n_p}$ , implies  $S_{k_{p-1}+1} - T_{k_{p-1}+1} \equiv (\beta_{n_p} - \alpha_{n_p}) \chi_{n_p}$  on  $\Delta^*(i_{p-1}, k_{p-1})$ . But  $\chi_{n_p}$  changes signs in that punctured interval while the left-hand side of the above identity is constant in that punctured interval. The only possibility, then, is that  $\beta_{n_p} = \alpha_{n_p}$  which in turn forces  $S_{k_{p-1}+1} - T_{k_{p-1}+1} \equiv 0$  on  $\Delta^*(i_{p-1}, k_{p-1})$ . Since both partial sums are constant throughout  $\Delta(i_{p-1}, k_{p-1})$ , this statement and (22) are incompatible; consequently (27) does hold.

Let  $j'_p \neq j_p$  with  $j'_p = 1$  or  $2$ . Then, by (21),  $\Delta(j_p, n_p) \cup \Delta(j'_p, n_p) = \Delta^*(i_{p-1}, k_{p-1})$ ; and, by (22),  $S_{k_{p-1}+1} \neq T_{k_{p-1}+1}$  on  $\Delta(i_{p-1}, k_{p-1})$ . Hence by the contrapositive of (18),

$$|\sigma_{k_{p-1}}(S; x)| \leq M_0 \quad \text{for } x \in \Delta(i_{p-1}, k_{p-1}).$$

Note also by (22) and the fact that  $S$  and  $S^*$  are identical outside  $\mathcal{D}$ , we have

$$\sigma_n(S; x) \equiv \sigma_n(S^*; x) \quad \text{for } x \in \Delta(i_{p-1}, k_{p-1})$$

whenever  $n = k_{p-1}, k_{p-1} + 1, \dots, n_{p+1} - 1$ . Consequently, the above inequality becomes

$$(28) \quad |\sigma_{k_{p-1}}(S^*; x)| \leq M_0 \quad \text{for } x \in \Delta(i_{p-1}, k_{p-1}).$$

Using (27) and the contrapositive of (18) we can also conclude that on at least one of the intervals  $\Delta(j_p, n_p), \Delta(j'_p, n_p)$ ,

$$|\sigma_{n_p}(S^*; x)| \leq M_0.$$

But

$$\begin{aligned} \sigma_{n_p}(S^*; x) &= \sum_{k=0}^{n_p-1} \left(1 - \frac{k}{n_p+1}\right) \alpha_k \chi_k(x) + \frac{\alpha_{n_p} \chi_{n_p}(x)}{n_p+1} \\ &\equiv \Sigma_1(x) + \frac{\alpha_{n_p} \chi_{n_p}(x)}{n_p+1}. \end{aligned}$$

Hence by (20) and the triangle inequality,

$$|\Sigma_1(x)| \leq |\sigma_{n_p}(S^*; x)| + |\alpha_{n_p} \chi_{n_p}(x)/(n_p+1)| \leq M_0 + M_0 = 2M_0$$

on at least one of the intervals  $\Delta(j_p, n_p), \Delta(j'_p, n_p)$ . But  $\Sigma_1$  is constant throughout the union of both these intervals, so  $|\Sigma_1(x)| \leq 2M_0$  for  $x \in \Delta(i_{p-1}, k_{p-1})$ .

Applying (20) and the triangle inequality again, we conclude that

$$(29) \quad |\sigma_{n_p}(S^*; x)| \leq 3M_0 \quad \text{for } x \in \Delta(i_{p-1}, k_{p-1}).$$

We shall now complete the proof of step V by showing that (28), (29) and (26) are incompatible with (27).

The definition of  $(C, 1)$  sums and the construction of  $S^*$  allow us to write the following equations for any  $x \in \Delta(i_{p-1}, k_{p-1})$ .

$$(30) \quad \sigma_L(S^*; x) = \frac{n_p+1}{L+1} \sigma_{n_p}(S^*; x) + \frac{L-n_p+1}{L+1} S_{n_p+1}^*(x);$$



$$\begin{aligned}
 \sigma_L(S^*; x) &= \frac{k_{p-1} + 1}{L + 1} \sigma_{k_{p-1}}(S^*; x) + \frac{L - k_{p-1} + 1}{L + 1} S_{n_p}^*(x) \\
 (31) \quad &+ \frac{L - n_p + 1}{L + 1} \alpha_{n_p} \chi_{n_p}(x); \\
 (32) \quad \sigma_{n_p}(S^*; x) &= \frac{k_{p-1} + 1}{n_p + 1} \sigma_{k_{p-1}}(S^*; x) + \frac{n_p - k_{p-1} + 1}{n_p + 1} S_{n_p}^*(x) + \frac{\alpha_{n_p} \chi_{n_p}(x)}{n_p + 1}.
 \end{aligned}$$

Since  $\chi_{n_p}$  is constant on  $\Delta(j_p, n_p)$  there are two possible cases:

$$\begin{aligned}
 (34) \quad &\frac{L - n_p + 1}{L + 1} |\alpha_{n_p} \chi_{n_p}(x)| > M_0 \quad \text{for } x \in \Delta(j_p, n_p) \\
 \text{or} \\
 (35) \quad &\frac{L - n_p + 1}{L + 1} |\alpha_{n_p} \chi_{n_p}(x)| \leq M_0 \quad \text{for } x \in \Delta(j_p, n_p).
 \end{aligned}$$

If (34) holds then we use (29) and (26) on (30) to conclude that

$$\frac{L - n_p + 1}{L + 1} |S_{n_p+1}^*(x)| > 10M_0 \quad \text{on } \Delta(j_p, n_p).$$

But  $S_{n_p+1}^* = S_{n_p}^* + \alpha_{n_p} \chi_{n_p}$ , so (34) implies

$$\frac{L - n_p + 1}{L + 1} |S_{n_p}^*| > 10M_0 - M_0 = 9M_0 \quad \text{on } \Delta(j_p, n_p).$$

Since  $S_{n_p}^*$  is constant on  $\Delta(i_{p-1}, k_{p-1})$  this inequality must hold throughout the larger interval:

$$(36) \quad \frac{L - n_p + 1}{L + 1} |S_{n_p}^*(x)| > 9M_0 \quad \text{for } x \in \Delta(i_{p-1}, k_{p-1}).$$

On the other hand, if (35) holds, then using (28) on (31) implies

$$\left| \frac{L - k_{p-1} + 1}{L + 1} S_{n_p}^*(x) + \frac{L - n_p + 1}{L + 1} \alpha_{n_p} \chi_{n_p}(x) \right| > 12M_0.$$

on  $\Delta(i_{p-1}, k_{p-1})$ , which by (35) guarantees

$$\frac{L - k_{p-1} + 1}{L + 1} |S_{n_p}^*| > 11M_0 > 9M_0 \quad \text{on } \Delta(j_p, n_p).$$

This shows that (36) holds in any case.

Now,  $2n_p$  is an element of the sequence  $n_1, n_2, \dots$  by (27). Hence by the choice of  $n_p$  relative to  $L$ ,  $L < 2n_p$ . The construction of the sequence  $k_j$  shows that  $2k_{p-1} \leq n_p$ . These two inequalities together yield  $3(n_p - k_{p-1} + 1) > L - k_{p-1} + 1$ . Consequently, applying (20), (28), and (36) to (32), we conclude  $|\sigma_{n_p}(S^*; x)| > (1/3)9M_0 - M_0 - M_0 = M_0$  for  $x \in \Delta(i_{p-1}, k_{p-1})$ . This, together with (18) and (21), implies  $S_{n_p+1} \equiv T_{n_p+1}$  on  $\Delta^*(i_{p-1}, k_{p-1})$ . By (27) this is impossible.

This final contradiction completes the proof of step V which in turn completes the proof of this lemma.

4. **The proof of the theorem.** Let  $\{Z_1, Z_2, \dots\}$  be the set of points in  $[0, 1]$  where

$$\limsup_{n \rightarrow \infty} |\sigma_n(S; x)| = +\infty.$$

Suppose that  $T$  is the D-Haar Fourier series of  $f$ , but that the theorem is false. Choose  $k_0$  least so that the  $k_0$ th Fourier coefficient of  $f$  is different from  $\alpha_{k_0}$ . Clearly, then,  $S_{k_0+1} \neq T_{k_0+1}$ . Since  $T$  satisfies Condition G (Lemma 3 of [2]) the series  $S - T$  satisfies the hypotheses of Lemma 1. The series  $S$  and  $T$  also satisfy the hypotheses of Lemma 4.

Applying Lemmas 1 and 4 countably many times, we can thus choose a sequence of intervals  $\Delta(i_1, k_1), \dots, \Delta(i_N, k_N), \dots$  of the form (1) or (2) such that

$$(37) \quad Z_n \text{ does not lie in the closure of } \Delta(i_N, k_N),$$

$$(38) \quad \text{the closure of } \Delta(i_N, k_N) \text{ is a subset of } \Delta(i_{N+1}, k_{N+1}),$$

and

$$(39) \quad |\sigma_{k_N}(S; x)| > N \quad \text{for } x \in \Delta(i_N, k_N), \text{ for } N = 1, 2, \dots.$$

By (38),  $\bigcap_{N=1}^{\infty} \Delta(i_N, k_N)$  is not empty; let  $\xi$  be in this intersection.

By (37),  $\xi \notin \{Z_1, Z_2, \dots\}$  which, by the definition of this sequence, implies

$$(40) \quad \limsup_{n \rightarrow \infty} |\sigma_n(S; \xi)| < \infty.$$

Yet by (39), since  $\xi \in \Delta(i_N, k_N)$  for all  $N$ ,  $\limsup_{n \rightarrow \infty} |\sigma_n(S; \xi)| = \infty$ . This being incompatible with (40) completes the proof of the theorem by contradiction.

#### BIBLIOGRAPHY

1. G. Alexits, *Konvergenzprobleme der Orthogonalreihen*, Akad. Kiadó, Budapest, 1960; English transl., Internat. Series of Monographs in Pure and Appl. Math., vol. 20, Pergamon Press, New York, 1961. MR 28 #5292; MR 36 #1911.
2. F. G. Arutjunjan, *Reconstruction of the coefficients of series with respect to Haar and Walsh systems convergent to functions integrable in the sense of Denjoy*, Izv. Akad. Nauk SSSR Ser. Mat. 30 (1966), 325–344. (Russian) MR 33 #1650.
3. W. R. Wade, *A uniqueness theorem for Haar and Walsh series*, Trans. Amer. Math. Soc. 141 (1969), 187–194. MR 39 #4587.
4. ———, *Uniqueness theory for Cesaro summable Haar series*, Duke Math. J. 38 (1971), 221–227.
5. A. Zygmund, *Trigonometrical series*. Vol. II, Cambridge Univ. Press, New York, 1959. MR 21 #6498.