

# ON THE EXTENSION OF MAPPINGS IN STONE-WEIERSTRASS SPACES<sup>(1)</sup>

BY

ANTHONY J. D'ARISTOTLE

Dedicated to Orrin Frink, Jr.

**ABSTRACT.** N. Veličko generalized the well-known result of A. D. Taïmanov on the extension of continuous functions by showing that Taïmanov's theorem holds when  $Y$  (the image space) is  $H$ -closed and Urysohn and the mapping  $f$  is weakly  $\theta$ -continuous. We obtain, in a more direct fashion, an even stronger generalization of this theorem.

We proceed to show that the class of all SW spaces is not reflective in the category of all completely Hausdorff spaces and continuous mappings. However, an epi-reflective situation is achieved by suitably enlarging the class of admissible morphisms.

We conclude by establishing a number of results about SW extension spaces.

**1. Preliminaries.** A subset  $A$  of a space  $X$  is said to be a zero-set of  $X$  if there exists a function  $f$  in  $C(X)$  (the set of all continuous, real-valued functions on  $X$ ) such that  $A = f^{-1}(\{0\}) = \{x \in X : f(x) = 0\}$ . Complements of zero-sets are called cozero-sets. If  $C^*(X)$  is the set of all bounded functions in  $C(X)$ , the subset  $A$  of  $X$  is said to be  $C^*$ -embedded in  $X$  if every function in  $C^*(A)$  can be extended to a function in  $C^*(X)$  [11]. If  $X$  and  $Y$  are spaces, let  $C(X, Y)$  denote the set of all continuous functions from  $X$  to  $Y$ .

A mapping  $f$  of the space  $X$  into the space  $Y$  is said to be  $\theta$ -continuous (weakly  $\theta$ -continuous) if for an arbitrary point  $x \in X$  and an arbitrary open set  $V$  of  $Y$  containing  $y = f(x)$ , there exists an open set  $U$  of  $X$  with  $x \in U$  and  $f(\text{cl}_X U) \cap V \neq \emptyset$  ([8], [19]). The point  $x$  is a member of the  $\theta$ -closure of the set  $S$  in  $X$  if and only if  $(\text{cl}_X G) \cap S \neq \emptyset$  for all open sets  $G$  containing  $x$ . The set  $S$  is said to be  $\theta$ -closed if it is equal to its  $\theta$ -closure [25].

A space  $X$  is said to be quasicompact if every family of zero-sets of  $X$  with the finite intersection property has a nonempty intersection [9]. The space  $X$  is said to be Urysohn if distinct points of  $X$  are contained in disjoint closed neighborhoods, and is said to be completely Hausdorff if for

---

Received by the editors August 16, 1972 and, in revised form, March 11, 1974.  
AMS (MOS) subject classifications (1970). primary 54C20, 54C30, 54C45.

(1) Portions of this paper were presented to the Second Pittsburgh International Conference on General Topology and its Applications, December 18–22; 1972.

every pair  $x, y$  of distinct points there is a function  $f$  in  $C(X)$  such that  $f(x) \neq f(y)$ . The word *space*, unqualified, shall henceforth mean a completely Hausdorff space. As in [21], the space  $X$  is said to be a Stone-Weierstrass space (or briefly, an SW space) if every point-separating subalgebra of  $C^*(X)$  which contains the constants is uniformly dense in  $C^*(X)$ .

A mapping  $f: X \rightarrow Y$ , where  $X, Y$  are arbitrary spaces, is said to be cozero-set continuous if  $f^{-1}(C)$  is open for all cozero-sets  $C$  of  $Y$ . Cozero-set continuous functions will be referred to as  $c$ -maps. If  $X$  is a space, let  $\tilde{X}$  be the space obtained by taking on the same set  $X$  the weak topology relative to  $C(X)$ . As noted in [3],  $X$  is SW if and only if  $\tilde{X}$  is compact. Moreover, a map  $f$  from an arbitrary space  $X$  to  $Y$  is a  $c$ -map if and only if  $\gamma_Y f: X \rightarrow \tilde{Y}$  is continuous, where  $\gamma_Y: Y \rightarrow \tilde{Y}$  is the identity map.

A Hausdorff space  $X$  is said to be absolutely closed, or simply  $H$ -closed, if it is closed in every Hausdorff space in which it can be embedded. This concept is a generalization of a property of compact Hausdorff spaces, and was introduced in 1924 by Alexandroff and Urysohn [1]. In [15], Katětov showed that any Hausdorff space  $X$  could be densely embedded in an  $H$ -closed space  $\kappa X$ , now referred to as the Katětov extension of  $X$ , having the property that  $X$  is a  $C^*$ -embedded subset. For a construction of  $\kappa X$ , the reader is referred to [18].

A filter  $\mathcal{F}$  on a space  $X$  is said to be completely regular if  $\mathcal{F}$  has a base  $\mathcal{B}$  of open sets such that for each set  $A \in \mathcal{B}$  there exist a set  $B \in \mathcal{B}$  contained in  $A$  and a function  $f \in C(X)$  which is equal to 0 on  $B$  and 1 on  $X \setminus A$  [5]. The filter  $\mathcal{F}$  is said to be free or fixed according as the intersection of all its members is empty or nonempty.

The space  $Y$  is said to be an extension of the space  $X$  if there exists a homeomorphism  $h$  from  $X$  into  $Y$  such that  $h(X)$  is dense in  $Y$ . If  $h$  is the identity map, the reference to  $h$  is omitted. The extensions  $Y$  and  $Z$  of  $X$  are said to be isomorphic if there is a homeomorphism of  $Y$  onto  $Z$  which leaves  $X$  pointwise fixed.

An arbitrary topological space  $X$  is said to be realcompact if every real maximal ideal in  $C(X)$  is fixed [6].

An open filter is a filter in the lattice of open sets. The open filter  $\mathcal{Q}$  is said to have the countable closure intersection property (abbreviated c.c.i.p.) provided that for each countable subset  $\mathcal{C}$  of  $\mathcal{Q}$ ,  $\bigcap \{cl_X C: C \in \mathcal{C}\} \neq \emptyset$ . An open ultrafilter is a maximal open filter.

A Hausdorff space  $X$  is said to be almost realcompact if every open ultrafilter with the c.c.i.p. converges.

2. Extension of maps. The following theorem is proved by Taĭmanov in [23].

(2.1) Let  $A$  be a dense subspace of an arbitrary space  $X$ , and let  $f: A \rightarrow Y$  be a continuous mapping of  $A$  into the compact Hausdorff space  $Y$ . The mapping  $f$  has a continuous extension from  $X$  to  $Y$  if and only if, for every pair  $F_1, F_2$  of closed disjoint subsets of  $Y$ , we have  $\text{cl}_X f^{-1}(F_1) \cap \text{cl}_X f^{-1}(F_2) = \emptyset$ .

It is easily verified that in the above theorem we may replace "closed disjoint subsets" by "disjoint zero-sets".

**Lemma 2.2.** Let  $f: X \rightarrow Y$  be a map from an arbitrary space  $X$  to an  $H$ -closed Urysohn space  $Y$ . Then the following conditions are equivalent:

- (a)  $f$  is  $\theta$ -continuous,
- (b)  $f$  is weakly  $\theta$ -continuous, and
- (c)  $f$  is a  $c$ -map.

**Proof.** That (a) implies (b) is obvious. A weakly  $\theta$ -continuous function is always a  $c$ -map and therefore it remains to prove that (c) implies (a).

Suppose  $p \in X$  and that  $f(p) \in V$ , where  $V$  is open in  $Y$ . An  $H$ -closed Urysohn space is an SW space [18] and hence completely Hausdorff. There is for each  $q \in Y \setminus \{f(p)\}$  a function  $h_q$  in  $C(Y)$  with  $h_q(q) = 0$  and  $h_q(f(p)) = 1$ . It is clear that  $C_q = \{y \in Y: h_q(y) < 1/2\}$  is open in  $Y$ ,  $D_q = \{y \in Y: h_q(y) \leq 1/2\}$  is a zero-set of  $Y$ ,  $f(p) \notin D_q$ , and  $Y \subseteq \bigcup \{C_q: q \in Y \setminus \{f(p)\}\} \cup V$ . The space  $Y$  is  $H$ -closed, and so there exist elements  $q_1, q_2, q_3, \dots, q_n$  of  $Y \setminus \{f(p)\}$  with  $Y \subseteq \bigcup_{i=1}^n \text{cl}_Y C_{q_i} \cup \text{cl}_Y V = \bigcup_{i=1}^n D_{q_i} \cup \text{cl}_Y V$  [16]. Since  $\bigcup_{i=1}^n f^{-1}(D_{q_i}) = E$  is a zero-set of  $X$  and  $p \in X \setminus E$ , there is an open set  $U$  of  $X$  with  $p \in U \subseteq \text{cl}_X U \subseteq X \setminus E$ . Clearly  $f(\text{cl}_X U) \subseteq f(X \setminus E) \subseteq \text{cl}_Y V$  and thus  $f$  is  $\theta$ -continuous.

The following theorem generalizes and follows from (2.1).

**Theorem 2.3.** Let  $A$  be a dense subspace of an arbitrary space  $X$ , and let  $f: A \rightarrow Y$  be a  $c$ -map from  $A$  to the SW space  $Y$ . The mapping  $f$  has a  $c$ -extension from  $X$  to  $Y$  if and only if, for every pair  $F_1, F_2$  of disjoint zero-sets of  $Y$ , we have  $\text{cl}_X f^{-1}(F_1) \cap \text{cl}_X f^{-1}(F_2) = \emptyset$ .

**Proof.** That the condition is necessary follows from the inclusion  $\text{cl}_X f^{-1}(F_i) \subseteq g^{-1}(F_i)$  where  $g: X \rightarrow Y$  is a  $c$ -extension of  $f$ . To see that

the condition is sufficient, we note that disjoint zero-sets of  $Y$  are disjoint zero-sets of  $\tilde{Y}$  and hence  $\gamma_Y f: A \rightarrow \tilde{Y}$  extends to a continuous map  $l: X \rightarrow \tilde{Y}$  by (2.1). The function  $h: X \rightarrow Y$ , defined by  $\gamma_Y h = l$ , is a  $c$ -map and  $h|A = f$ .

**Corollary 2.4.** *Let  $A$  be a dense subset of an arbitrary space  $X$ , and let  $f: A \rightarrow Y$  be weakly  $\theta$ -continuous where  $Y$  is  $H$ -closed and Urysohn. Then the following conditions are equivalent:*

- (a)  *$f$  has a weakly  $\theta$ -continuous extension from  $X$  to  $Y$ ,*
- (b) *for any pair  $F_1, F_2$  of  $\theta$ -closed disjoint subsets of  $Y$ , we have*  
 $\text{cl}_X f^{-1}(F_1) \cap \text{cl}_X f^{-1}(F_2) = \emptyset$ ,
- (c) *for any pair  $F_1, F_2$  of disjoint zero-sets of  $Y$ , we have  $\text{cl}_X f^{-1}(F_1)$   
 $\cap \text{cl}_X f^{-1}(F_2) = \emptyset$ .*

**Proof.** (a)  $\Rightarrow$  (b): Suppose  $g$  is the weakly  $\theta$ -continuous extension of  $f$ . If  $p \in \text{cl}_X f^{-1}(F_i)$ , then  $g(p)$  is, easily, a member of the  $\theta$ -closure of  $F_i$ .

(b)  $\Rightarrow$  (c): We need merely observe that zero-sets are  $\theta$ -closed.

(c)  $\Rightarrow$  (a): The function  $f$  is a  $c$ -map by Lemma 2.2 and therefore has a  $c$ -extension  $g$  from  $X$  to  $Y$  by Theorem 2.3. Again, by Lemma 2.2,  $g$  is weakly  $\theta$ -continuous.

In the above corollary, there are simple examples showing that we may not replace "weakly  $\theta$ -continuous" by "continuous". On the other hand, by Lemma 2.2, "weakly  $\theta$ -continuous" may be replaced by " $\theta$ -continuous" or " $c$ -map".

Veličko generalized Taĭmanov's theorem by showing that, under the hypothesis of Corollary 2.4, (a) is equivalent to (b) [25]. Stephenson [21] has shown that there exists a noncompact regular SW space  $Y$ . Since a regular absolutely closed space is compact,  $Y$  cannot be absolutely closed. Thus,  $Y$  is seen to be an example of an SW space which is not  $H$ -closed and so Theorem 2.3 covers a wider class of spaces than Veličko's result.

Almost realcompact spaces were defined and studied by Frolík in [10], where he showed that in many instances they behave much like realcompact spaces. In [7], Engelking gave the analogue of Taĭmanov's theorem for completely regular realcompact spaces: Let  $A$  be a dense subspace of an arbitrary topological space  $X$ , and let  $f: A \rightarrow Y$  be a continuous function of  $A$  into the completely regular realcompact space  $Y$ . The mapping  $f$  has a continuous extension from  $X$  to  $Y$  if and only if, for any sequence  $\{F_i\}_{i=1}^{\infty}$  of closed subsets of  $Y$  such that  $\bigcap_{i=1}^{\infty} F_i = \emptyset$ , we have  $\bigcap_{i=1}^{\infty} \text{cl}_X f^{-1}(F_i) = \emptyset$ .

It is a simple matter to establish the counterpart of Theorem 2 for completely Hausdorff realcompact spaces. In the same vein, we have

**Theorem 2.5.** *Suppose  $Y$  is almost realcompact,  $\kappa Y$  is an SW space,  $A$  is dense in the arbitrary space  $X$ , and  $f: A \rightarrow Y$  is weakly  $\theta$ -continuous. The mapping  $f$  has a weakly  $\theta$ -continuous extension from  $X$  to  $Y$  if and only if for any sequence  $\{F_i\}_{i=1}^\infty$  of  $\theta$ -closed subsets of  $Y$  such that  $\bigcap_{i=1}^\infty F_i = \emptyset$ , we have  $\bigcap_{i=1}^\infty \text{cl}_X f^{-1}(F_i) = \emptyset$ .*

**Proof.** To prove that the condition is necessary, we need merely argue as in (a)  $\Rightarrow$  (b) of Corollary 2.4. We now establish the sufficiency of the condition and first note that  $f$  may be regarded as a weakly  $\theta$ -continuous function from  $A$  to  $\kappa Y$ . If  $H_1$  and  $H_2$  are disjoint  $\theta$ -closed subsets of  $\kappa Y$ , then  $H_1 \cap Y$  and  $H_2 \cap Y$  are disjoint  $\theta$ -closed subsets of  $Y$ , and therefore  $\text{cl}_X f^{-1}(H_1) \cap \text{cl}_X f^{-1}(H_2) = \text{cl}_X f^{-1}(H_1 \cap Y) \cap \text{cl}_X f^{-1}(H_2 \cap Y) = \emptyset$ . By virtue of Corollary 2.4,  $f$  has a weakly  $\theta$ -continuous extension  $g$  from  $X$  to  $\kappa Y$ . It remains to show that  $g(X) \subseteq Y$ .

Suppose, on the contrary, that there is a point  $p$  of  $X \setminus A$  such that  $g(p) = \mathcal{Q} \in \kappa Y \setminus Y$ . Now  $\mathcal{Q}$  is an open ultrafilter on  $Y$  which does not have the c.c.i.p., and so there exists a countable subset  $\mathcal{C} = \{G_i\}_{i=1}^\infty$  of  $\mathcal{Q}$  with  $\bigcap_{i=1}^\infty \text{cl}_Y G_i = \emptyset$ . Since  $\kappa Y$  is  $H$ -closed and Urysohn,  $\text{cl}_Y G_i$  is  $\theta$ -closed for all positive integers  $i$  [24], and letting  $F_i = \text{cl}_Y G_i$  it follows that  $\bigcap_{i=1}^\infty \text{cl}_X f^{-1}(F_i) = \emptyset$ . There exists a positive integer  $j$  such that  $p \in X \setminus \text{cl}_X f^{-1}(F_j)$ , and we observe that  $G_j \cup \{\mathcal{Q}\}$  is open in  $\kappa Y$  and contains  $\mathcal{Q}$ . If  $U$  is an arbitrary open subset of  $X$  which contains  $p$ , choose a point  $s$  of  $[X \setminus \text{cl}_X f^{-1}(F_j)] \cap U \cap A$ . Now  $s \notin \text{cl}_X f^{-1}(F_j)$  and hence  $s \notin f^{-1}(\text{cl}_Y G_j)$  so that  $g(s) = f(s) \notin \text{cl}_Y G_j$ . It follows that  $g(s) \in Y \setminus \text{cl}_Y G_j$ , an open subset of  $\kappa Y$  which does not intersect  $G_j \cup \{\mathcal{Q}\}$ . Hence  $g(U) \not\subseteq \text{cl}_{\kappa Y}[G_j \cup \{\mathcal{Q}\}]$ , and so  $g$  is not weakly  $\theta$ -continuous at  $p$ .

**Remarks.** Porter and Thomas have given necessary and sufficient conditions on  $X$  for  $\kappa X$  to be an SW space [18].

It is easily verified that Lemma 2.2 is still valid if it is required merely that  $\kappa Y$  be Urysohn; thus in the above theorem, "weakly  $\theta$ -continuous" may be replaced by " $\theta$ -continuous" or "c-map".

**3. Reflectiveness of SW spaces.** Many extensions such as the Stone-Čech compactification, the Hewitt realcompactification, and the Banaschewski zero-dimensional compactification have, on account of their similar

mapping properties, been studied from a categorical standpoint and classified as epi-reflections in appropriate categories. For a thorough discussion of this theory, the reader is referred to [13].

We will not distinguish among isomorphic objects of any category, and for any category,  $1_X$  will denote the identity morphism for the object  $X$ . For categorical notions not specifically defined, the reader should consult Mitchell [17].

**Definition.** If  $\mathcal{U}$  is a full subcategory of a category  $\mathcal{B}$  and if for each object  $X$  in  $\mathcal{B}$  there exist an object  $X_{\mathcal{U}}$  in  $\mathcal{U}$  and a morphism (resp. epimorphism)  $r: X \rightarrow X_{\mathcal{U}}$  such that for each object  $Y$  in  $\mathcal{U}$  and each morphism  $f: X \rightarrow Y$ , there exists a unique morphism  $\bar{f}: X_{\mathcal{U}} \rightarrow Y$  such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{r} & X_{\mathcal{U}} \\ & \searrow f & \swarrow \bar{f} \\ & Y & \end{array}$$

is commutative, then  $\mathcal{U}$  is said to be a reflective (resp. an epi-reflective) subcategory of  $\mathcal{B}$  and  $r$  is called a reflection morphism (resp. epimorphism) from  $X$  to  $X_{\mathcal{U}}$ .

In establishing our next theorem, the techniques employed in Theorem 1 of [12] were most useful. We will furthermore lean heavily upon the following modification of Niemytski's classic example [11, 3K]. Let  $X = I^2 = \{(x, y): 0 \leq x \leq 1, 0 \leq y \leq 1\}$  be the unit square with the usual topology  $\tau_1$  and let  $A = \{(x, 0): (x, 0) \in X\}$ . To each  $(x, 0) \in A$  define  $V_x = \{(x, 0)\} \cup \{(u, v) \in X: v > 0 \text{ and } (u - x)^2 + v^2 < (1/4)^2\}$ . Let  $\tau_2$  be the topology on  $X$  generated by the collection of sets  $\tau_1 \cup (V_x)_{x \in I}$  as a subbase. It is easily verified that  $(X, \tau_2)$  is  $H$ -closed and Urysohn and hence SW, and that  $A$  with the induced topology is discrete.

**Theorem 3.1.** *Let  $\mathcal{B}$  be the category of all completely Hausdorff spaces and continuous functions, and let  $\mathcal{U}$  be its full subcategory of all SW spaces. Then  $\mathcal{U}$  is not reflective in  $\mathcal{B}$ .*

**Proof.** Suppose, on the contrary, that  $\mathcal{U}$  is reflective in  $\mathcal{B}$ . Consider the space  $(X, \tau_2)$  described above, and let  $r$  be the reflection morphism from  $A$  to the SW space  $A_{\mathcal{U}}$ . If  $i: A \rightarrow X$  is the identity map, there is a unique morphism  $\bar{i}: A_{\mathcal{U}} \rightarrow X$  such that  $i = \bar{i} \circ r$ . Since  $i$  is a homeomorphism into, it follows readily that  $r$  is a homeomorphism into.

Let  $D_2$  be the discrete space composed of two elements, 1 and 2, let  $P = X \times D_2$ , and for  $n = 1, 2$ , let  $j_n: X \rightarrow P$  be the map defined by  $j_n(y) = (y, n)$ . For each  $y \in i(A)$  identify  $j_1(y)$  and  $j_2(y)$ , let  $Q$  be the corresponding quotient space, and let  $\chi$  be the quotient map from  $P$  to  $Q$ . Now  $X$  and  $D_2$  are  $H$ -closed and it follows that  $P$  and hence  $Q$  is  $H$ -closed. One can easily check that  $Q$  is Urysohn and therefore SW.

It is now possible to proceed exactly as Herrlich and Strecker have done in [12] to show that  $A$  is homeomorphic to the SW space  $A_{\mathfrak{U}}$ : this is a contradiction since  $A$  is not quasicompact.

Let  $\alpha_X$  be the set of all nonisomorphic SW extensions  $Y$  of  $X$  such that  $X$  is  $C^*$ -embedded in  $Y$ , and let  $e_X$  denote the set of all those members of  $\alpha_X$  with the property that each trace filter is completely regular.

Stephenson [21] introduced and studied a particular member of  $e_X$ , which we denote by  $\sigma X$ . Specifically, if  $\mathfrak{M}$  is the set of all free maximal completely regular filters on  $X$ , then  $\sigma X$  is the space whose points are the elements of  $X \cup \mathfrak{M}$  and whose topology is generated by all sets  $V^*$  of the form  $V \cup \{\mathcal{F} \in \mathfrak{M} | V \in \mathcal{F}\}$  for  $V$  open in  $X$ . The space  $\sigma X$  enjoys many of the properties of the Stone-Čech compactification of a Tychonoff space and is, in fact, homeomorphic to  $\beta X$  in case  $X$  is completely regular.

In [20], Raha has described an extension of a space  $X$ , which we denote by  $\delta X$ , whose points are again the elements of  $X \cup \mathfrak{M}$  and whose topology is similar, in construction, to the topology of the Katětov extension. In particular, any set, open in  $X$ , is also open in  $\delta X$ , and if  $\mathcal{F} \in \mathfrak{M}$ , basic neighborhoods of  $\mathcal{F}$  are sets of the form  $G \cup \{\mathcal{F}\}$  for  $G \in \mathcal{F}$ . It is easily verified that  $\delta X \in e_X$ .

**Lemma 3.2.** *If  $Y \in \alpha_X$  and  $f: X \rightarrow Z$  is a  $c$ -map from  $X$  to the SW space  $Z$ , then there exists a unique  $c$ -map  $g: Y \rightarrow Z$  with  $g|X = f$ .*

**Proof.** Let  $F_1$  and  $F_2$  be disjoint zero-sets of  $Z$ . The sets  $A_1 = f^{-1}(F_1)$  and  $A_2 = f^{-1}(F_2)$  are disjoint zero-sets of  $X$ , and so there is an element  $l$  of  $C^*(X)$  which is 0 on  $A_1$  and 1 on  $A_2$  [11]. Now  $l$  can be extended to a function in  $C^*(Y)$  which implies that  $\text{cl}_Y A_1 \cap \text{cl}_Y A_2 = \emptyset$ . By Theorem 2.3, there is a  $c$ -map  $g: Y \rightarrow Z$  with  $g|X = f$ . The uniqueness of  $g$  follows from the uniqueness of  $\gamma_Z g$ .

**Theorem 3.3.** *Let  $\mathfrak{B}^*$  be the category of all spaces and  $c$ -maps. If  $\mathfrak{U}^* \subseteq \mathfrak{B}^*$  is the full subcategory of all SW spaces, then the natural mappings  $r: X \rightarrow \sigma X$  and  $r_1: X \rightarrow \delta X$  are reflection epimorphisms for  $\mathfrak{U}^*$ .*

**Proof.** The composition of  $c$ -maps is a  $c$ -map, the identity function is a  $c$ -map, and therefore  $\mathfrak{B}^*$  is a category. Since  $X$  is  $C^*$ -embedded in  $\sigma X$  and  $\delta X$  ([21], [20]), the proof now follows directly from Lemma 3.2.

**4. Projective extrema.** If  $Y$  is an extension space of  $X$ , the trace filters of  $Y$  are the filters  $\mathcal{N}(y)$ ,  $y \in Y \setminus X$ , where  $\mathcal{N}(y)$  is the filter on  $X$  generated by the traces  $U \cap X$  of the open sets  $U$  of  $Y$  which contain  $y$ . If  $X$  is Tychonoff, then the trace filters  $\mathcal{N}(y)$  of  $\beta X$  are precisely the free maximal completely regular filters on  $X$  ([2], [5]).

The extension  $Y$  of  $X$  is said to be projectively larger than the extension  $Z$  of  $X$ , denoted  $Y \geq Z$ , if there exists a continuous surjection  $f: Y \rightarrow Z$  which leaves  $X$  pointwise fixed. If  $\eta$  is a class of extensions of  $X$ , an element  $Y$  of  $\eta$  is said to be a projective maximum (resp. projective minimum) if  $Y \geq Z$  (resp.  $Z \geq Y$ ) for all  $Z$  in  $\eta$ . Projective maximums (resp. projective minimums), if they exist, are unique [4].

We are now in a position to give a simple proof of the following theorem due to Stephenson [21, Theorem 4(vii)].

**Theorem 4.1 (Stephenson).** *The projective minimum of  $e_X$  is  $\sigma X$ .*

**Proof.** If  $Y \in e_X$ , let  $\tilde{\eta}(y)$  denote the trace filter of  $\tilde{Y}$  corresponding to  $y \in \tilde{Y} \setminus \tilde{X}$ . Noting that  $\beta\tilde{X} = \tilde{Y}$  and that  $\tilde{\eta}(y)$  is a subset of the completely regular filter  $\eta(y)$ , we must have  $\eta(y) = \tilde{\eta}(y)$ . Let  $g: Y \rightarrow \sigma X$  be the function defined by  $g(x) = x$  for  $x \in X$  and  $g(y) = \eta(y) \in \mathfrak{M}$  for  $y \in Y \setminus X$ . If  $V \cup \{\mathcal{G} \in \mathfrak{M}: V \in \mathcal{G}\} = T$  is a basic open set of  $\sigma X$ , then  $g^{-1}(T) = V \cup \{y \in Y \setminus X: V \in \eta(y)\}$  which is open in  $Y$  by Lemma 4.1 of [18]. Thus,  $g$  is continuous.

It is clear that  $g(Y)$  is quasicompact and thus SW [3]. It follows that  $g(Y)$  is closed in  $\sigma X$  [21], and therefore  $g$  is onto.

Stephenson also proved that the function  $g$  in the above theorem is 1-1. From the manner in which we have defined  $g$ , this follows immediately from the fact that  $Y$  is completely Hausdorff.

Porter and Thomas [18] and Liu [16] have shown that the Katětov extension is a projective maximum in the class of  $H$ -closed extensions of a Hausdorff space  $X$ . In view of its affinity with the Katětov extension, it is natural to inquire about the role of  $\delta X$  as a projective maximum.

**Theorem 4.2.** *If  $Y \in \alpha_X$ , then  $\delta X \geq Y$  if and only if  $Y \in e_X$ .*

**Proof.** If  $\delta X \geq Y$ , then for any  $y \in Y \setminus X$  there is an  $\mathcal{Q} \in \mathfrak{M}$  such that



$\eta(y) \subseteq \mathcal{Q}$ . Since  $\tilde{\eta}(y)$  is a maximal completely regular filter and  $\tilde{\eta}(y) \subseteq \eta(y)$ , it follows that  $\eta(y) = \mathcal{Q}$ . Hence  $Y \in e_X$ .

On the other hand, suppose  $Y \in e_X$ . The identity map  $i: X \rightarrow Y$  is a  $c$ -map,  $\delta X$  is a reflection epimorphism for  $\mathcal{U}^*$ , and therefore  $i$  can be extended to a  $c$ -map  $f: \delta X \rightarrow Y$ . We claim that  $f(\mathcal{M}) \subseteq Y \setminus X$ . For if  $f(\mathcal{Q}) = x \in X$ , let  $\mathcal{D}$  be the class of all cozero-sets of  $X$  which contain  $x$ . Since  $f$  is a  $c$ -map and  $X$  is  $C^*$ -embedded in  $Y$ , it follows that  $\mathcal{D} \subseteq \mathcal{Q}$ . Now  $\mathcal{D}$  is a base for a fixed maximal completely regular filter, and so  $\mathcal{Q}$  is fixed which is a contradiction.

It is clear that  $f$  is continuous at each point of  $x$ . If  $f(\mathcal{Q}) = y \in Y \setminus X$ , let  $U$  be an open set  $Y$  which contains  $y$ . Since  $\eta(y)$  is completely regular,  $\tilde{\eta}(y) = \eta(y)$  which entails the existence of a cozero-set  $C$  of  $Y$  containing  $y$  with  $C \cap X \subseteq U$ . Since  $f$  is a  $c$ -map, there is a member  $G$  of  $\mathcal{Q}$  with  $f(G \cup \{\mathcal{Q}\}) \subseteq C$ . Clearly,  $f(G \cup \{\mathcal{Q}\}) \subseteq U$  and so  $f$  is continuous at  $y$ . That  $f$  is onto follows as in Theorem 4.1. Thus  $\delta X \geq Y$ .

The following corollary is analogous to a result of Banaschewski in [4].

**Corollary 4.3.** *If  $X$  is a space, then  $\sigma X \leq Y \leq \delta X$  for all  $Y$  in  $e_X$ .*

**Remarks.** The set  $e_X$  (and hence also  $\alpha_X$ ) may have substantial cardinality. We shall make use of the fact that any Hausdorff extension  $Y \supseteq X$  such that  $\sigma X \leq Y \leq \delta X$  belongs to  $e_X$ . Let  $N$  be the space of positive integers with the discrete topology, let  $Y = N \cup \mathcal{M}$ , and if  $\mathcal{Q} \in \mathcal{M}$ , let  $\tau_{\mathcal{Q}}$  be the topology on  $Y$  generated by the topology of  $\sigma N$  together with  $\{N \cup \{\mathcal{Q}\}\}$ . Clearly  $\sigma N \leq (Y, \tau_{\mathcal{Q}}) \leq \delta N$  and so  $(Y, \tau_{\mathcal{Q}}) \in e_N$ . If  $\mathcal{Q}$  and  $\mathcal{B}$  are distinct members of  $\mathcal{M}$ , a routine argument shows that  $(Y, \tau_{\mathcal{Q}})$  and  $(Y, \tau_{\mathcal{B}})$  are non-isomorphic extensions of  $N$ . Finally, since  $\sigma N$  is the Stone-Ćech compactification of  $N$ , it follows that  $2^{\mathfrak{c}} = \text{card } \mathcal{M} \leq \text{card } e_N$  [11].

It is natural to ask if  $\sigma X$  is a projective minimum in  $\alpha_X$ . We shall answer this question negatively, but we will first need to describe another SW extension of a given completely Hausdorff space  $X$ . Let  $\mathcal{U}$  be the set of all free zero-set ultrafilters (see [11]) on  $X$ , and let  $\pi X = X \cup \mathcal{U}$ . We define a topology for  $\pi X$  by taking as a base for the open sets the family of all sets of the form  $G \cup \{\mathcal{Q} \in \mathcal{U} : \exists A \in \mathcal{Q} \text{ with } A \subseteq G\}$  where  $G$  is any open set of  $X$ . It is readily verified that  $\pi X$  is SW and that  $X$  is a dense,  $C^*$ -embedded subset.

Let  $I = [0, 1]$ , let  $\tau$  be the usual topology on  $I$ , let  $J$  be the subset of  $I$  consisting of all irrational numbers, and choose disjoint dense subsets  $J_1$ ,

$J_2$  of  $(I, \tau)$  such that  $J = J_1 \cup J_2$ . Let  $I_1 = I \setminus J_2$ , let  $\tau_1$  be the topology on  $I_1$  induced by  $\tau$ , let  $\delta_1$  be the topology on  $I_1$  generated by  $\tau_1 \cup \{J_1\}$ , and denote the space  $(I_1, \delta_1)$  by  $P$ . A routine argument shows that  $(I_1, \tau_1)$  and  $P$  have the same continuous functions, and so  $\tau_1$  is the collection of cozero-sets of both  $(I_1, \tau_1)$  and  $P$ .

Suppose  $g \in C(\pi P, \sigma P)$  and  $g|_X$  is the identity map. Since no cozero-set of  $P$  is contained in  $J_1$ , the set  $J_1$  is open in  $\sigma P$ . However, if  $p \in U \subseteq J_1$  where  $U \in \delta_1$ , then  $U$  contains a member of a free zero-set ultrafilter on  $P$ , and hence  $g^{-1}(J_1) = J_1$  is not open in  $\pi P$ . This is a contradiction, and it is now evident that  $\sigma P$  is not a projective minimum in  $\alpha_P$ .

Clearly  $\pi P \in \alpha_P \setminus e_P$ , and it follows from Theorem 4.2 that  $\delta P \not\geq P$ . Therefore  $\delta X$  is not necessarily the projective maximum of  $\alpha_X$ , and this disproves a theorem of Raha [20].

Although  $\alpha X$  is not in general a projective minimum in  $\alpha_X$ , we do have the following result.

**Theorem 4.4.** *If  $X$  is Tychonoff, then  $\alpha X$  is the projective minimum in  $\alpha_X$ .*

**Proof.** If  $Y \in \alpha_X$ , then  $\tilde{Y} = \beta X = \alpha X$ ; hence  $\gamma_Y: Y \rightarrow \tilde{Y}$  is the desired map.

However, even if  $X$  is Tychonoff,  $\delta X$  need not be the projective maximum of  $\alpha_X$ . For in [21], Stephenson has described a Tychonoff space  $X$  with a one-point noncompact SW extension  $Y$ . Moreover,  $X$  is  $C^*$ -embedded in  $Y$ . If  $Y = X \cup \{y\}$  and  $\eta(y)$  were completely regular, then one could check the various cases to show that  $Y$  would be Tychonoff and hence compact ([3], [14]). Hence  $Y \in \alpha_X \setminus e_X$ .

**5. Acknowledgement.** The author gratefully acknowledges the substantial help given to him by the referee in the form of five pages of constructive criticism. These comments led to new proofs of Theorems 2.3, 3.3, and 4.4, replacing proofs of my own which were either excessively long, unnatural, or both. In addition, the referee made many other valuable suggestions.

#### REFERENCES

1. P. Alexandroff and P. Urysohn, *Zur theorie der topologischen Räume*, Math. Ann. 92 (1924), 258–266.
2. P. Alexandroff, *Bikompakte Erweiterungen topologischer Räume*, Mat. Sb. (N. S.) 5 (47) (1939), 403–423. (Russian) MR 1, 318.
3. B. Banaschewski, *On the Weierstrass-Stone approximation theorem*, Fund. Math. 44 (1957), 249–252. MR 19, 1182.
4. ———, *Extensions of topological spaces*, Canad. Math. Bull. 7 (1964), 1–22. MR 28 #4501.
5. N. Bourbaki, *Eléments de mathématique*. Part 1. *Les structures fondamentales de l'analyse*. Livre III: *Topologie générale*. Chaps. 4, 5, 6, 7, 8, 9, 10, Actua-

- alirés Sci. Indust., nos. 1029, 1045, 1084, Hermann, Paris, 1947, 1958, 1961; English transl., Hermann, Paris; Addison-Wesley, Reading, Mass., 1966. MR 9, 261; 30 #3439; 26 #6918; 34 #5044b.
6. R. F. Dickman, Jr., *Compactifications and realcompactifications of arbitrary topological spaces*, Virginia Polytechnic Institute and State University (unpublished).
7. R. Engelking, *Remarks on realcompact spaces*, Fund. Math. 55 (1964), 303–308. MR 31 #4000.
8. S. Fomin, *Extensions of topological spaces*, Ann. of Math. (2) 44 (1943), 471–480. MR 5, 45.
9. Z. Frolík, *Generalisations of compact and Lindelöf spaces*, Czechoslovak Math. J. 9 (84) (1959), 172–217. (Russian) MR 21 #3821.
10. ———, *A generalization of realcompact spaces*, Czechoslovak Math. J. 13 (88) (1963), 127–138. MR 27 #5224.
11. L. Gillman and M. Jerison, *Rings of continuous functions*, University Ser. in Higher Math., Van Nostrand, Princeton, N. J., 1960. MR 22 #6994.
12. H. Herrlich and G. E. Strecker, *H-closed spaces and reflective subcategories*, Math. Ann. 177 (1968), 302–309. MR 38 #2744.
13. H. Herrlich, *On the concept of reflections in general topology*, Contributions to Extension Theory of Topological Structures (Proc. Sympos., Berlin, 1967), Deutsch. Verlag Wissensch., Berlin, 1969, pp.105–114. MR 44 #2210.
14. E. Hewitt, *Certain generalizations of the Weierstrass approximation theorem*, Duke Math. J. 14 (1947), 410–427. MR 9, 95.
15. M. Katětov, *Über H-abgeschlossene und bikompakte Räume*, Časopis Pěst. Mat. Fys. 69 (1940), 36–49. (German) MR 1, 317.
16. C.-T. Liu, *Absolutely closed spaces*, Trans. Amer. Math. Soc. 130 (1968), 86–104. MR 36 #2107.
17. B. Mitchell, *Theory of categories*, Pure and Appl. Math., vol. 17, Academic Press, New York, 1965. MR 34 #2647.
18. J. Porter and J. Thomas, *On H-closed and minimal Hausdorff spaces*, Trans. Amer. Math. Soc. 138 (1969), 159–170. MR 38 #6544.
19. J. Porter and C. Votaw, *H-closed extensions. II*, University of Kansas, 1973 (preprint).
20. A. B. Raha, *On completely Hausdorff-completion of a completely Hausdorff space*, Pacific J. Math. 38 (1971), 161–166. MR 46 #6299.
21. R. M. Stephenson, Jr., *Spaces for which the Stone-Weierstrass theorem holds*, Trans. Amer. Math. Soc. 133 (1968), 537–546. MR 37 #3337.
22. ———, *Product spaces for which the Stone-Weierstrass theorem holds*, Proc. Amer. Math. Soc. 21 (1969), 284–288. MR 40 #3499.
23. A. D. Taĭmanov, *On extension of continuous mappings of topological spaces*, Mat. Sb. (N. S.) 31 (73) (1952), 459–463. (Russian) MR 14, 395.
24. N. V. Veličko, *H-closed topological spaces*, Mat. Sb. 70 (112) (1966), 98–112; English transl., Amer. Math. Soc. Transl. (2) 78 (1968), 103–118. MR 33 #6576.
25. ———, *On extension of mappings of topological spaces*, Sibirsk. Mat. Ž. 6 (1965), 64–69; English transl., Amer. Math. Soc. Transl. (2) 92 (1970), 41–47. MR 31 #4010; 42 #3.

DEPARTMENT OF MATHEMATICS, BROOKLYN COLLEGE (CUNY), BROOKLYN, NEW YORK 11210

*Current address:* Departamento de Matemáticas, Universidad Simon Bolívar, Sartanejas-Baruta 5354, Caracas, Venezuela