

BRANCH POINT STRUCTURE OF COVERING MAPS ONTO NONORIENTABLE SURFACES

BY

CLOYD L. EZELL¹

ABSTRACT. Let $f: M \rightarrow N$ be a degree n branched cover onto a compact, connected nonorientable surface with branch points y_1, y_2, \dots, y_m in N , and let the multiplicities at points in $f^{-1}(y_i)$ be $\mu_{i1}, \mu_{i2}, \dots, \mu_{ik_i}$. The *branching array* of f , designated by B , is the following array of numbers:

$$\begin{array}{c} \mu_{11}, \mu_{12}, \dots, \mu_{1k_1} \\ \mu_{21}, \mu_{22}, \dots, \mu_{2k_2} \\ \vdots \\ \mu_{m1}, \mu_{m2}, \dots, \mu_{mk_m} \end{array}$$

We show that the numbers in the branching array must always satisfy the following conditions:

- (1) $\sum \{ \mu_{ij} + 1 \mid j = 1, 2, \dots, k_i \} = n$,
- (2) $\sum \{ \mu_{ij} \mid i = 1, 2, \dots, m; j = 1, 2, \dots, k_i \}$ is even.

Furthermore, if B is any array of numbers satisfying these conditions, and if N is not the projective plane, then there is a branched cover onto N with B as its branching array.

1. Introduction. Let $f: M \rightarrow N$ be a continuous map between 2-dimensional manifolds. An open set $U \subset N$ is *evenly covered* if $f^{-1}(U)$ is a union of disjoint open sets on each of which f is topologically equivalent to the complex map z^n , for some n . If every point $y \in N$ has an evenly covered neighborhood, then f is called a *branched cover* and the pair (f, M) a *branched covering space* of N . If f is equivalent to z^n at x , then the *local degree* of f at x is n and the *multiplicity* of x is $n - 1$. A point $x \in M$ with positive multiplicity is a *critical point* and a point $y \in N$ with at least one preimage a critical point is a *branch point*. If y and z in N are not branch points, then $|f^{-1}(y)|$ (the cardinality of $f^{-1}(y)$) equals $|f^{-1}(z)|$ and we may define the *degree* of f , $\deg f$, to be $|f^{-1}(y)|$, for any y not a branch point. Two branched covers f_i :

Presented to the Society, November 17, 1976; received by the editors December 10, 1976 and, in revised form, July 23, 1977.

AMS (MOS) subject classifications (1970). Primary 55A10; Secondary 30A90.

Key words and phrases. Branched cover, branching array, multiplicity of a branched cover, nonorientable surface.

¹The results presented in this paper form a part of the author's Ph. D thesis, written at Vanderbilt University under the direction of Professor Morris L. Marx.

© American Mathematical Society 1978

$M_i \rightarrow N, i = 1, 2$, are *equivalent* if there exists a homeomorphism $h: M_1 \rightarrow M_2$ such that $f_1 = f_2 h$.

In this paper we will be concerned with branched covers between compact, closed (no boundary), connected surfaces, either orientable or nonorientable. In this case the number of branch points, the number of critical points, and the degree are all finite. Let $f: M \rightarrow N$ be such a map with branch point set $Y = \{y_1, y_2, \dots, y_m\}$ in N and let $\mu_{i1}, \mu_{i2}, \dots, \mu_{ik_i}, i = 1, \dots, m$, be the multiplicities of the preimages of y_i . The *branching array* of f , designated by B , is the following array of numbers:

$$B = \begin{bmatrix} \mu_{11}, \mu_{12}, \dots, \mu_{1k_1} \\ \mu_{21}, \mu_{22}, \dots, \mu_{2k_2} \\ \vdots \\ \mu_{m1}, \mu_{m2}, \dots, \mu_{mk_m} \end{bmatrix}$$

We say an array of numbers B is *admissible* if there is a branched cover having B as its branching array. The sum, designated by μ , of all of the entries in the branching array of some map is called the *multiplicity* of the map. A well-known fact concerning the multiplicities of branched covers is the Riemann-Hurwitz formula; namely, if $f: M \rightarrow N$ is a degree n branched cover with multiplicity μ , then $\mu = n \cdot \chi(N) - \chi(M)$, where χ represents the Euler characteristic of a surface.

Many of the problems concerning branching arrays of branched covers fall into one of two categories: enumeration problems and branch point classification problems. In an enumeration problem, one considers a class of maps F and an array B which is admissible for some map in F . The problem is to count the number of topologically distinct maps in F with branching array B . In a classification problem, one considers the set \mathfrak{B} of all branching arrays corresponding to maps in some class F . The problem is to find necessary and sufficient conditions for an array of numbers B to be a member of \mathfrak{B} .

An example of an enumeration problem appears in the work of Hurwitz [4]. Hurwitz enumerated all n -sheeted Riemann surfaces over the sphere with $2n - 2$ simple branch points. Translating to our language, Hurwitz enumerated all degree n branched covers over the sphere having the branching array:

$$\left[\begin{array}{cccccc} 1 & 0 & 0 & \dots & 0 \\ 1 & 0 & 0 & \dots & 0 \\ & & \vdots & & \\ 1 & 0 & 0 & \dots & 0 \end{array} \right] \begin{array}{l} \\ \\ \\ \end{array} \left. \vphantom{\begin{array}{c} 1 \\ 1 \\ \vdots \\ 1 \end{array}} \right\} 2n - 2 \text{ terms}$$

$$\underbrace{\hspace{10em}}_{n - 1 \text{ terms}}$$

Note that it follows from the Riemann-Hurwitz formula that the covering surface must also be the sphere.

Examples of classification problems appear in papers by Thom [7] and Shepardson [6]. In Thom's problem, the class of maps were the degree n polynomials $p: S^2 \rightarrow S^2$, where S^2 is the Riemann sphere. If $B = \{\mu_{ij} | i = 1, \dots, m; j = 1, \dots, k_i\}$ is the branching array of a degree n polynomial, then the following conditions must be satisfied:

- (1) $k_i = 1$, for some i (∞ is a branch point with one preimage);
- (2) $\sum \{\mu_{ij} + 1 | j = 1, \dots, k_i\} = n$;
- (3) $\mu = 2n - 2$.

Thom showed that any array of numbers satisfying (1)–(3) is the branching array of some degree n polynomial.

In Shepardson's problem, the maps were the degree n branched covers onto an arbitrary orientable surface N with $\chi(N) \leq 0$ (i.e. N can be any orientable surface except the sphere). Shepardson showed that the branching arrays corresponding to members in this class are characterized by the conditions:

- (1) $\sum \{\mu_{ij} + 1 | j = 1, \dots, k_i\} = n$;
- (2) μ is an even number.

In this paper we extend Shepardson's result to include the nonorientable surfaces; i.e., we show that the above two conditions characterize the branching arrays of degree n branched covers onto any surface N , orientable or nonorientable, as long as $\chi(N) \leq 0$.

The branching arrays for maps onto the sphere and projective plane have not been characterized, although some observations on certain admissible arrays for maps onto the sphere have been made by Endler [1]. Also, Francis [2] has defined a combinatorial invariant, called an assemblage, with which one can determine certain admissible arrays for maps from bordered surfaces into the sphere.

The facts on branched covers that we use to obtain these results are known but not readily available in the literature. Hence a summary of these is given in §2 of this paper. The actual classification problem is done in §3.

2. Representation of branched covers. Let $f: M \rightarrow N$ be a degree n branched cover between compact, closed surfaces with N connected. (M is not necessarily connected.) Let $Y = \{y_1, y_2, \dots, y_m\}$ be a set of points in N containing the branch points, \bar{N} the punctured surface $\bar{N} = N - Y$, and $\Pi(\bar{N}, x) = \Pi$ the fundamental group of \bar{N} based at some point $x \in \bar{N}$. The map f induces a homomorphism $\rho: \Pi \rightarrow S_n$, where S_n is the symmetric group of permutations on the letters $\{1, 2, \dots, n\}$, in the following way. Let the preimages of x be x_1, \dots, x_n . For each element $\omega \in \Pi$, define $\rho(\omega) = \rho_\omega$ as follows: $\rho_\omega(i) = j$ if the lift of some path in ω to x_i terminates at x_j . Since

each such path in ω terminates at the same point, ρ_ω is well defined, and since there are no branch points in \bar{N} , ρ_ω is one-one and a permutation in S_n . The map ρ is easily seen to be a homomorphism.

The homomorphism ρ determined by f is not unique. In fact f determines an entire equivalence class of homomorphisms as follows. Define homomorphisms ρ and δ to be equivalent if there exists a permutation $\sigma \in S_n$ such that for every $\omega \in \Pi$, $\rho_\omega = \sigma^{-1}\delta_\omega\sigma$. Now if f induces ρ and if ρ is related to δ , a relabeling of the subscripts of $f^{-1}(x)$ by σ causes f to induce δ . Conversely, any relabeling of the subscripts causes f to induce another homomorphism related to ρ . We now observe that two branched covers which determine the same class of homomorphisms are themselves equivalent.

Let $f: M \rightarrow N$ and $g: L \rightarrow N$ be degree n branched covers with branch points in the set $Y = \{y_1, \dots, y_m\}$ and suppose f and g induce equivalent homomorphisms ρ and δ , respectively. Let x_1, \dots, x_n and z_1, \dots, z_n be the preimages of x in M and L and suppose x_1, \dots, x_{n_1} are the preimages of x in some connected component of M , say M_1 . Then the existence of σ implies that $z_{\sigma(1)}, \dots, z_{\sigma(n_1)}$ are the preimages of x in some component of L , say L_1 . We now observe that the covering spaces (\bar{f}_1, \bar{M}_1) and (\bar{g}_1, \bar{L}_1) of \bar{N} (where \bar{M} and \bar{L} are the punctured surfaces obtained by removing the x_i 's and z_i 's, and \bar{f}_1 and \bar{g}_1 are the restrictions of f and g to these surfaces) are equivalent.

If $\bar{f}_1^\#: \Pi(\bar{M}_1, x_1) \rightarrow \Pi$ and $\bar{g}_1^\#: \Pi(\bar{L}_1, z_{\sigma(1)}) \rightarrow \Pi$ are the usual monomorphisms from the fundamental groups of the covering spaces (see [5, p. 154]), then we have

$$\begin{aligned} \bar{f}_1^\# \left(\Pi(\bar{M}_1, x_1) \right) &= \{ \omega \in \Pi | \rho_\omega(1) = 1 \} = \{ \omega \in \Pi | \sigma^{-1}\delta_\omega\sigma(1) = 1 \} \\ &= \{ \omega \in \Pi | \delta_\omega(\sigma(1)) = \sigma(1) \} = \bar{g}_1^\# \left(\Pi(\bar{L}_1, z_{\sigma(1)}) \right). \end{aligned}$$

Hence the covering spaces are equivalent (see [5, p. 159]), and by replacing the excised points, we have that the branched covers f_1 and g_1 (the restrictions of f and g to M_1 and L_1) are equivalent. Similarly, one can show that there is a one-one correspondence between the components of M and L such that the restrictions of f and g to corresponding components are equivalent branched covers.

Conversely, it is easily shown that equivalent branched covers determine the same equivalence class of homomorphisms.

If we let \mathcal{F} be the set of all equivalence classes of degree n branched covers over N with branch set in Y , and \mathcal{P} the set of equivalence classes of homomorphisms $\rho: \Pi \rightarrow S_n$, then we have established a one-one map Γ from \mathcal{F} to \mathcal{P} . We will now observe, using some construction techniques of Hurwitz, that this map is actually onto all of \mathcal{P} .

Now suppose we are given a closed, connected surface N , a finite set of

points $Y = \{y_1, \dots, y_m\}$ in N , and a homomorphism $\rho: \Pi \rightarrow S_n$. We can construct a degree n branched cover $f: M \rightarrow N$, from some surface M , which induces ρ as described below.

Consider a system of generators of Π . For convenience, we choose the generators in the standard way; namely, Π is generated by $\alpha_1, \dots, \alpha_r, \beta_1, \dots, \beta_m$, where the α_i 's are generators of $\Pi(N, x)$ and the β_i 's are path classes represented by simple closed curves about the points y_i . Then Π is represented by these generators with the single relation $W(\beta)W(\alpha) = 1$ (the identity in Π), where $W(\beta)$ is the word $\beta_1\beta_2 \dots \beta_m$ and $W(\alpha)$ is the word $\alpha_1\alpha_2\alpha_1^{-1}\alpha_2^{-1} \dots \alpha_{r-1}\alpha_r\alpha_{r-1}^{-1}\alpha_r^{-1}$, $r = 2p$, for N orientable of genus p , and $\alpha_1^2\alpha_2^2 \dots \alpha_r^2$, $r = p$, for N nonorientable of genus p .

Let a_1, \dots, a_r be closed curves intersecting only at x representing the path classes $\alpha_1, \dots, \alpha_r$, and let b_1, \dots, b_m be a set of nonintersecting line segments connecting x and the points y_i . See below.




FIGURE 1

Decompose N by cutting along the curves a_i and the lines b_i to obtain a closed disk Z with oriented boundary edges corresponding to the letters in the word $W(\beta)W(\alpha)$; i.e., $b_1b'_1 \dots b_mb'_ma_1a_2a'_1a'_2 \dots a_{r-1}a_ra'_{r-1}a'_r$ for N orientable and $b_1b'_1 \dots b_mb'_ma_1a'_1a_2a'_2 \dots a_ra'_r$ for N nonorientable. See below.

Construct the covering surface M by taking n copies of Z , Z_1, \dots, Z_n , and identifying the oriented edges as follows. If w (either an a_i or b_i) is an edge on the boundary of Z for which $\rho(w)$ (where w is the corresponding α_i or β_i) in S_n maps k to j , attach Z_k to Z_j by identifying edges w^k and $(w^j)'$. (The superscripts indicate the disk containing the edge.) Define a degree n branched cover $f: M \rightarrow N$ by taking homeomorphisms $f_k: Z_k \rightarrow Z$, $k = 1, \dots, n$, and extending in the obvious way. One can label the preimages of x




FIGURE 2

so that f does induce the homomorphism ρ from which it was constructed.

By the construction, it is clear that the only possible branch points are the y_i and x . The multiplicities of the preimages of y_i are determined by the cycles in the permutation $\rho(\beta_i)$ as follows. If $\rho(\beta_i)$ has a cycle $(c_1 c_2 \cdots c_r)$, then by the construction, the oriented boundary of a small disk about y_i lifts to arcs in sheets c_1, c_2, \dots, c_r , respectively. Hence y_i has a preimage at which f is locally t -to-one. Thus if $\rho(\beta_i)$ has k_i cycles of lengths $\mu_{i1} + 1, \dots, \mu_{ik_i} + 1$, then y_i has k_i preimages with multiplicities $\mu_{i1}, \dots, \mu_{ik_i}$. Since ρ is a homomorphism, $W(\rho(\beta))W(\rho(\alpha)) = 1$ (the identity in S_n), and it follows that x has n preimages and cannot be a branch point. Hence all branch points are in Y .

If the target surface N is orientable, then it is well-known that any covering surface must be orientable. (Note that this is readily checked from the construction above.) For nonorientable targets, the orientability of the covering surface can be determined by examination of the generators of Π as described below.

Suppose N is nonorientable. A curve in M avoiding the preimages of x and crossing the lifts of the a_i 's and b_j 's transversely is orientation reversing if and only if it crosses the lifts of the a_i 's an odd number of times. Hence M is nonorientable if and only if there exists such a curve in M . Suppose $W(\alpha, \beta) = \omega_1 \omega_2 \cdots \omega_q$ is a word in Π with an odd number of occurrences of α_i 's and suppose the permutation $W(\rho(\alpha), \rho(\beta))$ (i.e. the permutation $\rho(\omega_q) \cdots \rho(\omega_2) \rho(\omega_1)$, multiplying from right to left) fixes some number $k \in \{1, \dots, n\}$. Let $w_i, i = 1, \dots, q$, be the edges in N corresponding to the ω_i 's. (ω_i is either $\alpha_i^{\pm 1}$ or $\beta_i^{\pm 1}$ and w_i is the corresponding a_i or b_i .) Construct a path by beginning at some point $z \in Z_k$, cross edge w_1^k to sheet $Z_{\rho(\omega_1)(k)}$, cross $w_2^{\rho(\omega_1)(k)}$ to sheet $Z_{\rho(\omega_2)\rho(\omega_1)(k)}$, etc. Since $\rho(\omega_q) \cdots \rho(\omega_2) \rho(\omega_1)(k) = k$, this

process produces a path ending at the original sheet Z_k . Terminate the path at the starting point $z \in Z_k$. Since there are an odd number of α terms in $W(\alpha, \beta)$, we cross the edges a_i an odd number of times, and the path is reversing. Similarly, the existence of such a path implies the existence of a word $W(\alpha, \beta)$ with an odd number of α terms such that $W(\rho(\alpha), \rho(\beta))$ fixes some letter k .

Finally, we note that if $\rho(\Pi)$ acts transitively on $\{1, \dots, n\}$, then we can construct a path in M from Z_k to Z_j , for any choice of k and j , and thus M must be connected. Conversely, the existence of such a path implies $\rho(\Pi)$ acts transitively. More generally, if we let i be related to j if there exists a permutation $P \in \rho(\Pi)$ such that $P(i) = j$, and if S_1, \dots, S_k are the equivalence classes which partition the set $\{1, \dots, n\}$ under this relation, then M has k connected components and the restrictions of f, f_1, \dots, f_k , to these components have degrees $|S_1|, \dots, |S_k|$.

We summarize all of the above in the following theorem. As before, N is some connected, closed surface, $Y = \{y_1, \dots, y_m\}$ are selected points in N , $\bar{N} = N - Y$, \mathcal{F} is the set of all equivalence classes of degree n branched covering spaces of N with branch point set in Y , and \mathcal{P} is the set of all equivalence classes of homomorphisms $\rho: \Pi \rightarrow S_n$. We assume Π is represented in the standard way as described earlier.

2.1. THEOREM. *There is a one-one correspondence $\Gamma: \mathcal{F} \rightarrow \mathcal{P}$ such that if $\Gamma(F) = H$, $(f, M) \in F$, and $\rho \in H$, then:*

(1) *There is a one-one correspondence between the connected components of M, M_1, \dots, M_k and the equivalence classes S_1, \dots, S_k of $\{1, \dots, n\}$ determined by the action of $\rho(\Pi)$ on $\{1, \dots, n\}$ such that the degree of f restricted to M_i equals $|S_i|$. Specifically, M is connected if and only if $\rho(\Pi)$ acts transitively on $\{1, \dots, n\}$.*

(2) *The point $y_i \in N$ is a branch point whose preimages have multiplicities $\mu_{i1}, \dots, \mu_{ik_i}$, if and only if $\rho(\beta_i)$ is a nonidentity permutation with cycles of lengths $\mu_{i1} + 1, \dots, \mu_{ik_i} + 1$.*

(3) *If N is nonorientable, then M is nonorientable if and only if there exists a word $W(\alpha, \beta) \in \Pi$ with an odd number of α terms such that the permutation $W(\rho(\alpha), \rho(\beta))$ fixes k , for some $k \in \{1, \dots, n\}$.*

3. The classification problem. In this section we give necessary and sufficient conditions for an array B to be the branching array for a degree n branched cover onto any surface N with $\chi(N) \leq 0$. Since the conditions are necessary for maps onto any surface but sufficient only for the ones with $\chi(N) \leq 0$, we do the necessity and sufficiency parts in different theorems.

3.1. THEOREM. *Let $f: M \rightarrow N$ be a degree n branched cover between connected, closed surfaces with branch points y_1, \dots, y_m in N where the multiplicities*

at points in $f^{-1}(y_i)$ are $\mu_{i1}, \dots, \mu_{ik}$. Then:

(1) $\sum \{ \mu_{ij} + 1 \mid j = 1, \dots, k_i \} = n$;

(2) $\mu = \sum \{ \mu_{ij} \mid i = 1, \dots, m; j = 1, \dots, k_i \}$ is an even number.

Furthermore, if N is nonorientable and n is odd, then M is nonorientable.

PROOF. (1) This condition is well known and can be shown directly from the definition of a branched cover. Note, however, that it follows immediately from 2.1(2) since the sum of the cycle lengths of a permutation in S_n is n .

(2) Let ρ be any homomorphism associated with f by 2.1 and let Π have the standard generators α_i, β_j . We say the parity of a permutation is 0 if it is even and 1 if it is odd. Since each $\alpha_i, i = 1, \dots, r$, occurs twice in the word $W(\alpha)$, the parity of the permutation $W(\rho(\alpha))$ is 0. Since ρ is a homomorphism, $W(\rho(\beta))W(\rho(\alpha))$ is the identity in S_n and the parity of $W(\rho(\beta))$ is also 0. By 2.1(2), the j th cycle c_{ij} in $\rho(\beta_i)$ has length $\mu_{ij} + 1$, hence the parity of the cycle c_{ij} , $\text{par}(c_{ij})$, is congruent to μ_{ij} modulo 2. But $\text{par}(W(\rho(\beta)))$ equals the mod 2 sum of the parities of all the cycles in $W(\rho(\beta))$, hence $0 = \text{par}(W(\rho(\beta))) \equiv \mu \pmod{2}$, and μ is even. Note that if N is orientable, then this condition follows immediately from the Riemann-Hurwitz formula since the Euler characteristic of an orientable surface is even.

Now suppose N is nonorientable and n is odd. Then $\rho(\alpha_1)$ is a permutation of odd length and has at least one odd cycle, say $(c_1 c_2 \dots c_p)$. Thus the word $\alpha_1^p \in \Pi$ has an odd number of occurrences of α terms and $\rho_{\alpha_1}^p(c_1) = c_1$. Thus M is nonorientable by 2.1(3). \square

The following lemmas are used for the sufficiency part of the classification theorem. The first, Lemma 3.2, is done in [6]. For completeness, a proof is also given here.

3.2. LEMMA. *Every even permutation is the product of two cycles.*

PROOF. Let P be a given permutation and expand P as a product of disjoint cycles labeling the first letter of each cycle as follows:

$$P = (x_1 X_1)(x_2 X_2) \dots (x_j X_j)(y_1 Y_1)(y_2 Y_2) \dots (y_k Y_k),$$

where the X_i 's and Y_i 's are (possibly empty) strings of the remaining letters in the cycles. The x 's designate cycles of odd length and the y 's cycles of even length. Let \underline{X}_i (\underline{Y}_i) be the string consisting of the even terms (i.e. the 2nd, 4th, 6th, etc.) in X_i (Y_i) and let \bar{X}_i (\bar{Y}_i) be the string consisting of the odd terms. Let Q be the cycle $(x_1 \dots x_j y_1 \dots y_k \underline{X}_1 \dots \underline{X}_j \underline{Y}_1 \dots \underline{Y}_k)$ and observe that the product QP (multiplying from the right) is the permutation

$$\begin{aligned} & (x_1 \underline{X}_1 \dots x_j \underline{X}_j y_1 \underline{Y}_1 \bar{Y}_2 y_3 \underline{Y}_3 \bar{Y}_4 y_5 \dots \bar{Y}_{k-1} y_k \underline{Y}_k) \\ & \cdot (\bar{X}_1 \dots \bar{X}_j \bar{Y}_1 y_2 \underline{Y}_2 \bar{Y}_3 y_4 \underline{Y}_4 \bar{Y}_5 \dots \bar{Y}_k) \end{aligned}$$

for k odd and

$$\begin{aligned} (x_1 \underline{X}_1 \cdots x_j \underline{X}_j y_1 \underline{Y}_1 \bar{Y}_2 y_3 \underline{Y}_3 \bar{Y}_4 y_5 \cdots \bar{Y}_k \bar{X}_1 \\ \cdots \bar{X}_j \bar{Y}_1 y_2 \underline{Y}_2 \bar{Y}_3 y_4 \cdots \bar{Y}_{k-1} y_k \underline{Y}_k) \end{aligned}$$

for k even. If P is an even permutation then k must be even and the product QP is a cycle. Hence $P = (Q^{-1})(QP)$ factors as claimed. \square

3.3. LEMMA. *If P and Q are cyclic permutations, then there is a permutation R having a cycle of length one such that $P = R^{-1}QR$.*

PROOF. Let $P = (c_1 c_2 \dots c_n)$ and $Q = (d_1 d_2 \dots d_n)$. If $c_i = d_j$, take R to be

$$\begin{pmatrix} c_1 c_2 \cdots c_i \cdots c_n \\ \cdots d_{j-1} d_j d_{j+1} \cdots \end{pmatrix}.$$

See [3, p. 55]. \square

3.4. THEOREM. *Let N be any surface with $\chi(N) \leq 0$ and let B be an array of nonnegative integers, $B = \{\mu_{ij} | i = 1, \dots, m; j = 1, \dots, k_i\}$ (each row containing some nonzero entry) such that:*

$$(1) \sum \{\mu_{ij} + 1 | j = 1, \dots, k_i\} = n;$$

$$(2) \mu = \sum \{\mu_{ij} | i = 1, \dots, m; j = 1, \dots, k_i\} \text{ is an even number.}$$

Then there exists a degree n branched cover $f: M \rightarrow N$ from some connected surface M with B as its branching array. Furthermore, if N is nonorientable and n even, then M can be chosen nonorientable.

PROOF. The proof in the case where N is orientable is done in [6]. Suppose N is nonorientable and choose points y_1, \dots, y_m in N . Let $\alpha_1, \dots, \alpha_r, \beta_1, \dots, \beta_m$ be the generators of Π . We define a homomorphism $\rho: \Pi \rightarrow S_n$ by defining ρ on the generators and extending to all of Π . Choose permutations $P_{\beta_1}, \dots, P_{\beta_m}$ such that P_{β_i} has cycles of lengths $\mu_{i1} + 1, \dots, \mu_{ik_i} + 1$. It follows from the fact that μ is even that the product $P_{\beta_m} \cdots P_{\beta_1}$ is an even permutation. Hence by 3.1 there is a permutation Q such that $QP_{\beta_m} \cdots P_{\beta_1}$ is a cycle. Since Q^{-1} and $QP_{\beta_m} \cdots P_{\beta_1}$ are cycles, they are conjugates and there is a permutation P_{α_2} such that $QP_{\beta_m} \cdots P_{\beta_1} = P_{\alpha_2} Q^{-1} P_{\alpha_2}^{-1}$; or equivalently,

$$P_{\alpha_2} Q P_{\alpha_2}^{-1} Q P_{\beta_m} \cdots P_{\beta_1} = 1.$$

Insert $P_{\alpha_2} P_{\alpha_2}^{-1}$ on the left to obtain

$$P_{\alpha_2} P_{\alpha_2} P_{\alpha_2}^{-1} Q P_{\alpha_2}^{-1} Q P_{\beta_m} \cdots P_{\beta_1} = 1;$$

or equivalently,

$$(P_{\alpha_2})^2 (P_{\alpha_2}^{-1} Q)^2 P_{\beta_m} \cdots P_{\beta_1} = 1.$$

Let $P_{\alpha_1} = P_{\alpha_2}^{-1} Q$. Define $\rho(\beta_i) = P_{\beta_i}$, $i = 1, \dots, m$, $\rho(\alpha_1) = P_{\alpha_1}$, $\rho(\alpha_2) = P_{\alpha_2}$,

and $\rho(\alpha_i) = 1$, for $i = 3, \dots, r$. Hence $\rho^2(\alpha_r) \dots \rho^2(\alpha_1)\rho(\beta_m) \dots \rho(\beta_1) = 1$ and ρ extends to a homomorphism. Let (f, M) be any branched covering space associated with ρ by 2.1. Since $P_{\alpha_1}P_{\alpha_2} (= P_{\alpha_2}^{-1}QP_{\alpha_2})$ is a conjugate of Q and hence a cycle, $\rho(\Pi)$ acts transitively on $\{1, \dots, n\}$ and M is connected by 2.1(1). The cover f has B as its branching array by 2.1(2) and the choice of the permutations P_{β_i} , $i = 1, \dots, m$. By 3.3, P_{α_2} can be chosen to have a cycle of length one. If this is done then M is nonorientable by 2.1(3). (Take $W(\alpha, \beta)$ to be α_2 .) \square

The proof in the case where N is orientable is similar. Choose Q and the P_{β_i} as before. Let P_{α_2} be the permutation such that $P_{\alpha_2}^{-1}QP_{\alpha_2} = QP_{\beta_m} \dots P_{\beta_1}$. Take $P_{\alpha_1} = Q$, $P_{\alpha_i} = 1$, for $i = 3, \dots, r$, and we have

$$P_{\alpha_r}^{-1}P_{\alpha_{r-1}}^{-1}P_{\alpha_r}P_{\alpha_{r-1}} \dots P_{\alpha_2}^{-1}P_{\alpha_1}^{-1}P_{\alpha_2}P_{\alpha_1}P_{\beta_m} \dots P_{\beta_1} = 1.$$

Now ρ defined as before extends to a homomorphism.

We have shown that given an array B satisfying conditions (1) and (2) above and a nonorientable surface N (not the projective plane), there is always a degree n branched cover having B as its branching array. We observed that if n is odd, the covering surface is always nonorientable, and if n is even, there is always a nonorientable covering surface with the prescribed branching. However, if n is even, there is not always an orientable covering surface with the prescribed branching. For example, there is no 4-1 orientable cover of the Klein bottle having one branch point with multiplicities of 0 and 2. If there were, there would be permutations P_{β_1} , P_{α_1} , and P_{α_2} with $P_{\alpha_2}^2P_{\alpha_1}^2P_{\beta_1} = 1$, P_{β_1} having one cycle of length one and one of length three, and the permutations P_{α_1} and P_{α_2} having no odd cycles. But no such permutations exist in S_4 .

Finally, we note that if N is the sphere S^2 or the projective plane P^2 , then the Riemann-Hurwitz formula places a third restriction on branching arrays of maps onto N ; namely,

$$(3) \mu \geq 2n - 2, \text{ for } N = S^2;$$

$$(3') \mu \geq n - 2, \text{ for } N = P^2.$$

However, these conditions combined with (1) and (2) of 3.4 are not enough to characterize the branching arrays of maps onto S^2 and P^2 . This is shown in the following examples, the first of which appears in [6].

3.5. EXAMPLE. The array

$$B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \\ 0 & 2 \end{bmatrix}$$

satisfies conditions (1), (2), and (3), but there is no degree 4 branched cover onto S^2 with branching array B . If there were, there would be permutations P_{β_1} , P_{β_2} , and P_{β_3} in S_4 with P_{β_1} and P_{β_2} having two 2-cycles and P_{β_3} having a

1-cycle and a 3-cycle such that $P_{\beta_3}P_{\beta_2}P_{\beta_1} = 1$. But no such permutations exist in S_4 .

3.6. EXAMPLE. The array $B = [1 \ 3]$ satisfies conditions (1), (2), and (3'), but there is no 6-1 branched cover of P^2 with branching array B . If there were, there would be permutations P_{β_1} and P_{α_1} in S_6 with P_{β_1} having a 2-cycle and a 4-cycle such that $P_{\alpha_1}P_{\beta_1} = 1$. But there are no such permutations in S_6 .

REFERENCES

1. O. Endler, *Compact Riemann surfaces with prescribed ramifications and Puiseux series*, Bol. Soc. Brasil. Mat. **2** (1971), 61–64.
2. G. Francis, *Assembling Riemann surfaces with given boundary curves and branch points*, Illinois J. Math. **20** (1976), 198–217.
3. M. Hall, *The theory of groups*, Macmillan, New York, 1959.
4. A. Hurwitz, *Riemann'sche Flächen mit gegebenen Verzweigungspunkten*, Math. Ann. **39** (1891), 1–61.
5. W. S. Massey, *Algebraic topology: An introduction*, Harcourt, Brace, and World, New York, 1967.
6. C. B. Shepardson, *Hurwitz-Riemann formulas*, Ph.D. thesis, Syracuse Univ., 1973.
7. R. Thom, *L'équivalence d'une fonction différentiable et d'une polynôme*, Topology **3** (1965), suppl. 2, 297–307.

DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE, TENNESSEE 37235

Current address: Department of Mathematics, Stetson University, DeLand, Florida 32720