

AN EQUIVARIANT WALL OBSTRUCTION THEORY

BY

JENNY A. BAGLIVO

ABSTRACT. Let G be a finite group. For a certain class of CW-complexes with a G -action which are equivariantly dominated by a finite complex we define algebraic invariants to decide when the space is equivariantly homotopy or homology equivalent to a finite complex.

1. Introduction. A connected CW-complex X is a G -complex if the group G acts on X as a group of homeomorphisms which permute the cells of X . The question we study here is: Suppose that G is a finite group, that X is a G -complex and that K is a finite G -complex. Suppose there exist equivariant cellular maps $\varphi: K \rightarrow X$ and $s: X \rightarrow K$ and an equivariant cellular homotopy $\varphi s \simeq 1_X$. (K G -dominates X .) Is X "equivalent" to a finite G -complex?

Our main result is the following theorem.

THEOREM (5.1). *Let G be a finite group. Let X be a G -complex such that for each subgroup H of G , X^H is connected, and such that $X^G \neq \emptyset$. Suppose that X is G -dominated by a finite G -complex. Then*

(i) *there exists an invariant $E_1(X, G)$ such that X is G -equivalent to a finite G -complex if and only if $E_1(X, G)$ vanishes;*

(ii) *there exists an invariant $E_2(X, G)$ such that X is G -homology-equivalent to a finite G -complex where the equivalence is G -2-connected if and only if $E_2(X, G)$ vanishes.*

The space X^H is the subcomplex of X fixed by the subgroup H . A G -equivalence is an equivariant homotopy equivalence. As the term suggests, a G -homology-equivalence is an algebraic concept. (See §5.) The notions are not equivalent but each is a homotopy equivalence.

The elements $E_1(X, G)$ and $E_2(X, G)$ are Wall-type invariants [15], [16] of equivariant homotopy type. They are defined by considering chain complexes related to the cellular chains of the universal cover of X , $C(\tilde{X})$. (See §5.)

In §2 we discuss lifting group actions to regular covers of a G -complex X . §3 then shows how to factor $C(\tilde{X})$ as needed for the main theorem. Equivariant cell-attaching is discussed in §4, where we also show that if X is G -dominated by a finite G -complex, then it is G -equivalent to a complex of

Presented to the Society, January 29, 1977; received by the editors May 31, 1977.

AMS (MOS) subject classifications (1970). Primary 55G99, 57E99, 54E60.

Key words and phrases. G -complex, Wall obstruction, G -equivalence, G -homology equivalence.

© 1979 American Mathematical Society

finite type. §5 is devoted to a proof of Theorem (5.1). Lastly, in §6, we study the notion of G -homology-equivalence further. The results we obtain here are more algebraic in flavor than in the rest of the paper.

Most of this work is the author's thesis, written under the direction of Professor Douglas Anderson of Syracuse University. I wish to thank him for suggesting the problem to me, and for his advice and support.

2. Definition of G -complex; lifting group actions. Let X be a connected CW-complex with fixed cellular decomposition and let $\mathcal{C}(X)$ denote the group of cellular isomorphisms f of X with the property that if f leaves a cell of X invariant, then f fixes the cell pointwise. Let G be a group which we write multiplicatively.

DEFINITION 1. A *cellular action* of G on X is a homomorphism $\alpha: G \rightarrow \mathcal{C}(X)$, denoted by $\alpha(g) \equiv g$. The space X with a fixed cellular action of G is called a G -complex. The action is said to be *effective* if $\ker \alpha = \{1\}$.

Whenever possible we suppress mentioning the homomorphism α . If H is a subgroup of G and X is a G -complex, then X is an H -complex by simply restricting the homomorphism α to H .

DEFINITION 2. Let X and Y be G -complexes. A map $\varphi: X \rightarrow Y$ is said to be a G -map if φ is a cellular map which respects the action of G , that is, such that $\varphi(gx) = g\varphi(x)$, $x \in X$ and $g \in G$.

We follow the notation and terminology of [2]. Thus, in particular, if $S \subseteq G$ we let X^S denote the *fixed-point set* of S in X .

In Lemma 3 we list some facts which follow easily from the definitions. If H is a subgroup of G we let NH denote the *normalizer* of H in G . We use the notation $H \leq G$ if H is a subgroup of G , and $H < G$ if H is a proper subgroup of G .

LEMMA 3. Let X be a G -complex and let H be a subgroup of G .

(i) The subspace X^H is NH -invariant. In particular, if H is normal in G then X^H is invariant under the action of G .

(ii) There is an induced action of NH/H on X^H . Thus X^H is also an NH/H -complex.

(iii) The restriction of g to X^H yields a homeomorphism $g: X^H \rightarrow X^{gHg^{-1}}$, for each $g \in G$.

(iv) Let $X^{>H} = \bigcup_{H' > H} X^{H'}$. Then $X^{>H}$ is an NH -complex, and an NH/H -complex. \square

We will be interested in considering group actions defined on a regular cover of X , \hat{X} , which lift the action of G on X . If an action exists the group involved is, in general, an extension of $G/\ker \alpha$ by the group of deck transformations of the cover, and it is a split extension of these groups if there is a fixed point of G in X . (See [2].) For technical reasons we will include here

actions which are not effective. If G and π are groups, we use the notation $\pi x_\alpha G$ to represent the *semidirect product* of π and G with respect to an *action* $\alpha: G \rightarrow \mathcal{Q}(\pi)$, where α is a homomorphism whose range is the *group of automorphisms* of π .

Let X be a connected G -complex and let $e^0 \in X^G$. Then G acts on $\pi_1(X, e^0)$ by setting $g \cdot \gamma = \pi_1(g)(\gamma)$. We denote this action by α . Let $p: \hat{X} \rightarrow X$ be a regular cover of X . If $\text{im}(\pi_1(p))$ is invariant under the action of G , then there is an induced action (also denoted by α) on $\pi_1(X, e^0)/\text{im}(\pi_1(p))$. Let $\hat{G} = \pi_1(X, e^0)/\text{im}(\pi_1(p))x_\alpha G$.

LEMMA 4. *In the situation described above, the space \hat{X} is a \hat{G} -complex.*

PROOF. Fix a point $\hat{e}^0 \in \hat{X}$ over e^0 and use the Lifting Map Theorem to construct $\hat{g}: (\hat{X}, \hat{e}^0) \rightarrow (\hat{X}, \hat{e}^0)$ such that $p\hat{g} = gp$, for $g \in G$.

The action of \hat{G} on \hat{X} is defined by $(\gamma, g)x \equiv \gamma(\hat{g}(x))$, $\gamma \in \pi_1(X, e^0)/\text{im}(\pi_1(p))$, $g \in G$, $x \in \hat{X}$. \square

Notice that the action of G depends on the choices of $e^0 \in X^G$ and $\hat{e}^0 \in \hat{X}$.

As a special case of Lemma 4, the universal cover of X , \tilde{X} , is a \tilde{G} -complex, with $\tilde{G} = \pi_1(X)x_\alpha G$. If $A \subset X$ is a connected subcomplex, then there is a regular cover \hat{A} of A in \tilde{X} . We next discuss extensions of actions to \hat{A} .

LEMMA 5. *Let X be a connected G -complex such that $X^G \neq \emptyset$. Let $A \subset X$ be a connected G -invariant subcomplex such that $A^G \neq \emptyset$. Let $\hat{A} \subset \tilde{X}$ be one component of $p^{-1}(A)$.*

(i) *The space A with regular cover $p|_A: \hat{A} \rightarrow A$ satisfies Lemma 4. Thus, \hat{A} is a \hat{G} -complex where $\hat{G} = \pi_1(A)/\text{im}(\pi_1(p|_A))x_\alpha G$.*

(ii) *Let $\tilde{G} = \pi_1(X)x_\alpha G$. Then \hat{G} is a subgroup of \tilde{G} and \hat{A} is a \hat{G} -invariant subspace of \tilde{X} .*

PROOF. (i) We need to show that $\text{im}(\pi_1(p|_A)) \subset \pi_1(A)$ is G -invariant. (We assume that the basepoint is some $e^0 \in A^G$.) But $\text{im}(\pi_1(p|_A)) = \ker(\pi_1(i))$, where $i: A \hookrightarrow X$ is the inclusion map. Since A is G -invariant, i is a G -map. The group $\ker(\pi_1(i))$ is, therefore, a G -invariant subgroup.

(ii) We assume that the basepoints for the fundamental groups are $e^0 \in A^G \subset X^G$ and $\hat{e}^0 \in \hat{A} \subset \tilde{X}$. Then

$$\pi_1(A)/\text{im}(\pi_1(p|_A)) = \pi_1(A)/\ker(\pi_1(i)) \cong \text{im}(\pi_1(i)) \leq \pi_1(X);$$

and thus $\hat{G} \leq \tilde{G}$ under this identification. The rest of the lemma follows from the construction in Lemma 4. \square

In particular we can apply Lemma 5 to fixed-point sets of a subgroup H of G in X . Suppose that X^H and $X^{>H} = \bigcup_{H' > H} X^{H'}$ are connected. These complexes are NH -invariant, and thus by Lemma 5, X^H and $X^{>H}$ are NH -complexes which are NH -invariant subcomplexes of \tilde{X} .

3. Permutation complexes; the structure of the cellular chain complex $C(\tilde{X})$. Let X be a connected G -complex and $p: \tilde{X} \rightarrow X$ be its universal cover. Suppose that $X^G \neq \emptyset$. By the discussion in §2, \tilde{X} is a \tilde{G} -complex, where $\tilde{G} = \pi_1(X)x_\alpha G$. Then the cellular chain complex $C(\tilde{X})$ is a complex over the group ring $\mathbb{Z}\tilde{G}$. The purpose of this section is to study the structure of this chain complex further. In particular, we prove the following proposition.

PROPOSITION 1. *Let X be a connected G -complex such that for every subgroup H of G , X^H is connected. Suppose also that $X^G \neq \emptyset$. Then the following statements hold:*

- (i) *The complex $C(\tilde{X})$ is a $\mathbb{Z}\tilde{G}$ -permutation complex.*
- (ii) *Let \tilde{X}^H represent the universal cover of X^H . The subcomplex*

$$C(p^{-1}(X^H)) \cong \mathbb{Z}(\pi_1(X)x_\alpha NH) \bigotimes_{\mathbb{Z}(\pi_1(X^H)x_\alpha NH)} C(\tilde{X}^H),$$

as complexes of $\mathbb{Z}(\pi_1(X)x_\alpha NH)$ -modules.

We will first state the relevant definitions, facts and conventions. Recall that if S is a set and $\mathfrak{B}(S)$ is the group of bijections of S , an *action* of G on S as a group of permutations is a homomorphism $\alpha: G \rightarrow \mathfrak{B}(S)$. We use the usual notation of G_x for the *isotropy group* of a point x , and $G(x)$ for its orbit. Let R be a ring with unit.

DEFINITION 2. An *RG -permutation module*, M , is a left RG -module which is free as an R -module with a distinguished basis S on which G acts as a group of permutations, and whose RG -module structure extends this action on S linearly. We distinguish several R -free submodules of M , with $H \leq G$ a subgroup:

- M^H is R -free with basis $S^H = \{x \in S: G_x \geq H\}$;
- $M^{>H}$ is R -free with basis $S^{>H} = \{x \in S: G_x > H\}$;
- M_H is R -free with basis $S_H = \{x \in S: G_x = H\}$.

If the distinguished basis of M is $G(x)$, the orbit of one element $x \in S$, we can identify $G(x)$ with G/H , the set of cosets of $H = G_x$, with action defined by left multiplication. This identification depends on the choice of $x \in S$. However, once a choice is made, we can write $M = R(G/H)$. In fact, all RG -permutation modules are RG -direct sums of such modules. We can write $M = \sum_{H \leq G} \oplus_R M_H = \bigcup_{H \leq G} M^H$.

As an R -module M^H decomposes into $M^{>H} \oplus M_H$. The modules $M^H, M_H, M^{>H}$ are not in general RG -invariant submodules of M . They are, however, invariant submodules of M under the action of the normalizer of H in G .

LEMMA 3. *Let M be an RG -permutation module.*

- (i) *The modules M^H, M_H and $M^{>H}$ are $R(NH)$ - and $R(NH/H)$ -permutation modules. The module M_H is a free $R(NH/H)$ -module;*

(ii) The modules $\bigcup_{g \in G} M^{gHg^{-1}}$, $\bigcup_{g \in G} M_{gHg^{-1}}$ and $\bigcup_{g \in G} M^{>gHg^{-1}}$ are RG -invariant submodules of M ;

(iii) For each $g \in G$, the action of g induces an R -isomorphism $g: M^H \rightarrow M^{gHg^{-1}}$, so that $g(M_H) = M_{gHg^{-1}}$ and $g(M^{>H}) = M^{>gHg^{-1}}$;

(iv) For each $g \in G$, the group isomorphism $g: NH \rightarrow NgHg^{-1}$ defined by $k \rightarrow gkg^{-1}$ extends linearly to an isomorphism of the group rings. We can represent

$$M^{gHg^{-1}} \cong R(NgHg^{-1}) \otimes_{R(NH)} M^H,$$

and we can represent $M_{gHg^{-1}}$ and $M^{>gHg^{-1}}$ similarly.

(v) For each $g \in G$, the group isomorphism $g: NH/H \rightarrow NgHg^{-1}/gHg^{-1}$ induced from the isomorphism in (iv) extends linearly to an isomorphism of the group rings. We can represent

$$M^{gHg^{-1}} \cong R(NgHg^{-1}/gHg^{-1}) \otimes_{R(NH/H)} M^H,$$

and we can represent the other distinguished submodules similarly.

PROOF. All parts are straightforward applications of the definitions. \square

DEFINITION 4. Let M and M' be RG -permutation modules. An RG -homomorphism $\varphi: M \rightarrow M'$ is said to be an RG -permutation homomorphism if $\text{im}(\varphi|_{M^H}) \subset M'^H$ for each subgroup H of G . We let φ^H denote $\varphi|_{M^H}$, $\varphi^{>H} = \varphi|_{M^{>H}}: M^{>H} \rightarrow M'^{>H}$ and $(\varphi_H, \varphi_H^{>H}): M_H \rightarrow M'_H \oplus M'^{>H} = M'^H$.

The reader should note that RG -homomorphisms are not in general RG -permutation homomorphisms. As an example of this, consider the trivial permutation module R and the free RG -module RG . Let G be a finite group. Define an R -homomorphism $h: R \rightarrow RG$ by $1 \mapsto \sigma = \sum_{g \in G} g$. Since $g'\sigma = \sigma$ for every $g' \in G$, h is also an RG -homomorphism. Now, $h(R^G) = h(R) = R\{\sigma\} \subsetneq (RG)^G = 0$.

The following facts are easily checked using the definitions and Lemma 3.

LEMMA 5. Let $\varphi: M \rightarrow M'$ be an RG -permutation homomorphism.

(i) The homomorphisms φ^H , $\varphi^{>H}$ and φ_H are $R(NH)$ - and $R(NH/H)$ -permutation homomorphisms.

(ii) We can represent $\varphi^{gHg^{-1}} = R(NgHg^{-1}/gHg^{-1}) \otimes_{R(NH/H)} \varphi^H$, and we can represent $\varphi^{>gHg^{-1}}$ and $\varphi_{gHg^{-1}}$ similarly.

(iii) We can represent $\varphi^{sHg^{-1}} = R(NgHg^{-1}) \otimes_{R(NH)} \varphi^H$, and we can represent $\varphi^{>sHg^{-1}}$ and $\varphi_{sHg^{-1}}$ similarly. \square

DEFINITION 6. An RG -chain complex C is an RG -permutation complex if each C_i and ∂_i is an RG -permutation module and homomorphism, respectively. If $\varphi: D \rightarrow C$ is an RG -chain map between permutation complexes and each φ_i is a permutation homomorphism, φ is an RG -permutation chain map.

The R -subcomplexes C^H , $C^{>H}$, and quotient C_H are defined in the obvious way and are clearly seen to be $R(NH)$ - and $R(NH/H)$ -permutation complexes. These complexes satisfy the short exact sequence

$$C^{>H} \twoheadrightarrow C^H \twoheadrightarrow C_H,$$

but C^H is not, in general, a direct sum of the subcomplexes $C^{>H}$ and C_H since ∂^H is not, in general, a direct sum of homomorphisms.

The proof of Proposition 1 will follow from Lemmas 7–9. Since the proofs of these lemmas are straightforward they will be omitted.

LEMMA 7. *Let X be a G -complex. The cellular chain complex $C(X)$ is a $\mathbb{Z}G$ -permutation complex. \square*

Notice that, as subcomplexes of $C(X)$, $C(X)^H = C(X^H)$ and

$$C(X)_H = C\left(X^H, \bigcup_{H' > H} X^{H'}\right) = C(X^H, X^{>H}).$$

LEMMA 8. *Let X be a connected G -complex such that $e^0 \in X^G$. Let $p: \tilde{X} \rightarrow X$ be a regular cover such that $\text{im}(\pi_1(p)) \subset \pi_1(X, e^0)$ is invariant under the action of G . Let $\hat{G} = \pi_1(X)/\text{im}(\pi_1(p))x_\alpha G$.*

Then $C(X) \cong \mathbb{Z}G \otimes_{\mathbb{Z}\hat{G}} C(\tilde{X})$ as $\mathbb{Z}G$ -complexes. \square

LEMMA 9. *Let X be a connected G -complex such that $X^G \neq \emptyset$ and let $p: \tilde{X} \rightarrow X$ be its universal cover. Let A be a connected G -invariant subcomplex of X such that $A^G \neq \emptyset$ and let $\hat{A} \subset \tilde{X}$ be a regular cover. Let $\tilde{G} = \pi_1(X)x_\alpha G$ and $\hat{G} = \text{im } \pi_1(i)x_\alpha G$. Then there is an isomorphism of the chain complexes $C(p^{-1}(A)) \cong \mathbb{Z}\tilde{G} \otimes_{\mathbb{Z}\hat{G}} C(\hat{A})$ as $\mathbb{Z}\tilde{G}$ -complexes. \square*

PROOF OF PROPOSITION 1. The first part follows from (2.4) and Lemma 7.

The given representation of $C(p^{-1}(X^H))$ follows from Lemmas 8 and 9 above. \square

4. G -equivalence and G -domination. Our objectives in this section are to show how to extend the “ G -connectivity” of a G -domination map by an equivariant cell-attaching technique (Lemma 5), and to prove

PROPOSITION 1. *Let X be a G -complex and G a finite group. If X is G -dominated by a finite complex then X is G -equivalent to a complex of finite type, Y . If n is the dimension of the dominating complex, then the complex Y is G -dominated by its n -skeleton.*

These results generalize the constructions in [15]. Note that a complex of finite type is one containing a finite number of cells in each dimension.

If φ is a G -map and $H \leq G$ is a subgroup, we let φ^H denote the restriction $\varphi|_{K^H}: K^H \rightarrow X^H$ and let $M(\varphi^H)$ denote its mapping cylinder. The following definitions will be needed.

DEFINITION 2. A G -map $\varphi: K \rightarrow X$ is said to be G - n -connected if $\pi_i(M(\varphi^H), K^H) = 0$ for all $i \leq n$, for all $H \leq G$, and for all choices of basepoint.

DEFINITION 3. A G -map $\varphi: K \rightarrow X$ is said to be a G -domination if there exists a G -map $s: X \rightarrow K$ and G -homotopy $F_1: \varphi s \simeq 1_X$. In addition, if there exists a G -homotopy $F_2: s\varphi \simeq 1_K$, φ is said to be a G -equivalence, and K and X are said to be of the same G -homotopy-type.

If X is a G -complex, $X \times I$ is a G -complex by defining $g(x, t) \equiv (gx, t)$, $x \in X$, $t \in I$. Thus, a G -homotopy $F: f_0 \simeq f_1$ between G -maps $f_0, f_1: X \rightarrow Y$ is a G -map $F: X \times I \rightarrow Y$ such that

$$F(x, i) = f_i(x), \quad i = 0, 1.$$

Also, if $\varphi: K \rightarrow X$ is a G -map there is an induced action of G on the mapping cylinder $M(\varphi)$, and this action has the property that $M(\varphi)^H = M(\varphi^H)$.

From the definitions it is clear that if φ is a G -equivalence then φ^H is a homotopy equivalence for each H . Thus, φ is G - n -connected for every n . The converse, which is an equivariant version of a theorem of Whitehead, is also true.

PROPOSITION 4. Let $\varphi: K \rightarrow X$ be a G -map. If φ is G - n -connected for every n then φ is a G -equivalence.

PROOF. See [10]. \square

We assume from now on that G is a finite group. Assume that X is G -dominated by a finite complex K , with G -dominating map $\varphi: K \rightarrow X$, and right inverse $s: X \rightarrow K$; and that both K and X are connected. By assumption, $\varphi^H: K^H \rightarrow X^H$ is a dominating map for each $H \leq G$. Although K^H and X^H may have more than one component each, since K is a finite complex both have a finite number of components.

We show that we can increase the G -connectivity of the map by successively adding a finite number of cells in each dimension. In the following lemma, let any G -dominating map be " G - (-1) -connected".

LEMMA 5. Let X be G -dominated by the finite complex K , with dominating map $\varphi: K \rightarrow X$. Suppose φ is G -($i-1$)-connected. Then there exists a finite G -complex L containing K and a G - i -connected dominating map $\psi: L \rightarrow X$ extending φ .

PROOF. If $\psi: L \rightarrow X$ extends φ , and we let $j: K \rightarrow L$ represent the inclusion, then js is a homotopy right inverse of ψ . Thus, it is sufficient to construct a G - i -connected extension of φ .

To do this we proceed by induction on the order $|H|$ of the subgroup H of G . To simplify notation, assume that $\varphi: K \rightarrow X$ is G -($i-1$)-connected, and $\varphi^{H'}: K^{H'} \rightarrow X^{H'}$ is i -connected for all H' where $|H'| > |H|$. In the induction

steps below, we add cells to K^H and to $K^{gHg^{-1}}$ for all conjugates to extend the connectivity of φ^H and all $\varphi^{gHg^{-1}}$. There are three cases to consider.

Case 1 ($i = 0$). By the domination $\pi_0(\varphi^H): \pi_0(K^H) \rightarrow \pi_0(X^H)$ is an epimorphism. If φ^H is one-to-one on components then it is already 0-connected and there is nothing to do. Suppose not, and suppose components K_1^H and K_2^H map to X_0^H , that is, $\varphi^H: K_1^H \cup K_2^H \rightarrow X_0^H$. Let e_1, e_2 be zero-cells in K_1^H and K_2^H respectively. Let $\omega: I \rightarrow X_0^H$ be a path from $\varphi^H(e_1)$ to $\varphi^H(e_2)$ along the one-skeleton of X_0^H . For each class $g \in G/H$ attach a one-cell e_g so that the characteristic map on the boundary $c_g: \dot{I} \rightarrow K^{gHg^{-1}}$ is $c_g(0) = ge_1$ and $c_g(1) = ge_2$. (This definition is independent of choice of representative for the class in G/H since H fixes K_1^H and K_2^H .) The cells are permuted by $g'e_g = e_{g'g}$. To extend the map φ to ψ define $\psi_0^H|_{e_g} = g\omega c_g^{-1}$.

Case 2 ($i = 1$). Consider $\varphi^H: K^H \rightarrow X^H$. Since φ is 0-connected K^H and X^H have the same number of components. Let $\varphi_0^H: K_0^H \rightarrow X_0^H$ be the restriction to corresponding components. Now since φ_0^H is a domination there exists a short exact sequence

$$1 \rightarrow \pi_1(M(\varphi_0^H), K_0^H) \rightarrow \pi_1(K_0^H) \rightarrow \pi_1(X_0^H) \rightarrow 1.$$

Also, $\pi_1(X_0^H)$ is finitely presented, and $\pi_1(M(\varphi_0^H), K_0^H)$ is finitely normally generated in $\pi_1(K_0^H)$ [15, Proof of Lemma 1.3]. Let $\{\alpha\}$ be a finite set of generators for $\pi_2(M(\varphi_0^H), K_0^H)$. Let $\alpha = \langle (\beta_\alpha, \gamma_\alpha) \rangle$ where

$$(\beta_\alpha, \gamma_\alpha): (D^2, S^1) \rightarrow (M(\varphi_0^H), K_0^H).$$

We attach 2-cells $e_{(g, \alpha)}$ to K via $g \circ \gamma_\alpha: S^1 \rightarrow K^{gHg^{-1}}$. To extend the action over the cells define $g'e_{(g, \alpha)} \equiv e_{(g'g, \alpha)}$. To extend the map φ let $\psi^H|_{e_{(g, \alpha)}} = g \circ \beta_\alpha$.

Case 3 ($i > 1$). By [15, Proof of Theorem A] $\pi_i(M(\varphi_0^H), K_0^H)$ is a finitely-generated $\mathbb{Z}\pi_1(K_0^H) = \mathbb{Z}(\pi_1(X_0^H))$ -module. Let $\{\alpha\}$ be a finite set of $\mathbb{Z}\pi_1(X_0^H)$ -generators and proceed as in Case 2. \square

PROOF OF PROPOSITION 1. Let $\varphi: K \rightarrow X$ be a G -dominating map with right homotopy inverse s . Using Lemma 5 inductively we can assume that φ is G - n -connected, where n is the dimension of K . By continuing the construction of Lemma 5 and by Proposition 4 we obtain a G -equivalence $\psi: Y \rightarrow X$, where Y is a G -complex of finite type and the n -skeleton of Y is K . Let s' be a homotopy inverse of ψ . Then the inclusion $K \hookrightarrow Y$ is a G -domination with right homotopy inverse $s'\varphi s\psi$. (Since all maps are cellular the image of $s'\varphi s\psi$ lies in $Y^n = K$.) \square

5. The two G -homotopy type invariants. In this section we prove Theorem (5.1), as stated in the introduction.

The definitions of the invariants $E_1(X, G)$ and $E_2(X, G)$ will require a description of some relative Wall invariants. We first describe briefly the Wall invariant as it will be needed below. See also [15], [16], [6], and [12]. Let R be

a ring with unit, and let $\tilde{K}_0(R)$ be the *reduced projective class group* of R . Let C be an R -free chain complex such that C_i is finitely-generated (f.g.) for each i . Let $C^{(n)}$ be the n -skeleton of C . If the inclusion $C^{(n)} \hookrightarrow C$ is an R -domination of C then

$$H(C, C^{(n)}) = H_{n+1}(C, C^{(n)})$$

is a f.g. projective R -module. The *Wall invariant* of C is defined by

$$w(C) = (-1)^{n+1} [H_{n+1}(C, C^{(n)})] \in \tilde{K}_0(R).$$

This invariant vanishes if and only if C is R -chain equivalent to a finite R -free complex.

Now let X satisfy the hypotheses of Theorem 1. Let $\varphi: K \rightarrow X$ be a G -dominating map where K is a finite G -complex. By (4.5) and (4.1) we may assume that X is of finite-type, and that the inclusion $K \hookrightarrow X$, where $K = X^n$ and $n \geq 2$, is a G -domination. (We must assume that $n \geq 2$ in order that $\pi_1(K^H) \cong \pi_1(X^H)$ for every H .) If H is a subgroup of G , then the inclusion $K^H \hookrightarrow X^H$ is an NH/H -domination. Let $p_H: \tilde{X}^H \rightarrow X^H$ be the universal cover. The spaces \tilde{K}^H and \tilde{X}^H are $\widetilde{NH/H} = \pi_1(X^H) \backslash NH/H$ -complexes by (2.4), the cellular chain complexes $C(\tilde{K}^H)$ and $C(\tilde{X}^H)$ are $\mathbb{Z}(\widetilde{NH/H})$ -permutation complexes by (3.1), and the inclusion $C(K^H) \hookrightarrow C(X^H)$ is a $\mathbb{Z}(\widetilde{NH/H})$ -domination in which all maps are permutation maps. It easily follows that the induced map

$$C(\tilde{K}^H, p_H^{-1}(K^{>H})) \hookrightarrow C(\tilde{X}^H, p_H^{-1}(X^{>H}))$$

is a $\mathbb{Z}(\widetilde{NH/H})$ -domination for the free $\mathbb{Z}(\widetilde{NH/H})$ -chain complexes. Since $C(\tilde{K}^H, p_H^{-1}(K^{>H}))$ is finite, $w(C(\tilde{X}^H, p_H^{-1}(X^{>H})))$ is defined.

DEFINITIONS. Let X satisfy the hypotheses of Theorem 1.

2. The *relative Wall invariant* of X with respect to H is defined by

$$w_H(X) \equiv w(C(\tilde{X}^H, p_H^{-1}(X^{>H}))) \in \tilde{K}_0(\mathbb{Z}(\widetilde{NH/H})).$$

3. Define $E_1(X, G) = \sum_{H < G} w_H(X) \in \sum_{H < G} \tilde{K}_0(\mathbb{Z}(\widetilde{NH/H}))$.

DEFINITION 4. Let K and X be G -complexes. Let $\varphi: K \rightarrow X$ be a G -map. The map φ is called a *G -homology-equivalence* if

$$\varphi_*^H: H(q^{-1}(K^H)) \rightarrow H(p^{-1}(X^H))$$

is an isomorphism for each subgroup H of G , where $q: \hat{K} \rightarrow K$ is the pullback via φ of the universal cover of X .

The invariant for G -equivalence used quotients of the chain complexes $C(X^H)$. For G -homology-equivalence, the invariant will be defined using quotients of the chain complexes $\tilde{C}(X^H)$, where $\tilde{X}^H \subset \tilde{X}$ is the regular cover of X^H which contains the fixed basepoint $e^0 \in p^{-1}(X^G)$.

Assume as before that X is of finite type and the inclusion $K \hookrightarrow X$, where $K = X^n$ and $n \geq 2$, is a G -domination. Let $i_H: X^H \hookrightarrow X$ be the inclusion and

let $\pi_1(i_H): \pi_1(X^H) \rightarrow \pi_1(X)$ be the induced homomorphism. Since $\text{im}(\pi_1(i_H)) \subseteq \pi_1(X)$ has a trivial H -action, there is an induced action of NH/H on $\text{im}(\pi_1(i_H))$. Hence by (2.4), $\widehat{X^H}$ is an $\widehat{NH/H} = \text{im } \pi_1(i_H) \times_\alpha NH/H$ -complex. Analogous to the discussion above, the inclusion

$$C(\widehat{K^H}, p^{-1}(K^{>H}) \cap \widehat{K^H}) \hookrightarrow C(\widehat{X^H}, p^{-1}(X^{>H}) \cap \widehat{X^H})$$

is a $\mathbf{Z}(\widehat{NH/H})$ -domination for the free $\mathbf{Z}(\widehat{NH/H})$ -complexes. Hence, the invariant $w(C(\widehat{X^H}, p^{-1}(X^{>H}) \cap \widehat{X^H}))$ is defined.

DEFINITION 5. Let X satisfy the hypotheses of Theorem 1. We define

$$\begin{aligned} E_2(X, G) &= \sum_{H < G} w(C(\widehat{X^H}, p^{-1}(X^{>H}) \cap \widehat{X^H})) \\ &\in \sum_{H < G} \oplus \tilde{K}_0(\mathbf{Z}(\widehat{NH/H})). \end{aligned}$$

The invariants $E_1(X, G)$ and $E_2(X, G)$ are related.

LEMMA 6. Let X satisfy the hypotheses of Theorem 1. Then

(i) There is a homomorphism $\sigma_H: \widehat{NH/H} \rightarrow \widehat{NH/H}$, and therefore an induced homomorphism

$$\sigma_* = \sum_{H < G} \sigma_H: \sum_{H < G} \oplus \tilde{K}_0(\mathbf{Z}(\widehat{NH/H})) \rightarrow \sum_{H < G} \oplus \tilde{K}_0(\mathbf{Z}(\widehat{NH/H})).$$

(ii) The invariants $E_1(X, G)$ and $E_2(X, G)$ satisfy the relation

$$E_2(X, G) = \sigma_* E_1(X, G).$$

In particular, if each $\pi_1(i_H)$ is an isomorphism then $E_2(X, G) = E_1(X, G)$.

PROOF. (i) The homomorphism σ_H is defined by $(\gamma, g) \mapsto (\pi_1(i_H)\gamma, g)$, $\gamma \in \pi_1(X^H)$ and $g \in NH/H$.

(ii) By a relative form of (3.8)

$$C(\widehat{X^H}, p^{-1}(X^{>H}) \cap \widehat{X^H}) \cong \mathbf{Z}(\widehat{NH/H}) \otimes_{\mathbf{Z}(\widehat{NH/H})} C(\widehat{X^H}, p_H^{-1}(X^{>H})).$$

Hence $w(C(\widehat{X^H}, p^{-1}(X^{>H}) \cap \widehat{X^H})) = \sigma_{H_*} w_H(X)$. By summing over all subgroups H we obtain the desired result. \square

It is clear that if $\varphi: K \rightarrow X$ is a G -equivalence then it is a G -homology-equivalence as well. In the context above, $E_1(X, G) = 0$ implies $E_2(X, G) = 0$. At the end of this section we will give an example of a G -complex X with the property that $E_1(X, G) \neq 0$ but $E_2(X, G) = 0$, which implies that we can construct a G -homology-equivalence between X and a finite complex when a G -equivalence does not exist.

Also note that since $g \in G$ is a homeomorphism it induces isomorphisms between projective class groups and $g_* w_H(X) = w_{gHg^{-1}}(X)$ and $g_* \sigma_* w_H(X) = \sigma_* w_{gHg^{-1}}(X)$. We could have defined both invariants as sums over conjugacy classes of subgroups instead of as sums over all subgroups.

PROOF OF THEOREM 1 (i). The element $E_1(X, G)$ is an invariant of G -homotopy type since each $w_H(X)$ is an invariant of $\mathbf{Z}(\widetilde{NH/H})$ -chain homotopy type.

To prove the condition is necessary, suppose that $\varphi: K \rightarrow X$ is a G -equivalence. There is an induced $\mathbf{Z}(\widetilde{NH/H})$ -chain equivalence

$$\varphi_*^H: C(\widetilde{K^H}, q_H^{-1}(K^{>H})) \rightarrow C(\widetilde{X^H}, p_H^{-1}(X^{>H})),$$

where $q_H: \widetilde{K^H} \rightarrow K^H$ is the universal cover.

Thus, $w_H(X) = 0$ for each H . By definition $E_1(X, G) = 0$.

To prove the sufficiency we first assume as above that X is a G -complex of finite type which is G -dominated by its n -skeleton, K , and that $n \geq 2$. By induction, we assume that K is an at most $(n+1)$ -dimensional subcomplex of X and we assume that the inclusion map $i_{H'}: K^{H'} \hookrightarrow X^{H'}$ is a homotopy equivalence for all subgroups H' such that $|H'| > |H|$. We let H represent a conjugacy class of subgroups.

Consider the short exact sequence of $\mathbf{Z}(\widetilde{NH/H})$ -complexes

$$\begin{aligned} C(p_H^{-1}(X^{>H}), p_H^{-1}(K^{>H})) &\rightarrowtail C(\widetilde{X^H}, \widetilde{K^H}) \\ &\twoheadrightarrow C(\widetilde{X^H}, p_H^{-1}(X^{>H}) \cup \widetilde{K^H}). \end{aligned}$$

By a relative form of (3.9),

$$C(p_H^{-1}(X^{>H}), p_H^{-1}(K^{>H})) \cong \mathbf{Z}(\widetilde{NH/H}) \otimes_R C(\widetilde{X^{>H}}, \widetilde{K^{>H}})$$

where $R = \mathbf{Z}(\pi_1(X^{>H})_{x_\alpha} NH/H)$. By the inductive hypothesis the inclusion $i_{>H}: K^{>H} \hookrightarrow X^{>H}$ is a homotopy equivalence. Then $C(\widetilde{X^{>H}}, \widetilde{K^{>H}})$ is acyclic and, therefore, by the universal coefficient theorem, the complex

$$C(p_H^{-1}(X^{>H}), p_H^{-1}(K^{>H}))$$

is acyclic. Thus, $H(\widetilde{X^H}, \widetilde{K^H}) \cong H(\widetilde{X^H}, p_H^{-1}(X^{>H}) \cup \widetilde{K^H})$.

By assumption, the only nonzero homology group for either complex is in dimension $(n+1)$. Since $w_H(X) = 0$ by the isomorphism above

$$[H_{n+1}(\widetilde{X^H}, \widetilde{K^H})] = 0 \in \tilde{K}_0(\mathbf{Z}(\widetilde{NH/H})).$$

Thus, there exist finitely-generated free $\mathbf{Z}(\widetilde{NH/H})$ -modules, F_1 and F_2 , such that

$$F_1 \cong H_{n+1}(\widetilde{X^H}, \widetilde{K^H}) \oplus F_2.$$

In the following construction we will attach a finite number of cells to the subspace K^H and to its homeomorphic copies $K^{gHg^{-1}}$, $g \in G$. We then replace X by a G -equivalent space of finite-type with an $(n+1)$ -subcomplex L satisfying: $i_{H'}: L^{H'} \hookrightarrow X^{H'}$ is a homotopy equivalence for all subgroups H' such that either $|H'| > |H|$ or H' is conjugate to H .

Step 1. We define a complex $K' = K \vee (\bigvee_{(i,g)} S_{(i,g)}^n)$ where $\{i\}$ represents a finite set of generators of F_2 , and g ranges over some set of coset representatives of G/H . We extend the action of G to K' by defining $g'S_{(i,g)}^n \equiv S_{(i,g'g)}^n$ where g is a representative of G/H and $g' \in G$.

Define a G -map $\varphi': K' \rightarrow X$ by $\varphi'|_K = \text{inclusion}$ and $\varphi'|_{S_{(i,g)}^n} = x_0$ where x_0 is the basepoint in X . Then φ' is clearly a G -domination extending the inclusion. By (4.1) we can replace X by a G -equivalent space X' of finite type in such a way that K'^H is a subcomplex of X'^H . For this new space we have the short exact sequence

$$H_{n+1}(\widetilde{X'^H}, \widetilde{K'^H}) \twoheadrightarrow H_{n+1}(\widetilde{X'^H}, \widetilde{K'^H}) \twoheadrightarrow H_n(\widetilde{K'^H}, \widetilde{K'^H})$$

which splits. Since the first term is clearly isomorphic to $H_{n+1}(\widetilde{X'^H}, \widetilde{K'^H})$ we have the direct sum decomposition

$$\begin{aligned} H_{n+1}(\widetilde{X'^H}, \widetilde{K'^H}) &\cong H_{n+1}(\widetilde{X'^H}, \widetilde{K'^H}) \oplus H_n(\widetilde{K'^H}, \widetilde{K'^H}) \\ &\cong H_{n+1}(\widetilde{X'^H}, \widetilde{K'^H}) \oplus F_2 \cong F_1. \end{aligned}$$

Step 2. For each generator t of F_1 associate an element $\sigma_t \in \pi_{n+1}(X'^H, K'^H)$ using the following sequence of isomorphisms

$$F_1 \cong H_{n+1}(\widetilde{X'^H}, \widetilde{K'^H}) \xleftarrow{h} \pi_{n+1}(\widetilde{X'^H}, \widetilde{K'^H}) \xrightarrow{p_H} \pi_{n+1}(X'^H, K'^H),$$

where h represents the Hurewicz map. (These are all $\mathbf{Z}(NH/H)$ -maps by functorial properties.) Let $(\beta_t, \gamma_t) \in \sigma_t$.

Define a complex L by $L = K' \cup \bigcup_{(i,g)} e_{(i,g)}^{n+1}$ where the $(n+1)$ -cell $e_{(i,g)}^{n+1}$ has attaching map $g \circ \gamma_t$, g ranges over representatives of G/H , and t ranges over generators of F_1 . (This definition is independent of the choice of representatives in G/H since H fixes K'^H .) We extend the G -action on K' to L by defining $g'e_{(i,g)}^{n+1} = e_{(i,g'g)}^{n+1}$. The map $\psi: L \rightarrow X'$, extending the inclusion on K' by setting $\psi|_{e_{(i,g)}^{n+1}} = g \circ \beta_t$, is a G -dominating map.

By construction, the Hurewicz theorem and Whitehead's theorem, ψ^H is a homotopy equivalence. Likewise, since each g is a homeomorphism, it follows that $\psi^{gHg^{-1}}$ is a homotopy equivalence. Since the fixed-point sets for H' , where $|H'| > |H|$, were not changed in the replacement of X by X' in Step 1, then $\psi^{H'} = i_{H'}: L^{H'} \hookrightarrow X'^{H'}$ is a homotopy equivalence.

Finally, by (4.1) we can now assume that L is a subcomplex of X , and our induction step is complete.

The induction will end in a finite number of steps since G is finite. When $H = 1$ Steps 1 and 2 and (4.4) imply that $\psi: L \rightarrow X$, is a G -equivalence. \square

PROOF OF THEOREM 1 (ii). It is clear that the element $E_2(X, G)$ is an invariant of G -homotopy type.

To prove necessity, suppose that $\varphi: K \rightarrow X$ is a G -2-connected, G -homology-equivalence, where K is a finite G -complex. Let H be a subgroup of G . Considering X as an NH -complex, by (2.5) and (3.9)

$$C(p^{-1}(X^H)) \cong \mathbf{Z}(\widehat{NH}) \otimes_{\mathbf{Z}(\widehat{NH})} C(\widehat{X^H})$$

where $\widehat{NH} = \pi_1(X)x_\alpha\widehat{NH}$ and $NH = \text{im}(\pi_1(i_H))x_\alpha NH$. Since $\mathbf{Z}(\widehat{NH})$ is a free $\mathbf{Z}(\widehat{NH})$ -module, the universal coefficient theorem implies that

$$H(p^{-1}(X^H)) \cong \mathbf{Z}(\widehat{NH}) \otimes_{\mathbf{Z}(\widehat{NH})} H(\widehat{X^H}).$$

A similar argument shows that $H(p^{-1}(X^{>H})) \cong \mathbf{Z}(\widehat{NH}) \otimes_{\mathbf{Z}(\widehat{NH})} H(\widehat{X^{>H}})$.

By replacing X by the mapping cylinder $M(\varphi)$ we can consider K^H as a subspace of X^H . Applying the above analysis we see that

$$0 = H(p^{-1}(X^H), p^{-1}(K^H)) \cong \mathbf{Z}(\widehat{NH}) \otimes_{\mathbf{Z}(\widehat{NH})} H(\widehat{X^H}, \widehat{K^H}).$$

Since the last term is a direct sum, we have $H(\widehat{X^H}, \widehat{K^H}) = 0$. Similarly, $H(\widehat{X^{>H}}, \widehat{K^{>H}})$ is zero.

Since $\widehat{X^H} \cap p^{-1}(X^{>H})$ is a union of components each homeomorphic to $\widehat{X^{>H}}$, it follows that

$$H(\widehat{K^H}, p^{-1}(K^{>H}) \cap \widehat{K^H}) \cong H(\widehat{X^H}, p^{-1}(X^{>H}) \cap \widehat{X^H})$$

as $\mathbf{Z}(\widehat{NH})$ -modules and as $\mathbf{Z}(\widehat{NH/H}) = \mathbf{Z}(\text{im}(\pi_1(i_H))x_\alpha NH/H)$ -modules. Since these chain complexes are $\mathbf{Z}(\widehat{NH/H})$ -free they are also chain equivalent. Hence, $w(C(\widehat{X^H}, p^{-1}(X^{>H}) \cap \widehat{X^H})) = 0$. It then follows that $E_2(X, G) = 0$.

Now assume that $E_2(X, G) = 0$. Assume that X is of finite type and the inclusion $K \hookrightarrow X$ is a G -domination, where $K = X^n$ and $n \geq 2$. Since the proof that X is G -homology-equivalent to a finite complex is similar in part to the proof of G -equivalence from part (i) of this theorem, we will omit many details.

The induction hypothesis states: K is a subcomplex of X of dimension (at most) $(n+1)$, and $i_{H'}: K^{H'} \hookrightarrow X^{H'}$ induces a homology isomorphism

$$H(p^{-1}(K^{H'})) \cong H(p^{-1}(X^{H'}))$$

for all subgroups H' such that $|H'| > |H|$. We let H represent a conjugacy class of subgroups.

From our inductive hypothesis and the previous discussion,

$$C(p^{-1}(X^{>H}) \cap \widehat{X^H}, p^{-1}(K^{>H}) \cap \widehat{K^H})$$

is acyclic. Thus, $H(\widehat{X^H}, \widehat{K^H}) \cong H(\widehat{X^H}, \widehat{K^H} \cup (p^{-1}(X^{>H}) \cap \widehat{X^H}))$. Since $E_2(X, G) = 0$ there exist f.g. free $\mathbf{Z}(\widehat{NH/H})$ -modules, F_1 and F_2 , such that $F_1 \cong H_{n+1}(\widehat{X^H}, \widehat{K^H}) \oplus F_2$.

In our construction we attach orbits of cells to $\bigcup_g K^{gHg^{-1}}$. We then replace X by a G -equivalent space of finite type containing an $(n+1)$ -subcomplex L

satisfying: the inclusion induces isomorphisms

$$H(p^{-1}(L^{H'})) \cong H(p^{-1}(X^{H'}))$$

for H' conjugate to H or $|H'| > |H|$.

Step 1. Follow Step 1 of part (i). The last line becomes $H_{n+1}(\widetilde{X'^H}, \widetilde{K'^H}) \cong H_{n+1}(\widetilde{X^H}, \widetilde{K^H}) \oplus F$, where F is $\mathbf{Z}(\widehat{NH/H})$ -free and has the same rank as F_2 . But now applying $[\mathbf{Z}(\widehat{NH/H}) \otimes_{\mathbf{Z}(NH/H)}]$ to both sides yields $H_{n+1}(\widetilde{X'^H}, \widetilde{K'^H}) \cong F_1$.

Step 2. The Hurewicz homomorphism

$$h: \pi_{n+1}(\widehat{X'^H}, \widehat{K'^H}) \rightarrow H_{n+1}(\widehat{X'^H}, \widehat{K'^H})$$

is onto by the Hurewicz theorem. Hence there is a $\mathbf{Z}(\widehat{NH/H})$ -homomorphism $\sigma: F_1 \rightarrow \pi_{n+1}(\widehat{X'^H}, \widehat{K'^H})$. Use this correspondence in Step 2 of part (i) to construct L and $\psi: L \rightarrow X$. Assume $L \subset X$ by (4.1). By construction we have the required homology isomorphisms. \square

EXAMPLE 7. We will construct a \mathbf{Z}_2 -complex X of finite type which is \mathbf{Z}_2 -dominated by its 2-skeleton. For this complex $E_1(X, \mathbf{Z}_2) \neq 0$ but $E_2(X, \mathbf{Z}_2) = 0$. Thus X is not \mathbf{Z}_2 -equivalent to a finite complex, but there does exist $\psi: L \rightarrow X$ where L is finite and ψ is a G -homology-equivalence.

The group $\tilde{K}_0(\mathbf{Z}(\mathbf{Z}_{23})) \cong \mathbf{Z}_3$ [12, p. 30]. Let w_0 be a generator. Define a 2-complex $K = S^1 \cup_{23} e^2$, by adjoining a 2-cell to S^1 by a map of degree 23. Then $\pi_1(K) \cong \mathbf{Z}_{23}$. By [15, Theorem F] there exists a complex Y of finite type such that $K \subset Y^2$, $Y^1 = K^1$, Y^2 dominates Y and $w(Y) = w(C(\tilde{Y})) = w_0 \in \tilde{K}_0(\mathbf{Z}(\mathbf{Z}_{23}))$. The space Y will be $X^{\mathbf{Z}_2}$ in our example.

Let $X = Y \cup e_1^2 \cup e_2^2$ where the 2-cells are attached via $1: S^1 \rightarrow S^1 \subset Y$. Clearly $\pi_1(X) = 1$. Let \mathbf{Z}_2 act on X by switching the 2-cells.

Let $j: Y^2 \hookrightarrow Y$ be the inclusion, let $s: Y \rightarrow Y^2$ be a right homotopy inverse such that $s|_{Y^1}$ is the identity and let $F: js \cong 1_Y(\text{rel } Y^1)$ be a homotopy. We will extend these to \mathbf{Z}_2 -maps as follows: extend s to $s': X \rightarrow X^2$ be defining it to be the identity on the 2-cells; extend F to $F': X \times I \rightarrow X$ by $(x, t) \mapsto x$, $x \in e_1^2 \cup e_2^2$ and $t \in I$. Let $i: X^2 \hookrightarrow X$. Then s' is a right \mathbf{Z}_2 -homotopy inverse of i . Therefore, X^2 \mathbf{Z}_2 -dominates X .

The invariant

$$E_1(X, \mathbf{Z}_2) = w_{\mathbf{Z}_2}(X) + w_1(X) = w(Y) + w_1(X) = w_0 \neq 0.$$

($w_1(X) = w(C(X, Y)) \in \tilde{K}_0(\mathbf{Z}(\mathbf{Z}_2)) = 0$.) The invariant $E_2(X, \mathbf{Z}_2) = w(C(X^{\mathbf{Z}_2})) + w(C(X, X^{\mathbf{Z}_2})) \in \tilde{K}_0(\mathbf{Z}) \oplus \tilde{K}_0(\mathbf{Z}(\mathbf{Z}_2)) = 0$. Hence, $E_2(X, \mathbf{Z}_2) = 0$. \square

The results of this section suggest a related problem: Let G be a finite group. Define a Wall invariant in the category of RG -permutation complexes and maps. We note here that such an invariant can be defined, but we omit the details.

6. Further results. Let $\varphi: K \rightarrow X$ be a G -dominating map where K is a finite G -complex. In the previous sections we have studied the problems of replacing φ by either a G -homotopy equivalence or a G -homology equivalence $\psi: L \rightarrow X$ where L is a finite G -complex. Since we can always assume that a G -domination is G -2-connected, a positive solution to either of these problems would show that it is possible to replace φ by a G -map $\psi: L \rightarrow X$ which is an ordinary homotopy equivalence.

It is the object of this section to investigate the problem of the existence of a G -homology equivalence further, with the additional necessary assumption that there exists a G -map which is an ordinary homotopy equivalence. We note that, by [8, Theorem 1.1] any finite complex whose reduced homology with \mathbb{Z}_p -coefficients is trivial, can be the fixed-point set of finite contractible \mathbb{Z}_p -complex. Therefore, a G -map which is an ordinary homotopy equivalence (in this case a contraction) need not be a G -homology-equivalence.

Our main results are Theorems 1 and 2.

THEOREM 1. *Let X be a G -complex such that X^H is connected and $X^G \neq \emptyset$. Suppose that $\pi_1(X)$ is finite and that either G acts semi-freely on X or G is solvable. Let $\varphi: K \rightarrow X$ be a G -dominating map where K is a finite G -complex, the dimension of K is n , and $n \geq 2$. Then the following statements are equivalent:*

- (i) *The invariant $E_2(X, G) = 0$.*
- (ii) *There exists a G - n -connected, G -dominating map $\psi: L \rightarrow X$ where L is a finite complex and $\dim(L) = n + 1$ such that ψ is an ordinary homotopy equivalence.*

In Theorem 2 we let $\widehat{\varphi}^G: \widehat{K}^G \rightarrow \widehat{X}^G$ be the restriction of $\tilde{\varphi}: \tilde{K} \rightarrow \tilde{X}$ to the regular cover of K^G in \tilde{K} . Also, we let $\widehat{\widehat{\varphi}}^G: \widehat{\widehat{K}}^G \rightarrow \widehat{\widehat{X}}^G$ be the restriction to the complete inverse image of K^G in \tilde{K} . The inclusion of X^G into X is denoted by i_G .

THEOREM 2. *Let X be a G -complex and let G act semi-freely on X . Suppose that X^G is connected and not empty. Let $\varphi: K \rightarrow X$ be a G -2-connected G -domination which is also a homotopy equivalence.*

- (i) *If each $H_i(M(\widehat{\varphi}^G), \widehat{K}^G)$ has a finite $\mathbb{Z}(\text{im}(\pi_1(i_G)))$ -free resolution of length 2, then $E_2(X, G) = 0$.*
- (ii) *If each $H_i(M(\widehat{\widehat{\varphi}}^G), \widehat{\widehat{K}}^G)$ has a finite $\mathbb{Z}\tilde{G}$ resolution of length 2, then $E_2(X, G) = w(C(\widehat{X}^G))$. In addition, if X is simply connected then $E_2(X, G) = 0$.*

Some special cases of Theorem 2 are of interest.

COROLLARY 3. *Let X , G and $\varphi: K \rightarrow X$ be defined as in Theorem 2. Suppose that $\pi_1(X)$ is finite. Then the finite resolutions of $H_i(M(\varphi^G), K^G)$ and $H_i(M(\varphi^G), K^G)$ in parts (i) and (ii) may be taken to be of any finite length.*

PROOF. It is sufficient to show that either module has a finite resolution of length 2.

Let π be a finite group and $F_n \rightarrow F_{n-1} \rightarrow \cdots \rightarrow F_0 \twoheadrightarrow M$ be a finite $\mathbb{Z}\pi$ -free resolution of M . There exists a finite projective resolution of M of length 2, $P_1 \rightarrow P_0 \twoheadrightarrow M$, by [13, Theorem 4.12]. We may assume that P_0 is free. There exists a chain map $\varphi: P \rightarrow F$ such that the algebraic mapping cone $M(\varphi)$ is acyclic. By Schanuel's Lemma [1, p. 36] applied to $M(\varphi)$ and the zero complex, it follows that P_1 is stably-free. Thus, there exist f.g. free modules F'_1 and F'_2 such that $F'_2 \cong F'_1 \oplus P_1$. Then $F'_1 \oplus P'_1 \rightarrow F'_1 \oplus P_0 \twoheadrightarrow M$ is the desired resolution. \square

COROLLARY 4. *Let X be a \mathbb{Z}_n -complex which is simply connected. Let K be a finite \mathbb{Z}_n -complex and $\varphi: K \rightarrow X$ be a \mathbb{Z}_n -dominating 2-connected map which is a homotopy equivalence. Suppose that $H(M(\varphi^{\mathbb{Z}_n}), K^{\mathbb{Z}_n})$ is finite and prime to n . Then X is \mathbb{Z}_n -homology equivalent to a finite complex.*

PROOF OF COROLLARY 4. By [8, Lemma 1.1] each $H_i(M(\varphi^{\mathbb{Z}_n}), K^{\mathbb{Z}_n})$ has a finite free $\mathbb{Z}(\mathbb{Z}_n)$ -resolution of length 2. The rest follows from Corollary 3 and (6.1). \square

The proofs of the main theorems will follow from Lemmas 5–7.

LEMMA 5. *Let G be a finite group. Then G is solvable if and only if for each subgroup H of G there exists a maximal proper subgroup M of H which is normal in H .*

PROOF. Recall that a finite group is solvable if and only if the factor groups in a composition series are cyclic of prime order [7, p. 139].

Let G be a solvable group. Since each subgroup of a solvable group is solvable, it suffices to prove that a solvable group has a maximal proper subgroup which is normal. Let $G = G_0 \triangleright G_1 \triangleright \cdots \triangleright G_n = 1$ be a composition series for G . Then G/G_1 is cyclic of prime order and thus G_1 is maximal in G .

Suppose that G is not solvable. Let $G \triangleright G_1 \triangleright G_2 \triangleright \cdots \triangleright G_n = 1$ be a composition series for G . Then there exists an i such that the order of G_i/G_{i+1} is the form pq where p is prime and $q \neq 1$. Hence G_i/G_{i+1} contains a subgroup of order p . It follows that although G_{i+1} is maximal among the normal subgroups of G_i , it is not maximal among the subgroups of G_i . \square

LEMMA 6. *Let G be a finite group. Let P be a finitely-generated projective $\mathbb{Z}G$ -module, and T be a finitely-generated free abelian group with trivial $\mathbb{Z}G$ -module structure. If $P \cong T$ as $\mathbb{Z}G$ -modules, then $P = T = 0$.*

PROOF. Let $\hat{H}(G, M)$ represent the complete derived homology and cohomology sequence of G with coefficients in the $\mathbf{Z}G$ -module M [4, Chapter XII]. For a projective P , $\hat{H}(G, P) = 0$ [4, p. 236]. If the trivial module $T \neq 0$, then there are finite direct sums $T = \Sigma \oplus \mathbf{Z}$ and $H(G, T) \cong \Sigma \oplus H(G, \mathbf{Z}) \neq 0$ by [4, p. 237] and thus P and T are not isomorphic as $\mathbf{Z}G$ -modules. \square

LEMMA 7. *Let P be a finitely-generated projective $\mathbf{Z}\pi$ -module where π is a finite group. If $\mathbf{Z} \otimes_{\mathbf{Z}\pi} P = 0$, then $P = 0$.*

PROOF. Suppose $P \neq 0$. By a theorem of Swan [5, p. 556] $P \cong F \oplus P_0$ where F is a free $\mathbf{Z}\pi$ -module and P_0 is a projective ideal in $\mathbf{Z}\pi$. Since the tensor product commutes with direct sums it follows that if $\mathbf{Z} \otimes_{\mathbf{Z}\pi} P = 0$ then $F = 0$ and $P = P_0$.

By [5, p. 557] $P \otimes_{\mathbf{Z}} \mathbf{Q} \cong \mathbf{Z}\pi \otimes_{\mathbf{Z}} \mathbf{Q} \cong \mathbf{Q}\pi$ where \mathbf{Q} is the field of rational numbers. Now we have

$$0 = (\mathbf{Z} \otimes_{\mathbf{Z}\pi} P) \otimes_{\mathbf{Z}} \mathbf{Q} \cong \mathbf{Z} \otimes_{\mathbf{Z}\pi} (P \otimes_{\mathbf{Z}} \mathbf{Q}) \cong \mathbf{Z} \otimes_{\mathbf{Z}\pi} \mathbf{Q}\pi \supset \mathbf{Z} \otimes_{\mathbf{Z}\pi} \mathbf{Z}\pi \cong \mathbf{Z}.$$

Thus, we have a contradiction. \square

PROOF OF THEOREM 1. (i) \Rightarrow (ii) by the proof of (5.1). (ii) \Rightarrow (i): By (4.1) we assume that $\varphi: K \rightarrow X$ is inclusion, $K = X^n$ and $n \geq 2$.

Suppose first that G is solvable.

By assumption $C(\tilde{X}, \tilde{K})$ is acyclic. Suppose that for all H' such that $|H'| < |H|$, $C(\widehat{X^{H'}}, \widehat{K^{H'}})$ and $C(X^{H'}, K^{H'})$ are acyclic, and let H be a subgroup of G . By Lemma 5 there exists a proper normal subgroup M of H such that H/M is a cyclic group of prime order. Thus H/M acts semi-freely on X^M . Note that its fixed-point set is $(X^M)^{H/M} = X^H$. To simplify notation let $\widehat{X^H} = p^{-1}(X^H) \cap \widehat{X^M}$.

(1) Consider the short exact sequence of $\mathbf{Z}(\text{im}(\pi_1(i_M)))$ -modules. (We ignore the action of H/M on the spaces in this step.)

$$C(\widehat{X^H}, \widehat{K^H}) \twoheadrightarrow C(\widehat{X^M}, \widehat{K^M}) \rightarrow C(\widehat{X^M}, \widehat{X^H} \cup \widehat{K^M}).$$

By the inductive assumption $C(\widehat{X^M}, \widehat{K^M})$ is acyclic. Thus $H_{i+1}(\widehat{X^M}, \widehat{X^H} \cup \widehat{K^M}) \cong H_i(\widehat{X^H}, \widehat{K^H})$ for all i . By assumption φ^H is $(n-1)$ -connected, and thus $H_i(\widehat{X^H}, \widehat{K^H}) = 0 = H_i(\widehat{X^H}, \widehat{K^H})$ for $i < (n-1)$, where $n = \dim(K)$. Since $C(\widehat{K^M}, \widehat{K^H})$ dominates $C(\widehat{X^M}, \widehat{X^H})$,

$$H_i(\widehat{X^M}, \widehat{X^H} \cup \widehat{K^M}) = 0$$

for $i > n+1$. Thus, the only nonzero groups are $H_{n+1}(\widehat{X^M}, \widehat{X^H} \cup \widehat{K^M}) \cong H_n(\widehat{X^H}, \widehat{K^H})$.

The module $H_{n+1}(\widehat{X^M}, \widehat{X^H} \cup \widehat{K^M})$ is finitely-generated $\mathbf{Z}(\text{im}(\pi_1(i_M)))$ -projective.

(2) By relative forms of (3.8) and (3.9) $C(X^H, K^H) \cong \mathbb{Z} \otimes C(\widehat{X^H}, \widehat{K^H})$, where we tensor over $\mathbb{Z}(\text{im } \pi_2(i_M))$. By Step 1 the only nonzero homology group of $C(\widehat{X^H}, \widehat{K^H})$ is a f.g. projective. Hence, by the universal coefficient theorem, $H_n(X^H, K^H) \cong \mathbb{Z} \otimes H_n(\widehat{X^H}, \widehat{K^H})$, which is a f.g. \mathbb{Z} -free module.

(3) Consider the sequence of $\mathbb{Z}(H/M)$ -complexes

$$C(X^H, K^H) \twoheadrightarrow C(X^M, K^M) \rightarrow C(X^M, X^H \cup K^M).$$

By the inductive assumption and Step 2, the only nonzero groups are $H_{n+1}(X^M, X^H \cup K^M) \cong H_n(X^H, K^H)$. Since $C(K^M, K^H)$ dominates $C(X^M, X^H)$ as free $\mathbb{Z}(H/M)$ -complexes, $H_{n+1}(X^M, X^H \cup K^M)$ is a finitely-generated projective $\mathbb{Z}(H/M)$ -module.

(4) Since the action of H/M on $C(X^H, K^H)$ is necessarily trivial, Lemma 6 implies that

$$H_{n+1}(X^M, X^H \cup K^M) = H_n(X^H, K^H) = 0.$$

By Lemma 7, since

$$H_n(X^H, K^H) \cong \mathbb{Z} \otimes H_n(\widehat{X^H}, \widehat{K^H}) \quad \text{and} \quad H_n(X^H, K^H) = 0, \\ H_n(\widehat{X^H}, \widehat{K^H}) = 0 \quad \text{and thus} \quad H_{n+1}(\widehat{X^M}, \widehat{X^H} \cup \widehat{K^M}) = 0.$$

The induction step is complete.

Now suppose that G acts semi-freely on X . Since we only need to consider $C(\tilde{X}, \widehat{\tilde{X}^G} \cup \tilde{K})$ and $C(\widehat{\tilde{X}^G}, \widehat{\tilde{K}^G})$ to check for G -homology equivalence, just consider G and $M = 1$ in the above induction step. \square

The proof of Theorem 1 can be applied more generally if we know something about the action of G on X . Namely, if for each nontrivial subgroup H of G there exists a subgroup M of G such that M is normal in H , $M \neq H$ and H/M acts semi-freely on X^H , then the proof of Theorem 1 goes through. Lemma 5 shows that the proof cannot be extended to any larger class of subgroups without some extra conditions on the action.

The next lemma is used in the proof of Theorem 2.

LEMMA 8. *Let C be a free R -complex, and let $\varphi: D \rightarrow C$ be an R -dominating map, where D is a finite free R -complex. Suppose that each $H_i(M(\varphi))$ has a finite free R -resolution of length 2. Then C is R -chain equivalent to a finite free R -complex.*

PROOF. We assume by induction that the map $\varphi: D \rightarrow C$ has the property that $H_j(M(\varphi)) = 0$ for $j < i$. To prove the lemma it is sufficient to construct a finite free chain complex E and chain map $\psi: E \rightarrow C$ such that $H_j(M(\psi)) = 0$ for $j < i + 1$ and $H_j(M(\psi)) = H_j(M(\varphi))$ for $j > i + 1$. The construction will terminate in a finite number of steps since φ is a domination and D is finite.

The construction will be in two steps. Let $F' \twoheadrightarrow F \xrightarrow{h} H_{i+1}(M(\varphi))$ be a finite free resolution of $H_{i+1}(M(\varphi))$.

(1) Use the map $h: F \twoheadrightarrow H_{i+1}(\overline{M}(\varphi))$ and [3, Lemma 3] to construct a finite free complex \overline{D} and map $\overline{\varphi}: \overline{D} \rightarrow C$. Then $H_j(M(\overline{\varphi})) \cong H_j(M(\varphi))$ for $j \neq i+1, i+2$. By construction, $H_{i+1}(M(\overline{\varphi})) = 0$ and $H_{i+2}(M(\overline{\varphi})) \cong H_{i+2}(M(\varphi)) \oplus F'$.

(2) Use the map $h': F' \hookrightarrow H_{i+2}(M(\varphi)) \oplus F' \cong H_{i+2}(M(\overline{\varphi}))$ and [3, Lemma 3] to construct a finite complex E and map $\psi: E \rightarrow C$ which satisfy the properties listed above. \square

PROOF OF THEOREM 2. Since G acts semi-freely on X ,

$$E_2(X, G) = w(C(\widehat{X^G})) + w(C(\tilde{X}, p^{-1}(X^G))).$$

By assumption $C(M(\tilde{\varphi}), \tilde{K})$ is acyclic and

$$H_0(M(\widehat{\varphi^G}), \widehat{K^G}) = H_0(M(\tilde{\varphi}), M(\widehat{\varphi^G}) \cup \tilde{K}) = 0.$$

We also have the following isomorphisms:

$$\begin{aligned} H_{i+1}(M(\tilde{\varphi}), M(\widehat{\varphi^G}) \cup \tilde{K}) &\cong H_i(M(\widehat{\varphi^G}), p^{-1}(K^G)) \\ &\cong \mathbb{Z}\tilde{G} \otimes H_i(M(\widehat{\varphi^G}), \widehat{K^G}). \end{aligned}$$

In the last term we tensor over $\mathbb{Z}(\text{im } \pi_1(i_G))$. Let $\varphi_{\#}^G: C(\widehat{K^G}) \rightarrow C(\widehat{X^G})$ and $\varphi_{\#}: C(\tilde{K}, p^{-1}(K^G)) \rightarrow C(\tilde{X}, p^{-1}(X^G))$ be induced chain maps, and let $M(\varphi_{\#}^G)$ and $M(\varphi_{\#})$ be the algebraic mapping cones. Then $H(M(\varphi_{\#}^G)) \cong H(M(\widehat{\varphi^G}), \widehat{K^G})$ and $H(M(\varphi_{\#})) \cong H(M(\tilde{\varphi}), M(\widehat{\varphi^G}) \cup \tilde{K})$.

(i) By assumption and Lemma 8, $w(C(\widehat{X^G})) = 0$.

Let $F_{i1} \twoheadrightarrow F_{i0} \twoheadrightarrow H_i(M(\varphi_{\#}^G))$ be a finite $\mathbb{Z}(\text{im}(\pi_1(i_G)))$ -resolution. By tensoring with $\mathbb{Z}\tilde{G}$ over $\mathbb{Z}(\text{im}(\pi_1(i_G)))$ we obtain an induced $\mathbb{Z}\tilde{G}$ -resolution

$$F'_{i1} \twoheadrightarrow F'_{i0} \twoheadrightarrow H_{i+1}(M(\varphi_{\#})).$$

Since $H_0(M(\varphi_{\#})) = 0$ we can apply Lemma 8 to obtain $w(C(\tilde{X}, p^{-1}(X^G))) = 0$.

(ii) By the assumption, the isomorphisms above and Lemma 8, $w(C(\tilde{X}, p^{-1}(X^G))) = 0$. Hence, $E_2(X, G) = w(C(\widehat{X^G}))$.

If X is simply-connected, then $w(C(\widehat{X^G})) = w(C(X^G)) \in \tilde{K}_0(\mathbb{Z}) = 0$. Thus, $E_2(X, G) = 0$. \square

BIBLIOGRAPHY

1. H. Bass, *Algebraic K-theory*, Benjamin, New York, 1968.
2. G. Bredon, *Introduction to compact transformation groups*, Academic Press, New York, 1972.
3. K. Brown, *Homological criteria for finiteness*, Comment. Math. Helv **50** (1975), 129–135.
4. H. Cartan and S. Eilenberg, *Homological algebra*, Princeton Univ. Press, Princeton, N. J., 1956.
5. C. Curtis and I. Reiner, *Representation theory of finite groups and associative algebras*, Wiley, New York, 1966.
6. S. M. Gersten, *A product formula for Wall's obstruction*, Amer. J. Math. **88** (1966), 337–346.
7. M. Hall, *The theory of groups*, Macmillan, New York, 1959.

8. L. Jones, *The converse to the fixed point theorem of P. A. Smith. I*, Ann. of Math. **94** (1971), 52–68.
9. A. Lundell and S. Weingram, *The topology of CW complexes*, Van Nostrand Reinhold, New York, 1969.
10. T. Matumoto, *On G-CW complexes and a theorem of J. H. C. Whitehead*, J. Fac. Sci. Univ. Tokyo Sect. I A Math. **18** (1971), 363–374.
11. J. Milnor, *Introduction to algebraic K-theory*, Princeton Univ. Press, Princeton, N. J., 1971.
12. ———, *Uses of the fundamental group*, Colloquium Lectures, 73rd Summer Meeting of the Amer. Math. Soc., University of Wisconsin, 1968.
13. D. Rim, *Modules over finite groups*, Ann. of Math. **69** (1959), 700–712.
14. E. H. Spanier, *Algebraic topology*, McGraw-Hill, New York, 1966.
15. C. T. C. Wall, *Finiteness conditions for CW complexes*, Ann. of Math. **81** (1965), 55–69.
16. ———, *Finiteness conditions for CW complexes. II*, Proc. Roy. Soc. London Ser. A **295** (1966), 129–139.

DEPARTMENT OF MATHEMATICS, FAIRFIELD UNIVERSITY, FAIRFIELD, CONNECTICUT 06430