

ASYMPTOTIC COMPLETENESS FOR CLASSES OF TWO, THREE, AND FOUR PARTICLE SCHRÖDINGER OPERATORS¹

BY

GEORGE A. HAGEDORN²

ABSTRACT. Formulas for the resolvent $(z - H)^{-1}$ are derived, where $H = H_0 + \sum_{i < j} \lambda_{ij} V_{ij}$ is an N particle Schrödinger operator with the center of mass motion removed. For a large class of two-body potentials and generic couplings $\{\lambda_{ij}\}$, these formulas are used to prove asymptotic completeness in the $N < 4$ body problem when the space dimension is $m > 3$. The allowed potentials belong to a space of dilation analytic multiplication operators which fall off more rapidly than $r^{-2-\epsilon}$ at ∞ . In particular, Yukawa potentials, generalized Yukawa potentials, and potentials of the form $(1 + r)^{-2-\epsilon}$ are allowed.

I. Introduction. The scattering theory of two particle nonrelativistic quantum mechanics is reasonably well understood for short range potentials. The wave operators are asymptotically complete, and the singular continuous spectrum is empty (see Agmon [1], Enss [42], and Reed and Simon [24]). Much less is known about general N particle scattering.

For $N = 3$ and space dimension $m > 3$, Faddeev [6] first proved completeness of the wave operators for a large class of potentials and almost all coupling constants. These results have been simplified and extended by Ginibre and Moulin [7], Thomas [36], and Howland [11]. All of these authors require the potentials to fall off faster than $r^{-2-\epsilon}$ at infinity and obtain Kuroda completeness for almost all couplings. Mourre [19] has extended these results to allow potentials which fall faster than $r^{-1-\epsilon}$, as long as the potentials are repulsive for large r .

For general N , Balslev and Combes [3] have proved the absence of singular continuous spectrum for dilation analytic potentials. Asymptotic completeness for potentials falling off faster than $r^{-2-\epsilon}$ has been proved for small couplings by Iorio and O'Carroll [13], and for repulsive potentials falling off faster than $r^{-5/2-\epsilon}$ by Lavine [18]. (These are single channel results only.) Hepp [10] and Sigal [26] have reduced general N -body asymptotic completeness to the verification of properties of certain operators. They conjecture that these hold for generic potentials in certain classes.

Simultaneous to our announcement [9] of results, Sigal [28] announced a proof of asymptotic completeness for generic elements of a space of dilation analytic potentials and almost all couplings. More recently, van Winter [43] announced a

Received by the editors May 23, 1978.

AMS (MOS) subject classifications (1970). Primary 47A40, 81A48.

¹Based on a Princeton University Doctoral Thesis.

²Partially supported by an NSF Graduate Fellowship.

proof of completeness for some analytic Hilbert-Schmidt potentials. Also, Enss [42] has obtained some many body results by beautiful, time dependent methods.

Our main results are the following:

THEOREM I.1. *Let $m \geq 3$ and $N \leq 4$. Let*

$$\tilde{H} = - \sum_{i=1}^N (2m_i)^{-1} \Delta_i + \sum_{i < j} \lambda_{ij} V_{ij} \quad \text{on } L^2(\mathbf{R}^{Nm})$$

be the Schrödinger operator for N particles moving in m dimensions. Let $H = H_0 + \sum_{i < j} \lambda_{ij} V_{ij}$ denote the Schrödinger operator on $L^2(\mathbf{R}^{(N-1)m})$ for the same system with the center of mass motion removed. Assume each V_{ij} may be factored as $V_{ij} = U_{ij} W_{ij}$ so that

- (i) *each U_{ij} and W_{ij} is dilation analytic in some strip,*
 - (ii) *$(1 + x_{ij}^2)^\gamma U_{ij}(x_{ij})$ and $(1 + x_{ij}^2)^\gamma W_{ij}$ belong to $L^p(\mathbf{R}^m) + L^\infty(\mathbf{R}^m)$ for some $p > m$ and $\gamma > \frac{1}{2}$,*
 - (iii) *bound state energies of three body subsystems are nonpositive when $N = 4$.*
- Then for generic couplings $\{\lambda_{ij}\}$, asymptotic completeness holds.*

THEOREM I.2. *The set of generic couplings for Theorem I.1 is large in the sense that its complement is a closed set of Lebesgue measure zero.*

Remarks. (1) Hypothesis (ii) of Theorem I.1 implies that $U_{ij}(-\Delta_{ij} + 1)^{-1/2}$ and $W_{ij}(-\Delta_{ij} + 1)^{-1/2}$ are compact on $L^2(\mathbf{R}^m)$. Thus, $H = H_0 + \sum \lambda_{ij} V_{ij}$ is understood in the sense of quadratic forms.

(2) Balslev [2] and Simon [32] have given sufficient conditions for Hypothesis (iii) to hold. See Theorem II.10. Yukawa potentials, generalized Yukawa potentials, and potentials of the form $(1 + r)^{-2-\epsilon}$ obey these conditions. Therefore, for these potentials, Theorems I.1 and I.2 establish asymptotic completeness for generic couplings.

(3) For the generic couplings, H has finitely many thresholds. Sigal [27] has also obtained this result.

(4) The generic couplings are precisely those for which no cluster Hamiltonian has a threshold resonance or threshold bound state.

(5) Theorem I.1 will be proved under the assumption that all the masses are finite. If one of the masses is infinite, the result still holds, but several lemmas from §V must be modified. Various ρ functions must be changed, and in many instances, Case 2 of Lemma II.3 must be used in place of Case 1.

Our proof of Theorems I.1 and I.2 is organized as follows: In §II, we collect various technical results from the literature. Little is new here, except that weighted spaces L^2_δ are replaced by more general L^p_δ spaces. In §III, N -body Kuroda completeness is reduced to the study of the resolvent $(z - H)^{-1}$. Formulas for the N -body resolvent are derived in §IV. Equation (IV.4) is a generalization of Faddeev's three body formula [6] to the N -body case. It is much simpler than the Yakubovskii formula [41], but we have not been able to prove that it has all the desirable properties of the Yakubovskii equation. A second, more complicated

formula (equation (IV.5)), is derived from equation (IV.4). This formula is equivalent to a formula of Narodetskii and Yakubovskii [20]. Equation (IV.5) is modified for $N \leq 4$, in order to study scattering. In §V, these equations are used to prove properties of $(z - H)^{-1}$ for $N \leq 4$ under certain hypotheses. The validity of these hypotheses is studied in §VI.

Given the absence of singular continuous spectrum (Theorem II.8) and the existence of wave operators (Theorem II.12), Theorems I.1 and I.2 follow immediately from Theorem III.1, Propositions V.1, 2, and 3, and Proposition VI.3.

REMARK. The principal reason we do not have results for $N \geq 5$ is that we have been unable to prove the absence of spurious zeros in equation (IV.10).

ACKNOWLEDGEMENTS. I am very grateful to Professor Barry Simon for introducing me to many body Schrödinger operators and for his advice throughout this work. In addition, I would like to thank the ETH and IHES for their hospitality during the Spring and Summer of 1977, and Dr. I. M. Sigal for pointing out an error in a rough draft.

Preliminary definitions. The Schrödinger operator for a system of N particles moving in m dimensions is $\tilde{H} = -\sum_{i=1}^N (2m_i)^{-1} \Delta_i + \sum_{i < j} V_{ij}$ on $L^2(\mathbb{R}^{Nm})$. The mass of the i th particle is m_i , Δ_i is the Laplacian in the i th variable, and the potential energy between particles i and j is the multiplication operator $V_{ij}(r_i - r_j)$. We remove the trivial center of mass motion (see [23]) from \tilde{H} , to obtain $H = H_0 + \sum_{i < j} V_{ij}$ on $\mathcal{H} = L^2(\mathbb{R}^{(N-1)m})$.

A *cluster decomposition* $D = \{C_i\}_{i=1}^k$ is a partition of the set $\{1, 2, \dots, N\}$ into k disjoint clusters C_i . The Hamiltonian H_D is defined as $H_0 + V_D$, where V_D is the sum of all V_{ij} with i and j in the same cluster. The Hilbert space \mathcal{H} may be decomposed as $\mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \dots \otimes \mathcal{H}_k \otimes \mathcal{H}(D)$, so that

$$H_D = h_1 \otimes 1 \otimes \dots \otimes 1 + 1 \otimes h_2 \otimes \dots \otimes 1 + \dots \\ + 1 \otimes \dots \otimes h_k \otimes 1 + 1 \otimes \dots \otimes 1 \otimes K_D.$$

The Hamiltonian h_i corresponds to the energy of the particles in cluster C_i alone. K_D is the kinetic energy of the centers of mass of the clusters in D .

For each cluster Hamiltonian h_i , we choose eigenfunctions $\eta_j^{(i)}$ of h_i , so that $\{\eta_j^{(i)}\}$ is an orthonormal basis for the subspace of \mathcal{H}_i generated by the eigenfunctions of h_i .

A *channel* α is a cluster decomposition $D(\alpha)$, together with an eigenfunction $\eta^{(i)} \in \{\eta_j^{(i)}\}$ for each h_i . We define $E_\alpha = \sum_{i=1}^k E_i$, where $h_i \eta^{(i)} = E_i \eta^{(i)}$, and let $P_\alpha: \mathcal{H} \rightarrow \mathcal{H}$ denote the orthogonal projection onto all vectors of the form $\eta^{(1)} \otimes \eta^{(2)} \otimes \dots \otimes \eta^{(k)} \otimes \phi$, where $\phi \in \mathcal{H}(D(\alpha))$ is arbitrary. Let $T_\alpha = 1 \otimes 1 \otimes \dots \otimes 1 \otimes K_{D(\alpha)} + E_\alpha$, so that $H_{D(\alpha)} P_\alpha = T_\alpha P_\alpha$.

The *channel wave operators* are defined as

$$\Omega_\alpha^\mp = \text{strong-} \lim_{t \rightarrow \pm \infty} e^{itH} e^{-itT_\alpha} P_\alpha$$

if the limits exist. The wave operators are *asymptotically complete* if

$$\mathcal{H} = \bigoplus_\alpha \text{Ran } \Omega_\alpha^+ \oplus \mathcal{H}_{\text{bound}} = \bigoplus_\alpha \text{Ran } \Omega_\alpha^- \oplus \mathcal{H}_{\text{bound}},$$

where $\mathcal{H}_{\text{bound}}$ is the span of the eigenfunctions for H .

Each Ω_α^\pm is an isometry, so the projection onto $\text{Ran } \Omega_\alpha^\pm$ is $\Omega_\alpha^\pm \Omega_\alpha^\pm^*$. Since the channel wave operators have orthogonal ranges, the projection onto $\bigoplus_\alpha \text{Ran } \Omega_\alpha^\pm$ is $\sum_\alpha \Omega_\alpha^\pm \Omega_\alpha^\pm^*$.

The ranges of the wave operators are contained in $\mathcal{H}_{\text{a.c.}}$, the absolutely continuous subspace for H . Thus, $\bigoplus \text{Ran } \Omega_\alpha^\pm \subseteq \mathcal{H}_{\text{a.c.}}$. A weaker form of completeness is the condition $P_{\text{a.c.}} = \sum_\alpha \Omega_\alpha^\pm \Omega_\alpha^\pm^*$. This is called *Kuroda completeness*. Kuroda completeness implies asymptotic completeness if H has no singular continuous spectrum.

II. Technical devices. In this section we collect several results which are required for the other sections.

A. Jacobi coordinates (see Reed and Simon [23]).

We will use two types of coordinate systems for $\mathbf{R}^{(N-1)m}$. Both of these systems have the property that H_0 is of the form $-\sum_{i=1}^{N-1} (2\mu_i)^{-1} \Delta_i$, where $\mu_i > 0$ and Δ_i is the Laplacian in the i th coordinate. The particle positions r_i and their differences $r_i - r_j$ are not simple in these coordinates.

To obtain *Jacobi coordinates* $\{\xi_i\}$, we begin by labelling the particles by the numbers $1, 2, \dots, N$. Then, for $1 \leq j \leq N-1$, we let ξ_j denote the vector from the center of mass particles $1, 2, \dots, j$ to particle $j+1$. Explicitly, $\xi_j = r_{j+1} - (\sum_{i \leq j} m_i)^{-1} \sum_{i \leq j} m_i r_i$, where r_i denotes the position of particle i .

The second type of coordinate system is used to put K_D , the free Hamiltonian for the centers of mass of the clusters of D , into the form $-\sum_{i=1}^k (2\mu_i)^{-1} \Delta^{(k)}$. For each cluster C_i , we choose Jacobi coordinates $\xi_1^i, \xi_2^i, \dots, \xi_{n(i)-1}^i$ for the $n(i)$ particles labelled by the elements of C_i . Then we choose Jacobi coordinates $\xi_1, \xi_2, \dots, \xi_{k-1}$ for the centers of mass of the clusters. The resulting coordinates

$$\xi_1^1, \dots, \xi_{n(1)-1}^1, \quad \xi_1^2, \dots, \xi_{n(2)-1}^2, \quad \xi_1, \dots, \xi_{k-1}$$

are called *clustered Jacobi coordinates* for the cluster decomposition D .

B. Quadratic form techniques.

Quadratic forms are used to define the Hamiltonian H and to make sense out of various expressions involving resolvents, potentials, and square roots of the potentials.

The free Hamiltonian H_0 is a positive selfadjoint operator on \mathcal{H} . We let \mathcal{H}_{+1} denote the operator domain of $H_0^{1/2}$ with inner product $\langle \phi, \psi \rangle_{+1} = \langle \phi, \psi \rangle + \langle H_0^{1/2} \phi, H_0^{1/2} \psi \rangle$. Since $\mathcal{H}_{+1} \subseteq \mathcal{H}$, each $\phi \in \mathcal{H}$ defines a bounded linear functional on \mathcal{H}_{+1} by $\psi \rightarrow \langle \phi, \psi \rangle$. Thus, \mathcal{H} may be viewed as a subspace of the dual space \mathcal{H}_{-1} of \mathcal{H}_{+1} .

For each pair i, j , $L^2(\mathbf{R}^{(N-1)m})$ decomposes into $L^2(\mathbf{R}^m) \otimes L^2(\mathbf{R}^{(N-2)m})$, where the first factor denotes functions of x_{ij} . Under this decomposition, $V_{ij} = v_{ij} \otimes 1$. We assume $v_{ij} = u_{ij} w_{ij}$ such that $u_{ij}(-\Delta_{ij} + 1)^{-1/2}$ and $w_{ij}(-\Delta_{ij} + 1)^{-1/2}$ are compact on $L^2(\mathbf{R}^m)$.

We denote $u_{ij} \otimes 1$ by U_{ij} and $w_{ij} \otimes 1$ by W_{ij} . The assumptions on u_{ij} and w_{ij} imply that $U_{ij}: \mathcal{H} \rightarrow \mathcal{H}_{-1}$ and $W_{ij}: \mathcal{H}_{+1} \rightarrow \mathcal{H}$ are bounded. Moreover [22], for each $\varepsilon > 0$, there exists $a > 0$, such that

$$|\langle \psi, V_{ij}\psi \rangle| = |\langle U_{ij}^* \psi, W_{ij}\psi \rangle| \leq \varepsilon \|\psi\|_{+1}^2 + a \|\psi\|^2$$

for all $\psi \in \mathcal{H}_{+1}$.

Given this bound, the *KLMN* Theorem [22] implies that there is a unique selfadjoint operator H with $D(H) \subseteq \mathcal{H}_{+1}$, such that

$$\langle \psi, H\psi \rangle = \left\langle \psi, \left(H_0 + \sum_{i < j} V_{ij} \right) \psi \right\rangle$$

for $\psi \in \mathcal{H}_{+1}$. For each cluster decomposition D , the Hamiltonian H_D is defined by the same method.

If $z \notin \sigma(H_D)$, then $(z - H_D)^{-1}$ is bounded from \mathcal{H}_{-1} into \mathcal{H}_{+1} . Hence, products of the form $W_{ij}(z - H_D)^{-1}U_{kl}$ are bounded on \mathcal{H} .

C. Estimates related to Kato smoothness.

Although Kato smoothness [15] does not play a central role in our asymptotic completeness proof, some estimates from the theory of smooth perturbations will be used.

At least seven equivalent definitions of Kato smoothness are available (see [15], [24]). For our purposes, the following one is the most convenient.

DEFINITION. A closed operator F is called *Kato smooth* with respect to a selfadjoint operator A , if

$$\|F\|_A^2 = (2\pi)^{-1} \sup_{\substack{z \notin \mathbf{R}, \phi \in D(F^*) \\ \|\phi\|=1}} |\langle F^* \phi, [(z - A)^{-1} - (\bar{z} - A)^{-1}] F^* \phi \rangle| < \infty.$$

LEMMA II.1. Let $\phi_1: \mathbf{R} \rightarrow \mathcal{H}$ and $\phi_2: \mathbf{R} \rightarrow \mathcal{H}$ belong to $L^2(\mathbf{R}, \mathcal{H})$, where \mathcal{H} is a Hilbert space; if

$$\hat{\phi}_j(\lambda) = (2\pi)^{-1/2} \text{l.i.m.} \int_{\mathbf{R}} \phi_j(t) e^{-it\lambda} dt \quad \text{for } j = 1, 2,$$

then

$$\int_{\mathbf{R}} \langle \phi_1(t), \phi_2(t) \rangle dt = \int_{\mathbf{R}} \langle \hat{\phi}_1(\lambda), \hat{\phi}_2(\lambda) \rangle d\lambda.$$

PROOF. The usual Plancherel theorem shows that Fourier transform $\hat{\cdot}: L^2(\mathbf{R}) \rightarrow L^2(\mathbf{R})$ is unitary. Consequently, $\hat{\cdot} \otimes 1: L^2(\mathbf{R}) \otimes \mathcal{H} \rightarrow L^2(\mathbf{R}) \otimes \mathcal{H}$ is unitary. The lemma follows from this by the natural identification of $L^2(\mathbf{R}) \otimes \mathcal{H}$ with $L^2(\mathbf{R}, \mathcal{H})$. \square

REMARK. In the following lemma we prove an inequality. Equality can be proved [15], [24], but we will only use the inequality.

LEMMA II.2. Suppose F is Kato smooth with respect to A . Then

$$\|F\|_A^2 = \pi^{-1} \sup_{\substack{z \notin \mathbf{R}, \phi \in D(F^*) \\ \|\phi\|=1}} |\text{Im } z| \|(z - A)^{-1} F^* \phi\|^2$$

and

$$\|F\|_A^2 \geq (2\pi)^{-1} \sup_{\|\phi\|=1} \int_{-\infty}^{\infty} \|F e^{-itA} \phi\|^2 dt.$$

PROOF.

$$\begin{aligned}
 (2\pi)^{-1} |\langle F^* \phi, [(z - A)^{-1} - (\bar{z} - A)^{-1}] F^* \phi \rangle| \\
 &= (2\pi)^{-1} |\bar{z} - z| |\langle F^* \phi, (\bar{z} - A)^{-1} (z - A)^{-1} F^* \phi \rangle| \\
 &= \pi^{-1} |\operatorname{Im} z| \|(z - A)^{-1} F^* \phi\|^2.
 \end{aligned}$$

The first statement of the lemma follows by taking the supremum over $z \notin \mathbf{R}$ and $\phi \in D(F^*)$ with $\|\phi\| = 1$.

The operator

$$(2\pi i)^{-1} [(\bar{z} - A)^{-1} - (z - A)^{-1}] = \pi^{-1} (\operatorname{Im} z) (z - A)^{-1} (\bar{z} - A)^{-1}$$

is positive when $\operatorname{Im} z > 0$. Let $K(z)$ be its positive square root. Since F is A -smooth, $\|K(z) F^* \phi\|^2 \leq \|F\|_A^2 \|\phi\|^2$ for all $\phi \in D(F^*)$. Therefore, $\operatorname{Ran} K(z) \subseteq D(F)$ and $\|FK(z)\| \leq \|F\|_A$.

If $\phi(\lambda) = [(\lambda + i\varepsilon - A)^{-1} - (\lambda - i\varepsilon - A)^{-1}] \phi$, then explicit computation shows $\hat{\phi}(t) = i\varepsilon^{-1} e^{-\varepsilon|t|} e^{-itA} \phi$. Thus, by Lemma II.1,

$$\begin{aligned}
 (2\pi)^{-1} \int_{-\infty}^{\infty} \|Fe^{-itA} \phi\|^2 e^{-2\varepsilon|t|} dt \\
 &= (2\pi)^{-2} \int_{-\infty}^{\infty} \|F[(\lambda + i\varepsilon - A)^{-1} - (\lambda - i\varepsilon - A)^{-1}] \phi\|^2 d\lambda \\
 &\leq \int_{-\infty}^{\infty} \|FK(\lambda + i\varepsilon)\|^2 \|K(\lambda + i\varepsilon)\phi\|^2 d\lambda \\
 &\leq \|F\|_A^2 \int_{-\infty}^{\infty} \|K(\lambda + i\varepsilon)\phi\|^2 d\lambda \\
 &= (2\pi i)^{-1} \|F\|_A^2 \int_{-\infty}^{\infty} \langle \phi, [(\lambda - i\varepsilon - A)^{-1} - (\lambda + i\varepsilon - A)^{-1}] \phi \rangle d\lambda \\
 &= \|F\|_A^2 \frac{\varepsilon}{\pi} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{1}{(\lambda - \mu)^2 + \varepsilon^2} d\langle \phi, E(\mu) \phi \rangle d\lambda \\
 &= \|F\|_A^2 \|\phi\|^2.
 \end{aligned}$$

The lemma follows by taking ε to zero. \square

DEFINITION. For $1 < p < \infty$ and $\delta \in \mathbf{R}$, $L_\delta^p(\mathbf{R}^m)$ is the Banach space of functions $f: \mathbf{R}^m \rightarrow \mathbf{C}$, such that $(1 + x^2)^{\delta/2} f(x) \in L^p(\mathbf{R}^m)$. The L_δ^p -norm of f is the L^p -norm of $(1 + x^2)^{\delta/2} f(x)$.

REMARKS. (1) Lemma II.3 extends results of Iorio and O'Carroll [13].

(2) If the operators F_1 and F_2 are equal in Lemma II.3, then we conclude that $F_1^* = F_2^*$ is H_D -smooth.

LEMMA II.3. Let D be a cluster decomposition with at least two clusters. Let $\delta > 1$, $m \geq 3$, $p_1 > m$, and $p_2 > m$. Suppose $\{\xi_1, \dots, \xi_{k-1}\}$ and $\{\xi_1, \dots, \xi_{k-1}\}$ are two (possibly identical) choices for the intercluster clustered Jacobi coordinates corresponding to the cluster decomposition D . Suppose $F_1 \in L_\delta^{p_1}(\mathbf{R}^m)$ and $F_2 \in L_\delta^{p_2}(\mathbf{R}^m)$.

Let F_1 and F_2 denote multiplication by $F_1(\xi_1)$ and $F_2(\xi_1)$, respectively. Since $\{\xi_i\}$ and $\{\xi_i\}$ are bases for the same space, $\xi_1 = \sum \alpha_i \xi_i$.

Case 1. If $\alpha_1 \neq 0$, then $F_1^*(z - H_D)^{-1}F_2$ is uniformly bounded and analytic for $z \in \mathbb{C} \setminus \sigma(H_D)$, with norm continuous boundary values as z approaches $\sigma(H_D)$ from above and below.

Case 2. If $\alpha_1 = 0$, then the same conclusions hold, except that the boundary values are only strongly continuous.

PROOF. For $\text{Im } z > 0$,

$$(z - H_D)^{-1} = -i \int_0^\infty e^{-itH_D} e^{itz} dt.$$

$e^{-itH_D} = e^{-itA} \otimes e^{-itB}$ on $L^2(\mathbb{R}^m) \otimes L^2(\mathbb{R}^{(N-2)m})$, where the first factor denotes functions of ξ_1 . A is a negative multiple of the Laplacian, and e^{-itA} has an explicitly known integral kernel [22]. From the form of the integral kernel, an interpolation argument shows that e^{-itA} is bounded from $L^r(\mathbb{R}^m)$ to $L^s(\mathbb{R}^m)$ when $1 < r \leq 2$ and $r^{-1} + s^{-1} = 1$. Moreover, the norm of e^{-itA} is dominated by $(ct)^{-m(r^{-1}-2^{-1})}$.

Case 1. Since e^{-itB} is unitary and commutes with F_1^* ,

$$\|F_1^* e^{-itH_D} F_2\| = \|F_1^* e^{-itA} F_2\|.$$

To compute $\|F_1^* e^{-itA} F_2 \phi\|$, fix $\xi_2, \xi_3, \dots, \xi_{k-1}$, and compute the L^2 -norm in the ξ_1 variable first.

Hölder's inequality shows that F_1 and F_2 belong to $L^q(\mathbb{R}^m)$ for $m - \varepsilon < q < m + \varepsilon$. With ξ_2, \dots, ξ_{k-1} fixed, $F_2(\sum \alpha_i \xi_i) \phi(\xi_1, \dots, \xi_{k-1})$ belongs to $L^r(\mathbb{R}^m, d\xi_1)$, where $r^{-1} = q^{-1} + \frac{1}{2}$. e^{-itA} maps this into $L^s(\mathbb{R}^m, d\xi_1)$, where $s = \frac{1}{2} - q^{-1}$. Multiplication by $\overline{F_1(\xi_1)}$ then maps this back to $L^2(\mathbb{R}^m, d\xi_1)$. The result is

$$\begin{aligned} & \int |(\overline{F_1} e^{-itA} F_2 \phi)(\xi_1, \dots, \xi_{k-1})|^2 d\xi_1 \\ & \leq (ct)^{-2m/q} \|F_1\|_q^2 \|F_2\|_q^2 \alpha_1^{-2m/q} \int |\phi(\xi_1, \dots, \xi_{k-1})|^2 d\xi_1. \end{aligned}$$

By integrating over ξ_2, \dots, ξ_{k-1} , we obtain

$$\|F_1^* e^{-itA} F_2 \phi\| \leq (\alpha_1 ct)^{-m/q} \|F_1\|_q \|F_2\|_q \|\phi\|_2.$$

Since this is valid for $q \in (m - \varepsilon, m + \varepsilon)$, $\|F_1^* e^{-itA} F_2\|$ is an L^1 function of t .

For $\text{Im } z > 0$,

$$F_1^*(z - H_D)^{-1} F_2 = -i \int_0^\infty F_1^* e^{-itH_D} F_2 e^{itz} dt.$$

Thus,

$$\|F_1^*(z - H_D)^{-1} F_2\| \leq \int_0^\infty \|F_1^* e^{-itA} F_2\| dt \leq C.$$

Also,

$$\begin{aligned} & \|F_1[(z - H_D)^{-1} - (w - H_D)^{-1}]F_2\| \\ & \leq \int_0^\infty \|F_1^* e^{-itA} F_2\| |e^{itz} - e^{itw}| dt \end{aligned}$$

when $\text{Im } z > 0$ and $\text{Im } w > 0$. The norm Hölder continuity now follows by the bound: $|e^{itz} - e^{itw}| \leq \text{Min}(2, t|z - w|)$.

Case 2. It follows from Case 1 that $F_1^*(z - H_D)^{-1}F_1$ and $F_2^*(z - H_D)^{-1}F_2$ are uniformly bounded. Thus, F_1^* and F_2^* are H_D -smooth.

Since $\alpha_1 = 0$,

$$e^{-itH_D} = e^{-itA} \otimes e^{-itB} \otimes e^{-itC}$$

on $L^2(\mathbb{R}^m, d\xi_1) \otimes L^2(\mathbb{R}^m, d\xi_2) \otimes L^2(\mathbb{R}^{(N-3)m})$. A and B are negative multiples of the Laplacian; C is selfadjoint.

For $\phi, \psi \in \mathcal{H}$ and $\text{Im } z > 0$,

$$\begin{aligned} \langle \phi, F_1^*(z - H_D)^{-1}F_2\psi \rangle &= -i \int_0^\infty \langle \phi, F_1^* e^{-itH_D} F_2\psi \rangle e^{itz} dt \\ &= -i \int_0^\infty \langle e^{itB}\phi, F_1^* e^{-itA} e^{-itC} F_2\psi \rangle e^{itz} dt \\ &= -i \int_0^\infty \langle F_2^* e^{itB}\phi, F_1^* e^{-itA} e^{-itC}\psi \rangle e^{itz} dt. \end{aligned}$$

Hence,

$$\begin{aligned} |\langle \phi, F_1^*(z - H_D)^{-1}F_2\psi \rangle| &\leq \left(\int_0^\infty \|F_2^* e^{itB}\phi\|^2 dt \right)^{1/2} \left(\int_0^\infty \|F_1^* e^{-itA}\psi\|^2 dt \right)^{1/2} \\ &= \left(\int_0^\infty \|F_2^* e^{itH_D}\phi\|^2 dt \right)^{1/2} \left(\int_0^\infty \|F_1^* e^{-itH_D}\psi\|^2 dt \right)^{1/2} \\ &\leq \|F_2^*\|_{H_D} \|F_1^*\|_{H_D} \|\phi\| \|\psi\|. \end{aligned}$$

Here we have used the Schwarz and Hölder inequalities and Lemma II.1.

To prove the strong continuity, the same type of computation shows:

$$\begin{aligned} & |\langle \phi, F_1^*[(z - H_D)^{-1} - (w - H_D)^{-1}]F_2\psi \rangle| \\ & \leq \left(\int_0^\infty \|F_2^* e^{itH_D}\phi\|^2 dt \right)^{1/2} \left(\int_0^\infty \|F_1^* e^{-itH_D}\psi\|^2 |e^{itz} - e^{itw}|^2 dt \right)^{1/2} \\ & \leq \|F_2^*\|_{H_D} \|\phi\| \left(\int_0^\infty \|F_1^* e^{-itH_D}\psi\|^2 |e^{itz} - e^{itw}|^2 dt \right)^{1/2}. \end{aligned}$$

Hence,

$$\begin{aligned} & \|F_1^*[(z - H_D)^{-1} - (w - H_D)^{-1}]F_2\psi\| \\ & \leq \|F_2^*\|_{H_D} \left(\int_0^\infty \|F_1^* e^{-itH_D}\psi\|^2 |e^{itz} - e^{itw}|^2 dt \right)^{1/2}. \end{aligned}$$

By the dominated convergence theorem, the integral tends to zero as z approaches w .

The proof for $\text{Im } z < 0$ is similar. \square

LEMMA II.4. Suppose F_1 and F_2 satisfy the hypotheses of Lemma II.3. Then

$$\begin{aligned} F_1^*[(\lambda + i0 - H_D)^{-1} - (\lambda - i0 - H_D)^{-1}]F_2 \\ = F_1^*[(\lambda + i0 - H_D)^{-1} - (\lambda - i0 - H_D)^{-1}]E^D(a, b)F_2 \end{aligned}$$

for any $\lambda \in (a, b)$. $E^D(a, b)$ denotes the spectral projection for H_D corresponding to the interval (a, b) .

PROOF. If F_1 and F_2 are bounded, the result is trivial because

$$[(\lambda + i0 - H_D)^{-1} - (\lambda - i0 - H_D)^{-1}](1 - E^D(a, b)) = 0.$$

More general functions F_1 and F_2 are limits of bounded functions. By taking such limits, the lemma follows from the bounds of Lemma II.3. \square

D. Fourier restriction theorems (Agmon [1], Kuroda [17]).

DEFINITION. For $k > 0$, define $\pi(k): \mathcal{S}(\mathbf{R}^m) \rightarrow L^2(S^{m-1}, d\omega)$ by $(\pi(k)f)(\omega) = k^{1/2(m-1)}\hat{f}(k\omega)$, where ω belongs to the unit sphere S^{m-1} , \hat{f} is the Fourier transform of f , and $d\omega$ is the invariant surface measure on $S^{m-1} \subseteq \mathbf{R}^m$.

LEMMA II.5. Suppose $1 < p \leq 2$, $m \geq 3$, and $1/2 < \delta < m/2$. If $\delta(p) = 2\delta(1 - p^{-1})$, then $\pi(k)$ extends to a bounded mapping of $L_{\delta(p)}^p(\mathbf{R}^m)$ into $L^2(S^{m-1}, d\omega)$. The norm of this mapping is bounded by

$$\text{Min}\{c_1 k^{(m-1)(p^{-1}-1/2)}, c_2 k^{[(m-1)(p^{-1}-1/2) + (\delta-1/2)(2-2p^{-1})]}\},$$

where c_1 and c_2 depend only on p . Moreover, if $p > 1$, then $k \mapsto \pi(k)$ is norm Hölder continuous.

PROOF. When $p = 2$, these results may be found in [7] or [8].

When $p = 1$, norm continuity in k does not hold, but the other results are trivial because \hat{f} is continuous and $\|\hat{f}\|_\infty \leq c\|f\|_1$.

The results now follow by interpolation (see Stein [35]) between L^1 and L_δ^2 . Note that Hölder continuity of order θ on L_δ^2 and boundedness on L^1 imply Hölder continuity of order $2\theta(1 - p^{-1})$ on $L_{\delta(p)}^p$ for $1 < p \leq 2$. \square

PROPOSITION II.6. Suppose $m_1 \geq 3$, $\delta > 1$, $1 < p \leq 2$, and $\delta(p) = 2\delta(1 - p^{-1})$. $\pi(k)$ extends to a bounded mapping of $L_{\delta(p)}^p(\mathbf{R}^{m_1}) \otimes L^2(\mathbf{R}^{m_2})$ into $L^2(S^{m_1+m_2-1}, d\omega)$. The mapping $k \mapsto \pi(k)$ is strongly continuous and uniformly bounded in norm when k is restricted to a bounded subset of $(0, \infty)$.

PROOF (GINIBRE AND MOULIN [7]). Choose $f \in \mathcal{S}(\mathbf{R}^{m_1+m_2})$, and let p_1 and p_2 denote the Fourier conjugate variables to $x_1 \in \mathbf{R}^{m_1}$ and $x_2 \in \mathbf{R}^{m_2}$. Decompose p_1 into radial and angular variables $|p_1|$ and ω_1 .

If $k_1 = (k^2 - |p_2|^2)^{1/2}$ and $m = m_1 + m_2$, then

$$\begin{aligned} \|\pi(k)f\|_{L^2(S^{m-1})}^2 &= k^{(m-1)} \int_{S^{m-1}} |\hat{f}(k\omega)|^2 d\omega \\ &= \int_{\mathbf{R}^m} \delta(k - |p|) |\hat{f}(p)|^2 dp. \end{aligned}$$

Change variables to $|p_1|$, ω_1 , and p_2 , and compute the $|p_1|$ integral:

$$\begin{aligned}\|\pi(k)f\|_{L^2(S^{m-1})}^2 &= k \int_{|p_2| < k} dp_2 k_1^{(m-2)} \int_{S^{m-1}} |\hat{f}(k_1 \omega_1, p_2)|^2 d\omega_1 \\ &= k \int_{|p_2| < k} dp_2 k_1^{-1} \|\pi_1(k_1) \tilde{f}(\cdot, p_2)\|_{L^2(S^{m-1})}^2,\end{aligned}$$

where the last norm is computed for fixed p_2 , \tilde{f} denotes partial Fourier transform in the second variable, and $\pi_1(k_1)$ denotes the Fourier restriction map in the first variable.

By Lemma II.5, $k_1^{-1} \|\pi_1(k_1)\|^2$ is uniformly bounded for $k_1 \in (0, k)$ from $L_{\delta(p)}^P(\mathbf{R}^{m_1})$ into $L^2(S^{m-1}, d\omega_1)$, so

$$\begin{aligned}\|\pi(k)f\|_{L^2(S^{m-1})}^2 &\leq k \int_{|p_2| < k} dp_2 k_1^{-1} \|\pi_1(k_1)\|^2 \|f(\cdot, p_2)\|_{L_{\delta(p)}^2(\mathbf{R}^{m_1})}^2 \\ &\leq kC(k) \int_{|p_2| < k} dp_2 \|\tilde{f}(\cdot, p_2)\|_{L_{\delta(p)}^2}^2 \\ &= kC(k) \int dx_2 \|f(\cdot, x_2)\|_{L_{\delta(p)}^2}^2 \\ &= kC(k) \|f\|_{L_{\delta(p)}^2 \otimes L^2}^2.\end{aligned}$$

This implies the uniform boundedness for k restricted to a bounded subset of $(0, \infty)$. The strong continuity of $\pi(k)$ follows from the norm continuity of $\pi_1(k_1)$.

□

E. *Dilation analyticity* (Balslev and Combes [3], see also van Winter [38], [39]).

The unitary group of dilations on $L^2(\mathbf{R}^n)$ is given by $(U(\theta)f)(x) = e^{n\theta/2} \psi(e^\theta x)$. A simple computation shows $H_0(\theta) = U(\theta)H_0U(-\theta) = e^{-2\theta}H_0$ for all $\theta \in \mathbf{R}$. $H_0(\theta)$ clearly extends to an analytic family $H_0(\theta)$ for all $\theta \in \mathbf{C}$.

Following Simon's treatment [30] of dilation analyticity for quadratic forms, we make the following definitions.

DEFINITION. Suppose $\alpha > 0$. A compact operator $v: \mathcal{H}_{+1} \rightarrow \mathcal{H}_{-1}$ belongs to the class \mathcal{F}_α , if and only if the family of operators $v(\theta) = U(\theta)vU(-\theta)$ has an analytic continuation to $\{\theta: |\operatorname{Im} \theta| < \alpha\}$ as operators from \mathcal{H}_{+1} into \mathcal{H}_{-1} . If $v(\theta): \mathcal{H}_{+1} \rightarrow \mathcal{H}_{-1}$ is norm continuous in $\{\theta: |\operatorname{Im} \theta| < \alpha\}$ and analytic in $\{\theta: |\operatorname{Im} \theta| < \alpha\}$, then v belongs to the class $\overline{\mathcal{F}}_\alpha$.

If $v \in \mathcal{F}_\alpha$ for some $\alpha > 0$, then v is called *dilation analytic*.

If $H = H_0 + \sum_{i < j} V_{ij}$, where $V_{ij} = v_{ij} \otimes 1$ and each $v_{ij} \in \mathcal{F}_\alpha$, then $H(\theta) = H_0(\theta) + \sum_{i < j} V_{ij}(\theta)$ may be defined by quadratic form methods. If D is a cluster decomposition, then $H_D(\theta)$ and the cluster Hamiltonians, $h_i(\theta)$ may be defined.

Let D be a cluster decomposition, with cluster Hamiltonians h_i ($1 \leq i \leq k$). If the potentials are dilation analytic, then we define $\Sigma_D(\theta) = \{E_1 + E_2 + \cdots + E_k: E_i \text{ is an eigenvalue of } h_i(\theta)\}$ and $\Sigma(\theta) = \{0\} \cup (\bigcup_{\#(D) \geq 2} \Sigma_D(\theta))$ is the set of *thresholds* of $H(\theta)$.

THEOREM II.7 (BALSLEV AND COMBES [3], SIMON [30]). *Suppose the two body potentials $V_{ij} = v_{ij} \otimes 1$ have $v_{ij} \in \mathfrak{F}_\alpha$ and are selfadjoint. The essential spectrum of $H(\theta)$ is $\sigma_{\text{ess}}(H(\theta)) = \{z + e^{-2\theta}r : z \in \Sigma(\theta), r \in [0, \infty)\}$. If $\mu = \inf \Sigma(0)$, then the discrete spectrum of $H(\theta)$ is contained in $\mathbf{R} \cup \{\mu + z : -2\text{Im } \theta < \arg z < 0\}$ if $\text{Im } \theta > 0$, and $\mathbf{R} \cup \{\mu + z : 0 < \arg z < -2\text{Im } \theta\}$ if $\text{Im } \theta < 0$. The discrete spectrum of $H(\theta)$ can accumulate only at points of $\Sigma(\theta)$. The nonthreshold eigenvalues of H are the real nonthreshold eigenvalues of $H(\theta)$, and $\Sigma(\theta) \cap \mathbf{R} = \Sigma(0)$ whenever $|\text{Im } \theta| < \pi/2$.*

THEOREM II.8 (BALSLEV AND COMBES [3]). *Suppose $H = H_0 + \sum_{i < j} V_{ij}$, where the two body potentials are dilation analytic. Then H has empty singular continuous spectrum.*

The following theorem proves exponential fall-off of bound states corresponding to nonthreshold eigenvalues. Without dilation analyticity, O'Connor [21] has proven this fact for bound states corresponding to discrete eigenvalues.

THEOREM II.9 (COMBES AND THOMAS [4]). *Suppose $H = H_0 + \sum_{i < j} V_{ij}$ is selfadjoint, with dilation analytic potentials. If $H\psi = E\psi$ and $E \notin \Sigma(0)$, then $\psi(x)$ lies in the domain of multiplication by $e^{a|x|}$ for some $a > 0$. Moreover, $e^{a|x|}\psi(x)$ belongs to the operator domain of $H_0^{1/2}$ for some $a > 0$.*

PROOF. Combes and Thomas [4] prove all but the last statement. Simon [34] proves $e^{a|x|}\psi(x) \in D(H_0^{1/2})$. \square

THEOREM II.10 (BALSLEV [2], SIMON [31], [32]). *Suppose $H = H_0 + \sum_{i < j} V_{ij}$ is selfadjoint. If $N = 2$ and $V \in \mathfrak{F}_\alpha$ for some $\alpha > 0$, or if $N \geq 3$ and each $V_{ij} = v_{ij} \otimes 1$, with $v_{ij} \in \mathfrak{F}_{\pi/2}$, then H has no positive eigenvalues.*

LEMMA II.11. *Suppose $M(z, \theta)$ is an analytic operator valued function in the region $\{(z, \theta) \in \mathbf{C}^2 : |\text{Im } \theta| < \alpha, z \in \mathbf{C} \setminus \sigma_{\text{ess}}(H(\theta))\}$, with norm continuous boundary values as z approaches $\sigma_{\text{ess}}(H(\theta))$. Suppose $U(\phi)M(z, \theta)U(-\phi) = M(z, \theta + \phi)$ for $\phi \in \mathbf{R}$, and assume $(M(z, \theta))^2$ is compact. If $E \in \mathbf{R}$, then $1 \in \sigma(M(E + i0, 0))$, if and only if $1 \in \sigma(M(E, \theta))$ for all θ with $0 < \text{Im } \theta < \text{Min}(\alpha, \pi/2)$.*

PROOF. Since $(M(z, \theta))^2$ is compact, $1 \in \sigma(M(z, \theta))$ if and only if $1 \in \sigma((M(z, \theta))^n)$ for all $n \geq 2$. Thus, it suffices to prove $1 \in \sigma((M(E + i0, 0))^n)$ if and only if $1 \in \sigma((M(E, \theta))^n)$ for all $n \geq 2$.

If $1 \in \sigma((M(E, \theta))^n)$ whenever $0 < \text{Im } \theta < \text{Min}(\alpha, \pi/2)$, then $1 \in \sigma((M(E + i0, 0))^n)$ by continuity.

Next, suppose $1 \in \sigma((M(E + i0, 0))^n)$ but $1 \notin \sigma((M(E, \phi))^n)$ for some $n \geq 2$ and ϕ with $\text{Im } \phi$ small and positive. Since $M(E + i0, \theta) = U(\theta)M(E + i0, 0)U(-\theta)$ for $\theta \in \mathbf{R}$, $1 \in \sigma((M(E + i0, \theta))^n)$ for all $\theta \in \mathbf{R}$.

Since $1 \notin \sigma((M(E, \phi))^n)$, the set of $\theta \in \mathbf{R}$ such that $1 \in \sigma((M(E + i0, \theta))^n)$ must be a closed set of measure zero by a theorem of Kuroda [16] (see also Simon [29, p. 127]).

This contradiction proves the lemma. \square

F. Existence of wave operators.

The existence of channel wave operators has been proved by Schechter [25] and Simon [33] for the types of potentials which concern us. Simon's proof is an extension of a simple method due to Cook [5].

THEOREM II.12 (SIMON [33]). *Suppose the two body potentials V_{ij} belong to $L^p_\delta(\mathbf{R}^m) + L^\infty_\delta(\mathbf{R}^m)$ for some $p > m/2$ and $\delta > 2$. Then the channel wave operators*

$$\Omega^\mp_\alpha = \text{strong-} \lim_{t \rightarrow \pm \infty} e^{itH} e^{-itT_\alpha} P_\alpha$$

exist.

III. Many-body Kuroda completeness as a stationary problem. This section is devoted to proving Theorem III.1, which reduces the many-body Kuroda completeness problem to the question of finding an appropriate formula for the resolvent $(z - H)^{-1}$. The idea of this theorem is not new. Implicitly, it appears in the works of Faddeev [6], Hepp [10], Ginibre and Moulin [7], Thomas [36], and Sigal [26]; explicitly, it may be found in Howland [11].

DEFINITION. Suppose H is a many-body Hamiltonian. The *multiparticle limiting absorption principle* holds for H , if

$$(z - H)^{-1} = \sum_D (z - H_D)^{-1} P_D \sum_{l=1}^{L(D)} F_{l,D} Z_{l,D}(z),$$

where

(a) there exists δ_0 such that $\phi \in L^2_{\delta_0}(\mathbf{R}^{(N-1)m})$ implies $Z_{l,D}(z)\phi$ is an $L^2(\mathbf{R}^{(N-1)m})$ -valued meromorphic function in $\mathbb{C} \setminus \sigma_{\text{ess}}(H)$, with continuous extensions to $\sigma_{\text{ess}}(H)$ from above and below, in the complement of a closed set \mathcal{E} of measure zero; and

(b) for each l and D , $F_{l,D}$ maps $L^2(\mathbf{R}^{(N-1)m})$ into $L^p_\mu(\mathbf{R}^m) \otimes L^2(\mathbf{R}^{(N-2)m})$, for some $\mu > 1$ and $p \in (1, 2]$, where the first factor denotes functions of a Jacobi coordinate for the motion of the centers of mass of the clusters of D .

THEOREM III.1. *Suppose the multiparticle limiting absorption principle holds for H on $L^2(\mathbf{R}^{(N-1)m})$, where $m \geq 3$. If the channel wave operators exist, then they are complete in the sense that $P_{\text{a.c.}} = \sum_\alpha \Omega^\pm_\alpha \Omega^\pm_\alpha^*$.*

As a first step toward the proof of this theorem, we prove several lemmas related to the orthogonality of channels. Next, using the limiting absorption principle, we obtain a formula for $(\Omega^\pm_\alpha \phi)^\wedge$, when $\phi \in L^2_{\delta_0}(\mathbf{R}^{(N-1)m})$. This and Stone's formula for spectral projections are then used to prove Theorem III.1.

DEFINITION. Let H_1 and H_2 be selfadjoint operators on \mathcal{H} . For each $\varepsilon > 0$ and each Borel set $A \subseteq \mathbf{R}$, the operators $\Omega^\pm(\varepsilon, H_2, H_1, A)$ are defined by

$$\langle \phi, \Omega^\mp(\varepsilon, H_2, H_1, A) \psi \rangle = \frac{\varepsilon}{\pi} \int_A \langle (\lambda \pm i\varepsilon - H_2)^{-1} \phi, (\lambda \pm i\varepsilon - H_1)^{-1} \psi \rangle d\lambda.$$

LEMMA III.2 (HOWLAND [11]). *Let H_1 and H_2 be selfadjoint on \mathcal{H} . $\Omega^\mp(\varepsilon, H_2, H_1, A)$ are well-defined contractions on \mathcal{H} for all Borel sets $A \subseteq \mathbf{R}$ and all $\varepsilon > 0$. If H_1 has purely absolutely continuous spectrum and spectral projections $E_1(\cdot)$, then $E_1(A)\psi = 0$ implies $\lim_{\varepsilon \downarrow 0} \Omega^\mp(\varepsilon, H_2, H_1, A)\psi = 0$.*

PROOF. Let $\delta_\varepsilon(t) = (\varepsilon/\pi)/(t^2 + \varepsilon^2)$.

The Schwarz inequality shows that the absolute value of the integrand appearing in the definition of $\Omega^\mp(\varepsilon, H_2, H_1, A)$ is bounded by

$$\begin{aligned} & (\varepsilon/\pi) \|(\lambda \pm i\varepsilon - H_2)^{-1}\phi\| \|(\lambda \pm i\varepsilon - H_1)^{-1}\psi\| \\ &= \left((\varepsilon/\pi) \|(\lambda \pm i\varepsilon - H_2)^{-1}\phi\|^2 \right)^{1/2} \left((\varepsilon/\pi) \|(\lambda \pm i\varepsilon - H_1)^{-1}\psi\|^2 \right)^{1/2} \\ &= (\langle \phi, \delta_\varepsilon(\lambda - H_2)\phi \rangle)^{1/2} (\langle \psi, \delta_\varepsilon(\lambda - H_1)\psi \rangle)^{1/2}. \end{aligned}$$

So, by the Schwarz inequality and Fubini's theorem,

$$\begin{aligned} & |\langle \phi, \Omega^\mp(\varepsilon, H_2, H_1, A)\psi \rangle|^2 \\ & \leq \left(\int_{\mathcal{A}} \langle \phi, \delta_\varepsilon(\lambda - H_2)\phi \rangle d\lambda \right) \left(\int_{\mathcal{A}} \langle \psi, \delta_\varepsilon(\lambda - H_1)\psi \rangle d\lambda \right) \\ &= \left(\int_{\mathbf{R}} (\delta_\varepsilon * \chi_{\mathcal{A}})(\lambda) d\langle \phi, E_2(\lambda)\phi \rangle \right) \left(\int_{\mathbf{R}} (\delta_\varepsilon * \chi_{\mathcal{A}})(\lambda) d\langle \psi, E_1(\lambda)\psi \rangle \right). \quad (\text{III.1}) \end{aligned}$$

Since $\int_{\mathbf{R}} \delta_\varepsilon(\lambda) dt = 1$ and $\|\chi_{\mathcal{A}}\|_\infty < 1$, we have $0 < \delta_\varepsilon * \chi_{\mathcal{A}} < 1$. So, the right-hand side of (III.1) is bounded by $\|\phi\|^2 \|\psi\|^2$. The operators $\Omega^\mp(\varepsilon, H_2, H_1, A)$ are, therefore, well-defined contractions.

The measure $d\langle \psi, E_1(\cdot)\psi \rangle$ is absolutely continuous with respect to Lebesgue measure. As a consequence ([37, pp. 28–31]), $\lim_{\varepsilon \downarrow 0} \delta_\varepsilon * \chi_{\mathcal{A}} = \chi_{\mathcal{A}}$ a.e. with respect to $d\langle \psi, E_1(\cdot)\psi \rangle$. Moreover, $0 < \delta_\varepsilon * \chi_{\mathcal{A}} < 1$, so the dominated convergence theorem shows:

$$\lim_{\varepsilon \downarrow 0} \int_{\mathbf{R}} (\delta_\varepsilon * \chi_{\mathcal{A}})(\lambda) d\langle \psi, E_1(\lambda)\psi \rangle = \int_{\mathbf{R}} \chi_{\mathcal{A}}(\lambda) d\langle \psi, E_1(\lambda)\psi \rangle = \|E_1(A)\psi\|^2.$$

Since

$$0 \leq \int_{\mathbf{R}} (\delta_\varepsilon * \chi_{\mathcal{A}})(\lambda) d\langle \phi, E_2(\lambda)\phi \rangle \leq \|\phi\|^2,$$

(III.1) shows

$$\limsup_{\varepsilon \downarrow 0} \|\Omega^\mp(\varepsilon, H_2, H_1, A)\psi\|^2 \leq \|E_1(A)\psi\|^2.$$

Therefore, $E_1(A)\psi = 0$ implies $\lim_{\varepsilon \downarrow 0} \Omega^\mp(\varepsilon, H_2, H_1, A)\psi = 0$. \square

LEMMA III.3. Let H_1 and H_2 be selfadjoint on \mathcal{H} . For all $\varepsilon > 0$,

$$\Omega^\mp(\varepsilon, H_2, H_1, \mathbf{R})\psi = 2\varepsilon \int_0^{\pm\infty} e^{\mp 2\varepsilon t} e^{itH_2} e^{-itH_1} \psi dt.$$

PROOF. Consider only Ω^- ; the proof for Ω^+ is similar. Fix $\phi, \psi \in \mathcal{H}$ and define $\phi(\lambda) = (\varepsilon/\pi)^{1/2}(\lambda + i\varepsilon - H_2)^{-1}\phi$ and $\psi(\lambda) = (\varepsilon/\pi)^{1/2}(\lambda + i\varepsilon - H_1)^{-1}\psi$. Then, by explicit computation,

$$\hat{\phi}(t) = (2\varepsilon)^{1/2} i\chi_{(0,\infty)}(t) e^{-\varepsilon t} e^{-itH_2}\phi$$

and

$$\hat{\psi}(t) = (2\varepsilon)^{1/2} i\chi_{(0,\infty)}(t) e^{-\varepsilon t} e^{itH_1}\psi.$$

These formulas and Lemma II.1 show

$$\begin{aligned}
 \langle \phi, \Omega^-(\varepsilon, H_2, H_1, \mathbf{R})\psi \rangle &= \int_{\mathbf{R}} \langle \phi(\lambda), \psi(\lambda) \rangle d\lambda \\
 &= \int_{\mathbf{R}} \langle \hat{\phi}(t), \hat{\psi}(t) \rangle dt \\
 &= 2\varepsilon \int_0^\infty \langle \phi, e^{-2\varepsilon t} e^{itH_2} e^{-itH_1} \psi \rangle dt \\
 &= \langle \phi, 2\varepsilon \int_0^\infty e^{-2\varepsilon t} e^{itH_2} e^{-itH_1} \psi dt \rangle. \quad \square
 \end{aligned}$$

LEMMA III.4 (HOWLAND [11]). *Let H_1 and H_2 be commuting selfadjoint operators on \mathcal{H} . Assume H_1 has purely absolutely continuous spectrum and that $H = H_1 - H_2$ has kernel $\{0\}$. Then, for Borel sets $A \subseteq \mathbf{R}$,*

$$\text{weak-lim}_{\varepsilon \downarrow 0} \Omega^\mp(\varepsilon, H_2, H_1, A) = 0.$$

PROOF.

$$\begin{aligned}
 \Omega^\mp(\varepsilon, H_2, H_1, A) &= \Omega^\mp(\varepsilon, H_2, H_1, \mathbf{R})E_1(A) - \Omega^\mp(\varepsilon, H_2, H_1, \mathbf{R} \setminus A)E_1(A) \\
 &\quad + \Omega^\mp(\varepsilon, H_2, H_1, A)E_1(\mathbf{R} \setminus A),
 \end{aligned}$$

where H_1 has spectral projections $E_1(\cdot)$.

The last two terms in this expression converge strongly to zero as $\varepsilon \rightarrow 0$ by Lemma III.2. Thus, it suffices to show $\text{weak-lim}_{\varepsilon \downarrow 0} \Omega^\mp(\varepsilon, H_2, H_1, \mathbf{R}) = 0$.

By Lemma III.3,

$$\begin{aligned}
 \langle \phi, \Omega^\mp(\varepsilon, H_2, H_1, \mathbf{R})\psi \rangle &= 2\varepsilon \int_0^{\pm\infty} e^{\mp 2\varepsilon t} \langle \phi, e^{itH_2} e^{-itH_1} \psi \rangle dt \\
 &= 2\varepsilon \int_0^{\pm\infty} e^{\mp 2\varepsilon t} \langle \phi, e^{-itH} \psi \rangle dt \\
 &= \pm 2i\varepsilon \langle \phi, (H \mp 2i\varepsilon)^{-1} \psi \rangle.
 \end{aligned}$$

As ε tends to zero, this quantity converges to $-\langle \phi, P_0 \psi \rangle$, where P_0 is the orthogonal projection onto the kernel of H . Hence, the hypotheses imply it converges to zero. \square

LEMMA III.5. *Let H_1 and H_2 be commuting selfadjoint operators on \mathcal{H} , such that $H_1 - H_2$ has kernel $\{0\}$ and H_1 has purely absolutely continuous spectrum. Let F_1 and F_2 be operators on \mathcal{H} such that F_1^* and F_2^* are Kato smooth with respect to H_1 and H_2 , respectively. Suppose $\phi_\pm: D_\pm \rightarrow \mathcal{H}$ and $\psi_\pm: D_\pm \rightarrow \mathcal{H}$ are continuous functions, where $D_\pm = \{z = x \pm iy \in \mathbf{C}: a < x < b, 0 < y < c\}$. Then*

$$\begin{aligned}
 \lim_{\varepsilon \downarrow 0} \varepsilon \int_a^b \langle (\lambda \pm i\varepsilon - H_2)^{-1} F_2 \phi_\pm(\lambda \pm i\varepsilon), \\
 (\lambda \pm i\varepsilon - H_1)^{-1} F_1 \psi_\pm(\lambda \pm i\varepsilon) \rangle d\lambda = 0.
 \end{aligned}$$

PROOF. Assume, without loss of generality, that $F_1 \neq 0$, $F_2 \neq 0$, $\phi_{\pm} \neq 0$, $\psi_{\pm} \neq 0$, $a < b$, $c > 0$. For convenience, consider only the $+$ signs, with $\phi = \phi_+$, $\psi = \psi_+$, and $D = D_+$.

Let $\delta > 0$ be given, and choose M such that $\|\phi(z)\| < M$ and $\|\psi(z)\| < M$ for all $z \in D$. Since F_1 and F_2 have dense domains, there exist simple functions $f: [a, b] \rightarrow \mathcal{D}(F_1)$ and $g: [a, b] \rightarrow \mathcal{D}(F_2)$, such that $\|f(x)\| < M$, $\|g(x)\| < M$, $\|\phi(x) - g(x)\| < \delta$, and $\|\psi(x) - f(x)\| < \delta$, for all $x \in [a, b]$.

By the continuity of ϕ and ψ , and the compactness of D , there exists $\gamma > 0$, such that $0 < \varepsilon < \gamma$ implies $\|\phi(\lambda + i\varepsilon) - \phi(\lambda)\| < \delta$ and $\|\psi(\lambda + i\varepsilon) - \psi(\lambda)\| < \delta$ for all $\lambda \in [a, b]$.

Thus, $\|\phi(\lambda + i\varepsilon) - g(\lambda)\| < 2\delta$ and $\|\psi(\lambda + i\varepsilon) - f(\lambda)\| < 2\delta$ for all $\lambda \in [a, b]$, whenever $0 < \varepsilon < \gamma$.

Define:

$$\begin{aligned} A(\varepsilon) &= \varepsilon \int_a^b \langle (\lambda + i\varepsilon - H_2)^{-1} F_2 g(\lambda), (\lambda + i\varepsilon - H_1)^{-1} F_1 f(\lambda) \rangle d\lambda \\ B(\varepsilon) &= \varepsilon \int_a^b \langle (\lambda + i\varepsilon - H_2)^{-1} F_2 g(\lambda), \\ &\quad (\lambda + i\varepsilon - H_1)^{-1} F_1 [\psi(\lambda + i\varepsilon) - f(\lambda)] \rangle d\lambda \\ C(\varepsilon) &= \varepsilon \int_a^b \langle (\lambda + i\varepsilon - H_2)^{-1} F_2 [\phi(\lambda + i\varepsilon) - g(\lambda)], \\ &\quad (\lambda + i\varepsilon - H_1)^{-1} F_1 \psi(\lambda + i\varepsilon) \rangle d\lambda. \end{aligned}$$

By Lemma III.4, there exists $\gamma_1 < \gamma$, so that $\varepsilon < \gamma_1$ implies $|A(\varepsilon)| < \delta$.

By Hölder's inequality and Lemma II.2, $\varepsilon < \gamma$ implies

$$\begin{aligned} |B(\varepsilon)| &\leq \varepsilon \int_a^b \|(\lambda + i\varepsilon - H_2)^{-1} F_2 g(\lambda)\| \\ &\quad \times \|(\lambda + i\varepsilon - H_1)^{-1} F_1 [\psi(\lambda + i\varepsilon) - f(\lambda)]\| d\lambda \\ &\leq \left(\varepsilon \int_a^b \|(\lambda + i\varepsilon - H_2)^{-1} F_2 g(\lambda)\|^2 d\lambda \right)^{1/2} \\ &\quad \times \left(\varepsilon \int_a^b \|(\lambda + i\varepsilon - H_1)^{-1} F_1 [\psi(\lambda + i\varepsilon) - f(\lambda)]\|^2 d\lambda \right)^{1/2} \\ &\leq \left(\frac{b-a}{\pi} \|F_2\|_{H_2}^2 \left(\sup_{\lambda} \|g(\lambda)\| \right)^2 \right)^{1/2} \\ &\quad \times \left(\frac{b-a}{\pi} \|F_1\|_{H_1}^2 \left(\sup_{\lambda} \|\psi(\lambda + i\varepsilon) - f(\lambda)\| \right)^2 \right)^{1/2} \\ &\leq 2 \frac{b-a}{\pi} M \|F_1\|_{H_1} \|F_2\|_{H_2} \delta. \end{aligned}$$

Similarly, when $\varepsilon < \gamma$,

$$\begin{aligned}
 |C(\varepsilon)| &< \left(\varepsilon \int_a^b \left\| (\lambda + i\varepsilon - H_2)^{-1} F_2 [\phi(\lambda + i\varepsilon) - g(\lambda)] \right\|^2 d\lambda \right)^{1/2} \\
 &\quad \times \left(\varepsilon \int_a^b \left\| (\lambda + i\varepsilon - H_1)^{-1} F_1 \psi(\lambda + i\varepsilon) \right\|^2 d\lambda \right)^{1/2} \\
 &< \left(\frac{b-a}{\pi} \|F_2\|_{H_2} \left(\sup_{\lambda} \|\phi(\lambda + i\varepsilon) - g(\lambda)\| \right)^2 \right)^{1/2} \\
 &\quad \times \left(\frac{b-a}{\pi} \|F_1\|_{H_1} \left(\sup_{\lambda} \|\psi(\lambda + i\varepsilon)\| \right)^2 \right)^{1/2} \\
 &< 2 \frac{b-a}{\pi} M \|F_1\|_{H_1} \|F_2\|_{H_2} \delta.
 \end{aligned}$$

Therefore, $\varepsilon < \gamma_1$ implies

$$\begin{aligned}
 |A(\varepsilon) + B(\varepsilon) + C(\varepsilon)| &< |A(\varepsilon)| + |B(\varepsilon)| + |C(\varepsilon)| \\
 &< \left(1 + 4 \frac{b-a}{\pi} M \|F_1\|_{H_1} \|F_2\|_{H_2} \right) \delta.
 \end{aligned}$$

Thus, $\lim_{\varepsilon \downarrow 0} A(\varepsilon) + B(\varepsilon) + C(\varepsilon) = 0$, which is the desired result. \square

REMARKS. (1) Let α and β be channels. T_α and T_β are multiplication operators in momentum representation, so they commute. Moreover, if $D(\alpha) \neq D(\beta)$ or $E_\alpha \neq E_\beta$, then $T_\alpha - T_\beta$ has kernel $\{0\}$. To see this, notice that T_α and T_β are multiplication by different polynomials in momentum representation (because $D(\alpha) \neq D(\beta)$ or $E_\alpha \neq E_\beta$). Therefore, $T_\alpha - T_\beta$ multiplies by a nonzero polynomial $Q(p_1, \dots, p_{(N-1)m})$. Functions in the kernel of $T_\alpha - T_\beta$ must have support in $\{p: Q(p) = 0\}$. This set is a nontrivial algebraic variety, and consequently has measure zero. Thus, $T_\alpha - T_\beta$ has kernel $\{0\}$.

(2) Suppose α is a channel with cluster decomposition D . Then $H_D P_\alpha = T_\alpha P_\alpha = P_\alpha T_\alpha$ and $P_\alpha (z - H_D)^{-1} = P_\alpha (z - T_\alpha)^{-1}$. Also, if $P_\alpha F^*$ is Kato smooth with respect to H_D , then $P_\alpha F^*$ is Kato smooth with respect to T_α .

(3) If D is a cluster decomposition, then $P_D = \sum_{D(\alpha)=D} P_\alpha$. Consequently, $(z - H_D)^{-1} P_D = \sum_{D(\alpha)=D} (z - T_\alpha)^{-1} P_\alpha$. So, when the multiparticle limiting absorption principle holds for H ,

$$\begin{aligned}
 (z - H)^{-1} &= \sum_D (z - H_D)^{-1} P_D \sum_{l=1}^{L(D)} F_{l,D} Z_{l,D}(z) \\
 &= \sum_\alpha (z - T_\alpha)^{-1} P_\alpha \sum_{l=1}^{L(D(\alpha))} F_{l,D(\alpha)} Z_{l,D(\alpha)}(z).
 \end{aligned}$$

(4) In Propositions III.6 and III.7, we restrict some Fourier transforms to ellipsoids rather than spheres. This is because certain reduced masses are not equal. However, a linear change of coordinates shows that the results of §II.D apply to ellipsoids as well as spheres.

The ellipsoids which we use are the surfaces in momentum space corresponding to a given energy for a channel Hamiltonian T_α . In some clustered Jacobi coordinates, $T_\alpha = \sum_{l=1}^{k-1} (-2M_l)^{-1} \Delta_{\xi_l} + E_\alpha$, where M_l is a reduced mass. We choose the “unit ellipsoid”

$$\Gamma = \left\{ q \in \mathbf{R}^{(k-1)m}: \sum_{l=1}^{k-1} (2M_l)^{-1} |q_l|^2 = 1 \right\}$$

as the analogue of the unit sphere. The mapping $\pi_\alpha(k): L_\delta^P \otimes L^2 \rightarrow L^2(\Gamma)$ is defined as the analogue of $\pi(k): L_\delta^P \otimes L^2 \rightarrow L^2(S^{m-1})$ from Proposition II.6.

(5) For convenience we identify the range of P_α with $\mathcal{H}_\alpha = \mathcal{H}(D(\alpha))$, defined in the introduction. The identification is given by the mapping $\phi \otimes \psi_\alpha \rightarrow \phi$, where $\phi \otimes \psi_\alpha$ denotes an arbitrary element of $P_\alpha \mathcal{H}$.

PROPOSITION III.6. *Suppose the multiparticle limiting absorption principle holds for H , and let α be a channel. Let $(\xi_1, \dots, \xi_{k-1}, \xi_1^1, \dots, \xi_{n(k)-1}^k)$ be clustered Jacobi coordinates corresponding to the decomposition $D(\alpha)$, and let (q_1, \dots, q_{k-1}) be the Fourier conjugate variables to $(\xi_1, \dots, \xi_{k-1})$. Let $k_\alpha(q) = (\sum_{j=1}^{k-1} (2M_j)^{-1} |q_j|^2)^{1/2}$, where M_j is the reduced mass corresponding to the coordinate ξ_j . Suppose the wave operators Ω_α^\pm exist, and that $\phi \in L_{\delta_0}^2(\mathbf{R}^{(N-1)m})$.*

If $k^2 + E_\alpha$ does not belong to the exceptional set \mathcal{E} for H , then $((\Omega_\alpha^\pm)^ \phi)^\wedge$ may be restricted to the ellipsoid*

$$\Gamma(k) = \{ q \in \mathbf{R}^{(k-1)m}: k_\alpha(q) = k \},$$

and

$$\pi_\alpha(k)(\Omega_\alpha^\pm)^* \phi = \pi_\alpha(k) P_\alpha \sum_{l=1}^{L(D(\alpha))} F_{l,D(\alpha)} Z_{l,D(\alpha)} (k^2 + E_\alpha \mp i0) \phi.$$

PROOF. Choose an interval $[a, b]$ containing $k^2 + E_\alpha$, but which does not intersect the exceptional set \mathcal{E} . Choose $\psi_1 \in \mathcal{H}_\alpha = \mathcal{H}(D(\alpha))$ so that ψ_1 is C^∞ with support in $\bigcup_{\lambda \in [a, b]} \Gamma(\lambda)$.

Consider only Ω_α^- ; the proof for Ω_α^+ is similar.

The existence of Ω_α^- implies that Ω_α^- equals its Abelian limit (see [29]). So, if $\psi = \psi_1 \otimes \psi_\alpha$, then

$$\Omega_\alpha^- \psi = \lim_{\varepsilon \downarrow 0} 2\varepsilon \int_0^\infty e^{-2\varepsilon t} e^{itH} e^{-itT_\alpha} \psi \, dt.$$

Thus, by Lemma II.1,

$$\begin{aligned} \langle \phi, \Omega_\alpha^- \psi \rangle &= \lim_{\varepsilon \downarrow 0} 2\varepsilon \int_0^\infty \langle e^{-\varepsilon t} e^{-itH} \phi, e^{-\varepsilon t} e^{-itT_\alpha} \psi \rangle \, dt \\ &= \lim_{\varepsilon \downarrow 0} \frac{\varepsilon}{\pi} \int_{-\infty}^\infty \langle (\lambda + i\varepsilon - H)^{-1} \phi, (\lambda + i\varepsilon - T_\alpha)^{-1} \psi \rangle \, d\lambda. \end{aligned}$$

Lemma III.2 and the choice of ψ_1 now show

$$\langle \phi, \Omega_\alpha^- \psi \rangle = \lim_{\varepsilon \downarrow 0} \frac{\varepsilon}{\pi} \int_a^b \langle (\lambda + i\varepsilon - H)^{-1} \phi, (\lambda + i\varepsilon - T_\alpha)^{-1} \psi \rangle \, d\lambda.$$

By Remark (3) after Lemma III.5,

$$(z - H)^{-1} = \sum_{\beta} (z - T_{\beta})^{-1} P_{\beta} \sum_{l=1}^{L(D(\beta))} F_{l,D(\beta)} Z_{l,D(\beta)}(z).$$

Hence,

$$\langle \phi, \Omega_{\alpha}^{-} \psi \rangle = \lim_{\varepsilon \downarrow 0} \frac{\varepsilon}{\pi} \int_a^b \left\langle \sum_{\beta} \sum_l (\lambda + i\varepsilon - T_{\beta})^{-1} P_{\beta} F_{l,D(\beta)} Z_{l,D(\beta)}(z) \phi, (\lambda + i\varepsilon - T_{\alpha})^{-1} \psi \right\rangle d\lambda.$$

By Lemma III.5 and Remark (1) after it, all terms with $D(\alpha) \neq D(\beta)$ or $E_{\alpha} \neq E_{\beta}$ vanish as $\varepsilon \rightarrow 0$. All other terms with $\alpha \neq \beta$ vanish because $P_{\beta} P_{\alpha} = 0$ and $P_{\alpha} \psi = \psi$. Therefore,

$$\langle \phi, \Omega_{\alpha}^{-} \psi \rangle = \lim_{\varepsilon \downarrow 0} \frac{\varepsilon}{\pi} \int_a^b \left\langle (\lambda + i\varepsilon - T_{\alpha})^{-1} P_{\alpha} \sum_{l=1}^{L(D(\alpha))} F_{l,D(\alpha)} Z_{l,D(\alpha)}(\lambda + i\varepsilon) \phi, (\lambda + i\varepsilon - T_{\alpha})^{-1} \psi \right\rangle d\lambda.$$

For convenience put $F_l = F_{l,D(\alpha)}$, $L = L(D(\alpha))$, and $\phi_l(z) = Z_{l,D(\alpha)}(z)\phi$, so

$$\langle \phi, \Omega_{\alpha}^{-} \psi \rangle = \sum_{l=1}^L \lim_{\varepsilon \downarrow 0} \frac{\varepsilon}{\pi} \int_a^b \langle (\lambda + i\varepsilon - T_{\alpha})^{-1} P_{\alpha} F_l \phi_l(\lambda + i\varepsilon), (\lambda + i\varepsilon - T_{\alpha})^{-1} \psi \rangle d\lambda.$$

The identification of $P_{\alpha} \mathcal{H}$ with \mathcal{H}_{α} allows the inner product in this integral to be computed as an inner product on \mathcal{H}_{α} . Moreover, as elements of \mathcal{H}_{α} , ψ_1 and $P_{\alpha} F_l \phi_l(\lambda + i\varepsilon)$ have Fourier transforms which may be restricted to ellipsoids by Proposition II.6. So, using ellipsoidal coordinates on $\mathbb{R}^{(k-1)m}$ to compute the \mathcal{H}_{α} inner product, we have

$$\langle \phi, \Omega_{\alpha}^{-} \psi \rangle = \sum_{l=1}^L \lim_{\varepsilon \downarrow 0} \int_a^b \int_0^{\infty} \frac{\varepsilon/\pi}{(k^2 + E_{\alpha} - \lambda)^2 + \varepsilon^2} \times \langle \pi_{\alpha}(k) P_{\alpha} F_l \phi_l(\lambda + i\varepsilon), \pi_{\alpha}(k) \psi_1 \rangle_{L^2(\Gamma)} dk d\lambda,$$

where $\pi_{\alpha}(k): \mathcal{H}_{\alpha} \rightarrow L^2(\Gamma)$ restricts Fourier transforms to $\Gamma(k)$, and then does the scaling to yield a function on $\Gamma = \Gamma(1)$.

The inner product in the last integrand depends continuously on k and has support in $[a, b]$ in the k variable. Thus, Fubini's theorem and the dominated convergence theorem may be applied to yield:

$$\langle \phi, \Omega_{\alpha}^{-} \psi \rangle = \sum_{l=1}^L \int_a^b \lim_{\varepsilon \downarrow 0} \int_a^b \frac{\varepsilon/\pi}{(k^2 + E_{\alpha} - \lambda)^2 + \varepsilon^2} \times \langle \pi_{\alpha}(k) P_{\alpha} F_l \phi_l(\lambda + i\varepsilon), \pi_{\alpha}(k) \psi_1 \rangle_{L^2(\Gamma)} d\lambda dk.$$

The limit may now be computed explicitly (Ikebe [12], Simon [29]):

$$\langle \phi, \Omega_{\alpha}^{-} \psi \rangle = \int_a^b \left\langle \pi_{\alpha}(k) P_{\alpha} \sum_l F_l \phi_l(k^2 + E_{\alpha} + i0), \pi_{\alpha}(k) \psi_1 \right\rangle_{L^2(\Gamma)} dk.$$

Since ψ_1 is an arbitrary element of a dense subspace of $E^{\alpha}([a, b])\mathcal{H}_{\alpha}$, the proposition follows. \square

PROPOSITION III.7. *Suppose the multiparticle limiting absorption principle holds for H . Suppose Ω_α^\pm exist. Let $[a, b]$ be any compact interval disjoint from the exceptional set \mathcal{E} . Let $E[a, b]$ be the spectral projection for H on the interval $[a, b]$. Then, $\phi \in L_{\delta_0}^2(\mathbf{R}^{(N-1)m})$ implies*

$$\|E[a, b]\phi\|^2 = \sum_\alpha \int_{a < k_\alpha(q) + E_\alpha < b} |(\Omega_\alpha^\pm * \phi)^\sim(q)|^2 dq.$$

PROOF. Apply Stone's formula and the first resolvent formula:

$$\|E[a, b]\phi\|^2 = \lim_{\varepsilon \downarrow 0} \frac{\varepsilon}{\pi} \int_a^b \langle (\lambda \pm i\varepsilon - H)^{-1} \phi, (\lambda \pm i\varepsilon - H)^{-1} \phi \rangle d\lambda.$$

For simplicity take only the $+$ sign (this yields the result for Ω^-). Then use the multiparticle limiting absorption principle formula:

$$\begin{aligned} \|E[a, b]\phi\|^2 &= \sum_{\alpha, \beta} \sum_{l=1}^{L(D(\alpha))} \sum_{j=1}^{L(D(\beta))} \lim_{\varepsilon \downarrow 0} \frac{\varepsilon}{\pi} \\ &\quad \times \int_a^b \langle (\lambda + i\varepsilon - T_\alpha)^{-1} P_\alpha F_{l, D(\alpha)} \phi_{l, D(\alpha)}(\lambda + i\varepsilon), \\ &\quad (\lambda + i\varepsilon - T_\beta)^{-1} P_\beta F_{j, D(\beta)} \phi_{j, D(\beta)}(\lambda + i\varepsilon) \rangle d\lambda, \end{aligned}$$

where $\phi_{l, D(\alpha)}(z) = Z_{l, D(\alpha)}(z)\phi$.

By Lemma III.5 and Remark (1) after it, all terms with $D(\alpha) \neq D(\beta)$ or $E_\alpha \neq E_\beta$ vanish as $\varepsilon \rightarrow 0$. All other terms with $\alpha \neq \beta$ vanish because $P_\alpha P_\beta = 0$. Therefore,

$$\|E[a, b]\phi\|^2 = \sum_\alpha \lim_{\varepsilon \downarrow 0} \int_a^b \frac{\varepsilon}{\pi} \left\| \sum_{l=1}^L (\lambda + i\varepsilon - T_\alpha)^{-1} P_\alpha F_{l, D(\alpha)} \phi_{l, D(\alpha)}(\lambda + i\varepsilon) \right\|^2 d\lambda.$$

Proposition II.2 and the dominated convergence theorem show that the limit may be taken through the integral:

$$\begin{aligned} \|E[a, b]\phi\|^2 &= \sum_\alpha \int_a^b (2\pi i)^{-1} \sum_l \sum_j \\ &\quad \left[\langle \phi_{l, D(\alpha)}(\lambda + i0), (F_{l, D(\alpha)}^*(\lambda - i0 - T_\alpha)^{-1} P_\alpha F_{j, D(\alpha)}) \phi_{j, D(\alpha)}(\lambda + i0) \rangle \right. \\ &\quad \left. - \langle \phi_{l, D(\alpha)}(\lambda + i0), (F_{l, D(\alpha)}^*(\lambda + i0 - T_\alpha)^{-1} P_\alpha F_{j, D(\alpha)}) \phi_{j, D(\alpha)}(\lambda + i0) \rangle \right] d\lambda. \end{aligned}$$

Lemma II.4 shows that this may be rewritten as

$$\begin{aligned} \|E[a, b]\phi\|^2 &= \sum_\alpha \int_a^b \lim_{\varepsilon \downarrow 0} \frac{\varepsilon}{\pi} \left\| (\lambda + i\varepsilon - T_\alpha)^{-1} E^\alpha[a, b] P_\alpha \sum_l F_{l, D(\alpha)} \phi_{l, D(\alpha)}(\lambda + i\varepsilon) \right\|^2 d\lambda. \end{aligned}$$

The quantity inside the norm in the integrand belongs to $\text{Ran } P_\alpha$, so it may be lifted to \mathcal{H}_α by the identification of \mathcal{H}_α with $P_\alpha \mathcal{H}$. Ellipsoidal coordinates may then be used to evaluate the norm on \mathcal{H}_α :

$$\begin{aligned} \|E[a, b]\phi\|^2 &= \sum_{\alpha} \int_a^b \lim_{\varepsilon \downarrow 0} \int_{(a-E_{\alpha})^{1/2}}^{(b-E_{\alpha})^{1/2}} \frac{\varepsilon/\pi}{(k^2 + E_{\alpha} - \lambda)^2 + \varepsilon^2} \\ &\quad \times \left\| \pi_{\alpha}(k) P_{\alpha} \sum_l F_{l,D(\alpha)} \phi_{l,D(\alpha)}(\lambda + i\varepsilon) \right\|_{L^2(\Gamma)}^2 dk d\lambda. \end{aligned}$$

The limit may be evaluated explicitly (Ikebe [12], Simon [29]). Proposition III.6 and a change of variables yield the desired result:

$$\begin{aligned} \|E[a, b]\phi\|^2 &= \sum_{\alpha} \int_a^b \left\| \pi_{\alpha}(\nu) P_{\alpha} \sum_l F_{l,D(\alpha)} \phi_{l,D(\alpha)}(\nu^2 + E_{\alpha} + i0) \right\|_{L^2(\Gamma)}^2 d\nu \\ &= \sum_{\alpha} \int_{a < k_{\alpha}(q)^2 + E_{\alpha} < b} |(\Omega_{\alpha}^{\pm} * \phi)^{\wedge}(q)|^2 dq. \quad \square \end{aligned}$$

PROOF OF THEOREM III.1. The wave operators are partial isometries, so it suffices to prove

$$\|P_{\text{a.c.}}\phi\|^2 = \sum_{\alpha} \|\Omega_{\alpha}^{\pm} * \phi\|^2$$

for $\phi \in L_{\delta_0}^2(\mathbf{R}^{(N-1)m})$.

The complement of the exceptional set \mathcal{E} is open, so it is the countable union of compact intervals A_n , with rational end points, such that any two distinct intervals intersect in at most one point. Since \mathcal{E} has measure zero,

$$\|P_{\text{ac}}\phi\|^2 = \sum_n \|E(A_n)\phi\|^2,$$

where $E(A_n)$ is the spectral projection for H corresponding to the interval A_n .

Therefore, by Proposition III.7 and Fubini's theorem,

$$\begin{aligned} \|P_{\text{a.c.}}\phi\|^2 &= \sum_{n=1}^{\infty} \sum_{\alpha} \int_{k_{\alpha}(q)^2 + E_{\alpha} \in A_n} |(\Omega_{\alpha}^{\pm} * \phi)^{\wedge}(q)|^2 dq \\ &= \sum_{\alpha} \|(\Omega_{\alpha}^{\pm} * \phi)^{\wedge}\|^2 = \sum_{\alpha} \|\Omega_{\alpha}^{\pm} * \phi\|^2. \quad \square \end{aligned}$$

IV. Resolvent formulas. Having proved Theorem III.1, the problem of asymptotic completeness is reduced to the study of resolvent formulas which satisfy certain conditions. Derivation of such formulas for the two, three, and four particle cases is the goal of this section.

To begin with, the desired two-body formula is given in equation (IV.1). Next, a generalization of equation (IV.1) is obtained for the N -particle case. This is equation (IV.4), which involves $\frac{1}{2}N(N-1) \times \frac{1}{2}N(N-1)$ matrices. It coincides with the Faddeev equation [6] when $N = 3$, but does not have the form required by the multiparticle limiting absorption principle. Thus, various modifications are necessary.

A second N -body formula, equation (IV.5), is obtained from equation (IV.4), by increasing the size of the matrices so that the individual matrix entries become simpler. This formula is closely related, but not identical to a formula of Nardetskii and Yakubovskii [20]. However, as was the case with the first N -body formula, equation (IV.5) is not in a suitable form for scattering.

Consequently, for $N = 3$ and $N = 4$, a third formula, which does have the correct form, is derived from equation (IV.5). Ginibre and Moulin [7] have used this three-body equation, and their ideas have been generalized to give the four-body formula.

For $N > 5$ analogous equations, having the proper form, may be written down. However, it is not at all clear that the bounds required by the multiparticle limiting absorption principle can be proved. This is due to the fact that we have not been able to prove the absence of spurious zeros in the four-body case.

Throughout this section we assume that $V_{ij} = U_{ij}W_{ij}$, such that $U_{ij}(-\Delta_{ij} + 1)^{-1/2}$ and $W_{ij}(-\Delta_{ij} + 1)^{-1/2}$ are compact as operators on $L^2(\mathbb{R}^m)$, where Δ_{ij} is the Laplacian in the x_{ij} variable. These assumptions imply the relative form boundedness of the potentials with respect to H_0 , with relative bound zero. So, when $\text{Re } z$ is sufficiently large negative, the perturbation series

$$\begin{aligned} (z - H)^{-1} &= (z - H_0)^{-1} + (z - H_0)^{-1} \sum_{\alpha} V_{\alpha} (z - H_0)^{-1} \\ &\quad + (z - H_0)^{-1} \sum_{\alpha} V_{\alpha} (z - H_0)^{-1} \sum_{\beta} V_{\beta} (z - H_0)^{-1} + \dots \end{aligned}$$

is norm convergent.

When $N = 2$, this is particularly simple. Using geometric series, the perturbation expansion may be resummed in the form:


$$\begin{aligned} (z - H)^{-1} &= (z - H_0)^{-1} \\ &\quad + (z - H_0)^{-1} U [1 - W(z - H_0)^{-1} U]^{-1} W (z - H_0)^{-1}. \end{aligned} \quad (\text{IV.1})$$

This is the desired equation when $N = 2$.

For $N > 3$, the graphical symbolism of Weinberg [40] (see also Simon [29]) is used to identify each term of the perturbation series with a graph. For example, when $N = 4$, the term

$$(z - H_0)^{-1} V_{12} (z - H_0)^{-1} V_{23} (z - H_0)^{-1} V_{12} (z - H_0)^{-1} V_{24} (z - H_0)^{-1}$$

is assigned the graph



Using this symbolism, the sum of all nontrivial graphs is

$$(z - H)^{-1} - (z - H_0)^{-1}$$

Several definitions are now required:

DEFINITION. To each nontrivial graph

$$(z - H_0)^{-1} V_{\alpha_1} (z - H_0)^{-1} V_{\alpha_2} \cdots (z - H_0)^{-1} V_{\alpha_n} (z - H_0)^{-1},$$

there is associated a *truncated graph*

$$W_{\alpha_1} (z - H_0)^{-1} V_{\alpha_2} \cdots (z - H_0)^{-1} V_{\alpha_n} (z - H_0)^{-1}.$$

A truncated graph is *connected* (respectively *disconnected*) if its associated graph is connected (disconnected).

DEFINITION. An expression of the form

$$W_{\alpha_1} (z - H_0)^{-1} V_{\alpha_2} (z - H_0)^{-1} \cdots (z - H_0)^{-1} V_{\alpha_{n-1}} (z - H_0)^{-1} U_{\alpha_n}$$

is a *barely connected part* if

$$W_{\alpha_1} (z - H_0)^{-1} V_{\alpha_2} (z - H_0)^{-1} \cdots (z - H_0)^{-1} V_{\alpha_{n-1}} (z - H_0)^{-1} V_{\alpha_n} (z - H_0)^{-1}$$

is connected, but

$$W_{\alpha_1} (z - H_0)^{-1} V_{\alpha_2} (z - H_0)^{-1} \cdots (z - H_0)^{-1} V_{\alpha_{n-1}} (z - H_0)^{-1}$$

is disconnected.

DEFINITION. $\mathcal{Q}_\alpha(z) = (z - H_0)^{-1} U_\alpha$.

$\mathcal{M}_{\alpha,\beta}(z)$ is the sum of all barely connected parts whose initial factor is W_α and whose final factor is U_β .

$\mathcal{C}_\alpha(z)$ is the sum of all disconnected truncated graphs whose initial factor is W_α .

Let $G = W_{\alpha_1} (z - H_0)^{-1} V_{\alpha_2} (z - H_0)^{-1} \cdots V_{\alpha_n} (z - H_0)^{-1}$ be a truncated graph. Either G is disconnected or G is the product of a barely connected part $W_{\alpha_1} (z - H_0)^{-1} V_{\alpha_2} (z - H_0)^{-1} \cdots (z - H_0)^{-1} U_{\alpha_j}$ and a truncated graph $G' = W_{\alpha_j} (z - H_0)^{-1} V_{\alpha_{j+1}} (z - H_0)^{-1} \cdots V_{\alpha_n} (z - H_0)^{-1}$. By iterating this procedure G may be uniquely decomposed as a product of $k \geq 0$ barely connected parts times a disconnected truncated graph.

In order to compute the sum of all graphs, first sum all graphs containing precisely k barely connected parts, and then sum over k . The sum of all nontrivial graphs with no barely connected parts is the sum of all nontrivial disconnected graphs: $\sum_\alpha \mathcal{Q}_\alpha \mathcal{C}_\alpha$. The sum of all graphs containing one barely connected part is $\sum_{\alpha,\beta} \mathcal{Q}_\alpha \mathcal{M}_{\alpha,\beta} \mathcal{C}_\beta$. Similarly, the sum of all graphs containing k barely connected parts is

$$\sum_{\alpha, \beta_1, \beta_2, \dots, \beta_k} \mathcal{Q}_\alpha \mathcal{M}_{\alpha, \beta_1} \mathcal{M}_{\beta_1, \beta_2} \mathcal{M}_{\beta_2, \beta_3} \cdots \mathcal{M}_{\beta_{k-1}, \beta_k} \mathcal{C}_{\beta_k}.$$

Therefore, using matrix multiplication, $(z - H)^{-1} - (z - H_0)^{-1} = \mathcal{Q}(1 + \mathfrak{N} + \mathfrak{N}^2 + \dots)\mathcal{C}$.

When $\operatorname{Re} z$ is sufficiently large negative, the geometric series $1 + \mathfrak{N}(z) + \mathfrak{N}(z)^2 + \dots$ converges in norm to $(1 - \mathfrak{N}(z))^{-1}$ and

$$(z - H)^{-1} = (z - H_0)^{-1} + \mathcal{Q}(z)(1 - \mathfrak{N}(z))^{-1}\mathcal{C}(z).$$

To obtain explicit formulas for $\mathfrak{N}(z)$ and $\mathcal{C}(z)$ in terms of the resolvents $(z - H_D)^{-1}$, it is useful to introduce strings and make several definitions.

DEFINITION. Suppose $2 \leq k \leq N - 1$. A *string* is an ordered collection of cluster decompositions $S = (D_N, D_{N-1}, \dots, D_k)$, such that

- (i) D_l has l clusters, and
- (ii) D_{l+1} is a refinement of D_l , i.e., D_{l+1} is formed from D_l by splitting one cluster into two.

DEFINITION. A string $S = (D_N, D_{N-1}, \dots, D_k)$ is called *long* if $k = 2$. If $k \geq 3$, then S is a *short* string.

DEFINITION. Suppose $S = (D_N, D_{N-1}, \dots, D_k)$ is a string. Since D_{N-1} has exactly one cluster containing two elements, it may be identified with a pair α . Define $U_S = U_\alpha$ and

$$\begin{aligned} \tilde{C}_S(z) = & W_\alpha(z - H_{D_{N-1}})^{-1}(V_{D_{N-2}} - V_{D_{N-1}})(z - H_{D_{N-2}})^{-1} \\ & \dots (V_{D_k} - V_{D_{k-1}})(z - H_{D_k})^{-1}. \end{aligned} \quad (\text{IV.2})$$

DEFINITION. Let $S = (D_N, D_{N-1}, \dots, D_k)$ and $S' = (D'_N, D'_{N-1}, \dots, D'_l)$ be strings.

$$\tilde{M}_{S,S'}(z) = \begin{cases} 0 & \text{if } S \text{ is short,} \\ 0 & \text{if } S \text{ is long and } D'_{N-1} \text{ refines } D_2, \\ \tilde{C}_S U_{S'} & \text{if } S \text{ is long and } D'_{N-1} \text{ does not refine } D_2. \end{cases}$$

REMARK. $\tilde{M}_{S,S'}$ has been defined so that its nonzero entries are sums of barely connected parts. The partition D'_{N-1} refines D_2 if and only if $\tilde{C}_S U_{S'}$ is not a sum of barely connected parts.

Each disconnected truncated graph

$$G = W_{\alpha_1}(z - H_0)^{-1}V_{\alpha_2}(z - H_0)^{-1} \dots V_{\alpha_n}(z - H_0)^{-1}$$

has an associated string determined by the following procedure. For $j = 1, 2, \dots, n$, there is a unique cluster decomposition $D(j)$, with a maximal number of clusters, such that all the pairs $\alpha_1, \dots, \alpha_j$ refine $D(j)$. Since G is disconnected, the sequence $D(1), D(2), \dots, D(k)$ has the following properties:

- (i) $D(1)$ has $N - 1$ clusters,
- (ii) $D(n)$ has $k \geq 2$ clusters,
- (iii) either $D(i) = D(i + 1)$ or $D(i)$ has one cluster more than $D(i + 1)$, and
- (iv) $D(i)$ refines $D(j)$ whenever $i \leq j$.

As a consequence, removal of the repetitions in the sequence $D(1), D(2), \dots, D(n)$ yields a sequence $D_{N-1}, D_{N-2}, \dots, D_k$. The string associated with G is $S = (D_N, D_{N-1}, \dots, D_k)$.

At large negative $\text{Re } z$, each resolvent $(z - H_D)^{-1}$ equals its perturbation series. Substituting the series for each resolvent factor in $\tilde{C}_S(z)$, $\tilde{C}_S(z)$ becomes the sum of all truncated graphs whose string is S . Thus $\mathcal{C}_\alpha(z)$ may be written as the sum of all $\tilde{C}_S(z)$, such that $S = (D_N, D_{N-1}, \dots, D_k)$, and D_{N-1} is the decomposition determined by the pair α .

Similarly, $\mathfrak{N}_{\alpha,\beta}(z)$ is the sum of terms $\tilde{M}_{S,S'}(z)$, where S' is any string with D'_{N-1} identified with β , and the sum is over all S , with D_{N-1} identified with α .

With \mathfrak{N} and \mathcal{C} written in closed form, we have

$$(z - H)^{-1} = (z - H_0)^{-1} + \mathcal{Q}(z)(1 - \mathfrak{N}(z))^{-1}\mathcal{C}(z). \quad (\text{IV.4})$$

By the quadratic form methods of §IIB, $\mathcal{Q}(z)$, $\mathfrak{N}(z)$, and $\mathcal{C}(z)$ are analytic for $z \in \mathbb{C} \setminus \sigma_{\text{ess}}(H)$, and $\mathfrak{N}(z)$ is compact. Therefore, the uniqueness of analytic continuation shows that equation (IV.4) is valid for $z \in \mathbb{C} \setminus \sigma(H)$, except for an at most countable set, where the right-hand side may have removable singularities.

EXAMPLES. (1) When $N = 3$, equation (IV.4) is a modified form of the Faddeev equation [6]. This equation with $U_\alpha = |V_\alpha|^{1/2}$ and $W_\alpha = |V_\alpha|^{1/2} \text{sgn}(V_\alpha)$ may also be found in Ginibre and Moulin [7]. Explicitly

$$\begin{aligned} \mathcal{Q}(z) &= \begin{bmatrix} (z - H_0)^{-1}U_{12} & (z - H_0)^{-1}U_{13} & (z - H_0)^{-1}U_{23} \end{bmatrix}, \\ \mathfrak{N}(z) &= \begin{bmatrix} 0 & W_{12}(z - H_{12})^{-1}U_{13} & W_{12}(z - H_{12})^{-1}U_{23} \\ W_{13}(z - H_{13})^{-1}U_{12} & 0 & W_{13}(z - H_{13})^{-1}U_{23} \\ W_{23}(z - H_{23})^{-1}U_{12} & W_{23}(z - H_{23})^{-1}U_{13} & 0 \end{bmatrix}, \\ \mathcal{C}(z) &= \begin{bmatrix} W_{12}(z - H_{12})^{-1} \\ W_{13}(z - H_{13})^{-1} \\ W_{23}(z - H_{23})^{-1} \end{bmatrix}. \end{aligned}$$

(2) When $N = 4$, the matrices have six rows or columns, and the entries are more complicated. The following is a representative sample of the entries:

$$\begin{aligned} \mathcal{Q}_{12}(z) &= (z - H_0)^{-1}U_{12}, \\ \mathfrak{N}_{12,12}(z) &= 0, \\ \mathfrak{N}_{12,13}(z) &= W_{12}(z - H_{12})^{-1}(V_{14} + V_{24})(z - H_{124})^{-1}U_{13} \\ &\quad + W_{12}(z - H_{12})^{-1}V_{34}(z - H_{12,34})^{-1}U_{13}, \\ \mathfrak{N}_{12,34}(z) &= W_{12}(z - H_{12})^{-1}(V_{13} + V_{23})(z - H_{123})^{-1}U_{34} \\ &\quad + W_{12}(z - H_{12})^{-1}(V_{14} + V_{24})(z - H_{124})^{-1}U_{34}, \\ \mathcal{C}_{12}(z) &= W_{12}(z - H_{12})^{-1} + W_{12}(z - H_{12})^{-1}(V_{13} + V_{23})(z - H_{123})^{-1} \\ &\quad + W_{12}(z - H_{12})^{-1}(V_{14} + V_{24})(z - H_{124})^{-1} \\ &\quad + W_{12}(z - H_{12})^{-1}V_{34}(z - H_{12,34})^{-1}. \end{aligned}$$

For $N \geq 4$, several of the matrix entries appearing in equation (IV.4) contain many terms. In order to avoid certain combinatorial difficulties it is necessary to obtain matrices with less complicated entries. This can be done at the expense of using larger matrices.

Fix $N \geq 3$ and let l denote the number of strings which occur in the N body case. $\tilde{A}(z)$ is the $1 \times l$ matrix with $\tilde{A}_S(z) = (z - H_0)^{-1} U_S$. $\tilde{M}_{S,S'}(z)$ is the $l \times l$ matrix whose entries are given by equation (IV.3). $\tilde{C}(z)$ is the $l \times 1$ matrix whose entries are given by equation (IV.2).

Our second formula for the N -body resolvent is

$$(z - H)^{-1} = (z - H_0)^{-1} + \tilde{A}(z)(1 - \tilde{M}(z))^{-1} \tilde{C}(z). \quad (\text{IV.5})$$

The validity of this formula may be established by computing $\mathcal{Q}(z)(\mathcal{M}(z))^k \mathcal{C}(z)$, with the entries of $\mathcal{M}(z)$ expressed in terms of the operators $\tilde{M}_{S,S'}(z)$. The result of this computation is $\tilde{A}(z)(\tilde{M}(z))^k \tilde{C}(z)$. Equation (IV.5) then follows from equation (IV.4) by using geometric series.

An alternative proof may be given. Notice first that each barely connected part is the product of a disconnected truncated graph and a final factor U_β . Therefore, each barely connected part has an associated string determined by its initial truncated graph.

Next, recall that each graph G may be uniquely decomposed as $A_\alpha B_1 B_2 \dots B_k G'$, where B_1, B_2, \dots, B_k are barely connected parts and G' is disconnected. Let S_1, S_2, \dots, S_{k+1} be the strings associated to B_1, B_2, \dots, B_k , and G' , respectively. Then G determines the sequence S_1, S_2, \dots, S_{k+1} uniquely.

To sum all graphs, first sum all graphs with a given sequence of strings, and then sum over all sequences. Since $\tilde{M}_{S,S'}(z)$ is the sum of all barely connected parts with string S and final factor $U_{S'}$, the sum of all graphs with sequence S_1, S_2, \dots, S_{k+1} is: $\tilde{A}_{S_1} \tilde{M}_{S_1, S_2} \tilde{M}_{S_2, S_3} \dots \tilde{M}_{S_k, S_{k+1}} \tilde{C}_{S_{k+1}}$. Thus, the sum of all graphs with sequence of length $k + 1$ is $\tilde{A}(z)(\tilde{M}(z))^k \tilde{C}(z)$. Equation (IV.5) follows by using geometric series and the uniqueness of analytic continuation.

REMARK. Equation (IV.4) involves a $\frac{1}{2}N(N-1) \times \frac{1}{2}N(N-1)$ matrix in the N particle case. Equation (IV.5) contains a much larger matrix. For the N particle case, it is $l_N \times l_N$, where $l_3 = 3$ and $l_N = \frac{1}{2}N(N-1)(l_{N-1} + 1)$.

$$\begin{aligned} N = 3: & \quad l_N = 3, \\ N = 4: & \quad l_N = 24, \\ N = 5: & \quad l_N = 250, \\ N = 6: & \quad l_N = 3765. \end{aligned}$$

Next, we give the final resolvent formula for the three body case. The same formula may be found in Ginibre and Moulin [7]; similar equations may be found in Thomas [36] and Howland [11].

For the moment, the operators ρ_{12} , ρ_{13} , and ρ_{23} will not be specified. They will be chosen in §V so that certain bounds can be established. Algebraically, equation (IV.6) holds for any choice of these ρ functions.

Denoting $(z - H_0)^{-1}$ and $(z - H_\alpha)^{-1}$ by G_0 and G_α , respectively, the final three-body equation is:

$$(z - H)^{-1} = (z - H_0)^{-1} + A(z)(1 - M(z))^{-1}C(z), \quad (\text{IV.6})$$

where the matrices $A(z)$, $M(z)$ and $C(z)$ are given in Figure 1.

We view $A(z)$, $M(z)$, and $C(z)$ as 1×3 , 3×3 , and 3×1 matrices of 1×2 , 2×2 , and 2×1 blocks, respectively. In accordance with this viewpoint, we label the entries $A_{\alpha,i}$, $M_{\alpha,i;\beta,j}$, and $C_{\alpha,i}$, where α and β run over the pairs, while i and j run from 1 to 2.

Equation (IV.6) holds because

$$A(z)(M(z))^k C(z) = \mathcal{Q}(z)(\mathfrak{M}(z))^k \mathcal{C}(z) \quad (\text{IV.7})$$

for $k \geq 0$, and because geometric series may be used at large negative $\text{Re } z$ when the ρ_α are suitably chosen.

To prove equation (IV.7), consider a typical nonzero term in the expression for $\mathcal{Q} \mathfrak{M}^k \mathcal{C}$. It has the form

$$\begin{aligned} & \mathcal{Q}_{\alpha_1} \mathfrak{M}_{\alpha_1, \alpha_2} \dots \mathfrak{M}_{\alpha_k, \alpha_{k+1}} \mathcal{C}_{\alpha_{k+1}} \\ &= G_0 U_{\alpha_1} W_{\alpha_1} G_{\alpha_1} U_{\alpha_2} W_{\alpha_2} G_{\alpha_2} U_{\alpha_3} \dots W_{\alpha_k} G_{\alpha_k} U_{\alpha_{k+1}} W_{\alpha_{k+1}} G_{\alpha_{k+1}}, \end{aligned} \quad (\text{IV.8})$$

where $\alpha_i \neq \alpha_{i+1}$.

Let $\mathcal{Q}_{\alpha,1} = (1 - P_\alpha)$ and $\mathcal{Q}_{\alpha,2} = P_\alpha = (P_\alpha \rho_\alpha)(\rho_\alpha^{-1} P_\alpha)$, and replace each G_α factor by $G_\alpha \mathcal{Q}_{\alpha,1} + G_\alpha \mathcal{Q}_{\alpha,2}$ in equation (IV.8). The right side of equation (IV.8) becomes a sum of 2^{k+1} terms of the form:

$$\begin{aligned} & G_0 U_{\alpha_1} W_{\alpha_1} G_{\alpha_1} \mathcal{Q}_{\alpha_1, i_1} U_{\alpha_2} W_{\alpha_2} G_{\alpha_2} \mathcal{Q}_{\alpha_2, i_2} U_{\alpha_3} \dots W_{\alpha_k} G_{\alpha_k} \mathcal{Q}_{\alpha_k, i_k} U_{\alpha_{k+1}} W_{\alpha_{k+1}} G_{\alpha_{k+1}} \mathcal{Q}_{\alpha_{k+1}, i_{k+1}} \\ &= A_{\alpha_1, i_1} M_{\alpha_1, i_1; \alpha_2, i_2} M_{\alpha_2, i_2; \alpha_3, i_3} \dots M_{\alpha_k, i_k; \alpha_{k+1}, i_{k+1}} C_{\alpha_{k+1}, i_{k+1}}. \end{aligned} \quad (\text{IV.9})$$

Summing over all indices and using the identity $G_0 V_\alpha G_\alpha = (G_\alpha - G_0)$, equation (IV.7) is obtained.

Equations (IV.4) and (IV.5) are not suitable for scattering because $\mathfrak{M}(z)$, $\mathcal{C}(z)$, $\tilde{M}(z)$, and $\tilde{C}(z)$ are all singular as z approaches the essential spectrum of H . The advantage of equation (IV.6) is that $M(z)$ and $C(z)$ are reasonably well behaved as z approaches the essential spectrum, and the singularities all appear explicitly in the factor $A(z)$.

Equation (IV.6) was obtained from equation (IV.4) by the following procedure: First, the inverse term in equation (IV.4) was expanded using geometric series. Second, the factors $(1 - P_\alpha)$ and $P_\alpha = P_\alpha \rho_\alpha \rho_\alpha^{-1} P_\alpha$ were inserted after each G_α . Third, each term was decomposed as a product in a way which depended on the sequence of projections $(1 - P_\alpha)$ or P_α . Then, the result was resummed to give equation (IV.6).

The four body case is similar. Equations (IV.4) and (IV.5) are not suitable, so equation (IV.10) will be derived from equation (IV.5). Again geometric series will be used, and factors $(1 - P_{D_2})$, P_{D_2} , $(1 - P_{D_3})$, P_{D_3} will be inserted, along with ρ 's and ρ^{-1} 's. Next, we "symmetrize" certain terms, which does not change the sum of all terms. Resumming by geometric series, we obtain equation (IV.10).

$$\begin{aligned}
 A(z) &= \begin{bmatrix} G_{012} & (G_{12}-G_0)P_{12}P_{12} & G_{013} & (G_{13}-G_0)P_{13}P_{13} & G_{023} & (G_{23}-G_0)P_{23}P_{23} \end{bmatrix} \\
 M(z) &= \begin{bmatrix} 0 & 0 & W_{12}G_{12}(1-P_{12})U_{13} & W_{12}G_{12}(1-P_{12})V_{13}G_{13}P_{13}P_{13} & W_{12}G_{12}(1-P_{12})U_{23} & W_{12}G_{12}(1-P_{12})V_{23}G_{23}P_{23}P_{23} \\ 0 & 0 & \rho_{12}^{-1}P_{12}U_{13} & \rho_{12}^{-1}P_{12}V_{13}G_{13}P_{13}P_{13} & \rho_{12}^{-1}P_{12}U_{23} & \rho_{12}^{-1}P_{12}V_{23}G_{23}P_{23}P_{23} \\ W_{13}G_{13}(1-P_{13})U_{12} & W_{13}G_{13}(1-P_{13})V_{12}G_{12}P_{12}P_{12} & 0 & 0 & W_{13}G_{13}(1-P_{13})U_{23} & W_{13}G_{13}(1-P_{13})V_{23}G_{23}P_{23}P_{23} \\ \rho_{13}^{-1}P_{13}U_{12} & \rho_{13}^{-1}P_{13}V_{12}G_{12}P_{12}P_{12} & 0 & 0 & \rho_{13}^{-1}P_{13}U_{23} & \rho_{13}^{-1}P_{13}V_{23}G_{23}P_{23}P_{23} \\ W_{23}G_{23}(1-P_{23})U_{12} & W_{23}G_{23}(1-P_{23})V_{12}G_{12}P_{12}P_{12} & W_{23}G_{23}(1-P_{23})U_{13} & W_{23}G_{23}(1-P_{23})V_{13}G_{13}P_{13}P_{13} & 0 & 0 \\ \rho_{23}^{-1}P_{23}U_{12} & \rho_{23}^{-1}P_{23}V_{12}G_{12}P_{12}P_{12} & \rho_{23}^{-1}P_{23}U_{13} & \rho_{23}^{-1}P_{23}V_{13}G_{13}P_{13}P_{13} & 0 & 0 \end{bmatrix} \\
 C(z) &= \begin{bmatrix} W_{12}G_{12}(1-P_{12}) \\ \rho_{12}^{-1}P_{12} \\ W_{13}G_{13}(1-P_{13}) \\ \rho_{13}^{-1}P_{13} \\ W_{23}G_{23}(1-P_{23}) \\ \rho_{23}^{-1}P_{23} \end{bmatrix}
 \end{aligned}$$

FIGURE 1

REMARKS. (1) In the three body case, no symmetrization step was necessary because all strings contained only one nontrivial cluster decomposition. In the four body case, some terms require symmetrization. This involves the replacement of certain terms by a combinatorial factor times a sum of those terms.

(2) The symmetrization step allows us to prove asymptotic completeness when the subsystems have embedded (nonpositive) eigenvalues. The equations used by Sigal [26] are not symmetrized, and embedded eigenvalues are not treated by Sigal's asymptotic completeness proof.

(3) Equation (IV.10) is derived from equation (IV.5) by replacing the entries of the 24×24 matrix of equation (IV.5) by 3×3 blocks. As a result, equation (IV.10) has an unpleasantly large 72×72 matrix. An equation equivalent to equation (IV.10) can be derived from equation (IV.4). This formula has a 42×42 matrix, but the entries are complicated. So, we prefer to use equation (IV.10).

The final resolvent formula for the four body case is

$$(z - H)^{-1} = (z - H_0)^{-1} + A(z)(1 - M(z))^{-1}C(z), \quad (\text{IV.10})$$

where A , M , and C are the 1×72 , 72×72 , and 72×1 matrices indicated in the Appendix. A representative sample of the entries is given, and the number associated with each entry refers to the proposition of §V in which that entry is studied. A , M , and C should be viewed as 1×24 , 24×24 , and 24×1 matrices of 1×3 , 3×3 , and 3×1 blocks, respectively. The entries are denoted by $A_{S,i}$, $M_{S,i;S',j}$, and $C_{S,i}$, where S and S' denote strings, and i and j run from 1 to 3.

As in the three body case, equation (IV.10) involves some operators ρ_{ij} , $\rho_{ij,k}$, and $\rho_{ij,kl}$, which will be specified in §V. Also, equation (IV.10) holds because

$$A(z)(M(z))^k C(z) = \tilde{A}(z)(\tilde{M}(z))^k \tilde{C}(z) \quad (\text{IV.11})$$

for $k \geq 0$. Equation (IV.11) implies equation (IV.10) in the same way equation (IV.7) implied (IV.6). However, the proof of equation (IV.11) is more complicated than that of equation (IV.7).

To prove equation (IV.11), consider first the case $k = 0$, and define $Q_{D_3,1} = (1 - P_{D_3})$, $Q_{D_3,2} = P_{D_3}$, $Q_{D_3,3} = 1$, $Q_{D_2,1} = Q_{D_2,2} = (1 - P_{D_2})$, and $Q_{D_2,3} = P_{D_2}$. If $S = (D_4, D_3)$ is a short string, then

$$\begin{aligned} \tilde{A}_S \tilde{C}_S &= G_0 V_{D_3} G_{D_3} \\ &= G_0 V_{D_3} G_{D_3} Q_{D_3,1} + G_0 V_{D_3} G_{D_3} Q_{D_3,2} \\ &= A_{S,1} C_{S,1} + A_{S,2} C_{S,2} + 0 \\ &= \sum_{i=1}^3 A_{S,i} C_{S,i}. \end{aligned} \quad (\text{IV.12})$$

For long strings $S = (D_4, D_3, D_2)$, the situation is not so simple because $A_{S,3}$ has been symmetrized. For each decomposition D_2 with two clusters, let $\Lambda_{D_2} = \{S' = (D_4, D'_3, D'_2): D'_2 = D_2\}$, and let $l(D_2)$ denote the number of elements of Λ_{D_2} . Then

$$\begin{aligned}
\sum_{S \in \Lambda_{D_2}} \tilde{A}_S \tilde{C}_S &= \sum_{D_3 \text{ refines } D_2} G_0 V_{D_3} G_{D_3} (V_{D_2} - V_{D_3}) G_{D_2} \\
&= \sum_{D_3 \text{ refines } D_2} \sum_{i=1}^2 G_0 V_{D_3} G_{D_3} Q_{D_3,i} (V_{D_2} - V_{D_3}) G_{D_2} Q_{D_2,i} \\
&\quad + \sum_{D_3' \text{ refines } D_2} G_0 V_{D_3'} G_{D_3'} Q_{D_3',3} (V_{D_2} - V_{D_3'}) G_{D_2} Q_{D_2,3} \\
&= \sum_{S \in \Lambda_{D_2}} \sum_{i=1}^2 A_{S,i} C_{S,i} \\
&\quad + \sum_{D_3 \text{ refines } D_2} \left(\frac{1}{l(D_2)} \sum_{D_3' \text{ refines } D_2} G_0 V_{D_3'} G_{D_3'} Q_{D_3',3} (V_{D_2} - V_{D_3'}) G_{D_2} Q_{D_2,3} \right) \\
&= \sum_{S \in \Lambda_{D_2}} \sum_{i=1}^3 A_{S,i} C_{S,i}. \tag{IV.13}
\end{aligned}$$

Equation (IV.11), for $k = 0$, is now obtained from (IV.12) and (IV.13) by the following computation:

$$\begin{aligned}
\tilde{A}\tilde{C} &= \sum_S \tilde{A}_S \tilde{C}_S = \sum_{S \text{ short}} \tilde{A}_S \tilde{C}_S + \sum_{D_2} \sum_{S \in \Lambda_{D_2}} \tilde{A}_S \tilde{C}_S \\
&= \sum_{S \text{ short}} \sum_{i=1}^3 A_{S,i} C_{S,i} + \sum_{D_2} \sum_{S \in \Lambda_{D_2}} \sum_{i=1}^3 A_{S,i} C_{S,i} \\
&= \sum_S \sum_{i=1}^3 A_{S,i} C_{S,i} = AC.
\end{aligned}$$

When $k > 0$, the ideas are the same, but more sums are involved. For each sequence $D_2^{(1)}, D_2^{(2)}, \dots, D_2^{(k)}$, let $\Lambda_j = \Lambda_{D_2^{(j)}}$. Then, by explicit computation,

$$\begin{aligned}
&\sum_{S_1 \in \Lambda_1} \sum_{S_2 \in \Lambda_2} \dots \sum_{S_k \in \Lambda_k} \tilde{A}_{S_1} \tilde{M}_{S_1, S_2} \tilde{M}_{S_2, S_3} \dots \tilde{M}_{S_k, S_{k+1}} \tilde{C}_{S_{k+1}} \\
&= \sum_{S_1 \in \Lambda_1} \sum_{S_2 \in \Lambda_2} \dots \sum_{S_k \in \Lambda_k} \sum_{i_1=1}^3 \sum_{i_2=1}^3 \dots \\
&\quad \sum_{i_{k+1}=1}^3 A_{S_1, i_1} M_{S_1, i_1; S_2, i_2} \dots M_{S_k, i_k; S_{k+1}, i_{k+1}} C_{S_{k+1}, i_{k+1}} \tag{IV.14}
\end{aligned}$$

for any choice of S_{k+1} . This is the generalization of equation (IV.13). Equation (IV.12) does not need to be generalized because $M_{S, S'} = 0$ whenever S is short. Now equation (IV.11) follows from (IV.14).

$$\begin{aligned}
\tilde{A}\tilde{M}^k\tilde{C} &= \sum_{S_1} \sum_{S_2} \cdots \sum_{S_k} \sum_{S_{k+1}} \tilde{A}_{S_1} \tilde{M}_{S_1, S_2} \cdots \tilde{M}_{S_k, S_{k+1}} \tilde{C}_{S_{k+1}} \\
&= \sum_{S_1 \text{ long}} \sum_{S_2 \text{ long}} \cdots \sum_{S_k \text{ long}} \sum_{S_{k+1}} \tilde{A}_{S_1} \tilde{M}_{S_1, S_2} \cdots \tilde{M}_{S_k, S_{k+1}} \tilde{C}_{S_{k+1}} \\
&= \sum_{D_2^{(1)}} \sum_{D_2^{(2)}} \cdots \sum_{D_2^{(k)}} \sum_{S_1 \in \Lambda_1} \sum_{S_2 \in \Lambda_2} \cdots \\
&\quad \sum_{S_k \in \Lambda_k} \sum_{S_{k+1}} \tilde{A}_{S_1} \tilde{M}_{S_1, S_2} \cdots \tilde{M}_{S_k, S_{k+1}} \tilde{C}_{S_{k+1}} \\
&= \sum_{D_2^{(1)} \dots D_2^{(k)}} \sum_{S_1 \in \Lambda_1 \dots S_k \in \Lambda_k} \sum_{S_{k+1}} \sum_{i_1 \dots i_{k+1}}^3 A_{S_1, i_1} M_{S_1, i_1; S_2, i_2} \\
&\quad \dots M_{S_k, i_k; S_{k+1}, i_{k+1}} C_{S_{k+1}, i_{k+1}} \\
&= AM^kC.
\end{aligned}$$

This establishes equation (IV.11).

In the three body case, equations (IV.8) and (IV.9) show that individual terms in the expression for AM^kC correspond to products of blocks of A , M , and C . This is not true in the four body case, although something closely related is true. Products of certain blocks of \tilde{A} , \tilde{M} , and \tilde{C} (determined by the sequence $D_2^{(1)}, \dots, D_2^{(k)}$) are equal to products of corresponding blocks of A , M , and C . This is the content of equation (IV.14). Equation (IV.11) is then obtained by summing over the blocks.

Having established equation (IV.11), the validity of equation (IV.10) is proved. In §V, $(M(z))^2$ is shown to be compact, and the bounds required by the multiparticle limiting absorption principle are proved. However, $A(z)$ must first be put into the correct form. This is done by using the following identities:

$$\begin{aligned}
(1) \quad & G_0 V_{ij} G_{ij} = G_{ij} - G_0, \\
(2) \quad & G_0 (V_{ij} G_{ij} (V_{ik} + V_{jk}) + V_{ik} G_{ik} (V_{ij} + V_{jk}) + V_{jk} G_{jk} (V_{ij} + V_{ik})) G_{ijk} \\
&= G_{ijk} - (G_{ij} P_{ij} + G_{ik} P_{ik} + G_{jk} P_{jk}) - G_0 (1 - P_{ij} - P_{ik} - P_{jk}) \\
&\quad - G_0 (V_{ij} G_{ij} (1 - P_{ij}) + V_{ik} G_{ik} (1 - P_{ik}) + V_{jk} G_{jk} (1 - P_{jk})), \\
(3) \quad & G_0 (V_{ij} G_{ij} V_{kl} + V_{kl} G_{kl} V_{ij}) G_{ij, kl} \\
&= G_{ij, kl} - (G_{ij} P_{ij} + G_{kl} P_{kl}) - G_0 (1 - P_{ij} - P_{kl}) \\
&\quad - G_0 (V_{ij} G_{ij} (1 - P_{ij}) + V_{kl} G_{kl} (1 - P_{kl})). \tag{IV.15}
\end{aligned}$$

Notice that the symmetrization is necessary to put A in a form in which these identities may be used to give a formula of the type required in §III. The other reason for the symmetrization is to make $M(z)$ well behaved as z approaches $\sigma_{\text{ess}}(H)$.

We give in the Appendix a representative sample of the blocks of the matrices $A(z)$, $C(z)$, and $M(z)$.

V. Estimates. Theorem III.1 and the formulas of §IV reduce the problem of Kuroda completeness to the proof of certain operator bounds. These bounds are established in this section under the following hypotheses:

(1) $H = H_0 + \sum_{i < j} V_{ij}$ on $L^2(\mathbf{R}^{(N-1)m})$, where $m \geq 3$ and $N \leq 4$. Each two body potential V_{ij} is a selfadjoint multiplication operator $U_{ij}W_{ij}$, where U_{ij} and W_{ij} belong to $L^p_\delta(\mathbf{R}^m) + L^\infty_\delta(\mathbf{R}^m)$ for some $p > m$ and $\delta > 1$.

(2) $U_{ij}: \mathcal{H} \rightarrow \mathcal{H}_{-1}$ and $W_{ij}: \mathcal{H}_{+1} \rightarrow \mathcal{H}$ are dilation analytic.

(3) No cluster Hamiltonian has a bound state or resonance at any of its thresholds.

(4) When $N = 4$, we assume the three body cluster Hamiltonians have no positive eigenvalues.

(5) Each cluster Hamiltonian has exactly one bound state.

REMARKS. (1) Hypothesis (2) asserts dilation analyticity only. The boundedness of $U_{ij}: \mathcal{H} \rightarrow \mathcal{H}_{-1}$ and $W_{ij}: \mathcal{H}_{+1} \rightarrow \mathcal{H}$ follow from Hypothesis (1) and methods of §II.B.

(2) Bounds established for the cluster Hamiltonians are required to prove bounds for the full Hamiltonian. So, we need two body information to treat the three body case, and two and three body information to treat the four body case.

(3) The threshold set $\Sigma(h)$ for a cluster Hamiltonian h is defined as in §II.E. We prove $(z - h)^{-1} = (z - h_0)^{-1} + A(z)(1 - M(z))^{-1}C(z)$, where $M(z)$ has norm continuous boundary values $M(E \pm i0)$ as z approaches $E \in \mathbf{R}$ from above or below. Hypothesis (3) states that $1 \notin \sigma(M(E \pm i0))$ whenever $E \in \Sigma(h)$.

(4) By continuity, Hypothesis (3) implies $1 \notin \sigma(M(E \pm i0))$ for E in the neighborhood of $\Sigma(h)$. We prove $E \notin \Sigma(h)$ and $1 \notin \sigma(M(E \pm i0))$ imply E is not an eigenvalue of h . Thus, h has no eigenvalues near its thresholds. Hypotheses (1) and (2) imply that eigenvalues may accumulate only at thresholds (see Balslev and Combes [3]). So h has finitely many eigenvalues. Therefore, Hypotheses (1)–(3) imply H has finitely many channels.


(5) In §VI, we prove that Hypotheses (1) and (2) imply Hypothesis (3) for a dense open set of couplings.

(6) Hypothesis (5) has been introduced merely for convenience. We denote cluster eigenfunctions ψ_{12}, ψ_{123} , etc., and let their corresponding eigenvalues be E_{12}, E_{123} , etc., respectively. Define $\psi_{12,34} = \psi_{12} \otimes \psi_{34}$, $\psi_{13,24} = \psi_{13} \otimes \psi_{24}$, and $\psi_{14,23} = \psi_{14} \otimes \psi_{23}$. Similarly, let $E_{12,34} = E_{12} + E_{34}$, $E_{13,24} = E_{13} + E_{24}$, and $E_{14,23} = E_{14} + E_{23}$.

(7) Theorem II.10 gives sufficient conditions for Hypothesis (4) to hold. Yukawa potentials, generalized Yukawa potentials and potentials of the form $(1 + r)^{-2-\epsilon}$ are allowed.

Since there are finitely many potentials, Hypothesis (1) implies the existence of a $p > m$ and $\delta > \gamma > 1$, such that all U_{ij} and W_{ij} belong to $L^p_\delta(\mathbf{R}^m) + L^\infty_\delta(\mathbf{R}^m)$. Fix such a γ and p .

DEFINITION. Let ξ be the coordinate from the center of mass of particles i and j to particle k . Let ζ denote the coordinate from the center of mass of particles i, j , and k to particle l . Let η be the coordinate from the center of mass of particles i and j to the center of mass of particles k and l .



Define

$$\rho_{ij,k}(\xi) = (1 + \xi^2)^{-\gamma/2},$$

$$\rho_{ijk}(\zeta) = (1 + \zeta^2)^{-\gamma/2},$$

$$\rho_{ij,kl}(\eta) = (1 + \eta^2)^{-\gamma/2}.$$

In the three body problem, we let $\rho_{ij} = \rho_{ij,k}$ for convenience. In the four body problem we choose functions ρ_{ij} arbitrarily as $\rho_{12} = \rho_{12,3}$, $\rho_{13} = \rho_{13,2}$, $\rho_{23} = \rho_{23,1}$, $\rho_{34} = \rho_{34,1}$, $\rho_{14} = \rho_{14,3}$, and $\rho_{24} = \rho_{24,3}$.

These are the ρ functions introduced in §IV. The choice is certainly not unique, and we have not made the same choice as other authors. However, our ρ functions have the advantage of being dilation analytic.

The results of this section are summarized by the first three propositions. These depend on the technical lemmas which make up most of this section.

PROPOSITION V.1. *Assume Hypothesis (1). If $N = 2$, then the limiting absorption principle holds for H . If Hypothesis (2) also holds, then the exceptional set \mathfrak{E} is contained in $(-\infty, 0]$, and \mathfrak{E} consists of all eigenvalues of H and possibly the point 0.*

PROOF. Equation (IV.1) states

$$(z - H)^{-1} = (z - H_0)^{-1} + (z - H_0)^{-1}U(1 - W(z - H_0)^{-1}U)^{-1}W(z - H_0)^{-1}.$$

Let $\rho(x) = (1 + x^2)^{-\gamma/2}$. Define $F_1 = \rho$, $F_2 = U$, $Z_1(z) = \rho^{-1}$, and $Z_2(z) = (1 - W(z - H_0)^{-1}U)^{-1}W(z - H_0)^{-1}$. Then

$$(z - H)^{-1} = \sum_{i=1}^2 (z - H_0)^{-1}F_iZ_i(z).$$

Hölder's inequality shows that F_1 and F_2 map $L^2(\mathbf{R}^m)$ into $L_\gamma^q(\mathbf{R}^m) + L_\gamma^2(\mathbf{R}^m)$, where $q^{-1} = p^{-1} + 1/2$.

Repeating the proof of Lemma II.3, we see that $W(z - H_0)^{-1}U$ is uniformly bounded and analytic for $z \in \mathbb{C} \setminus [0, \infty)$, with norm continuous boundary values as z approaches $[0, \infty)$ from above or below. Similarly, $W(z - H_0)^{-1}$ has these properties as a mapping of $L_\gamma^2(\mathbf{R}^m)$ into $L^2(\mathbf{R}^m)$.

If U and W belong to $L^2(\mathbf{R}^m)$, then $We^{-iH_0t}U$ has an L^2 integral kernel for $t \neq 0$. Thus, $We^{-iH_0t}U$ is compact for $t \neq 0$. When U and W belong to $L_\gamma^p(\mathbf{R}^m) + L_\gamma^\infty(\mathbf{R}^m)$, we may approximate them by L^2 functions, and conclude the compactness

of $We^{-itH_0}U$ for $t \neq 0$, by taking limits. Taking Fourier transforms as in Lemma II.3, we see that $W(z - H_0)^{-1}U$ is compact, and $\|W(z - H_0)^{-1}U\| \rightarrow 0$ as $|\operatorname{Im} z| \rightarrow \infty$ or $\operatorname{Re} z \rightarrow -\infty$.

By the analytic Fredholm theorem and a theorem of Kuroda [16] (see Simon [29]), $(1 - W(z - H_0)^{-1}U)^{-1}$ exists for all z in the closed cut plane, except for some set \mathcal{E} . $\mathcal{E} \cap (\mathbb{C} \setminus [0, \infty))$ is discrete; $\mathcal{E} \cap [0, \infty)$ is closed and has Lebesgue measure 0.

If $w \in \mathcal{E} \cap (\mathbb{C} \setminus [0, \infty))$, then $\phi = W(w - H_0)^{-1}U\phi$ for some $\phi \in \mathcal{H}$. Since $w \notin \sigma(H_0)$, $(w - H_0)^{-1}U$ is bounded, and $\psi = (w - H_0)^{-1}U\phi$ belongs to \mathcal{H} . However, $\phi = W(w - H_0)^{-1}U\phi = W\psi$, and $\psi = (w - H_0)^{-1}V\psi$. Applying $(w - H_0)$ to both sides of this equation, we obtain $H\psi = w\psi$. Therefore, $\mathcal{E} \subseteq \mathbb{R}$ and $\mathcal{E} \cap (-\infty, 0)$ consists of eigenvalues of H .

If $w \pm i0 \in \mathcal{E} \cap [0, \infty)$, and Hypothesis (2) holds, then Lemma II.11 shows that $1 \in \sigma(W(\theta)(w - H_0(\theta))^{-1}U(\theta))$ for some θ with $\pm \operatorname{Im} \theta > 0$. If $w \neq 0$, then $\delta \notin \sigma(H_0(\theta))$, and we can construct ψ (as above), so that $H(\theta)\psi = w\psi$. Theorem II.7 shows that w is an eigenvalue of H . Since $w > 0$, this contradicts Theorem II.10. Therefore, $\mathcal{E} \subseteq (-\infty, 0]$ consists of eigenvalues of H and possibly the point 0. \square

PROPOSITION V.2 (GINIBRE AND MOULIN [7], THOMAS [36], HOWLAND [11]). *Assume Hypotheses (1)–(3); if $N = 3$, then the multiparticle limiting absorption principle holds for H . The exceptional set \mathcal{E} is at most countable, and consists of eigenvalues of H and possibly some thresholds of H .*

PROOF. For convenience, we also assume Hypothesis (5).

As in the proof of Proposition V.1, we begin by writing

$$(z - H)^{-1} = \sum_D (z - H_D)^{-1} P_D \sum_I F_{I,D} Z_{I,D}(z).$$

This expression is obtained from equation (IV.6). The $F_{I,D}$'s are U_{ij} 's, ρ_{ij} 's, and P_{ij} ρ_{ij} 's. The $Z_{I,D}(z)$'s are made up of components of $(1 - M(z))^{-1}C(z)$.

$C(z): L^2_\gamma(\mathbb{R}^{2m}) \rightarrow \bigoplus_{i=1}^6 L^2(\mathbb{R}^{2m})$ is analytic and uniformly bounded for $z \in \mathbb{C} \setminus \sigma_{\text{ess}}(H)$, with norm continuous boundary values as z approaches $\sigma_{\text{ess}}(H)$ from above or below. This is proved by Lemmas V.4 and V.8 below.

$M(z): \bigoplus_{i=1}^6 L^2(\mathbb{R}^{2m}) \rightarrow \bigoplus_{i=1}^6 L^2(\mathbb{R}^{2m})$ has all the properties mentioned above for $C(z)$. In addition, $(M(z))^2$ is compact and $\|(M(z))^2\| \rightarrow 0$ as $\operatorname{Re} z \rightarrow -\infty$. These facts are proved by Lemma V.13.

Let \mathcal{E} be the set of z , such that $(1 - M(z))^{-1}$ does not exist. If $z \in \mathcal{E} \cap (\mathbb{C} \setminus \sigma_{\text{ess}}(H))$, then the compactness of $(M(z))^2$ shows that $M(z)\phi = \phi$ for some $\phi \in \bigoplus_{i=1}^6 \mathcal{H}$;

$$\phi_{\alpha,i} = \sum_{\beta,j} M_{\alpha,i;\beta,j}(z) \phi_{\beta,j}.$$

Define $\eta_\alpha = \phi_{\alpha,1} + W_\alpha(z - H_\alpha)^{-1}P_\alpha\rho_\alpha\phi_{\alpha,2}$. Then by the relation of $M(z)$ and $\mathfrak{N}(z)$,

$$\begin{aligned}\eta_\alpha &= \sum_\beta \mathfrak{N}_{\alpha,\beta}(z)\eta_\beta = \sum_{\beta \neq \alpha} W_\alpha(z - H_\alpha)^{-1}U_\beta\eta_\beta \\ &= \sum_{\beta \neq \alpha} W_\alpha(z - H_0)^{-1}U_\beta\eta_\beta + \sum_{\beta \neq \alpha} W_\alpha(z - H_0)^{-1}V_\alpha(z - H_\alpha)^{-1}U_\beta\eta_\beta \\ &= \sum_{\beta \neq \alpha} W_\alpha(z - H_0)^{-1}U_\beta\eta_\beta + W_\alpha(z - H_0)^{-1}U_\alpha\eta_\alpha \\ &= \sum_\beta W_\alpha(z - H_0)^{-1}U_\beta\eta_\beta.\end{aligned}\tag{V.1}$$

Since $z \notin \sigma_{\text{ess}}(H)$, $(z - H_0)^{-1}U_\beta$ is bounded. Hence, $\psi = (z - H_0)^{-1}\sum_\beta U_\beta\eta_\beta$ belongs to \mathcal{H} . Moreover, equation (V.1) shows $\eta_\alpha = W_\alpha\psi$, so $\psi = (z - H_0)^{-1}(V_{12} + V_{13} + V_{23})\psi$. Therefore, $H\psi = z\psi$.

If $z \pm i0 \in \mathfrak{E} \cap \sigma_{\text{ess}}(H)$, and z is not a threshold, then Lemma II.11 shows that $z \in \mathfrak{E}$ implies $1 \in \sigma(M(z, \theta))$, for some θ with $\pm \text{Im } \theta > 0$. Since $z \notin \sigma_{\text{ess}}(H(\theta))$, equation (V.1) shows $H(\theta)\psi = z\psi$ for some $\psi \in \mathcal{H}$. Theorem II.7 shows that z is an eigenvalue of H .

Since $M(z)$ depends continuously on z , \mathfrak{E} is a closed set. Since eigenvalues of H may accumulate only at thresholds, \mathfrak{E} is at most countable, and has Lebesgue measure zero. \square

PROPOSITION V.3. *Assume Hypotheses (1)–(4). If $N = 4$, then the multiparticle limiting absorption principle holds for H .*

PROOF. Using equation (IV.10) and the identities (IV.15), we may write $(z - H)^{-1}$ in the required form.

$C(z): L^2_\gamma(\mathbf{R}^{3m}) \rightarrow \bigoplus_{i=1}^{72} L^2(\mathbf{R}^{3m})$ is analytic and uniformly bounded for $z \in \mathbf{C} \setminus \sigma_{\text{ess}}(H)$, with norm continuous boundary values as z approaches $\sigma_{\text{ess}}(H)$ from above or below. This is proved by Corollary V.5, Lemma V.8, and Propositions V.15, V.16, V.25, and V.26.

$M(z): \bigoplus_{i=1}^{72} L^2(\mathbf{R}^{3m}) \rightarrow \bigoplus_{i=1}^{72} L^2(\mathbf{R}^{3m})$ is analytic and uniformly bounded for $z \in \mathbf{C} \setminus \sigma_{\text{ess}}(H)$, with norm continuous boundary values as z approaches $\sigma_{\text{ess}}(H)$ from above or below. $(M(z))^2$ is compact, and $\|(M(z))^2\| \rightarrow 0$ as $\text{Re } z \rightarrow -\infty$. The compactness of M^2 follows from the fact that only the lower left corner entries in any of the 3×3 blocks for M may fail to be compact. Also, these are the only ones which do not approach zero as $\text{Re } z \rightarrow -\infty$.

These facts about $M(z)$ are proved by Lemma V.4 and Propositions V.15–V.20, V.25–V.32.

The exceptional set \mathfrak{E} is defined as the set of all real z such that $1 \in \sigma(M(z))$. The analytic Fredholm theorem and a theorem of Kuroda [16] (see also Simon [29]) show that \mathfrak{E} is a closed set of measure zero.

The only facts remaining to be proved are the boundedness, analyticity, and continuity up to $\sigma_{\text{ess}}(H)$ of certain factors which occur in $A(z)$. These facts are proved by Lemma V.8. \square

LEMMA V.4. Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct indices. Choose coordinates ξ, ζ , and η as in the definitions of $\rho_{ij,k}$, ρ_{ijk} , and $\rho_{ij,kl}$. Then, for all $\alpha, \beta, \delta \in \mathbf{R}$,

(a) $\rho_{ij,k}^{-1} P_{ij} U_{ik}$ is bounded from $L_{-\beta}^2(\mathbf{R}^m, dx_{ij}) \otimes L_{\delta}^2(\mathbf{R}^{2m}, d\xi d\xi^*)$ to $L_{\alpha}^2(\mathbf{R}^m, dx_{ij}) \otimes L_{\delta}^2(\mathbf{R}^{2m}, d\xi d\xi^*)$.

(b) $\rho_{ijk}^{-1} P_{ijk} U_{kl}$ is bounded from $L_{-\beta}^2(\mathbf{R}^{2m}, dx_{ij} d\xi) \otimes L_{\delta}^2(\mathbf{R}^m, d\xi^*)$ to $L_{\alpha}^2(\mathbf{R}^{2m}, dx_{ij} d\xi) \otimes L_{\delta}^2(\mathbf{R}^m, d\xi^*)$.

(c) $\rho_{ij,kl}^{-1} P_{ij,kl} U_{ik}$ is bounded from $L_{-\beta}^2(\mathbf{R}^{2m}, dx_{ij} dx_{kl}) \otimes L_{\delta}^2(\mathbf{R}^m, d\eta)$ to $L_{\alpha}^2(\mathbf{R}^{2m}, dx_{ij} dx_{kl}) \otimes L_{\delta}^2(\mathbf{R}^m, d\eta)$.

PROOF. First, we prove three inequalities involving coordinates:

$$\begin{aligned} & 2(1 + x_{ij}^2)(1 + x_{ik}^2) - (1 - \xi^2) \\ &= 1 + 2x_{ij}^2 x_{ik}^2 + x_{ik}^2 - 2m_j(m_i + m_j)^{-1} x_{ij} \cdot x_{ik} \\ & \quad + (2 - m_j^2(m_i + m_j)^{-2}) x_{ij}^2 \\ & \geq 1 + 2x_{ij}^2 x_{ik}^2 + (x_{ik} - m_j(m_i + m_j)^{-1} x_{ij})^2 \geq 0, \end{aligned}$$

where we have used $(2 - m_j^2(m_i + m_j)^{-2}) \geq 1 > m_j^2(m_i + m_j)^{-2}$. Therefore,

$$(1 + \xi^2) \leq 2(1 + x_{ij}^2)(1 + x_{ik}^2). \quad (\text{V.2})$$

The same computation (but with different masses) shows

$$(1 + \zeta^2) \leq 2(1 + x_{kl}^2)(1 + \xi^2). \quad (\text{V.3})$$

Once again, the same computation (with different masses again) shows: $(1 + \eta^2) \leq 2(1 + \xi^2)(1 + x_{kl}^2)$. Combining this with inequality (V.2), we obtain

$$(1 + \eta^2) \leq 4(1 + x_{ij}^2)(1 + x_{ik}^2)(1 + x_{kl}^2).$$

Since Hypotheses (1)–(3) are satisfied, Theorem II.9 shows that $\psi_{ij}(x_{ij})$, $\psi_{ijk}(x_{ij}, \xi)$, and $\psi_{ij,kl}(x_{ij}, x_{kl})$ fall off exponentially. It is therefore sufficient to consider only the case $\alpha = \beta = 0$. Also, it suffices to consider only $\delta = 0$, because $\rho_{ij,k}^{-1} P_{ij} U_{ik}$ commutes with multiplication by functions of ξ and ζ , $\rho_{ijk}^{-1} P_{ijk} U_{kl}$ commutes with multiplication by functions of ζ , and $\rho_{ij,kl}^{-1} P_{ij,kl} U_{ik}$ commutes with multiplication by functions of η .

$$\begin{aligned} \rho_{ij,k}^{-1} P_{ij} U_{ik} &= P_{ij} U_{ik} \rho_{ij,k}^{-1} \\ &= [P_{ij} (1 + x_{ij}^2)^{\delta/2} (1 - \Delta_{ij})^{1/2}] [(1 - \Delta_{ij})^{-1/2} U_{ik} (1 + x_{ik}^2)^{\delta/2}] \\ & \quad \times [(1 + x_{ik}^2)^{-\delta/2} (1 + x_{ij}^2)^{-\delta/2} \rho_{ij,k}^{-1}]. \end{aligned}$$

The first factor is bounded because $(1 + x_{ij}^2)^{\delta/2} \psi_{ij}$ belongs to the quadratic form domain of $-\Delta_{ij}$ (Theorem II.9). The third factor is bounded because of inequality (V.2).

The second factor is bounded if its absolute square, $(1 + x_{ik}^2)^{\delta/2} \bar{U}_{ik} (1 - \Delta_{ij})^{-1} U_{ik} (1 + x_{ik}^2)^{\delta/2}$, is bounded. To bound the absolute square, we write it as a

Laplace transform:

$$\int_0^\infty \overline{F(x_{ik})} e^{t\Delta_\eta} F(x_{ik}) e^{-t} dt,$$

where $F(x_{ik}) = (1 + x_{ik}^2)^{\delta/2} U_{ik}(x_{ik})$.

Suppose $\phi \in L^2(\mathbf{R}^{3m})$. To compute $\|F(x_{ik})e^{t\Delta_\eta}F(x_{ik})\phi\|$, we use coordinates x_{ij} , ξ , ζ , and first compute the L^2 -norm in the x_{ij} variable with ξ and ζ fixed. The operator $e^{t\Delta_\eta}$ has an explicitly known integral kernel [22] and $x_{ij} = c_1 x_{ij} + c_2 \xi$, with $c_1 \neq 0$, since all masses are finite: $F(x_{ik}) \in L^p(\mathbf{R}^m, dx_{ik}) + L^\infty(\mathbf{R}^m, dx_{ik})$, for some $p > m$. So, for each fixed ξ and ζ , $F(c_1 x_{ij} + c_2 \xi)\phi(x_{ij}, \xi, \zeta)$ belongs to $L^r(\mathbf{R}^m, dx_{ij}) + L^2(\mathbf{R}^m, dx_{ij})$, where $r^{-1} = p^{-1} + 1/2$. If $t > 0$, $e^{t\Delta_\eta}$ maps this into $L^2(\mathbf{R}^m, dx_{ij}) \cap L^\infty(\mathbf{R}^m, dx_{ij})$ (where we have used the fact that $r < 2$). Multiplication by $\overline{F(c_1 x_{ij} + c_2 \xi)}$ maps this into $L^2(\mathbf{R}^m, dx_{ij})$. By keeping track of the norms at each stage, we obtain:

$$\begin{aligned} & \int |e^{-t}(F(x_{ik})e^{t\Delta_\eta}F(x_{ik})\phi)(x_{ij}, \xi, \zeta)|^2 dx_{ij} \\ & \leq C_F f(t) \int |\phi(x_{ij}, \xi, \zeta)|^2 dx_{ij}, \end{aligned}$$

where C_F is a constant depending only on F and $\int_0^\infty (f(t))^{1/2} dt$ is finite.

Integrating over ξ and ζ , we see that

$$\| \overline{F(x_{ik})} e^{t\Delta_\eta} F(x_{ik}) e^{-t} \phi \| \leq C_F (f(t))^{1/2} \|\phi\|,$$

which belongs to $L^1((0, \infty), dt)$. The Laplace transform now yields part (a) of the lemma.

Parts (b) and (c) are proved by using the same methods, with inequalities (V.3) and (V.4) in place of (V.2). \square

COROLLARY V.5. *Let i, j, k , and l be distinct indices. Let $\delta \in \mathbf{R}$.*

(a) $\rho_{ij,k}^{-1}$, ρ_{ijk}^{-1} , and $\rho_{ij,kl}^{-1}$ are bounded from $L_{\delta+\gamma}^2(\mathbf{R}^{3m})$ to $L_\delta^2(\mathbf{R}^{3m})$.

(b) Assume Hypotheses (1)–(3), and (5), and let n equal j, k , or l .

For any α and β , the following operators are bounded:

$$\begin{aligned} W_{in} P_{ij} &: L_{-\beta}^2(\mathbf{R}^m, dx_{ij}) \otimes L_\delta^2(\mathbf{R}^{2m}, d\xi d\zeta) \rightarrow L_\alpha^2(\mathbf{R}^m) \otimes L_\delta^2(\mathbf{R}^{2m}, d\xi d\zeta), \\ W_{in} P_{ijk} &: L_{-\beta}^2(\mathbf{R}^{2m}, dx_{ij} d\xi) \otimes L_\delta^2(\mathbf{R}^m, d\zeta) \rightarrow L_\alpha^2(\mathbf{R}^{2m}, dx_{ij} d\xi) \otimes L_\delta^2(\mathbf{R}^m, d\zeta), \\ W_{in} P_{ij,kl} &: L_{-\beta}^2(\mathbf{R}^{2m}, dx_{ij} dx_{kl}) \otimes L_\delta^2(\mathbf{R}^m, d\eta) \rightarrow L_\alpha^2(\mathbf{R}^{2m}, dx_{ij} dx_{kl}) \otimes L_\delta^2(\mathbf{R}^m, d\eta). \end{aligned}$$

PROOF. Part (a) follows immediately from the definitions of the ρ 's and L_δ^2 -spaces.

If $n \neq j$, then Lemma V.4 yields the boundedness of $W_{in} P_{ij} = W_{in} P_{ij} \rho_{ij,n}^{-1} \rho_{ij,n}$ on the weighted spaces, because $\rho_{ij,n}$ is bounded. If $n = j$, we use the fact that the operator domain of W_{ij} contains the operator domain of $H_0^{1/2}$ (see §II.B). Theorem II.9 shows that $e^{a|x_\eta|} \psi_{ij}$ belongs to the domain of $H_0^{1/2}$, so $W_{ij} P_{ij}$ is bounded on the given weighted spaces.

The same methods apply to the remaining operators of part (b). \square

DEFINITION. Suppose $A(z): L_\alpha^2(\mathbf{R}^n) \rightarrow L_\beta^2(\mathbf{R}^n)$ is an operator valued function, which has the following properties:

- (i) $A(z)$ is analytic for $z \in \mathbf{C} \setminus \sigma_{\text{ess}}(H)$.
- (ii) $A(z)$ is uniformly bounded for $\text{Re } z \leq C$.
- (iii) $\|A(z)\| \rightarrow 0$ as $\text{Re } z \rightarrow -\infty$.
- (iv) $A(z)$ has norm continuous extensions $A(x \pm i0)$ to $\sigma_{\text{ess}}(H)$ from above and below ($A(x_0 + i0) = A(x_0 - i0)$ if $x_0 = \inf \sigma_{\text{ess}}(H)$). Then $A(z)$ is called *well behaved* from $L_\alpha^2(\mathbf{R}^n)$ to $L_\beta^2(\mathbf{R}^n)$. If $\alpha = \beta = 0$, then we say $A(z)$ is *well behaved*.

LEMMA V.6. Assume Hypotheses (1)–(3), and (5). Let i, j, k and l be distinct indices. Let D_1 and D_2 be cluster decompositions. The following are well behaved:

$$\begin{aligned} K_1(z) &= W_{ij}(z - H_0)^{-1}U_{ik}, & K_2(z) &= W_{ij}(z - H_{kl})^{-1}U_{ik}, \\ K_3(z) &= W_{ij}P_{D_1}(z - H_0)^{-1}U_{ik}, & K_4(z) &= W_{ij}P_{D_1}(z - H_{kl})^{-1}U_{ik}, \\ K_5(z) &= W_{ij}(z - H_0)^{-1}P_{D_2}U_{ik}, & K_6(z) &= W_{ij}(z - H_{kl})^{-1}P_{D_2}U_{ik}, \\ K_7(z) &= W_{ij}P_{D_1}(z - H_0)^{-1}P_{D_2}U_{ik}, & \text{and } K_8(z) &= W_{ij}P_{D_1}(z - H_{kl})^{-1}P_{D_2}U_{ik}. \end{aligned}$$

PROOF. Lemma II.3 yields the result for K_1 and K_2 . If P_{D_1} commutes with W_{ij} , then Lemma II.3 yields the result for K_3 and K_4 , also. If i and j belong to the same cluster of D_1 , then Corollary V.5 shows that $W_{ij}P_{D_1}(1 + x_{ij}^2)$ is bounded. However, Lemma II.3 applies to $(1 + x_{ij}^2)^{-1}(z - H_0)^{-1}U_{ik}$ and $(1 + x_{ij}^2)^{-1}(z - H_{kl})^{-1}U_{ik}$. So, it suffices to consider K_3 and K_4 when i and j belong to different clusters of D_1 , but W_{ij} and P_{D_1} do not commute. In that case, Lemma V.4 shows that the adjoint of $W_{ij}P_{D_1}\rho^{-1}$ is bounded into some tensor product of weighted spaces. Thus, $W_{ij}P_{D_1}\rho^{-1}$ is bounded. Lemma II.3 shows that $\rho P_{D_1}(z - H_0)^{-1}U_{ik}$ and $\rho P_{D_1}(z - H_{kl})^{-1}U_{ik}$ are well behaved. So, K_3 and K_4 are well behaved. Of course, the correct choice of ρ must be made.

The same methods control K_5, K_6, K_7 , and K_8 . \square

LEMMA V.7. Assume Hypotheses (1)–(3). Suppose $N \geq 3$, and α is any pair. Then $K(z) = W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1}U_\alpha$ is well behaved.

PROOF. We may decompose \mathcal{H} as a direct integral of Hilbert spaces for the two body problem determined by the pair α . $K(z)$ is fibered under this decomposition, and it suffices to control the operators on the individual fibers. So, we need only show $W_\alpha(1 - P_\alpha)(z - h_\alpha)^{-1}U_\alpha$ is well behaved on $L^2(\mathbf{R}^m)$:

$$\begin{aligned} W_\alpha(1 - P_\alpha)(z - h_\alpha)^{-1}U_\alpha &= W_\alpha(1 - P_\alpha)(z - h_0)^{-1}U_\alpha \\ &+ W_\alpha(1 - P_\alpha)(z - h_0)^{-1}U_\alpha(1 - W_\alpha(z - h_0)^{-1}U_\alpha)^{-1}W_\alpha(z - h_0)^{-1}U_\alpha. \end{aligned}$$

In view of Lemma V.6, it suffices to show that the factor of $(1 - P_\alpha)$ cancels the singularities of $(1 - W_\alpha(z - h_0)^{-1}U_\alpha)^{-1}$.

Let $M(z) = W_\alpha(z - h_0)^{-1}U_\alpha$. The proof of Proposition V.1 shows that $(1 - M(z))^{-1}$ is singular when z is a real negative eigenvalue of h_α , and possibly when $z = 0$. Hypothesis (3) implies that it is not singular at $z = 0$.

The poles of $(1 - M(z))^{-1}$ are first order poles. To prove this, let z_0 be an eigenvalue of h_α , and consider the Laurent series

$$(1 - M(z))^{-1} = \sum_{k=-n}^{\infty} (z - z_0)^k L_k,$$

where $L_{-n} \neq 0$. Since $L_{-n} \neq 0$, there is a vector ϕ in the domain of U_α , such that $L_{-n}\phi \neq 0$. $[(1 - M(z))^{-1} - 1]\phi = (1 - M(z))^{-1}M(z)\phi$ must have a pole of order n , so $\psi = M(z_0)\phi = C(z_0)(U_\alpha\phi)$ is a nonzero element of $\text{Ran } C(z)$ with $L_{-n}\psi \neq 0$. Similarly, using

$$[(1 - M(z))^{-1} - 1] = M(z)(1 - M(z))^{-1},$$

we see that every nonzero $\eta \in \text{Ran } L_n$ has $A(z_0)\eta \neq 0$. Therefore,

$$(z - h_\alpha)^{-1}(U_\alpha\phi) = (z - h_0)^{-1}(U_\alpha\phi) + A(z)(1 - M(z))^{-1}C(z)(U\phi)$$

has a pole of order n . Since h_α is selfadjoint, $n = 1$.

If $Q = L_{-1}$ is the residue at z_0 , then Q is a projection onto solutions of $M(z_0)\phi = \phi$ (see Kato [14]). If P is the orthogonal projection onto all eigenvectors of h_α with eigenvalue z_0 , then $PA(z_0) = A(z_0)Q$. (This is proved in the proof of Proposition V.1.) Therefore, for z near z_0 ,

$$\begin{aligned} (1 - P)A(z)(1 - M(z))^{-1} &= (1 - P)A(z_0)(1 - M(z))^{-1} - (z - z_0)(1 - P) \\ &\quad \times (z - h_0)^{-1}A(z_0)(1 - M(z))^{-1} \\ &= A(z_0)(1 - Q) \sum_{k=-1}^{\infty} (z - z_0)^k L_k - (1 - P) \\ &\quad \times (z - h_0)^{-1}A(z_0) \sum_{k=-1}^{\infty} (z - z_0)^{k+1} L_k \\ &= A(z_0)(1 - Q) \sum_{k=0}^{\infty} (z - z_0)^k L_k - (1 - P) \\ &\quad \times (z - h_0)^{-1}A(z_0) \sum_{k=-1}^{\infty} (z - z_0)^{k+1} L_k. \end{aligned}$$

Thus, $(1 - P)A(z)(1 - M(z))^{-1}$ has a removable singularity at z_0 . Since $(1 - P_\alpha) = (1 - P_\alpha)(1 - P)$, the factor $(1 - P_\alpha)$ cancels the singularities of $(1 - M(z))^{-1}$ and the lemma is proved. \square

LEMMA V.8. Assume Hypotheses (1)–(3), and (5). Let α , β , and δ be pairs such that α and δ have at least one index in common. Then $K_1(z) = W_\alpha(1 - P_\beta)(z - H_\beta)^{-1}U_\delta$ is well behaved, and $K_2(z) = W_\alpha(1 - P_\beta)(z - H_\beta)^{-1}$ is well behaved from $L^2_\gamma(\mathbb{R}^{(N-1)m})$ to $L^2(\mathbb{R}^{(N-1)m})$.

PROOF. Consider only K_1 ; proof for K_2 is similar.

If β has no index in common with α , or if β has no index in common with δ , then Lemma V.6 applies. Thus, we may assume β has indices in common with α and δ :

$$\begin{aligned}
 & W_\alpha(1 - P_\beta)(z - H_\beta)^{-1}U_\delta \\
 &= W_\alpha(1 - P_\beta)(z - H_0)^{-1}U_\delta + W_\alpha(1 - P_\beta)(z - H_\beta)^{-1}V_\beta(z - H_0)^{-1}U_\delta \\
 &= W_\alpha(1 - P_\beta)(z - H_0)^{-1}U_\delta + W_\alpha(z - H_\beta)^{-1}(1 - P_\beta)V_\beta(z - H_0)^{-1}U_\delta \\
 &= W_\alpha(1 - P_\beta)(z - H_0)^{-1}U_\delta \\
 &\quad + (W_\alpha(z - H_0)^{-1}(1 - P_\beta)U_\beta)(W_\beta(z - H_0)^{-1}U_\delta) \\
 &\quad + (W_\alpha(z - H_0)^{-1}U_\beta)(W_\beta(1 - P_\beta)(z - H_\beta)^{-1}U_\beta)(W_\beta(z - H_0)^{-1}U_\delta).
 \end{aligned}$$

Apply Lemmas V.6 and V.7. \square

LEMMA V.9. Assume Hypotheses (1)–(3), and (5). Let α , β , δ , and σ be pairs with $\alpha \neq \beta$ and $\beta \neq \sigma$. If α and β have no index in common, assume $\beta = \delta$ or that δ and σ have an index in common. Let D be any cluster decomposition, such that the indices of α belong to the same cluster and the indices of β belong to the same cluster. The following are well behaved:

$$\begin{aligned}
 K_1(z) &= W_\alpha(z - H_0)^{-1}V_\beta(z - H_0)^{-1}U_\sigma, \\
 K_2(z) &= W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1}V_\beta(z - H_0)^{-1}U_\sigma, \\
 K_3(z) &= W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1}V_\beta(1 - P_\delta)(z - H_\delta)^{-1}U_\sigma, \\
 K_4(z) &= W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1}V_\beta P_D(z - H_0)^{-1}U_\sigma, \\
 K_5(z) &= W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1}V_\beta P_D(1 - P_\delta)(z - H_\delta)^{-1}U_\sigma.
 \end{aligned}$$

PROOF (SEE IORIO AND O'CARROL [13]). Lemma II.3 shows that K_1 is well behaved if it has norm continuous boundary values. If α and β have an index in common and β and σ have an index in common, then Lemma II.3 shows K_1 is well behaved.

If α and β , or β and σ have no index in common, then, by taking adjoints if necessary, we may assume β and σ have no index in common. Then, for $\text{Im } z > 0$, $\text{Im } w > 0$,

$$\begin{aligned}
 & |\langle \phi(K_1(z) - K_1(w))\psi \rangle| \\
 &= \left| \int_0^\infty \int_0^\infty \langle \phi, W_\alpha e^{-itH_0} V_\beta e^{-isH_0} U_\sigma \psi \rangle (e^{i(t+s)z} - e^{i(t+s)w}) ds dt \right| \\
 &\leq \int_0^\infty \int_0^\infty |\langle \phi, W_\alpha e^{-itH_0} e^{-is(H_0 - H_\beta^\beta)} U_\sigma V_\beta e^{-isH_\beta^\beta} \psi \rangle| |e^{i(t+s)z} - e^{i(t+s)w}| ds dt \\
 &= \int_0^\infty \int_0^\infty |\langle U_\beta^* e^{itH_\beta^\beta} (U_\sigma^* e^{i(t+s)(H_0 - H_\beta^\beta)} W_\alpha^*) \phi, W_\beta e^{-isH_\beta^\beta} \psi \rangle| \\
 &\quad \times |e^{i(t+s)z} - e^{i(t+s)w}| ds dt.
 \end{aligned}$$

Change variables by replacing s with $r = t + s$.

$$\begin{aligned}
& |\langle \phi, (K_1(z) - K_1(w))\psi \rangle| \\
& \leq \int_0^\infty \int_0^\infty |\langle U_\beta^* e^{itH_\beta^\delta} (U_\sigma^* e^{ir(H_0 - H_\beta^\delta)} W_\alpha^*) \phi, W_\beta e^{-i(r-t)H_\beta^\delta} \psi \rangle| |e^{irz} - e^{irw}| dt dr \\
& \leq \int_0^\infty \int_0^\infty \|U_\beta^* e^{itH_\beta^\delta} (U_\sigma^* e^{ir(H_0 - H_\beta^\delta)} W_\alpha^*) \phi\| \\
& \quad \times \|W_\beta e^{-i(r-t)H_\beta^\delta} \psi\| |e^{irz} - e^{irw}| dt dr \\
& \leq \int_0^\infty \left(\int_0^\infty \|U_\beta^* e^{itH_\beta^\delta} (U_\sigma^* e^{ir(H_0 - H_\beta^\delta)} W_\alpha^*) \phi\|^2 dt \right)^{1/2} \\
& \quad \times \left(\int_0^\infty \|W_\beta e^{-i(r-t)H_\beta^\delta} \psi\|^2 dt \right)^{1/2} |e^{irz} - e^{irw}| dr \\
& \leq \int_0^\infty C \|U_\sigma^* e^{ir(H_0 - H_\beta^\delta)} W_\alpha \phi\| \|\psi\| |e^{irz} - e^{irw}| dr.
\end{aligned}$$

We have used the Schwarz inequality, Hölder inequality, and Kato smoothness of U_β^* and W_β with respect to H_0^β (see Lemma II.2).

Since $\alpha \neq \beta$ and β has no index in common with σ , α and σ have an index in common. Therefore, as in the proof of Lemma II.3, $\|U_\sigma^* e^{ir(H_0 - H_\beta^\delta)} W_\alpha^* \phi\|$ is dominated by an L^1 function of r which depends only on $\|\phi\|$. So, as in the proof of Lemma II.3, $|\langle \phi, (K_1(z) - K_1(w))\psi \rangle|$ tends to zero uniformly in ϕ and ψ as z approaches w .

$$\begin{aligned}
K_2(z) &= W_\alpha(1 - P_\alpha)(z - H_0)^{-1} V_\beta(z - H_0)^{-1} U_\sigma \\
&\quad + W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1} U_\alpha K_1(z).
\end{aligned}$$

The first term is controlled by the methods used for $K_1(z)$. The result for $K_1(z)$ and Lemma V.7 control the second term.

If β has indices in common with α and σ , then Lemma V.8 shows that $K_3(z)$ is well behaved. If β and σ have an index in common, but α and β do not, then there are two cases:

Case 1. $\beta = \delta$:

$$\begin{aligned}
K_3(z) &= W_\alpha(1 - P_\alpha)(z - H_0)^{-1} V_\beta(z - H_\beta)^{-1} (1 - P_\beta) U_\sigma \\
&\quad + (W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1} U_\alpha) \\
&\quad \times (W_\alpha(z - H_0)^{-1} V_\beta(z - H_\beta)^{-1} (1 - P_\beta) U_\sigma) \\
&= W_\alpha(1 - P_\alpha) [(z - H_\beta)^{-1} - (z - H_0)^{-1}] (1 - P_\beta) U_\sigma \\
&\quad + (W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1} U_\alpha) \\
&\quad \times (W_\alpha [(z - H_\beta)^{-1} - (z - H_0)^{-1}] (1 - P_\beta) U_\sigma).
\end{aligned}$$

Since $\beta \neq \sigma$ and α has no index in common with β , α and σ have an index in common. Lemmas V.6 and V.8 therefore apply to the terms which appear in the last expression.

Case 2. $\delta \neq \beta$:

$$K_3(z) = W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1}V_\beta(z - H_0)^{-1}(1 - P_\delta)U_\sigma \\ + (W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1}V_\beta(z - H_0)^{-1}U_\delta)(W_\delta(z - H_\delta)^{-1}(1 - P_\delta)U_\sigma).$$

The methods used to control $K_2(z)$ apply to the first term. The first factor of the second term has the same form as $K_2(z)$. Lemma V.8 applies to the final factor, since δ and σ are assumed to have an index in common.

If β has no indices in common with σ , then there are again two cases.

Case 3. $\beta = \delta$: Since β and σ have no common index and $\alpha \neq \beta$, α and σ must have a common index. Apply the methods of Case 1.

Case 4. $\beta \neq \delta$: Since β and σ have no index in common and $\beta \neq \delta$, δ and σ have a common index. Apply the methods of Case 2.

To show $K_4(z)$ and $K_5(z)$ are well behaved, first notice that $(1 + x_\alpha^2 + x_\beta^2)^{\delta/2}W_\beta P_D(1 + x_\alpha^2 + x_\beta^2)^{\delta/2}$ is bounded (by Corollary V.5). Moreover, σ must have an index in common with α or β . Using this information, $K_4(z)$ and $K_5(z)$ may be rewritten as products of factors which may be controlled by the methods of Lemmas V.6 and V.8, when U_β is bounded. A density argument then shows that the result holds for general U_β . \square

REMARK. Let ξ be the coordinate from the center of mass of particles i and j to particle k . With the inner product on \mathbf{R}^{3m} determined by the masses, ξ and x_{ij} are orthogonal. However, ξ is not orthogonal to any other x_α .

LEMMA V.10. Assume Hypotheses (1)–(3), and (5). Let i, j , and k be distinct indices, and choose pairs $\alpha \neq (i, j)$ and $\beta \neq (i, j)$, so that α and β have at least one common index. The following are well behaved:

$$K_1(z) = W_\alpha(z - H_{ij})^{-1}P_{ij}\rho_{ij,k}$$

and

$$K_2(z) = W_\alpha(z - H_{ij})^{-1}P_{ij}U_\beta.$$

PROOF. Notice first that $(z - H_{ij})^{-1}P_{ij}$ equals P_{ij} times a multiplication operator in the momentum representation. This fact, together with the above remark, show that $K_1(z)$ may be controlled by the methods of Lemma II.3 or Lemma V.6. These methods apply to $K_2(z)$ if β has no index in common with (i, j) .

If β has an index in common with (i, j) , then we may assume $\beta = (i, k)$. $K_2(z)$ may then be written as $K_2(z) = K_1(z)\rho_{ij,k}^{-1}P_{ij}U_{ik}$. Since $K_1(z)$ is well behaved, Lemma V.4 shows that $K_2(z)$ is well behaved. \square

LEMMA V.11. Assume Hypotheses (1)–(3), and (5). Let i, j , and k be distinct indices. Let $\alpha = (i, j)$, (i, k) , or (j, k) . Choose l such that $\alpha \cup \{l\} = \{i, j, k\}$. The following are well behaved:

$$\begin{aligned} K_1(z) &= \rho_{ij,k}^{-1} P_{ij} V_{ik} (z - H_\alpha)^{-1} P_\alpha \rho_{\alpha,l}, \\ K_2(z) &= \rho_{ij,k}^{-1} P_{ij} V_{ik} (1 - P_{ijk}) (z - H_\alpha)^{-1} P_\alpha \rho_{\alpha,l}, \\ K_3(z) &= \rho_{ij,k}^{-1} P_{ij} V_{ik} (z - H_0)^{-1} P_\alpha \rho_{\alpha,l}, \\ K_4(z) &= \rho_{ij,k}^{-1} P_{ij} V_{ik} (1 - P_{ijk}) (z - H_0)^{-1} P_\alpha \rho_{\alpha,l}. \end{aligned}$$

PROOF. If $\alpha \neq (i, k)$, then

$$K_1(z) = (\rho_{ij,k}^{-1} P_{ij} U_{ik}) (W_{ik} (z - H_\alpha)^{-1} P_\alpha \rho_{\alpha,l}).$$

Apply Lemmas V.4 and V.10.

If $\alpha = (i, k)$, then $(z - H_\alpha)^{-1} P_\alpha$ equals P_α times a multiplication operator in the momentum representation. This operator $(z - K_\alpha - E_\alpha)^{-1}$ commutes with W_{ik} , as does $\rho_{\alpha,l}$. Therefore,

$$K_1(z) = (\rho_{ij,k}^{-1} P_{ij} U_{ik} (1 + x_{ij}^2)^{\gamma/2}) ((1 + x_{ij}^2)^{-\gamma/2} (z - K_\alpha - E_\alpha)^{-1} \rho_{\alpha,l}) (W_{ik} P_\alpha).$$

Lemma V.4 bounds the first factor. The proof of Lemma II.3 controls the second. Corollary V.5 bounds the final factor.

$$\begin{aligned} K_2(z) &= K_1(z) - \rho_{ij,k}^{-1} P_{ij} V_{ik} P_{ijk} (z - H_\alpha)^{-1} P_\alpha \rho_{\alpha,l} \\ &= K_1(z) - (\rho_{ij,k}^{-1} P_{ij} U_{ik}) (W_{ik} P_{ijk} \rho_{\alpha,l}^{-1}) (\rho_{\alpha,l} (z - H_\alpha)^{-1} P_\alpha \rho_{\alpha,l}). \end{aligned}$$

Lemma V.4 bounds the first factor; Corollary V.5 bounds the second; and the proof of Lemma II.3 controls the third factor.

Proofs for $K_3(z)$ and $K_4(z)$ are similar. \square

LEMMA V.12. Assume Hypotheses (1)–(3), and (5). Let α, β, δ , and σ be pairs, with $\alpha \neq \beta$ and $\beta \neq \sigma$. Let k be any index not contained in the pair δ . Let D be any cluster decomposition, such that the indices of α belong to the same cluster and the indices of β belong to the same cluster. The following are well behaved:

$$\begin{aligned} K_1(z) &= W_\alpha (1 - P_\alpha) (z - H_\alpha)^{-1} V_\beta (z - H_\delta)^{-1} P_\delta \rho_{\delta,k}, \\ K_2(z) &= W_\alpha (1 - P_\alpha) (z - H_\alpha)^{-1} V_\beta (z - H_\delta)^{-1} P_\delta U_\sigma, \\ K_3(z) &= W_\alpha (1 - P_\alpha) (z - H_\alpha)^{-1} V_\beta (1 - P_D) (z - H_\delta)^{-1} P_\delta \rho_{\delta,k}, \\ K_4(z) &= W_\alpha (1 - P_\alpha) (z - H_\alpha)^{-1} V_\beta (1 - P_D) (z - H_\delta)^{-1} P_\delta U_\sigma, \\ K_5(z) &= W_\alpha (1 - P_\alpha) (z - H_\alpha)^{-1} V_\beta (1 - P_D) (z - H_0)^{-1} P_\delta \rho_{\delta,k}. \end{aligned}$$

PROOF. We consider only $K_1(z)$. The others may be treated in a similar fashion. There are three cases.

Case 1. $\beta = \delta$:

$$\begin{aligned} K_1(z) &= W_\alpha(1 - P_\alpha)(z - H_0)^{-1}V_\beta(z - H_\beta)^{-1}P_\beta\rho_{\beta,k} \\ &\quad + (W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1}U_\alpha)(W_\alpha(z - H_0)^{-1}V_\beta(z - H_\beta)^{-1}P_\beta\rho_{\beta,k}) \\ &= W_\alpha(1 - P_\alpha)[(z - H_\beta)^{-1} - (z - H_0)^{-1}]P_\beta\rho_{\beta,k} \\ &\quad + (W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1}U_\alpha)(W_\alpha[(z - H_\beta)^{-1} - (z - H_0)^{-1}]P_\beta\rho_{\beta,k}). \end{aligned}$$

Apply Lemmas II.3, V.8, V.10, and the remark preceding Lemma V.10.

Case 2. $\beta \neq \delta$; α and β have a common index.

$$K_1(z) = (W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1}U_\beta)(W_\beta(z - H_\delta)^{-1}P_\delta\rho_{\delta,k}).$$

Apply Lemmas V.8 and V.10.

Case 3. $\beta \neq \delta$; α and β have no common index.

$$\begin{aligned} K_1(z) &= W_\alpha(1 - P_\alpha)(z - H_0)^{-1}V_\beta(z - H_\delta)^{-1}P_\delta\rho_{\delta,k} \\ &\quad + (W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1}U_\alpha)(W_\alpha(z - H_0)^{-1}V_\beta(z - H_\delta)^{-1}P_\delta\rho_{\delta,k}). \end{aligned}$$

Proceeding as in Case 1, all the results may be proved, except for norm continuity of the boundary values. However, a density argument shows that it suffices to prove norm continuity under the additional assumption that V_β is bounded.

In that case,

$$\begin{aligned} K_1(z) &= (W_\alpha(1 - P_\alpha)(z - H_\alpha)^{-1}(1 + x_\delta^2)^{-\gamma/2}) \\ &\quad \times (V_\beta(1 + x_\delta^2)^{\gamma/2}P_\delta(z - H_\delta)^{-1}\rho_{\delta,l}). \end{aligned}$$

Since α and β have no common index and $\beta \neq \delta$, α and δ have a common index. So, Lemma V.6 applies to the first factor. Lemma V.10 and Theorem II.9 show that the second factor is well behaved. \square

LEMMA V.13. Assume Hypotheses (1)–(3). If $N = 3$, then $M(z)$ is well behaved and $(M(z))^2$ is compact.

PROOF. $M(z)$ is well behaved by Lemmas V.4, V.8, V.11, and V.12. $W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}U_{ik}$ is a norm convergent sum of connected graphs, when $\operatorname{Re} z$ is large negative. Standard methods (e.g. Ginibre and Moulin [7]) show that each term is compact. Similarly,

$$W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{ik}(z - H_{ik})^{-1}P_{ik}\rho_{ik,j}$$

and

$$\rho_{ij,k}^{-1}P_{ij}V_{ik}(z - H_{ik})^{-1}P_{ik}\rho_{ik,j}$$

are compact for large negative $\operatorname{Re} z$. Since these operator valued functions are well behaved, compactness holds for all z .

The entries of $(M(z))^2$ are sums of terms, each of which contains one of these compact factors. Thus $(M(z))^2$ is compact. \square

LEMMA V.14. Assume Hypotheses (1)–(3), and (5). Let i, j , and k be distinct indices, and let α and β be distinct pairs. The following are well behaved:

$$K_1(z) = \rho_{ij,k}^{-1} P_{ij} V_{ik} (z - H_0)^{-1} U_\beta,$$

$$K_2(z) = \rho_{ij,k}^{-1} P_{ij} V_{ik} (1 - P_{ijk})(z - H_0)^{-1} U_\beta,$$

$$K_3(z) = \rho_{ij,k}^{-1} P_{ij} V_{ik} (1 - P_\alpha)(z - H_\alpha)^{-1} U_\beta,$$

$$K_4(z) = \rho_{ij,k}^{-1} P_{ij} V_{ik} P_\alpha (z - H_\alpha)^{-1} U_\beta,$$

$$K_5(z) = \rho_{ij,k}^{-1} P_{ij} V_{ik} (1 - P_{ijk})(1 - P_\alpha)(z - H_\alpha)^{-1} U_\beta,$$

$$K_6(z) = \rho_{ij,k}^{-1} P_{ij} V_{ik} (1 - P_{ijk}) P_\alpha (z - H_\alpha)^{-1} U_\beta.$$

PROOF. If β and (i, k) have an index in common, $K_1(z)$ is well behaved by Lemmas V.4 and V.6. If β and (i, k) have no common index, then Lemmas II.3 and V.4 yield the results, except for the norm continuity of the boundary values. Given this, we need only prove the norm continuity when V_{ik} is bounded. In that case,

$$K_1(z) = (\rho_{ij,k}^{-1} P_{ij} V_{ik} (1 + x_{ij}^2)^{\gamma/2}) ((1 + x_{ij}^2)^{-\gamma/2} (z - H_0)^{-1} U_\beta).$$

Since β and (i, j) must have a common index, Lemma II.3 controls the boundary values of the second factor. Lemma V.4 bounds the first.

Proofs for the other operators are similar. \square

PROPOSITION V.15. Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct indices. Let $n = i, j$, or k . Then

$$K_1(z) = W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} U_n$$

is well behaved and compact. Moreover

$$K_2(z) = W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1}$$

is compact and well behaved from $L_\delta^2(\mathbb{R}^{3m})$ to $L^2(\mathbb{R}^{3m})$, when $\delta > 1$.

PROOF. Consider only $K_1(z)$; proof for $K_2(z)$ is similar.

Using equation (IV.6), we have $K_1(z) = \text{I}(z) + \text{II}(z) + \text{III}(z)$, where

$$\text{I}(z) = W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_0)^{-1} U_n,$$

$$\text{II}(z) = W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})A(z)C(z)U_n,$$

$$\begin{aligned} \text{III}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})A(z)(1 - M(z))^{-1} \\ &\quad \times M(z)C(z)U_n. \end{aligned}$$

Lemma V.9 shows $\text{I}(z)$ is well behaved.

$A(z)C(z)$ is a sum of terms, each of which contains only one resolvent. We substitute this sum in $\text{II}(z)$. $\text{II}(z)$ then becomes a sum of terms which are well behaved by Lemmas V.9 and V.12.

$W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})A(z)$ is well behaved by Lemmas V.9 and V.12. $M(z)$ is well behaved by Lemmas V.4, V.8, V.11, and V.12. $M(z)C(z)U_n$

is well behaved by Lemmas V.9, V.12, and V.14. So, $\text{III}(z)$ is well behaved if the factor $(1 - P_{ijk})$ in the operator

$$W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})A(z)$$

cancels the singularities of $(1 - M(z))^{-1}$.

To prove this, we mimic part of the proof of Lemma V.7. \mathcal{H} is a direct integral of spaces for the three body problem determined by (i, j, k) . $A(z)$, $M(z)$, and $C(z)$ are fibered under this decomposition, and we consider their restrictions (which we also denote by $A(z)$, $M(z)$, and $C(z)$) to the individual fibers.

Let z_0 be a point at which $(1 - M(z))^{-1}$ is singular on some fixed fiber. Hypothesis (3) and Lemma II.11 show that $(1 - M(z, \theta))^{-1}$ is singular at z_0 for some θ with $z_0 \notin \sigma_{\text{ess}}(H(\theta))$ and $|\text{Im } \theta|$ arbitrarily close to 0. $(1 - M(z, \theta))^{-1}$ may be written as a Laurent series $\sum_{k=-n}^{\infty} (z - z_0)^k L_k$ for z near z_0 . Mimicking the proof of Lemma V.7, we see that $L_{-n}C(z_0, \theta) \neq 0$ and that no $\phi \in \text{Ran } L_{-n}$ has $M(z_0, \theta)\phi = 0$. By using $G_\alpha = G_0 + G_\alpha V_\alpha G_0$, we see that $A(z_0, \theta)\psi = 0$ implies $M(z_0, \theta)\psi = 0$. Putting these facts together, $A(z_0, \theta)L_{-n}C(z_0, \theta) \neq 0$. Equation (IV.6) shows that $(z - H(\theta))^{-1}$ has a pole of order n at z_0 . Selfadjointness of H and standard dilation analyticity arguments show $n = 1$. The rest of the argument of Lemma V.7 now goes through, since equation (V.1) shows $(1 - P)A(z_0) = A(z_0)(1 - Q)$, where Q projects onto the eigenvalue 1 eigenvectors of $M(z_0)$. Thus $(1 - P_{ijk})$ cancels the singularities of $(1 - M(z))^{-1}$, and $\text{III}(z)$ is well behaved. Therefore, $K_1(z)$ is well behaved.

It suffices to prove compactness at large negative $\text{Re } z$. Write $K_1(z)$ as:

$$\begin{aligned} & W_{ij}(z - H_{ij})^{-1}(V_{ik} + V_{jk})(z - H_{ijk})^{-1}U_{nl} \\ & - W_{ij}P_{ij}(z - H_{ij})^{-1}(V_{ik} + V_{jk})(z - H_{ijk})^{-1}U_{nl} \\ & - W_{ij}(z - H_{ij})^{-1}(V_{ik} + V_{jk})(z - H_{ijk})^{-1}P_{ijk}U_{nl} \\ & + W_{ij}P_{ij}(z - H_{ij})^{-1}(V_{ik} + V_{jk})(z - H_{ijk})^{-1}P_{ijk}U_{nl}. \end{aligned}$$

Expand the resolvent factors as sums of graphs. Replace each $W_{ij}P_{ij}$ by

$$(W_{ij}P_{ij}(1 + x_{ij}^2))(1 + x_{ij}^2)^{-1}$$

and each $P_{ijk}U_{nl}$ by

$$(1 + x_{ij}^2 + x_{ik}^2 + x_{jk}^2)^{-1}((1 + x_{ij}^2 + x_{ik}^2 + x_{jk}^2)P_{ijk}U_{nl}).$$

Then apply Corollary V.5 and any of the usual compactness proofs to each term.

□

PROPOSITION V.16. *Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct indices, and let $n = i, j$, or k .*

$$K_1(z) = \rho_{ij}^{-1}P_{ij}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1}U_{nl}$$

is compact and well behaved.

$$K_2(z) = \rho_{ij,k}^{-1}P_{ij}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1}$$

is compact and well behaved from $L_\delta^2(\mathbf{R}^{3m})$ to $L^2(\mathbf{R}^{3m})$ whenever $\delta > 1$.

PROOF. Consider only $K_1(z)$; the proof for $K_2(z)$ is similar.

Using equation (IV.6), we have $K_1(z) = I(z) + II(z) + III(z)$, where

$$I(z) = \rho_{ij,k}^{-1} P_{ij}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_0)^{-1} U_{nl},$$

$$II(z) = \rho_{ij,k}^{-1} P_{ij}(V_{ik} + V_{jk})(1 - P_{ijk})A(z)C(z)U_{nl},$$

$$III(z) = \rho_{ij,k}^{-1} P_{ij}(V_{ik} + V_{jk})(1 - P_{ijk})A(z)(1 - M(z))^{-1}M(z)C(z)U_{nl}.$$

Lemma V.14 shows that $I(z)$ and $II(z)$ are well behaved. $M(z)$ is well behaved by Lemmas V.4, V.8, V.12, and V.14. $M(z)C(z)U_{nl}$ is well behaved by Lemmas V.9, V.12, and V.14. Lemmas V.11 and V.14 control $\rho_{ij,k}^{-1} P_{ij}(V_{ik} + V_{jk})(1 - P_{ijk})A(z)$.

As in Proposition V.15, $(1 - P_{ijk})$ removes the singularities of $(1 - M(z))^{-1}$. So, $K_1(z)$ is well behaved.

It suffices to prove compactness for large negative $\text{Re } z$. Furthermore, a density argument shows that we may assume all U 's and W 's lie in $\mathfrak{S}(\mathbb{R}^m)$.

Under this assumption,

$$\begin{aligned} & \rho_{ij,k}^{-1} P_{ij}(V_{ik} + V_{jk}) P_{ijk}(z - H_{ijk})^{-1} U_{nl} \\ &= (\rho_{ij,k}^{-1} P_{ij}(V_{ik} + V_{jk})) (P_{ijk}(z - H_{ijk})^{-1} \rho_{ijk}) (\rho_{ijk}^{-1} P_{ijk} U_{nl}). \end{aligned}$$

Lemma V.4 bounds the initial and final factors. The middle factor equals $P_{ijk}(z - K_{ijk} - E_{ijk})^{-1} \rho_{ijk}$, where K_{ijk} is the kinetic energy associated with the coordinate from the center of mass of particles i, j , and k to particle l . P_{ijk} is $(\text{compact}) \otimes 1$. $(z - K_{ijk} - E_{ijk})^{-1} \rho_{ijk}$ is $1 \otimes (\text{compact})$. So, the middle term is compact.

Hence, it suffices to prove $\rho_{ij,k}^{-1} P_{ij}(V_{ik} + V_{jk})(z - H_{ijk})^{-1} U_{nl}$ is compact when the U 's and W 's belong to $\mathfrak{S}(\mathbb{R}^m)$. Lemma V.4 shows that it suffices to prove that $(1 + x_{ij}^2)^{-\alpha} (W_{ik} + W_{jk})(z - H_{ijk})^{-1} U_{nl}$ is compact. We expand $(z - H_{ijk})^{-1}$ as a sum of graphs. The only graphs for which compactness could fail are those which do not connect i and j . It therefore suffices to prove the compactness of $(1 + x_{ij}^2)^{-\alpha} (W_{ik}(z - H_{ik})^{-1} + W_{jk}(z - H_{jk})^{-1}) U_{nl}$.

Expand $(1 + x_{ij}^2)^{-\alpha} W_{ik}(z - H_{ik})^{-1} U_{nl}$ as a sum of graphs. Use coordinates x_{ij} , x_{ik} , and x_{nl} , with conjugate momenta p_{ij} , p_{ik} , and p_{nl} , respectively. Following Reed and Simon [24, p. 128], write out the integral kernel for each graph in momentum representation. If z is real and large negative, then the replacement of each factor $(z - H_0)^{-1}$ by $(z - p_{ij}^2 - p_{ik}^2 - p_{nl}^2)^{-1}$ will increase the absolute value of the integral kernel at every point. The new integral kernel is square integrable, so the original one is, also. Consequently, each graph is Hilbert-Schmidt.

The proof for $(1 + x_{ij}^2)^{-\alpha} W_{jk}(z - H_{jk})^{-1} U_{nl}$ is similar. \square

PROPOSITION V.17. Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct indices; let $n = i, j$, or k ; let p be any index not equal to n or l ; and let α be the pair not containing n or l . The following are well behaved and compact:

$$\begin{aligned} K_1(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\ &\quad \times V_{nl}(z - H_{nl})^{-1} P_{nl} \rho_{nl,p} \end{aligned}$$

and

$$K_2(z) = W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\ \times V_{nl}(z - H_{nl})^{-1}P_{nl}U_\alpha.$$

PROOF. Consider only $K_1(z)$; the proof for $K_2(z)$ is essentially identical.

For large negative $\operatorname{Re} z$, $W_{nl}P_{nl}(z - H_{nl})^{-1}\rho_{nl,p}$ is bounded, so compactness follows from Proposition V.15. Similarly, it is clear that $\|K_1(z)\|$ tends to zero as $\operatorname{Re} z \rightarrow -\infty$. It is therefore sufficient to prove analyticity and the norm continuity of the boundary values.

$K_1(z) = \text{I}(z) + \text{II}(z) + \text{III}(z)$, where

$$\begin{aligned} \text{I}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_0)^{-1} \\ &\quad \times V_{nl}(z - H_{nl})^{-1}P_{nl}\rho_{nl,p}, \\ \text{II}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})A(z)C(z) \\ &\quad \times V_{nl}(z - H_{nl})^{-1}P_{nl}\rho_{nl,p}, \\ \text{III}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})A(z) \\ &\quad \times (1 - M(z))^{-1}M(z)C(z)V_{nl}(z - H_{nl})^{-1}P_{nl}\rho_{nl,p}. \end{aligned}$$

Lemma V.12 controls $\text{I}(z)$ after the factor $(z - H_0)^{-1}V_{nl}(z - H_{nl})^{-1}$ is replaced by $[(z - H_{nl})^{-1} - (z - H_0)^{-1}]$.

Since

$$\begin{aligned} A(z)C(z) &= (z - H_{ij})^{-1} + (z - H_{ik})^{-1} + (z - H_{jk})^{-1} - 3(z - H_0)^{-1} \\ &= [(z - H_{ij})^{-1}V_{ij} + (z - H_{ik})^{-1}V_{ik} + (z - H_{jk})^{-1}V_{jk}](z - H_0)^{-1}, \\ \text{II}(z) &= \sum_{\substack{\beta=(ij),(ik), \\ (jk)}} W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk}) \\ &\quad \times (1 - P_{ijk})(z - H_\beta)^{-1}V_\beta[(z - H_{nl})^{-1} - (z - H_0)^{-1}]P_{nl}\rho_{nl,p} \\ &= \sum_\beta W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_\beta)^{-1}V_\beta \\ &\quad \times (z - H_{nl})^{-1}P_{nl}\rho_{nl,p} \\ &\quad - \sum_\beta W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk}) \\ &\quad \times [(z - H_\beta)^{-1} - (z - H_0)^{-1}]P_{nl}\rho_{nl,p}. \end{aligned}$$

Using the remark before Lemma V.10, Lemma II.3, and Theorem II.9, we see that $(1 + x_{nl}^2)^\delta W_\beta(z - H_{nl})^{-1}P_{nl}\rho_{nl,p}$ is well behaved for each $\beta = (i, j)$ (i, k), or (j, k) . So, to control the first sum, it is sufficient to control

$$W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_\beta)^{-1}U_\beta(1 + x_{nl}^2)^{-\delta}.$$

This equals

$$\begin{aligned} & W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(1 - P_{\beta}) \\ & \times (z - H_{\beta})^{-1}U_{\beta}(1 + x_{nl}^2)^{-\delta} + W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk}) \\ & \times (1 - P_{ijk})P_{\beta}(z - H_{\beta})^{-1}U_{\beta}(1 + x_{nl}^2)^{-\delta}. \end{aligned}$$

Lemma V.9 controls the first of these terms. The second is well behaved by Corollary V.5 and Lemmas V.4 and V.12.

The second sum in the last expression for $\Pi(z)$ is controlled by Lemmas V.9 and V.12.

Next consider $\text{III}(z)$.

$$W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})A(z)(1 - M(z))^{-1}$$

has been studied in the proof of Proposition V.15. So, it suffices to control $M(z)C(z)V_{nl}(z - H_{nl})^{-1}P_{nl}\rho_{nl,p}$.

Two types of terms appear here:

$$W_{\delta}(1 - P_{\delta})(z - H_{\delta})^{-1}V_{\sigma}(z - H_{\sigma})^{-1}V_{nl}(z - H_{nl})^{-1}P_{nl}\rho_{nl,p},$$

and

$$\rho_{\delta,q}^{-1}P_{\delta}V_{\sigma}(z - H_{\sigma})^{-1}V_{nl}(z - H_{nl})^{-1}P_{nl}\rho_{nl,p}$$

where δ and σ have a common index. The first type appears in $\Pi(z)$, and has been studied. The second equals

$$\begin{aligned} & \rho_{\delta,q}^{-1}P_{\delta}V_{\sigma}(1 - P_{\sigma})(z - H_{\sigma})^{-1}V_{nl}(z - H_{nl})^{-1}P_{nl}\rho_{nl,p} \\ & + \rho_{\delta,q}^{-1}P_{\delta}V_{\sigma}P_{\sigma}(z - H_{\sigma})^{-1}V_{nl}(z - H_{nl})^{-1}P_{nl}\rho_{nl,p}. \end{aligned}$$

The first of these is well behaved by Lemmas V.4 and V.12. The second may be rewritten as:

$$\begin{aligned} & (\rho_{\delta,q}^{-1}P_{\delta}U_{\sigma}(1 + x_{\delta}^2)^{\gamma/2})(W_{\sigma}(1 + x_{\delta}^2)^{-\gamma/2}P_{\sigma}(z - H_{\sigma})^{-1}U_{nl}(1 + x_{\sigma}^2)^{\gamma/2}) \\ & \times ((1 + x_{\sigma}^2)^{-\gamma/2}P_{nl}(z - H_{nl})^{-1}\rho_{nl,p}). \end{aligned}$$

Lemma V.4 applies to the first factor. Lemma V.4 and the proof of Lemma II.3 control the others. \square

PROPOSITION V.18. *Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct; let $n = i, j$, or k ; let p be any index not equal to n or l ; let α be the pair not containing n or p . The following are well behaved and compact:*

$$\begin{aligned} K_1(z) &= \rho_{ij,k}^{-1}P_{ij}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1}V_{nl}P_{nl}(z - H_{nl})^{-1}\rho_{nl,p}, \\ K_2(z) &= \rho_{ij,k}^{-1}P_{ij}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1}V_{nl}P_{nl}(z - H_{nl})^{-1}U_{\alpha}. \end{aligned}$$

PROOF. The technique for extending the proof for Proposition V.16 to these operators is the same as the technique used to obtain the proof of Proposition V.17 from that of Proposition V.15. In some instances, Lemma V.14 must be used in place of Lemmas V.9 and V.12. \square

PROPOSITION V.19. Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct, and let $n = j$ or k . The following are compact and well behaved:

$$K_1(z) = \rho_{jk}^{-1} P_{ijk} V_{il} P_{il} (z - H_{il})^{-1} \rho_{il,n},$$

$$K_2(z) = \rho_{jk}^{-1} P_{ijk} V_{il} P_{il} (z - H_{il})^{-1} U_{jk},$$

$$K_3(z) = \rho_{ij,kl}^{-1} P_{ij,kl} V_{il} P_{il} (z - H_{il})^{-1} \rho_{il,n},$$

$$K_4(z) = \rho_{ij,kl}^{-1} P_{ij,kl} V_{il} P_{il} (z - H_{il})^{-1} U_{jk}.$$

PROOF. Consider only $K_1(z)$; proofs for the others are similar.

$$K_1(z) = (\rho_{jk}^{-1} P_{ijk} U_{il} (1 + x_{ij}^2 + x_{ik}^2 + x_{jk}^2)^\alpha) \\ \times ((1 + x_{ij} + x_{ik} + x_{jk})^{-\alpha} W_{il} P_{il} (z - H_{il})^{-1} \rho_{il,n}).$$

Lemma V.4 bounds the first factor. $P_{il}(z - H_{il})^{-1} = P_{il}(z - K_{il} - E_{il})^{-1}$, where K_{il} is a multiplication operator in the momentum representation. The methods of Lemma II.3 may be used to prove that the second factor is well behaved. For large negative $\text{Re } z$, $(1 + x_{ij}^2 + x_{ik}^2 + x_{jk}^2)^{-\alpha} W_{il}(z - H_{il})^{-1}$ is compact because $(1 + x_{ij}^2 + x_{ik}^2 + x_{jk}^2)^{-\alpha} W_{il}$ falls off in all directions in configuration space. This implies compactness. \square

PROPOSITION V.20. Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct; let $n = j$ or l ; and let $p = j$ or k . The following are well behaved and compact:

$$K_1(z) = \rho_{jk}^{-1} P_{ijk} V_{il} (z - H_{il})^{-1} (V_{ip} + V_{lp}) (z - H_{ilp})^{-1} P_{ilp} \rho_{ilp},$$

$$K_2(z) = \rho_{jk}^{-1} P_{ijk} V_{il} (z - H_{il})^{-1} V_{jk} (z - H_{il,jk})^{-1} P_{il,jk} \rho_{il,jk},$$

$$K_3(z) = \rho_{ij,kl}^{-1} P_{ij,kl} V_{ik} (z - H_{ik})^{-1} (V_{in} + V_{kn}) (z - H_{ikn})^{-1} P_{ikn} \rho_{ikn},$$

$$K_4(z) = \rho_{ij,kl}^{-1} P_{ij,kl} V_{ik} (z - H_{ik})^{-1} V_{jl} (z - H_{ik,jl})^{-1} P_{ik,jl} \rho_{ik,jl}.$$

PROOF. Consider only $K_1(z)$; proofs for the others are similar.

$$K_1(z) = \rho_{jk}^{-1} P_{ijk} V_{il} (z - H_{ilp})^{-1} P_{ilp} \rho_{ilp} \\ - \rho_{jk}^{-1} P_{ijk} V_{il} (1 - P_{il}) (z - H_{il})^{-1} P_{ilp} \rho_{ilp} \\ - \rho_{jk}^{-1} P_{ijk} V_{il} P_{il} (z - H_{il})^{-1} P_{ilp} \rho_{ilp}.$$

The methods used to prove Proposition V.19 may easily be modified to control the first and last terms. The middle term may be written as:

$$- (\rho_{jk}^{-1} P_{ijk} U_{il}) (W_{il} (1 - P_{il}) (z - H_{il})^{-1} (1 + x_{il}^2)^{-\gamma/2}) ((1 + x_{il}^2)^{\gamma/2} P_{ilp} \rho_{ilp}).$$

Lemma V.4 bounds the first factor. Theorem II.9 bounds the third. The proof of Lemma V.7 controls the middle factor.

Given the above results, it suffices to prove compactness when V_{ip} and V_{lp} are bounded. In that case,

$$K_1(z) = (\rho_{jk}^{-1} P_{ijk} V_{il} (z - H_{il})^{-1} (V_{ip} + V_{lp})) ((z - H_{ilp})^{-1} P_{ilp} \rho_{ilp}).$$

For large negative $\text{Re } z$, the first factor is bounded. In the second,

$$(z - H_{ilp})^{-1} P_{ilp} = (z - K_{ilp} - E_{ilp})^{-1} P_{ilp}.$$

P_{ilp} is (compact) $\otimes 1$; $(z - K_{ilp} - E_{ilp})^{-1} \rho_{ilp}$ is $1 \otimes$ (compact). This implies compactness. \square

In order to prove certain results involving $(z - H_{ij,kl})^{-1}$, a formula analogous to equation (IV.6) must be obtained. To motivate the formula we will use, we first give a simpler formula:

$$(z - H_{ij,kl})^{-1} = (z - H_0)^{-1} + \mathcal{Q}(z)(1 - \mathfrak{N}(z))^{-1}\mathcal{C}(z), \quad (\text{V.5})$$

where

$$\begin{aligned} \mathcal{Q}(z) &= \begin{bmatrix} (z - H_0)^{-1} U_{ij} & (z - H_0)^{-1} U_{kl} \\ 0 & W_{ij}(z - H_{ij})^{-1} U_{kl} \end{bmatrix}, \\ \mathfrak{N}(z) &= \begin{bmatrix} 0 & W_{ij}(z - H_{ij})^{-1} U_{kl} \\ W_{kl}(z - H_{kl})^{-1} U_{ij} & 0 \end{bmatrix}, \\ \mathcal{C}(z) &= \begin{bmatrix} W_{ij}(z - H_{ij})^{-1} \\ W_{kl}(z - H_{kl})^{-1} \end{bmatrix}. \end{aligned}$$

For large negative $\text{Re } z$, the inverse term may be expanded by geometric series. Then, using the diagram symbolism, the formula is easily checked. Using the uniqueness of analytic continuation, the formula is valid wherever both sides contain only bounded operators.

If the relative motion of the centers of mass of particles i and j and of particles k and l is removed, then the barely connected terms $W_{ij}(z - H_{ij})^{-1} U_{kl}$ and $W_{kl}(z - H_{kl})^{-1} U_{ij}$ are compact. So, the analytic Fredholm theorem shows that the inverse term of equation (V.5) exists for $z \notin \sigma_{\text{ess}}(H_{ij,kl})$ except at those z 's for which the system of equations

$$\phi_{ij} = W_{ij}(z - H_{ij})^{-1} U_{kl} \phi_{kl} \quad \text{and} \quad \phi_{kl} = W_{kl}(z - H_{kl})^{-1} U_{ij} \phi_{ij}$$

has a nontrivial solution. Mimicking equation (V.1), we see that this occurs only when z is an eigenvalue of $H_{ij,kl}$. Thus, equation (V.5) is valid for $z \in \mathbb{C} \setminus \sigma(H_{ij,kl})$.

As z approaches $\sigma_{\text{ess}}(H_{ij,kl})$, various terms in equation (V.5) are singular, unless $P_{ij} = P_{kl} = 0$. So, we require another formula. Because V_{kl} and P_{ij} commute, there is no analogue of the ρ functions in this case. However, a formula very much like equation (IV.6) is valid. Let $G_0 = (z - H_0)^{-1}$, $G_{ij} = (z - H_{ij})^{-1}$, etc. Then,

$$G_{ij,kl} = G_{ij} + G_{kl} - G_0 + A(z)(1 - M(z))^{-1}C(z), \quad (\text{V.6})$$

where $A(z)$, $M(z)$, and $C(z)$ are given in Figure 2.

$$\begin{aligned}
A(z) &= \begin{bmatrix} G_o U_{ij} & (G_{ij} - G_o) P_{ij} U_{kl} & G_o U_{kl} & (G_{kl} - G_o) P_{kl} U_{ij} \end{bmatrix} \\
M(z) &= \begin{bmatrix} 0 & 0 & W_{ij}(1-P_{ij})G_{ij}U_{kl} & W_{ij}(1-P_{ij})G_{ij}V_{kl}P_{kl}G_{kl}U_{ij} \\ 0 & 0 & P_{ij} & P_{ij}W_{kl}P_{kl}G_{kl}U_{ij} \\ W_{kl}(1-P_{kl})G_{kl}U_{ij} & W_{kl}(1-P_{kl})G_{kl}V_{ij}P_{ij}G_{ij}U_{kl} & 0 & 0 \\ P_{kl} & P_{kl}W_{ij}P_{ij}G_{ij}U_{kl} & 0 & 0 \end{bmatrix} \\
C(z) &= \begin{bmatrix} W_{ij}(1-P_{ij})G_{ij}V_{kl}G_{kl} \\ P_{ij}W_{kl}G_{kl} \\ W_{kl}(1-P_{kl})G_{kl}V_{ij}G_{ij} \\ P_{kl}W_{ij}G_{ij} \end{bmatrix}
\end{aligned}$$

FIGURE 2

To establish the validity of equation (V.6), we expand the inverse term of equation (V.5) and insert projections P_{ij} , $(1 - P_{ij})$, P_{kl} and $(1 - P_{kl})$ before the resolvent factors. We then sum the resulting series to obtain equation (V.6). This is completely analogous to the derivation of equation (IV.6) from equation (IV.4). The absence of spurious zeros for equation (V.6) is proved by mimicking the proof of Proposition V.2. Below, we will prove $M(z)$ is well behaved, and $(M(z))^2$ is compact (when the motion of the center of mass of particles i and j relative to the center of mass of particles k and l is removed). $A(z)$ and $C(z)$ are analytic for $z \in \mathbb{C} \setminus \sigma_{\text{ess}}(H_{ij,kl})$, so equation (V.6) is valid for $z \notin \sigma(H_{ij,kl})$.

In the two and three body cases (with finite masses), $M(z)$ was well behaved. We have not been able to prove this for equation (V.6) (or for the three body case with an infinite mass, in which case an equation like (V.6) is used in place of equation (IV.6)). However, $(M(z))^2$ is well behaved. The next few lemmas prove this.

LEMMA V.21. *Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct. $K(z) = W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}U_{kl}$ is analytic for $z \in \mathbb{C} \setminus \sigma_{\text{ess}}(H_{ij,kl})$; tends to zero as $\text{Re } z \rightarrow -\infty$; and has strongly continuous boundary values on $\sigma_{\text{ess}}(H_{ij,kl})$ from above and below. If the motion of the center of mass of particles i and j relative to the center of mass of particles k and l has been removed, then $K(z)$ is compact for $z \in \mathbb{C} \setminus \sigma_{\text{ess}}(H_{ij,kl})$.*

PROOF. Except for compactness, Lemma II.3 yields these results for the operator valued function $W_{ij}(1 - P_{ij})(z - H_0)^{-1}U_{kl}$:

$$\begin{aligned}
K(z) &= W_{ij}(1 - P_{ij})(z - H_0)^{-1}U_{kl} \\
&\quad + (W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}U_{ij})(W_{ij}(z - H_0)^{-1}U_{kl}).
\end{aligned}$$

Lemmas II.3 and V.7 apply to the various factors here.

To prove compactness, we expand the resolvent in $K(z)$ as a sum of graphs. The standard compactness proofs apply to each term at large negative $\text{Re } z$. \square

LEMMA V.22. Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct. $K(z) = P_{ij}W_{kl}P_{kl}(z - H_{kl})^{-1}U_{ij}$ is well behaved. Moreover, if the relative motion of the centers of mass of cluster $\{i, j\}$ and $\{k, l\}$ is removed, then $K(z)$ is compact.

PROOF. $K(z) = (W_{kl}P_{kl})(P_{kl}P_{ij}(1 + x_{ij}^2)^\alpha)((1 + x_{ij}^2)^{-\alpha}(z - K_{kl} - E_{kl})^{-1}U_{ij})$. Corollary V.5 bounds the first factor. The second is compact on $L^2(\mathbb{R}^{2m})$. Lemma II.3 controls the third. \square

LEMMA V.23. Assume Hypotheses (1)–(3), and (5). If the relative motion of the centers of mass of the clusters $\{i, j\}$ and $\{k, l\}$ is removed, then $M(z)$ in equation (V.6) is analytic in $\mathbb{C} \setminus \sigma_{\text{ess}}(H_{ij,kl})$, with strongly continuous boundary values on the cut. $(M(z))^2$ is compact and well behaved.

PROOF. All results, except for compactness and norm continuity of $(M(z))^2$ up to the cut, are consequences of Lemmas V.12, V.21, and V.22.

$(M(z))^2$ has four types of nonzero terms.

Type 1. $W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{kl}(z - H_{kl})^{-1}U_{ij}$.

Lemmas V.9 and V.12 imply the continuity. Compactness follows from Lemma V.21, since $U_{kl}(z - H_{kl})^{-1}U_{ij}$ is bounded for large negative $\text{Re } z$.

Type 2. $P_{ij}W_{kl}(z - H_{kl})^{-1}U_{ij}$.

This equals

$$P_{ij}W_{kl}P_{kl}(z - H_{kl})^{-1}U_{ij} + P_{ij}W_{kl}(1 - P_{kl})(z - H_{kl})^{-1}U_{ij}.$$

So, Lemmas V.21 and V.22 show that it suffices to prove the continuity when W_{kl} is bounded. In that case,

$$P_{ij}W_{kl}(z - H_{kl})^{-1}U_{ij} = (W_{kl}P_{ij}(1 + x_{ij}^2)^\alpha)((1 + x_{ij}^2)^{-\alpha}(z - H_{kl})^{-1}U_{ij}).$$

The first factor is bounded. The second is well behaved by Lemma II.3.

Type 3. $W_{ij}(1 - P_{ij})V_{kl}(z - H_{kl})^{-1}V_{ij}(z - H_{ij})P_{ij}U_{kl}$.

$G_{ij} - G_{kl} = G_{kl}(V_{ij} - V_{kl})G_{ij}$, so the operator in question equals

$$\begin{aligned} & W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{kl}(z - H_{ij})^{-1}P_{ij}U_{kl} \\ & - W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{kl}(z - H_{kl})^{-1}P_{ij}U_{kl} \\ & + W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{kl}(z - H_{kl})^{-1}V_{kl}(z - H_{ij})^{-1}P_{ij}U_{kl}. \end{aligned}$$

The first term here is zero, since the P_{ij} commutes with $(z - H_{ij})^{-1}V_{kl}(z - H_{ij})^{-1}$. By Lemmas V.21 and V.22, it suffices to prove the continuity for the remaining terms when V_{kl} and U_{kl} are bounded. The second term is controlled by Lemmas V.9 and V.12 after factorization as

$$- (W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{kl}(z - H_{kl})^{-1}(1 + x_{ij}^2)^{-\alpha})((1 + x_{ij}^2)^\alpha P_{ij}U_{kl}).$$

The third term equals

$$\begin{aligned} & (W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{kl}(z - H_{kl})^{-1}(1 + x_{ij}^2)^{-\alpha}) \\ & \times ((1 + x_{ij}^2)^\alpha P_{ij}U_{kl})(W_{kl}(z - H_{ij})^{-1}U_{kl}). \end{aligned}$$

Lemmas V.9 and V.12 control the first factor. The second factor is bounded. The third is controlled by Lemma II.3.

Type 4. $P_{ij}W_{kl}(z - H_{kl})^{-1}V_{ij}(z - H_{ij})^{-1}P_{ij}U_{kl}$.

This equals

$$P_{ij}W_{kl}(1 - P_{kl})(z - H_{kl})^{-1}V_{ij}(z - H_{ij})^{-1}P_{ij}U_{kl} \\ + (P_{ij}W_{kl}P_{kl}(z - H_{kl})^{-1}U_{ij})(P_{kl}W_{ij}P_{ij}(z - H_{ij})^{-1}U_{kl}).$$

Compactness follows from Lemmas V.21 and V.22. Lemmas V.12 and V.22 imply the norm continuity of the first and second terms, respectively. \square

REMARK. $C(z)$ in equation (V.6) is well behaved from $L^2_\delta(\mathbf{R}^{2m})$ to $L^2(\mathbf{R}^{2m})$. Also,

$$G_{ij} = G_{ij}P_{ij} + G_0(1 - P_{ij}) + G_0V_{ij}G_{ij}(1 - P_{ij})$$

and

$$G_{kl} = G_{kl}P_{kl} + G_0(1 - P_{kl}) + G_0V_{kl}G_{kl}(1 - P_{kl}).$$

These facts, together with Lemma V.23 and Theorem III.1 imply asymptotic completeness for $H_{ij,kl}$, when Hypotheses (1)–(3) hold. Much easier proofs of this fact exist, but this result is a simple by-product of our methods.

LEMMA V.24. *Suppose $A(z)$ is a strongly continuous bounded function with values in the bounded operators on \mathcal{H} . Suppose $B(z)$ is a norm continuous function with values in the compact operators on \mathcal{H} . Then $A(z)B(z)$ is norm continuous and compact.*

PROOF. The compactness of $A(z)B(z)$ is a well-known fact.

Suppose $\varepsilon > 0$ is given. Fix z , and let $N = \sup\|A(w)\|$. Choose a finite rank operator C , such that $\|B(z) - C\| < \varepsilon/6N$. The restriction of $A(w)$ to the range of C is norm continuous. Therefore, there exists $\delta > 0$, so that $|z - w| < \delta$ implies $\|(A(z) - A(w))C\| < \varepsilon/3N$.

If $|z - w| < \delta$, then by the triangle inequality,

$$\|A(z)B(z) - A(w)B(w)\| \\ \leq \|(A(z) - A(w))(B(z) - C)\| + \|(A(z) - A(w))C\| \\ + \|A(w)(B(z) - B(w))\| \\ < 2N\varepsilon/6N + \varepsilon/3 + N\varepsilon/3N = \varepsilon. \quad \square$$

PROPOSITION V.25. *Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct indices.*

$$K_1(z) = W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1}U_{ik}$$

is compact and well behaved.

$$K_2(z) = W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1}$$

is compact and well behaved from $L^2_\delta(\mathbf{R}^{3m})$ to $L^2(\mathbf{R}^{3m})$, for any $\delta > 1$.

PROOF. Consider only $K_1(z)$; the proof for $K_2(z)$ is similar.

Compactness for large negative $\operatorname{Re} z$ is easily proved by expanding the resolvents as sums of graphs. It therefore suffices to prove $K_1(z)$ is well behaved.

Using equation (V.6), we write $K_1(z)$ as a sum of four terms.

$$\begin{aligned} \text{I}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})(z - H_{ij})^{-1} U_{ik}, \\ \text{II}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})(z - H_{kl})^{-1} U_{ik}, \\ \text{III}(z) &= -W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})(z - H_0)^{-1} U_{ik}, \\ \text{IV}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})A(z)(1 - M(z))^{-1}C(z)U_{ik}. \end{aligned}$$

Lemmas V.9 and V.12 control $\text{I}(z)$, $\text{II}(z)$, and $\text{III}(z)$. Lemmas II.3 and V.21 show that $W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{kl}(1 - P_{ij,kl})A(z)$ is well behaved, except that the boundary values are only strongly continuous. Lemma V.23 shows that $M(z)$ has similar properties.

By mimicking part of the proof of Proposition V.15, we see that the factor of $(1 - P_{ij,kl})$ in $\text{IV}(z)$ removes the singularities of $(1 - M(z))^{-1}$. Thus,

$$W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})A(z)(1 - M(z))^{-1}$$

is well behaved, except that the boundary values are strongly continuous. Lemma V.24 now shows that it suffices to prove $C(z)U_{ik}$ is well behaved and compact.

The first and third entries of $C(z)U_{ik}$ are compact for large negative $\text{Re } z$, because they may be written as sums of connected graphs. The second and fourth entries are compact by the compactness argument used in the proof of Proposition V.16.

Lemmas V.9 and V.12 show that the first and third entries of $C(z)U_{ik}$ are well behaved. The second entry equals

$$P_{ij}W_{kl}(1 - P_{kl})(z - H_{kl})^{-1}U_{ik} + P_{ij}W_{kl}P_{kl}(z - H_{kl})^{-1}U_{ik}.$$

Lemma V.6 controls the first term. The second may be written as

$$(P_{ij}(1 + x_{ij}^2)^\alpha)(W_{kl}P_{kl})((1 + x_{ij}^2)^{-\alpha}(z - H_{kl})^{-1}U_{ik}).$$

Apply Lemma II.3, Theorem II.9, and Corollary V.5. This shows that the second entry of $C(z)U_{ik}$ is well behaved. The fourth entry is handled in the same way. \square

PROPOSITION V.26. *Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct.*

$$K_1(z) = P_{ij}W_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1}U_{ik}$$

is compact and well behaved.

$$K_2(z) = P_{ij}W_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1}$$

is compact and well behaved from $L_\delta^2(\mathbb{R}^{3m})$ to $L^2(\mathbb{R}^{3m})$, for any $\delta > 1$.

PROOF. Consider only $K_1(z)$; proof for $K_2(z)$ is similar.

P_{ij} and $W_{kl}(1 - P_{ij,kl})$ commute. $P_{ij}(z - H_{ij,kl})^{-1} = P_{ij}(z - E_{ij} - H'_{kl})^{-1}$, where $H'_{kl} = H_{kl} - H_0^{(ij)}$. Moreover, $(1 - P_{ij,kl})P_{ij} = (1 - P_{kl})P_{ij}$. Therefore,

$$K(z) = (P_{ij}W_{kl}(1 - P_{kl})(z - E_{ij} - H'_{kl})^{-1}\rho_{ij,k})(\rho_{ij,k}^{-1}P_{ij}U_{ik}).$$

Lemma V.4 bounds the second factor. P_{ij} is (compact) $\otimes 1$;

$$W_{kl}(1 - P_{kl})(z - E_{ij} - H'_{kl})^{-1} \rho_{ij,k} = 1 \otimes X(z).$$

$X(z)$ is compact for large negative $\operatorname{Re} z$. The proof of this is the same as the compactness of the three body operator $W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} U_{ik}$ on $L^2(\mathbb{R}^{2m})$. $X(z)$ is well behaved by the proof of Lemma V.8 and the remark before Lemma V.10. Thus, $K_1(z)$ is compact and well behaved. \square

PROPOSITION V.27. *Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct, and let $n = j$ or l . The following are well behaved and compact:*

$$\begin{aligned} K_1(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1} \\ &\quad \times V_{ik}(z - H_{ik})^{-1} P_{ik} U_{jl} \end{aligned}$$

and

$$\begin{aligned} K_2(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1} \\ &\quad \times V_{ik}(z - H_{ik})^{-1} P_{ik} \rho_{ik,n}. \end{aligned}$$

PROOF. Consider only $K_1(z)$; the proof for $K_2(z)$ is similar.

For large negative $\operatorname{Re} z$, $W_{ik}(z - H_{ik})^{-1} P_{ik} U_{jl}$ is bounded, so compactness follows from Proposition V.25. Hence, it suffices to prove $K_1(z)$ is well behaved.

Using equation (V.6), we write $K_1(z)$ as a sum of four terms:

$$\begin{aligned} \text{I}(z) &= -W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})(z - H_0)^{-1} \\ &\quad \times V_{ik}(z - H_{ik})^{-1} P_{ik} U_{jl}, \end{aligned}$$

$$\begin{aligned} \text{II}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})(z - H_{ij})^{-1} \\ &\quad \times V_{ik}(z - H_{ik})^{-1} P_{ik} U_{jl}, \end{aligned}$$

$$\begin{aligned} \text{III}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})(z - H_{kl})^{-1} \\ &\quad \times V_{ik}(z - H_{ik})^{-1} P_{ik} U_{jl}, \end{aligned}$$

$$\begin{aligned} \text{IV}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl}) A(z) (1 - M(z))^{-1} \\ &\quad \times C(z) V_{ik}(z - H_{ik})^{-1} P_{ik} U_{jl}. \end{aligned}$$

$$\text{I}(z) = W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl}) [(z - H_{ik})^{-1} - (z - H_0)^{-1}] P_{ik} U_{jl}.$$

This is well behaved by Lemma V.12.

$$\begin{aligned} \text{II}(z) &= -\text{I}(z) + W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})(z - H_{ij})^{-1} \\ &\quad \times V_{ij} [(z - H_{ik})^{-1} - (z - H_0)^{-1}] P_{ik} U_{jl} \\ &= -\text{I}(z) + (W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})(1 - P_{ij})(z - H_{ij})^{-1} U_{ij}) \\ &\quad \times (W_{ij} [(z - H_{ik})^{-1} - (z - H_0)^{-1}] P_{ik} U_{jl}). \end{aligned}$$

Apply Lemmas V.9, V.10, and II.3:

$$\begin{aligned}
 \text{III}(z) &= -\text{I}(z) + W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})(z - H_{kl})^{-1} \\
 &\quad \times V_{kl}[(z - H_{ik})^{-1} - (z - H_0)^{-1}] P_{ik} U_{jl} \\
 &= -\text{I}(z) + (W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})(z - H_{kl})^{-1} \\
 &\quad \times U_{kl}(1 + x_{ik}^2)^{-\alpha})((1 + x_{ik}^2)^\alpha W_{kl} P_{ik}(z - H_{ik})^{-1} U_{jl}) \\
 &\quad - W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl}) \\
 &\quad \times [(z - H_{kl})^{-1} - (z - H_0)^{-1}] P_{ik} U_{jl}.
 \end{aligned}$$

Lemmas V.9 and V.12 control the last term. Lemma V.10 and the fall off of Ψ_{ik} control the second factor of the middle term. If U_{kl} is bounded, then Lemmas V.4, V.9, and V.12 control the first factor. When U_{kl} is unbounded, a density argument shows that it suffices to prove uniform boundedness of the first factor. To do this, we rewrite this first factor as

$$\begin{aligned}
 &W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} V_{kl}(1 - P_{ij,kl})(z - H_{kl})^{-1} \\
 &\quad \times P_{kl} U_{kl}(1 + x_{ik}^2)^{-\alpha} + W_{ij}(1 - P_{ij})(z - H_{ij})^{-1} \\
 &\quad \times V_{kl}(1 - P_{ij,kl})(z - H_{kl})^{-1} (1 - P_{kl}) U_{kl}(1 + x_{ik}^2)^{-\alpha}.
 \end{aligned}$$

Lemmas V.8 and V.21 show the second term is uniformly bounded. Corollary V.5 and Lemmas V.4 and V.12 control the first term.

To control $\text{IV}(z)$, we proceed as in the proof of Proposition V.25. This shows that we need only prove $C(z) V_{ik}(z - H_{ik})^{-1} P_{ik} U_{jl}$ is compact and well behaved. Since $W_{ik}(z - H_{ik})^{-1} P_{ik} U_{jl}$ is bounded for large negative $\text{Re } z$, compactness follows from arguments in the proof of Proposition V.25. To prove $C(z) V_{ik}(z - H_{ik})^{-1} P_{ik} U_{jl}$ is well behaved, notice that the first and third entries are of the same form as $\text{III}(z)$, above. The second entry may be rewritten as

$$\begin{aligned}
 &P_{ij} W_{kl}[(z - H_{ik})^{-1} - (z - H_0)^{-1}] P_{ik} U_{jl} \\
 &\quad + P_{ij} W_{kl}(z - H_{kl})^{-1} V_{kl}(z - H_{ik})^{-1} P_{ik} U_{jl} \\
 &\quad - P_{ij} W_{kl}[(z - H_{kl})^{-1} - (z - H_0)^{-1}] P_{ik} U_{jl}.
 \end{aligned}$$

Replace each $(z - H_{kl})^{-1}$ in these expressions by $(z - H_{kl})^{-1} P_{kl} + (z - H_{kl})^{-1} (1 - P_{kl})$. The only terms which are not controlled by various lemmas are those containing the factor P_{kl} . To control them, notice that $W_{kl} P_{kl}$ and $P_{kl} U_{kl}$ are bounded. Then use Lemma V.4 and the fall off of Ψ_{ij} and Ψ_{ik} to obtain factors which are controlled by Lemma II.3. This controls the second entry of $C(z) V_{ik}(z - H_{ik})^{-1} P_{ik} U_{jl}$. The same method applies to the fourth entry. \square

PROPOSITION V.28. *Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct, and let $n = i$ or j . The following are well behaved and compact:*

$$K_1(z) = P_{ij} W_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1} V_{ik}(z - H_{ik})^{-1} P_{ik} U_{jl}$$

and

$$K_2(z) = P_{ij}W_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1}V_{ik}(z - H_{ik})^{-1}P_{ik}\rho_{ik,n}.$$

PROOF. Consider only $K_1(z)$; the proof for $K_2(z)$ is similar.

For large negative $\text{Re } z$, $W_{ik}(z - H_{ik})^{-1}P_{ik}U_{jl}$ is bounded, so compactness follows from Proposition V.26. It is therefore sufficient to show $K_1(z)$ is well behaved.

P_{ij} commutes with W_{kl} , and $P_{ij}(1 - P_{ij,kl}) = P_{ij}(1 - P_{kl})$. So,

$$\begin{aligned} K_1(z) &= P_{ij}W_{kl}(1 - P_{kl})(z - H_{kl})^{-1}V_{ik}(z - H_{ik})^{-1}P_{ik}U_{jl} \\ &\quad + P_{ij}W_{kl}(1 - P_{kl})(z - H_{kl})^{-1}V_{ij}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1} \\ &\quad \times V_{ik}(z - H_{ik})^{-1}P_{ik}U_{jl}. \end{aligned}$$

Lemma V.12 controls the first term. Proposition V.27 controls the second. \square

PROPOSITION V.29. Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct; let $n = i, j$, or k ; let p be any index not equal to n or l ; let q be the index not equal to n, l , or p . The following are well behaved and compact:

$$\begin{aligned} K_1(z) &= \rho_{ij,k}^{-1}P_{ij}(V_{ik} + V_{jk})(z - H_{ijk})^{-1}(1 - P_{ijk}) \\ &\quad \times [V_{nl}(z - H_{nl})^{-1}(V_{np} + V_{lp}) + V_{lp}(z - H_{lp})^{-1}(V_{np} + V_{nl})] \\ &\quad \times (z - H_{nlp})^{-1}P_{nlp}\rho_{nlp} \end{aligned}$$

and

$$\begin{aligned} K_2(z) &= \rho_{ij,k}^{-1}P_{ij}(V_{ik} + V_{jk})(z - H_{ijk})^{-1}(1 - P_{ijk}) \\ &\quad \times V_{nl}(z - H_{nl})^{-1}V_{pq}(z - H_{nl,pq})^{-1}P_{nl,pq}\rho_{nl,pq}. \end{aligned}$$

PROOF. Consider only $K_1(z)$; proof for $K_2(z)$ is similar.

For large negative $\text{Re } z$,

$$W_{nl}(z - H_{nl})^{-1}(V_{np} + V_{pl})(z - H_{nlp})^{-1}P_{nlp}\rho_{nlp}$$

and

$$W_{lp}(z - H_{lp})^{-1}(V_{np} + V_{nl})(z - H_{nlp})^{-1}P_{nlp}\rho_{nlp}$$

are uniformly bounded, so compactness and $\lim_{\text{Re } z \rightarrow -\infty} \|K_1(z)\| = 0$ follow from Proposition V.16.

Let $X(z) = \rho_{ij,k}^{-1}P_{ij}(V_{ik} + V_{jk})(z - H_{ijk})^{-1}(1 - P_{ijk})$.

$$\begin{aligned} K_1(z) &= X(z)(V_{nl} + V_{pl})(z - H_{nlp})^{-1}P_{nlp}\rho_{nlp} \\ &\quad - X(z)V_{nl}(z - H_{nl})^{-1}P_{nlp}\rho_{nlp} - X(z)V_{pl}(z - H_{pl})^{-1}P_{nlp}\rho_{nlp}. \end{aligned}$$

The second and third terms can each be broken into two terms which are controlled by Proposition V.18 or Lemma V.8 and Proposition V.16.

Since $\{i, j, k\} = \{n, p, q\}$, the first term equals:

$$\begin{aligned} & \rho_{ij,k}^{-1} P_{ij} (V_{ik} + V_{jk}) (1 - P_{ijk}) (z - H_{np})^{-1} (V_{nl} + V_{pl}) (z - H_{np})^{-1} \\ & \quad \times P_{nlp} \rho_{nlp} + \rho_{ij,k}^{-1} P_{ij} (V_{ik} + V_{jk}) (1 - P_{ijk}) (z - H_{ijk})^{-1} \\ & \quad \times (V_{nq} + V_{pq}) (z - H_{np})^{-1} (V_{nl} + V_{pl}) (z - H_{np})^{-1} P_{nlp} \rho_{nlp}. \end{aligned}$$

This may be rewritten as a sum of four terms:

$$\begin{aligned} \text{I}(z) &= \rho_{ij,k}^{-1} P_{ij} (V_{ik} + V_{jk}) (1 - P_{ijk}) (z - H_{np})^{-1} P_{nlp} \rho_{nlp}, \\ \text{II}(z) &= -\rho_{ij,k}^{-1} P_{ij} (V_{ik} + V_{jk}) (1 - P_{ijk}) (z - H_{np})^{-1} P_{nlp} \rho_{nlp}, \\ \text{III}(z) &= \rho_{ij,k}^{-1} P_{ij} (V_{ik} + V_{jk}) (1 - P_{ijk}) \\ & \quad \times (z - H_{ijk})^{-1} (V_{nq} + V_{pq}) (z - H_{np})^{-1} P_{nlp} \rho_{nlp}, \\ \text{IV}(z) &= -\rho_{ij,k}^{-1} P_{ij} (V_{ik} + V_{jk}) (1 - P_{ijk}) \\ & \quad \times (z - H_{ijk})^{-1} (V_{nq} + V_{pq}) (z - H_{np})^{-1} P_{nlp} \rho_{nlp}. \\ \text{I}(z) &= \rho_{ij,k}^{-1} P_{ij} (V_{ik} + V_{jk}) (z - H_{np})^{-1} P_{nlp} \rho_{nlp} \\ & \quad - \rho_{ij,k}^{-1} P_{ij} (V_{ik} + V_{jk}) P_{ijk} (z - H_{np})^{-1} P_{nlp} \rho_{nlp}. \end{aligned}$$

The first term in this expression equals

$$\begin{aligned} & \left[\rho_{ij,k}^{-1} P_{ij} U_{ik} (1 + x_{ij}^2)^\alpha \right] \left[(1 + x_{ij}^2)^{-\alpha} W_{ik} (z - K_{nlp} - E_{nlp})^{-1} P_{nlp} \rho_{nlp} \right] \\ & + \left[\rho_{ij,k}^{-1} P_{ij} U_{jk} (1 + x_{ij}^2)^\alpha \right] \left[(1 + x_{ij}^2)^{-\alpha} W_{jk} (z - K_{nlp} - E_{nlp})^{-1} P_{nlp} \rho_{nlp} \right]. \end{aligned}$$

Lemma V.4 bounds the first factors. The methods of Lemma II.3 control the second factors. The second term in the last expression for I(z) equals

$$\begin{aligned} & - (\rho_{ij,k}^{-1} P_{ij} U_{ik}) (W_{ik} P_{ijk}) (P_{ijk} P_{nlp} \rho_{nlp}^{-1}) (\rho_{nlp} (z - H_{nlp})^{-1} \rho_{nlp}) \\ & - (\rho_{ij,k}^{-1} P_{ij} U_{jk}) (W_{jk} P_{ijk}) (P_{ijk} P_{nlp} \rho_{nlp}^{-1}) (\rho_{nlp} (z - H_{nlp})^{-1} \rho_{nlp}). \end{aligned}$$

Lemma V.4 bounds the first factors. Corollary V.5 bounds the second factors. The proof of Lemma V.4 bounds the third factors. Lemma II.3 controls the final factors.

$$\begin{aligned} \text{III}(z) &= \rho_{ij,k}^{-1} P_{ij} (V_{ik} + V_{jk}) (1 - P_{ijk}) (z - H_{ijk})^{-1} V_{nq} (z - H_{nlp})^{-1} P_{nlp} \rho_{nlp} \\ & + \rho_{ij,k}^{-1} P_{ij} (V_{ik} + V_{jk}) (1 - P_{ijk}) (z - H_{ijk})^{-1} V_{pq} (z - H_{nlp})^{-1} P_{nlp} \rho_{nlp}. \end{aligned}$$

Since $(z - H_{nlp})^{-1} P_{nlp} = (z - K_{nlp} - E_{nlp})^{-1} P_{nlp}$, Lemma V.4 and Lemma II.3 show $(1 + x_{pl}^2 + x_{nl}^2 + x_{np}^2)^\alpha W_{nq} (z - H_{nlp})^{-1} P_{nlp} \rho_{nlp}$ is well behaved. Following the proof of Proposition V.16,

$$\rho_{ij,k}^{-1} P_{ij} (V_{ik} + V_{jk}) (1 - P_{ijk}) (z - H_{ijk})^{-1} U_{nq} (1 + x_{np}^2 + x_{nl}^2 + x_{pl}^2)^{-\alpha}$$

is well behaved. This controls the first term in the last expression for III(z). The same proof controls the second term.

Since $\{n, p, q\} = \{i, j, k\}$,

$$\text{II}(z) + \text{IV}(z) = -\rho_{ij,k}^{-1} P_{ij} (V_{ik} + V_{jk}) (1 - P_{ijk}) (z - H_{ijk})^{-1} P_{nlp} \rho_{nlp}.$$

The proof of Proposition V.16 shows this is well behaved. \square

PROPOSITION V.30. *Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct. The following are compact and well behaved:*

$$\begin{aligned}
 K_1(z) &= P_{ij} W_{kl} (1 - P_{ij,kl}) (z - H_{ij,kl})^{-1} \\
 &\quad \times \left[V_{ik} (z - H_{ik})^{-1} V_{jl} + V_{jl} (z - H_{jl})^{-1} V_{ik} \right] (z - H_{ik,jl})^{-1} P_{ij,kl} \rho_{ij,kl}, \\
 K_2(z) &= P_{ij} W_{kl} (1 - P_{ij,kl}) (z - H_{ij,kl})^{-1} \\
 &\quad \times \left[V_{ik} (z - H_{ik})^{-1} (V_{jk} + V_{ij}) + V_{jk} (z - H_{jk})^{-1} (V_{ik} + V_{ij}) \right] \\
 &\quad \times (z - H_{ijk})^{-1} P_{ijk} \rho_{ijk}, \\
 K_3(z) &= P_{ij} W_{kl} (1 - P_{ij,kl}) (z - H_{ij,kl})^{-1} \\
 &\quad \times \left[V_{il} (z - H_{il})^{-1} (V_{ik} + V_{lk}) + V_{ik} (z - H_{ik})^{-1} (V_{il} + V_{lk}) \right] \\
 &\quad \times (z - H_{ilk})^{-1} P_{ilk} \rho_{ilk}.
 \end{aligned}$$

PROOF. Compactness and the limiting behavior as $\text{Re } z \rightarrow -\infty$ follow from Proposition V.26 and simple bounds.

Let

$$\begin{aligned}
 X(z) &= P_{ij} W_{kl} (z - H_{ij,kl})^{-1} (1 - P_{ij,kl}) \\
 &= W_{kl} (z - H_{ij,kl})^{-1} P_{ij} (1 - P_{kl}); \\
 K_1(z) &= X(z) (V_{ik} + V_{jl}) (z - H_{ik,jl})^{-1} P_{ik,jl} \rho_{ik,jl} \\
 &\quad - X(z) V_{ik} (z - H_{ik})^{-1} P_{ik,jl} \rho_{ik,jl} - X(z) V_{jl} (z - H_{jl})^{-1} P_{ik,jl} \rho_{ik,jl}.
 \end{aligned}$$

Trivial extensions of Proposition V.28 control the last two terms. The first term equals:

$$\begin{aligned}
 &\left[P_{ij} W_{kl} (z - H_{ij,kl})^{-1} (1 - P_{ij,kl}) U_{ik} (1 + x_{ij}^2)^\alpha \right] \\
 &\quad \times \left[(1 + x_{ij}^2)^{-\alpha} W_{ik} (z - H_{ik,jl})^{-1} P_{ik,jl} \rho_{ik,jl} \right] \\
 &\quad + \left[P_{ij} W_{kl} (z - H_{ij,kl})^{-1} (1 - P_{ij,kl}) U_{jl} (1 + x_{ij}^2)^\alpha \right] \\
 &\quad \times \left[(1 + x_{ij}^2)^{-\alpha} W_{jl} (z - H_{ik,jl})^{-1} P_{ik,jl} \rho_{ik,jl} \right].
 \end{aligned}$$

Due to the fall off of Ψ_{ij} , the proof of Proposition V.26 can easily be extended to show the first factors are well behaved. The methods of Lemmas II.3 and V.4 show the second factors are well behaved.

$$\begin{aligned}
 K_2(z) &= X(z) (V_{ik} + V_{jk}) (z - H_{ijk})^{-1} P_{ijk} \rho_{ijk} \\
 &\quad - X(z) V_{ik} (z - H_{ik})^{-1} P_{ijk} \rho_{ijk} - X(z) V_{jk} (z - H_{jk})^{-1} P_{ijk} \rho_{ijk}.
 \end{aligned}$$

The last two terms can each be written as sums of two terms, which are well behaved by Proposition V.28 or Proposition V.26 and Lemma V.8. The first term

equals:

$$\begin{aligned} & P_{ij}W_{kl}(1 - P_{ij,kl})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(z - H_{ijk})^{-1}P_{ijk}\rho_{ijk} \\ & + P_{ij}W_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1}V_{kl}(z - H_{ij})^{-1}(V_{ik} + V_{jk}) \\ & \times (z - H_{ijk})^{-1}P_{ijk}\rho_{ijk}. \end{aligned}$$

We write this as a sum of four terms:

$$\begin{aligned} \text{I}(z) &= P_{ij}W_{kl}(1 - P_{ij,kl})(z - H_{ijk})^{-1}P_{ijk}\rho_{ijk}, \\ \text{II}(z) &= -P_{ij}W_{kl}(1 - P_{ij,kl})(z - H_{ij})^{-1}P_{ijk}\rho_{ijk}, \\ \text{III}(z) &= P_{ij}W_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1}V_{kl}(z - H_{ijk})^{-1}P_{ijk}\rho_{ijk}, \\ \text{IV}(z) &= -P_{ij}W_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1}V_{kl}(z - H_{ij})^{-1}P_{ijk}\rho_{ijk}, \\ \text{I}(z) &= (P_{ij}W_{kl}(1 - P_{ij,kl})P_{ijk}\rho_{ijk}^{-1})(\rho_{ijk}(z - H_{ijk})^{-1}\rho_{ijk}). \end{aligned}$$

Apply Lemmas V.4 and II.3 to control this.

$$\begin{aligned} \text{III}(z) &= (P_{ij}W_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1}U_{kl})(W_{kl}P_{ijk}\rho_{ijk}^{-1}) \\ &\times (\rho_{ijk}(z - H_{ijk})^{-1}\rho_{ijk}). \end{aligned}$$

The proof of Proposition V.26 easily generalizes to control the first factor; Lemma V.4 bounds the second factor; and Lemma II.3 controls the third.

$$\text{II}(z) + \text{IV}(z) = -P_{ij}W_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1}P_{ijk}\rho_{ijk}.$$

The proof of Proposition V.26 generalizes to control this term:

$$\begin{aligned} K_3(z) &= X(z)(V_{il} + V_{ik})(z - H_{ilk})^{-1}P_{ilk}\rho_{ilk} \\ &- X(z)V_{il}(z - H_{il})^{-1}P_{ikl}\rho_{ilk} - X(z)V_{ik}(z - H_{ik})^{-1}P_{ilk}\rho_{ilk}. \end{aligned}$$

The second and third terms are of a form studied in the proof for $K_2(z)$. The first term equals

$$\begin{aligned} & [P_{ij}W_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1}U_{il}(1 + x_{ij}^2)^\alpha] \\ & \times [(1 + x_{ij}^2)^{-\alpha}W_{il}(z - H_{ilk})^{-1}P_{ilk}\rho_{ilk}] \\ & + [P_{ij}W_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1}U_{ik}(1 + x_{ij}^2)^\alpha] \\ & \times [(1 + x_{ij}^2)^{-\alpha}W_{ik}(z - H_{ilk})^{-1}P_{ilk}\rho_{ilk}]. \end{aligned}$$

The first factors are of the same form as terms which have been controlled in the proof for $K_1(z)$. The second factors are controlled by methods of Lemmas V.4 and II.3. \square

PROPOSITION V.31. *Assume Hypotheses (1)–(3), and (5). Let i, j, k and l be distinct. The following are well behaved and compact:*

$$\begin{aligned}
 K_1(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\
 &\quad \times [V_{il}(z - H_{il})^{-1}(V_{jl} + V_{ij}) + V_{jl}(z - H_{jl})^{-1}(V_{il} + V_{ij})] \\
 &\quad \times (z - H_{ijl})^{-1}P_{ijl}\rho_{ijl}, \\
 K_2(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\
 &\quad \times V_{kl}(z - H_{kl})^{-1}V_{ij}(z - H_{ij,kl})^{-1}P_{ij,kl}\rho_{ij,kl}, \\
 K_3(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1} \\
 &\quad \times [V_{ik}(z - H_{ik})^{-1}(V_{jk} + V_{ij}) + V_{jk}(z - H_{jk})^{-1}(V_{ik} + V_{ij})] \\
 &\quad \times (z - H_{ijk})^{-1}P_{ijk}\rho_{ijk}, \\
 K_4(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1} \\
 &\quad \times [V_{ik}(z - H_{ik})^{-1}V_{jl} + V_{jl}(z - H_{jl})^{-1}V_{ik}](z - H_{ik,jl})^{-1}P_{ij,kl}\rho_{ij,kl}.
 \end{aligned}$$

PROOF. Consider only $K_1(z)$; the proofs for the others are similar.

Since

$$W_{il}(z - H_{il})^{-1}(V_{jl} + V_{ij})(z - H_{ijl})^{-1}P_{ijl}\rho_{ijl}$$

and

$$W_{jl}(z - H_{jl})^{-1}(V_{il} + V_{ij})(z - H_{ijl})^{-1}P_{ijl}\rho_{ijl}$$

are bounded for large negative $\text{Re } z$, compactness of $K_1(z)$ at large negative $\text{Re } z$ follows from Proposition V.15.

Let

$$X(z) = W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1}.$$

Then,

$$\begin{aligned}
 K_1(z) &= X(z)(V_{il} + V_{jl})(z - H_{ijl})^{-1}P_{ijl}\rho_{ijl} \\
 &\quad - X(z)V_{il}(z - H_{il})^{-1}P_{ijl}\rho_{ijl} - X(z)V_{jl}(z - H_{jl})^{-1}P_{ijl}\rho_{ijl}.
 \end{aligned}$$

The second and third terms can each be broken into two terms which are well behaved by Proposition V.17 or Lemma V.8 and Proposition V.15. The first term equals

$$\begin{aligned}
 &W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1}(V_{il} + V_{jl}) \\
 &\quad \times (z - H_{ijl})^{-1}P_{ijl}\rho_{ijl} \\
 &+ W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1}(V_{ik} + V_{jk}) \\
 &\quad \times (z - H_{ij})^{-1}(V_{il} + V_{jl})(z - H_{ijl})^{-1}P_{ijl}\rho_{ijl}.
 \end{aligned}$$

This may be written as a sum of four terms:

$$\begin{aligned}
 \text{I}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijl})^{-1}P_{ijl}\rho_{ijl}, \\
 \text{II}(z) &= -W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ij})^{-1}P_{ijl}\rho_{ijl}, \\
 \text{III}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\
 &\quad \times (V_{ik} + V_{jk})(z - H_{ijl})^{-1}P_{ijl}\rho_{ijl}, \\
 \text{IV}(z) &= -W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\
 &\quad \times (V_{ik} + V_{jk})(z - H_{ij})^{-1}P_{ijl}\rho_{ijl}. \\
 \text{I}(z) &= (W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}U_{ik})(W_{ik}P_{ijl}\rho_{ijl}^{-1})(\rho_{ijl}(z - H_{ijl})^{-1}\rho_{ijl}) \\
 &\quad + (W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}U_{jk})(W_{jk}P_{ijl}\rho_{ijl}^{-1})(\rho_{ijl}(z - H_{ijl})^{-1}\rho_{ijl}) \\
 &\quad - (W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}U_{ik})(W_{ik}P_{ijk}P_{ijl}\rho_{ijl}^{-1})(\rho_{ijl}(z - H_{ijl})^{-1}\rho_{ijl}) \\
 &\quad - (W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}U_{jk})(W_{jk}P_{ijk}P_{ijl}\rho_{ijl}^{-1})(\rho_{ijl}(z - H_{ijl})^{-1}\rho_{ijl}).
 \end{aligned}$$

Apply Lemmas V.4, V.8, and II.3.

III(z) and II(z) + IV(z) are well behaved for the same reasons that III(z) and II(z) + IV(z) in Proposition V.29 are well behaved. The only difference is that Proposition V.15 must be used in place of Proposition V.16. \square

PROPOSITION V.32. *Assume Hypotheses (1)–(3), and (5). Let i, j, k , and l be distinct.*

$$\begin{aligned}
 K_1(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\
 &\quad \times V_{il}(z - H_{il})^{-1}V_{jk}(z - H_{il,jk})^{-1}P_{il,jk}\rho_{il,jk}
 \end{aligned}$$

is well behaved and compact.

If Hypothesis (4) also holds, then the following are well behaved and compact:

$$\begin{aligned}
 K_2(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\
 &\quad \times [V_{il}(z - H_{il})^{-1}(V_{ik} + V_{lk}) + V_{kl}(z - H_{kl})^{-1}(V_{ik} + V_{il})] \\
 &\quad \times (z - H_{ilk})^{-1}P_{ilk}\rho_{ilk}. \\
 K_3(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{kl}(1 - P_{ij,kl})(z - H_{ij,kl})^{-1} \\
 &\quad \times [V_{il}(z - H_{il})^{-1}(V_{ik} + V_{lk}) + V_{ik}(z - H_{ik})^{-1}(V_{lk} + V_{il})] \\
 &\quad \times (z - H_{ilk})^{-1}P_{ilk}\rho_{ilk}.
 \end{aligned}$$

PROOF. Consider only $K_1(z)$; proofs for the others are similar. The extra assumption is required for $K_2(z)$ and $K_3(z)$ because they contain P_{ilk} rather than $P_{il,jk}$. Hypotheses (1)–(3), and (5) imply $E_{il,jk} < 0$; Hypotheses (3) and (4) require $E_{ilk} < 0$.

Compactness for $K_1(z)$ follows from Proposition V.15 for large negative $\text{Re } z$, since $W_{il}(z - H_{il})^{-1}V_{jk}(z - H_{il,jk})^{-1}P_{il,jk}\rho_{il,jk}$ is bounded for large negative $\text{Re } z$:

$$\begin{aligned}
K_1(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\
&\quad \times V_{il}(z - H_{iljk})^{-1}P_{iljk}\rho_{iljk} \\
&\quad - W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\
&\quad \times V_{il}(z - H_{il})^{-1}P_{iljk}\rho_{iljk}.
\end{aligned}$$

By Proposition V.17, the second term is well behaved. The first term equals

$$\begin{aligned}
&W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\
&\quad \times V_{il}(z - H_{iljk})^{-1}P_{iljk}\rho_{iljk} \\
&\quad + W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\
&\quad \times (V_{ij} + V_{ik})(z - H_{ijk})^{-1}V_{il}(z - H_{iljk})^{-1}P_{iljk}\rho_{iljk} \\
&= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{iljk})^{-1}P_{iljk}\rho_{iljk} \\
&\quad - W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1}P_{iljk}\rho_{iljk} \\
&\quad + W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\
&\quad \times (V_{ij} + V_{ik})(z - H_{iljk})^{-1}P_{iljk}\rho_{iljk} \\
&\quad - W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\
&\quad \times (V_{ij} + V_{ik})(z - H_{ijk})^{-1}P_{iljk}\rho_{iljk}.
\end{aligned}$$

This may be written as a sum of seven terms!

$$\begin{aligned}
\text{I}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{ik}(z - H_{iljk})^{-1}P_{iljk}\rho_{iljk}, \\
\text{II}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}V_{jk}(z - H_{iljk})^{-1}P_{iljk}\rho_{iljk}, \\
\text{III}(z) &= -W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})P_{ijk}(z - H_{iljk})^{-1}P_{iljk}\rho_{iljk}, \\
\text{IV}(z) &= -W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1}P_{iljk}\rho_{iljk}, \\
\text{V}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1} \\
&\quad \times (V_{ij} + V_{ik})(z - H_{iljk})^{-1}P_{iljk}\rho_{iljk}, \\
\text{VI}(z) &= -W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})^{-1}P_{iljk}\rho_{iljk}, \\
\text{VII}(z) &= W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}(V_{ik} + V_{jk})(1 - P_{ijk})(z - H_{ijk})P_{iljk}\rho_{iljk}, \\
\text{I}(z) &= (W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}U_{ik})(W_{ik}P_{iljk}\rho_{iljk}^{-1})(\rho_{iljk}(z - H_{iljk})^{-1}\rho_{iljk}).
\end{aligned}$$

Apply Lemmas V.4, V.8, and II.3:

$$\begin{aligned}
\text{II}(z) &= W_{ij}(1 - P_{ij})(z - H_0)^{-1}V_{jk}(z - H_{iljk})^{-1}P_{iljk}\rho_{iljk} \\
&\quad + (W_{ij}(1 - P_{ij})(z - H_{ij})^{-1}U_{ij}) \\
&\quad \times (W_{ij}(z - H_0)^{-1}V_{jk}(z - H_{iljk})^{-1}P_{iljk}\rho_{iljk}).
\end{aligned}$$

Lemma V.7 shows the first factor of the second term is well behaved.

$$(z - H_{il,jk})^{-1} P_{il,jk} = (z - E_{il,jk} - K_{il,jk})^{-1} P_{il,jk}$$

and $(z - E_{il,jk} - K_{il,jk})^{-1}$ commutes with V_{jk} . Moreover,

$$\begin{aligned} & (z - H_0)^{-1} (z - E_{il,jk} - K_{il,jk})^{-1} \\ &= \left[(z - H_0)^{-1} - (z - E_{il,jk} - K_{il,jk})^{-1} \right] (H_0^{(x_u)} + H_0^{(x_k)} - E_{il,jk})^{-1}. \end{aligned}$$

The second factor of the second term in the expression for $\text{II}(z)$ can now be written as:

$$\begin{aligned} & \left(W_{ij} \left[(z - H_0)^{-1} - (z - E_{il,jk} - K_{il,jk})^{-1} \right] \rho_{il,jk} \right) \\ & \quad \times \left((H_0^{(x_u)} + H_0^{(x_k)} - E_{il,jk})^{-1} U_{jk} \right) (W_{jk} P_{il,jk}). \end{aligned}$$

The methods of Lemma II.3 control the first factor; Corollary V.5 bounds the last factor; Theorem II.10 and Hypothesis (3) imply $E_{il,jk}$ is negative, so the middle factor is bounded (for K_2 and K_3 we need Hypothesis (4) at this point). The first term in the expression for $\text{II}(z)$ is treated by the same method.

$\text{III}(z)$ may be controlled by using Lemma V.4, the methods of Lemma II.3, and Corollary V.5.

$$\begin{aligned} \text{IV}(z) &= W_{ij} (1 - P_{ij}) (z - H_{ij})^{-1} (V_{ik} + V_{jk}) P_{ijk} (z - H_{jk})^{-1} P_{il,jk} \rho_{il,jk} \\ &\quad - W_{ij} (1 - P_{ij}) (z - H_{ij})^{-1} V_{ik} (z - H_{jk})^{-1} P_{il,jk} \rho_{il,jk} \\ &\quad - W_{ij} (1 - P_{ij}) (z - H_{ij})^{-1} V_{jk} (z - H_{jk})^{-1} P_{il,jk} \rho_{il,jk}. \end{aligned}$$

The first term may be controlled by the method used for $\text{III}(z)$, above. Lemma V.12 controls the remaining terms.

If Proposition V.16 is replaced by Proposition V.15, then the argument for $\text{III}(z)$ in Proposition V.29 shows that $\text{V}(z)$ is well behaved.

$\text{VI}(z)$ is well behaved by the proof of Proposition V.15.

$\text{VII}(z)$ is well behaved by Lemma V.12. \square

VI. Generic couplings. Hypotheses (1), (2), and (4) of §V hold for a large class of potentials (by Theorem II.10). However, Hypothesis (3) fails for some elements of any reasonable vector space of potentials. In this section, we show that Hypotheses (1) and (2) imply Hypothesis (3) for all couplings in the complement of a closed set of measure zero.

For the two body problem, Hypothesis (3) trivially holds, because there are no nontrivial cluster decompositions.

When $N = 3$, there are three nontrivial cluster decompositions, with cluster Hamiltonians $h_{ij} = h_0^{(i,j)} + \lambda_{ij} V_{ij}$. Hypothesis (3) states that 0 is not an eigenvalue or resonance energy for h_{ij} . This means 1 is not an eigenvalue of $\lambda_{ij} W_{ij} (0 - h_0)^{-1} U_{ij}$.

When $\lambda_{ij} = 0$, $\lambda_{ij} W_{ij} (0 - h_0)^{-1} U_{ij}$ is the zero operator. Moreover, $\lambda_{ij} W_{ij} (0 - h_0)^{-1} U_{ij}$ is a compact operator valued analytic function of λ_{ij} . So, the analytic Fredholm theorem shows that Hypothesis (3) fails only when λ_{ij} belongs to some discrete subset $\Lambda_{ij} \subseteq \mathbf{R}$.

Since $\{(\lambda_{12}, \lambda_{13}, \lambda_{23}) \in \mathbf{R}^3: \text{some } \lambda_{ij} \in \Lambda_{ij}\}$ is closed and has codimension 1, the set of couplings where Hypothesis (3) can fail is closed and has Lebesgue measure zero in \mathbf{R}^3 .

For $N = 4$, the situation is more complicated because Hypothesis (3) imposes the above conditions on the two body subsystems, as well as conditions on the three body subsystems. For each cluster Hamiltonian

$$h_{ijk} = h_0^{(i,j,k)} + \lambda_{ij} V_{ij} + \lambda_{ik} V_{ik} + \lambda_{jk} V_{jk},$$

we construct $M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, z)$, as in equation (IV.6). Hypothesis (3) requires that 1 not be an eigenvalue of $M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, z)$, whenever z is a threshold for h_{ijk} . $M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, z)$ is analytic in the λ variables, except when some λ_{nl} belongs to Λ_{nl} . At this point, M is discontinuous, and the argument used above for $N = 3$ is not valid.

For the nonzero thresholds (which must be negative), we can modify the argument so that it is valid in the four body case:

LEMMA VI.1. *Suppose $N = 4$ and Hypotheses (1) and (2) hold. Let $\Gamma_1 \subseteq \mathbf{R}^6$ be the set of couplings $\{\lambda_{ij}\}$, such that Hypothesis (3) fails for $H(\{\lambda_{ij}\}) = H_0 + \sum_{i < j} \lambda_{ij} V_{ij}$ at a negative energy threshold. Then the closure of Γ_1 has Lebesgue measure zero.*

PROOF. Since the two body subsystems have no negative thresholds, the conclusion of the lemma is a statement about the three body subsystems only.

If h_{ijk} is a three body cluster Hamiltonian, then a negative energy threshold E for h_{ijk} depends only on one coupling constant, which we may assume is λ_{ij} . Fix $\lambda_{ij} \notin \Lambda_{ij}$. The operator $M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, E(\lambda_{ij}) \pm i0)$ associated to h_{ijk} , has a compact square and is analytic for $(\lambda_{ik}, \lambda_{jk}) \in \Omega$, where

$$\Omega = \{(\lambda_{ik}, \lambda_{jk}) \in \mathbf{R}^2: \lambda_{ik} \notin \Lambda_{ik}, \lambda_{jk} \notin \Lambda_{jk}\}.$$

Ω is a union of open squares, and on each square Δ there are two possibilities:

- (i) $1 \notin \sigma(M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, E(\lambda_{ij}) \pm i0))$ for some $(\lambda_{ik}, \lambda_{jk}) \in \Delta$.
- (ii) $1 \in \sigma(M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, E(\lambda_{ij}) \pm i0))$ for all $(\lambda_{ik}, \lambda_{jk}) \in \Delta$.

When the first possibility occurs, the analytic Fredholm theorem shows that $1 \in \sigma(M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, E(\lambda_{ij}) \pm i0))$ only for those $(\lambda_{ik}, \lambda_{jk}) \in \Delta$, which belong to a measure zero union of analytic varieties. The union of the boundary of Δ and these varieties is closed in \mathbf{R}^2 .

Since h_{ijk} has finitely many thresholds (see Simon [29]), we see that the intersection of Γ_1 with any compact set is contained in a finite union of closed sets of measure zero, if possibility (ii) above, never occurs. Thus, it suffices to prove that there is no square Δ such that $M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, E(\lambda_{ij}) \pm i0)$ has eigenvalue 1 for all $(\lambda_{ik}, \lambda_{jk}) \in \Delta$.

$M(\lambda_{ij}, 0, 0, E(\lambda_{ij}) \pm i0) = 0$. So, if such a square Δ exists, we may assume (by changing squares if necessary) that Δ is adjacent to a square Δ_1 , on which possibility (i) occurs. Moreover, without loss, we may assume that it is possible to move from Δ_1 to Δ by increasing λ_{ik} .

Fix λ_{jk} so that $(\lambda_{ik}, \lambda_{jk}) \in \Delta_1$ for some λ_{ik} , and let $N(\lambda_{ik}, z) = M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, z)$, where λ_{ik} is chosen so that $(\lambda_{ik}, \lambda_{jk}) \in \Delta_1$. As we increase λ_{ik} , $N(\lambda_{ik}, z)$ is analytic

until λ_{ik} approaches $\omega \in \Lambda_{ik}$. At this point, $P_{ik}(\lambda_{ik})$ has a discontinuity. However, by representing $P_{ik}(\lambda_{ik})$ as a contour integral, we see that it has an analytic continuation $P'_{ik}(\lambda_{ik})$ from the region $\lambda_{ik} \in (\omega - \varepsilon_1, \omega)$ into the region $\lambda_{ik} \in (\omega - \varepsilon_1, \omega + \varepsilon_2)$, where $\varepsilon_1 > 0$, $\varepsilon_2 > 0$.

For $0 < \lambda_{ik} - \omega < \varepsilon_2$, $P_{ik}(\lambda_{ik}) - P'_{ik}(\lambda_{ik})$ is the projection onto the new eigenfunctions of $h_{ik}(\lambda_{ik}) \otimes 1$, which appear as λ_{ik} passes through ω . Let $\beta(\lambda_{ik})$ be the infimum of the corresponding new thresholds for h_{ijk} .

Replacing $P_{ik}(\lambda_{ik})$ by $P'_{ik}(\lambda_{ik})$ in Lemmas V.4, V.8, and V.11–V.13, we see that $N(\lambda_{ik}, z)$ has an analytic continuation into the region $\lambda_{ik} \in (\omega - \varepsilon_1, \omega - \varepsilon_2)$, as long as $z \notin [\beta(\lambda_{ik}), \infty)$. Since $E(\lambda_{ij}) < 0$ and $\beta(\omega) = 0$, $N(\lambda_{ik}, E(\lambda_{ij}) \pm i0)$ is analytic for $\omega - \varepsilon_1 < \lambda_{ik} < \omega + \varepsilon_3$, for some $\varepsilon_3 > 0$.

When $\omega < \lambda_{ik} < \omega + \varepsilon_3$, $(\lambda_{ik}, \lambda_{jk}) \in \Delta$. So, there exists a vector

$$\phi(\lambda_{ik}) = \begin{bmatrix} \phi_{ij,1}(\lambda_{ik}) \\ \phi_{ij,2}(\lambda_{ik}) \\ \phi_{ik,1}(\lambda_{ik}) \\ \phi_{ik,2}(\lambda_{ik}) \\ \phi_{jk,1}(\lambda_{ik}) \\ \phi_{jk,2}(\lambda_{ik}) \end{bmatrix},$$

such that $\phi(\lambda_{ik}) = M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, E(\lambda_{ij}) \pm i0)\phi(\lambda_{ik})$, whenever λ_{ik} belongs to some open subset of $(\omega, \omega + \varepsilon_3)$. Define

$$\psi(\lambda_{ik}) = \begin{bmatrix} \phi_{ij,1}(\lambda_{ik}) \\ \phi_{ij,2}(\lambda_{ik}) \\ [\phi_{ik,1}(\lambda_{ik}) + \lambda_{ik} W_{ik}(P_{ik}(\lambda_{ik}) \\ - P'_{ik}(\lambda_{ik})) G_{ik}(\lambda_{ik}) \rho_{ik} \phi_{ik,2}(\lambda_{ik})] \\ P'_{ik}(\lambda_{ik}) \phi_{ik,2}(\lambda_{ik}) \\ \phi_{jk,1}(\lambda_{ik}) \\ \phi_{jk,2}(\lambda_{ik}) \end{bmatrix}.$$

Then $\psi(\lambda_{ik}) = N(\lambda_{ik}, E(\lambda_{ij}) \pm i0)\psi(\lambda_{ik})$ for all λ_{ik} in the same open subset of $(\omega, \omega + \varepsilon_3)$.

The analytic Fredholm theorem now shows that $1 \in \sigma(N(\lambda_{ik}, E(\lambda_{ij}) \pm i0))$ for $\omega - \varepsilon_1 < \lambda_{ik} < \omega$. Since this is true for all choices of λ_{jk} , such that $(\lambda_{ik}, \lambda_{jk}) \in \Delta_1$ for some λ_{ik} , we conclude that $M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, E(\lambda_{ij}) \pm i0)$ has a constant eigenvalue 1 for all $(\lambda_{ik}, \lambda_{jk}) \in \Delta_1$. This contradicts the choice of Δ_1 , and the lemma is proved.

□

REMARK. The above argument fails for the zero energy threshold, because the new thresholds appear at zero when λ_{ik} passes ω . To avoid this problem, we move λ_{ik} around ω in the complex plane. The next lemma shows that we may do so without having the cuts of $\sigma_{\text{ess}}(h_{ijk})$ cross the negative real axis.

LEMMA VI.2. Assume Hypotheses (1) and (2) for the two body Hamiltonian $H(\lambda) = H_0 + \lambda V$. There is an open complex neighborhood X of the real axis, such that $\lambda \in X$ and $\text{Im } \lambda \neq 0$ imply that $H(\lambda)$ has no eigenvalues in the real interval $(-\infty, 0]$. Moreover, $\lambda \in X$ and $\text{Im } \lambda \neq 0$ imply $H(\lambda)$ has no zero energy resonances.

PROOF. The analytic Fredholm theorem shows that $1 \in \sigma(\lambda W(0 - H_0)^{-1}U)$ only when λ belongs to a discrete subset Λ of the complex plane. There exists an open neighborhood X of the real axis, such that $\Lambda \cap X \subseteq \mathbf{R}$. Since $H(\lambda)$ has a zero energy resonance or eigenvalue if and only if $\lambda \in \Lambda$, we need only show that $\text{Im } \lambda \neq 0$ implies $H(\lambda)$ has no eigenvalues in $(-\infty, 0)$.

Suppose $H(\lambda)\phi = E\phi$, where $\text{Im } \lambda \neq 0$ and $E < 0$.

$$\langle \phi, H_0 \phi \rangle + \lambda \langle \phi, V \phi \rangle = E \langle \phi, \phi \rangle < 0.$$

Since $\langle \phi, H_0 \phi \rangle \geq 0$, $\lambda \langle \phi, V \phi \rangle$ must be negative. Since $\langle \phi, V \phi \rangle \in \mathbf{R}$, $\text{Im } \lambda$ must be zero. This contradicts $\text{Im } \lambda \neq 0$. \square

PROPOSITION VI.3. Assume Hypotheses (1) and (2) for the $N \leq 4$ body Hamiltonian

$$H(\{\lambda_{ij}\}) = H_0 + \sum_{i < j} \lambda_{ij} V_{ij}.$$

Let Γ denote the set of all couplings $\{\lambda_{ij}\}$ such that Hypothesis (3) fails for $H(\{\lambda_{ij}\})$. Then Γ is closed and has Lebesgue measure zero.

PROOF. Γ is closed because eigenvalues and thresholds depend continuously on the couplings.

When $N = 2$ or 3 , the discussion at the beginning of this section proves the proposition. When $N = 4$, Lemma IV.1 shows that we need only study the threshold at zero energy.

The method of proof used for Lemma VI.1 may be applied to the zero energy threshold, with the following change. We cannot analytically continue $N(\lambda_{ik}, 0 \pm i0)$ through the point $\omega \in \Lambda_{ik}$. However, Lemmas V.4, V.8, and V.11–V.13, and VI.2 show that we may continue $N(\lambda_{ik}, 0 \pm i0)$ from the region $\omega - \varepsilon_1 < \lambda_{ik} < \omega$ ($\lambda_{ik} \in \mathbf{R}$) into the region

$$R = \{\lambda_{ik} = \omega + re^{i\theta} : 0 < r < \varepsilon, 0 < \arg \theta < 2\pi\}.$$

These lemmas also show that $M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, 0 + i0)$ and $M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, 0 - i0)$ have analytic continuations from the region $\omega < \lambda_{ik} < \omega + \delta$ into regions which intersect R in open subsets of the complex plane.

Using the fact that the analytically continued M operators have constant eigenvalue 1, we conclude that the analytically continued N operators have constant eigenvalue 1. (This is done as in Lemma VI.1, by explicitly constructing the eigenfunction of N from the eigenfunction of M , for λ_{ik} in the open set to which both operators have been continued.)

This argument shows that $M(\lambda_{ij}, \lambda_{ik}, \lambda_{jk}, 0 \pm i0)$ has constant eigenvalue 1 on the square Δ_1 , as in the proof of Lemma VI.1. This contradicts the choice of Δ_1 and proves the proposition. \square

REMARK. The coupling constants λ_{ij} and λ_{ik} are treated very differently in the proof of Lemma VI.1. So, our results say nothing about the case in which various coupling constants must be equal.

Appendix. Representative sample of the blocks of the matrices $A(z)$, $C(z)$, and $M(z)$ discussed in §IV.

The Matrix $A(z)$

$$A_{ij}(z) = \begin{bmatrix} G_o U_{ij} & G_o V_{ij} G_{ij} P_{ij} \rho_{ij} & 0 \end{bmatrix}$$

$$A_{ij,ijk}(z) = \begin{bmatrix} G_o U_{ij} & G_o V_{ij} G_{ij} P_{ij} \rho_{ij,k} & \frac{1}{3} G_o [V_{ij} G_{ij} (V_{ik} + V_{jk}) + V_{ik} G_{ik} (V_{ij} + V_{jk}) \\ & + V_{jk} G_{jk} (V_{ij} + V_{ik})] G_{ijk} P_{ijk} \rho_{ijk} \end{bmatrix}$$

$$A_{ij,(ij,k\ell)}(z) = \begin{bmatrix} G_o U_{ij} & G_o V_{ij} G_{ij} P_{ij} U_{k\ell} & \frac{1}{2} G_o [V_{ij} G_{ij} V_{k\ell} + V_{k\ell} G_{k\ell} V_{ij}] \\ & \times G_{ij,k\ell} P_{ij,k\ell} \rho_{ij,k\ell} \end{bmatrix}$$

The Matrix $C(z)$

$$C_{ij}(z) = \begin{bmatrix} W_{ij} G_{ij} (1 - P_{ij}) \\ \rho_{ij}^{-1} P_{ij} \\ 0 \end{bmatrix}$$

$$C_{ij,ijk}(z) = \begin{bmatrix} W_{ij} G_{ij} (1 - P_{ij}) (V_{ik} + V_{jk}) G_{ijk} (1 - P_{ijk}) \\ \rho_{ij,k}^{-1} P_{ij} (V_{ik} + V_{jk}) G_{ijk} (1 - P_{ijk}) \\ \rho_{ijk}^{-1} P_{ijk} \end{bmatrix}$$

$$C_{ij,(ij,k\ell)}(z) = \begin{bmatrix} W_{ij} G_{ij} (1 - P_{ij}) V_{k\ell} G_{ij,k\ell} (1 - P_{ij,k\ell}) \\ P_{ij} W_{k\ell} G_{ij,k\ell} (1 - P_{ij,k\ell}) \\ \rho_{ij,k\ell}^{-1} P_{ij,k\ell} \end{bmatrix}$$

12, 123; 14, 124

$w_{12} G_{12} (1-P_{12}) (V_{13}+V_{23}) G_{123} (1-P_{123}) U_{14}$ <p style="text-align: center;">V. 15</p>	$w_{12} G_{12} (1-P_{12}) (V_{13}+V_{23}) G_{123} \times$ $(1-P_{123}) V_{14} G_{14} P_{14} \rho_{14,2}$ <p style="text-align: center;">V. 17</p>	$\frac{1}{2} w_{12} G_{12} (1-P_{12}) (V_{13}+V_{23}) G_{123} (1-P_{123}) \times$ $[V_{14} G_{14} (V_{12}+V_{24}) + V_{24} G_{24} (V_{12}+V_{14})] \times$ $G_{124} P_{124} \rho_{124}$ <p style="text-align: center;">V. 31</p>
$\rho_{12,3}^{-1} P_{12} (V_{13}+V_{23}) G_{123} (1-P_{123}) U_{14}$ <p style="text-align: center;">V. 16</p>	$\rho_{12,3}^{-1} P_{12} (V_{13}+V_{23}) G_{123} (1-P_{123}) \times$ $V_{14} G_{14} P_{14} \rho_{14,2}$ <p style="text-align: center;">V. 18</p>	$\frac{1}{2} \rho_{12,3}^{-1} P_{12} (V_{13}+V_{23}) G_{123} (1-P_{123}) \times$ $[V_{14} G_{14} (V_{12}+V_{24}) + V_{24} G_{24} (V_{12}+V_{14})] \times$ $G_{124} P_{124} \rho_{124}$ <p style="text-align: center;">V. 29</p>
$\rho_{123}^{-1} P_{123} U_{14}$ <p style="text-align: center;">V. 4</p>	$\rho_{123}^{-1} P_{123} V_{14} G_{14} P_{14} \rho_{14,2}$ <p style="text-align: center;">V. 19</p>	$\rho_{123}^{-1} P_{123} V_{14} G_{14} (V_{12}+V_{24}) G_{124} P_{124} \rho_{124}$ <p style="text-align: center;">V. 20</p>

12, 123; 34, 134

$w_{12} G_{12} (1-P_{12}) (V_{13}+V_{23}) G_{123} (1-P_{123}) U_{34}$ <p style="text-align: center;">V. 15</p>	$w_{12} G_{12} (1-P_{12}) (V_{13}+V_{23}) G_{123} (1-P_{123}) \times$ $V_{34} G_{34} P_{34} \rho_{34,1}$ <p style="text-align: center;">V. 17</p>	$\frac{1}{2} w_{12} G_{12} (1-P_{12}) (V_{13}+V_{23}) G_{123} (1-P_{123}) \times$ $[V_{34} G_{34} (V_{13}+V_{14}) + V_{14} G_{14} (V_{13}+V_{34})] \times$ $G_{134} P_{134} \rho_{134}$ <p style="text-align: center;">V. 32</p>
$\rho_{12,3}^{-1} P_{12} (V_{13}+V_{23}) G_{123} (1-P_{123}) U_{34}$ <p style="text-align: center;">V. 16</p>	$\rho_{12,3}^{-1} P_{12} (V_{13}+V_{23}) G_{123} (1-P_{123}) \times$ $V_{34} G_{34} P_{34} \rho_{34,1}$ <p style="text-align: center;">V. 18</p>	$\frac{1}{2} \rho_{12,3}^{-1} P_{12} (V_{13}+V_{23}) G_{123} (1-P_{123}) \times$ $[V_{34} G_{34} (V_{13}+V_{14}) + V_{14} G_{14} (V_{13}+V_{34})] \times$ $G_{134} P_{134} \rho_{134}$ <p style="text-align: center;">V. 29</p>
$\rho_{123}^{-1} P_{123} U_{34}$ <p style="text-align: center;">V. 4</p>	$\rho_{123}^{-1} P_{123} V_{34} G_{34} P_{34} \rho_{34,1}$ <p style="text-align: center;">V. 19</p>	$\rho_{123}^{-1} P_{123} V_{34} G_{34} (V_{13}+V_{14}) G_{134} P_{134} \rho_{134}$ <p style="text-align: center;">V. 20</p>

12, 123; 14, 134

$W_{12}G_{12}(1-P_{12})(V_{13}+V_{23})G_{123}(1-P_{123})U_{14}$	$W_{12}G_{12}(1-P_{12})(V_{13}+V_{23})G_{123}(1-P_{123}) \times$ $V_{14}G_{14}P_{14}\rho_{14,3}$	$\frac{1}{2}W_{12}G_{12}(1-P_{12})(V_{13}+V_{23})G_{123}(1-P_{123}) \times$ $[V_{14}G_{14}(V_{13}+V_{34})+V_{34}G_{34}(V_{13}+V_{14})] \times$ $G_{134}P_{134}\rho_{134}$
V. 15	V. 17	V. 32
$\rho_{12,3}^{-1}P_{12}(V_{13}+V_{23})G_{123}(1-P_{123})U_{14}$	$\rho_{12,3}^{-1}P_{12}(V_{13}+V_{23})G_{123}(1-P_{123}) \times$ $V_{14}G_{14}P_{14}\rho_{14,3}$	$\frac{1}{2}\rho_{12,3}^{-1}P_{12}(V_{13}+V_{23})G_{123}(1-P_{123}) \times$ $[V_{14}G_{14}(V_{13}+V_{34})+V_{34}G_{34}(V_{13}+V_{14})] \times$ $G_{134}P_{134}\rho_{134}$
V. 16	V. 18	V. 29
$\rho_{123}^{-1}P_{123}U_{14}$	$\rho_{123}^{-1}P_{123}V_{14}G_{14}P_{14}\rho_{14,3}$	$\rho_{123}^{-1}P_{123}V_{14}G_{14}(V_{13}+V_{34})G_{134}P_{134}\rho_{134}$
V. 4	V. 19	V. 20

12, (12, 34); 13, (13, 24)

$W_{12}G_{12}(1-P_{12})V_{34}G_{12,34}(1-P_{12,34})U_{13}$	$W_{12}G_{12}(1-P_{12})V_{34}G_{12,34}(1-P_{12,34}) \times$ $V_{13}G_{13}P_{13}U_{24}$	$\frac{1}{2}W_{12}G_{12}(1-P_{12})V_{34}G_{12,34}(1-P_{12,34}) \times$ $[V_{13}G_{13}V_{24}+V_{24}G_{24}V_{13}] \times$ $G_{13,24}P_{13,24}\rho_{13,24}$
V. 25	V. 27	V. 31
$P_{12}W_{34}G_{12,34}(1-P_{12,34})U_{13}$	$P_{12}W_{34}G_{12,34}(1-P_{12,34})V_{13}G_{13}P_{13}U_{24}$	$\frac{1}{2}P_{12}W_{34}G_{12,34}(1-P_{12,34}) \times$ $[V_{13}G_{13}V_{24}+V_{24}G_{24}V_{13}] \times$ $G_{13,24}P_{13,24}\rho_{13,24}$
V. 26	V. 28	V. 30
$\rho_{12,34}^{-1}P_{12,34}U_{13}$	$\rho_{12,34}^{-1}P_{12,34}V_{13}G_{13}P_{13}U_{24}$	$\rho_{12,34}^{-1}P_{12,34}V_{13}G_{13}V_{24}G_{13,24}P_{13,24}\rho_{13,24}$
V. 4	V. 19	V. 20

12, 123; 14, (14,23)

$w_{12}G_{12}(1-P_{12})(V_{13}+V_{23})G_{123}(1-P_{123})U_{14}$	$w_{12}G_{12}(1-P_{12})(V_{13}+V_{23})G_{123}(1-P_{123}) \times$	$w_{12}G_{12}(1-P_{12})(V_{13}+V_{23})G_{123}(1-P_{123}) \times$
$V_{14}G_{14}P_{14}U_{23}$		$V_{14}G_{14}V_{23}G_{14,23}P_{14,23}P_{14,23}$
V. 15	V. 17	V. 32
$\rho_{12,3}^{-1}P_{12}(V_{13}+V_{23})G_{123}(1-P_{123})U_{14}$	$\rho_{12,3}^{-1}P_{12}(V_{13}+V_{23})G_{123}(1-P_{123}) \times$	$\rho_{12,3}^{-1}P_{12}(V_{13}+V_{23})G_{123}(1-P_{123}) \times$
$V_{14}G_{14}P_{14}U_{23}$		$V_{14}G_{14}V_{23}G_{14,23}P_{14,23}P_{14,23}$
V. 16	V. 18	V. 29
$\rho_{123}^{-1}P_{123}U_{14}$	$\rho_{123}^{-1}P_{123}V_{14}G_{14}P_{14}U_{23}$	$\rho_{123}^{-1}P_{123}V_{14}G_{14}V_{23}G_{14,23}P_{14,23}P_{14,23}$
V. 4	V. 19	V. 20

12, 123; 34, (12,34)

$w_{12}G_{12}(1-P_{12})(V_{13}+V_{23})G_{123}(1-P_{123})U_{34}$	$w_{12}G_{12}(1-P_{12})(V_{13}+V_{23})G_{123}(1-P_{123}) \times$	$w_{12}G_{12}(1-P_{12})(V_{13}+V_{23})G_{123}(1-P_{123}) \times$
	$V_{34}G_{34}P_{34}U_{12}$	$V_{34}G_{34}V_{12}G_{12,34}P_{12,34}P_{12,34}$
V. 15	V. 17	V. 31
$\rho_{12,3}^{-1}P_{12}(V_{13}+V_{23})G_{123}(1-P_{123})U_{34}$	$\rho_{12,3}^{-1}P_{12}(V_{13}+V_{23})G_{123}(1-P_{123}) \times$	$\rho_{12,3}^{-1}P_{12}(V_{13}+V_{23})G_{123}(1-P_{123}) \times$
	$V_{34}G_{34}P_{34}U_{12}$	$V_{34}G_{34}V_{12}G_{12,34}P_{12,34}P_{12,34}$
V. 16	V. 18	V. 29
$\rho_{123}^{-1}P_{123}U_{34}$	$\rho_{123}^{-1}P_{123}V_{34}G_{34}P_{34}U_{12}$	$\rho_{123}^{-1}P_{123}V_{34}G_{34}V_{12}G_{12,34}P_{12,34}P_{12,34}$
V. 4	V. 19	V. 20

12, (12, 34); 13, 123

$W_{12}G_{12}(1-P_{12})V_{34}G_{12,34}(1-P_{12,34})U_{13}$	$W_{12}G_{12}(1-P_{12})V_{34}G_{12,34}(1-P_{12,34}) \times$ $V_{13}G_{13}P_{13}^0_{13,2}$	$\frac{1}{2}W_{12}G_{12}(1-P_{12})V_{34}G_{12,34}(1-P_{12,34}) \times$ $[V_{13}G_{13}(V_{12}+V_{23})+V_{23}G_{23}(V_{12}+V_{13})] \times$ $G_{123}P_{123}^0_{123}$
V. 25	V. 27	V. 31
$P_{12}W_{34}G_{12,34}(1-P_{12,34})U_{13}$	$P_{12}W_{34}G_{12,34})V_{13}G_{13}P_{13}^0_{13,2}$	$\frac{1}{2}P_{12}W_{34}G_{12,34}(1-P_{12,34}) \times$ $[V_{13}G_{13}(V_{12}+V_{23})+V_{23}G_{23}(V_{12}+V_{13})] \times$ $G_{123}P_{123}^0_{123}$
V. 26	V. 28	V. 30
$\rho_{12,34}^{-1}P_{12,34}U_{13}$	$\rho_{12,34}^{-1}P_{12,34}V_{13}G_{13}P_{13}^0_{13,2}$	$\rho_{12,34}^{-1}P_{12,34}V_{13}G_{13}(V_{12}+V_{23})G_{123}P_{123}^0_{123}$
V. 4	V. 19	V. 20

12, (12, 34); 13, 134

$W_{12}G_{12}(1-P_{12})V_{34}G_{12,34}(1-P_{12,34})U_{13}$	$W_{12}G_{12}(1-P_{12})V_{34}G_{12,34}(1-P_{12,34}) \times$ $V_{13}G_{13}P_{13}^0_{13,4}$	$\frac{1}{2}W_{12}G_{12}(1-P_{12})V_{34}G_{12,34}(1-P_{12,34}) \times$ $[V_{13}G_{13}(V_{14}+V_{34})+V_{34}G_{34}(V_{14}+V_{13})] \times$ $G_{134}P_{134}^0_{134}$
V. 25	V. 27	V. 32
$P_{12}W_{34}G_{12,34}(1-P_{12,34})U_{13}$	$P_{12}W_{34}G_{12,34}(1-P_{12,34})V_{13}G_{13}P_{13}^0_{13,4}$	$\frac{1}{2}P_{12}W_{34}G_{12,34}(1-P_{12,34}) \times$ $[V_{13}G_{13}(V_{14}+V_{34})+V_{34}G_{34}(V_{14}+V_{13})] \times$ $G_{134}P_{134}^0_{134}$
V. 26	V. 28	V. 30
$\rho_{12,34}^{-1}P_{12,34}U_{13}$	$\rho_{12,34}^{-1}P_{12,34}V_{13}G_{13}P_{13}^0_{13,4}$	$\rho_{12,34}^{-1}P_{12,34}V_{13}G_{13}(V_{14}+V_{34})G_{134}P_{134}^0_{134}$
V. 4	V. 19	V. 20

12, 123; 14

$w_{12} G_{12} (1-P_{12}) (V_{13}+V_{23}) G_{123} (1-P_{123}) U_{14}$	$w_{12} G_{12} (1-P_{12}) (V_{13}+V_{23}) G_{123} (1-P_{123}) V_{14} G_{14} P_{14} \rho_{14}$	0
V.15	V.17	
$\rho_{12,3}^{-1} P_{12} (V_{13}+V_{23}) G_{123} (1-P_{123}) U_{14}$	$\rho_{12,3}^{-1} P_{12} (V_{13}+V_{23}) G_{123} (1-P_{123}) V_{14} G_{14} P_{14} \rho_{14}$	0
V.16	V.18	
$\rho_{123}^{-1} P_{123} U_{14}$	$\rho_{123}^{-1} P_{123} V_{14} G_{14} P_{14} \rho_{14}$	0
V.4	V.19	

12, 123; 34

$w_{12} G_{12} (1-P_{12}) (V_{13}+V_{23}) G_{123} (1-P_{123}) U_{34}$	$w_{12} G_{12} (1-P_{12}) (V_{13}+V_{23}) G_{123} (1-P_{123}) V_{34} G_{34} P_{34} \rho_{34}$	0
V.15	V.17	
$\rho_{12,3}^{-1} P_{12} (V_{13}+V_{23}) G_{123} (1-P_{123}) U_{34}$	$\rho_{12,3}^{-1} P_{12} (V_{13}+V_{23}) G_{123} (1-P_{123}) V_{34} G_{34} P_{34} \rho_{34}$	0
V.16	V.18	
$\rho_{123}^{-1} P_{123} U_{34}$	$\rho_{123}^{-1} P_{123} V_{34} G_{34} P_{34} \rho_{34}$	0
V.4	V.19	

$$\frac{1}{2}, \frac{(1}{2}, \frac{5}{4}); \frac{1}{2}, \frac{1}{2}$$

$W_{12}G_{12}(1-P_{12})V_{34}G_{12,34}(1-P_{12,34})U_{13}$	$W_{12}G_{12}(1-P_{12})V_{34}G_{12,34}(1-P_{12,34})V_{13}G_{13}P_{13}P_{13}$	0
V. 25	V. 27	
$P_{12}W_{34}G_{12,34}(1-P_{12,34})U_{13}$	$P_{12}W_{34}G_{12,34}(1-P_{12,34})V_{13}G_{13}P_{13}P_{13}$	0
V. 26	V. 28	
$\rho_{12,34}^{-1}P_{12,34}U_{13}$	$\rho_{12,34}^{-1}P_{12,34}V_{13}G_{13}P_{13}P_{13}$	0
V. 4	V. 19	

REFERENCES

1. S. Agmon, *Spectral properties of Schrödinger operators and scattering theory*, Ann. Scuola Norm. Sup. Pisa (IV) **2** (1975), 151–218.
2. E. Balslev, *Absence of positive eigenvalues of Schrödinger operators*, Arch. Rational Mech. Anal. **59** (1975), 343–357.
3. E. Balslev and J. M. Combes, *Spectral properties of many-body Schrödinger operators with dilatation-analytic interactions*, Comm. Math. Phys. **22** (1971), 280–294.
4. J. M. Combes and L. Thomas, *Asymptotic behavior of Eigenfunctions for multiparticle Schrödinger operators*, Comm. Math. Phys. **34** (1973), 251–270.
5. J. Cook, *Convergence of the Möller wave matrix*, J. Math. Phys. **36** (1957), 82–87.
6. L. D. Faddeev, *Mathematical aspects of the three-body problem in the quantum scattering theory*, Israel Program for Scientific Translations, Jerusalem, 1965.
7. J. Ginibre and M. Moulin, *Hilbert space approach to the quantum mechanical three-body problem*, Ann. Inst. H. Poincaré Sect. A **21** (1974), 97–145.
8. U. Greifenegger, K. Jörgens, J. Weidmann and M. Winkler, *Streutheorie für Schrödinger Operatoren*, preprint, 1972.
9. G. A. Hagedorn, *Asymptotic completeness for a class of four particle Schrödinger operators*, Bull. Amer. Math. Soc. **84** (1978), 155–156.
10. K. Hepp, *On the quantum mechanical N-body problem*, Helv. Phys. Acta **42** (1969), 425–458.
11. J. S. Howland, *Abstract stationary theory of multichannel scattering*, J. Functional Analysis **22** (1976), 250–282.
12. T. Ikebe, *Eigenfunctions associated with the Schrödinger operators and their applications to scattering theory*, Arch. Rational Mech. Anal. **5** (1960), 1–34.
13. R. J. Iorio and M. O'Carroll, *Asymptotic completeness for multi-particle Schrödinger Hamiltonians with weak potentials*, Comm. Math. Phys. **27** (1972), 137–145.
14. T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, Berlin-Heidelberg-New York, 1966.
15. ———, *Wave operators and similarity for some non-selfadjoint operators*, Math. Ann. **162** (1966), 258–279.

16. S. T. Kuroda, *An abstract stationary approach to perturbation of continuous spectra and scattering theory*, J. Analyse Math. **20** (1967), 57–117.
17. ———, *Scattering theory for differential operators. II. Self-adjoint elliptic operators*, J. Math. Soc. Japan **25** (1973), 222–234.
18. R. B. Lavine, *Commutators and scattering theory. I. Repulsive interactions*, Comm. Math. Phys. **20** (1971), 301–323.
19. E. Mourre, *Applications de la méthode de Lavine au problème à trois corps*, Ann. Inst. H. Poincaré Sect. A **26** (1977), 219–262.
20. I. M. Narodetskii and O. A. Yakubovskii, *Two cluster approximation in the integral equation for N particles*, Soviet J. Nuclear Phys. **14** (1972), 178–184.
21. A. J. O'Connor, *Exponential decay of bound state wave functions*, Comm. Math. Phys. **32** (1973), 319–340.
22. M. Reed and B. Simon, *Methods of modern mathematical physics, Vol. II: Fourier analysis, self-adjointness*, Academic Press, New York-London, 1975.
23. ———, *Methods of modern mathematical physics, Vol. III: Scattering theory*, Academic Press, New York-London, 1979.
24. ———, *Methods of modern mathematical physics, Vol. IV: Analysis of operators*, Academic Press, New York-London, 1978.
25. M. Schechter, *A new criterion for scattering theory*, Preprint, Yeshiva University, 1976.
26. I. M. Sigal, *Mathematical foundations of quantum scattering theory for multiparticle systems*, Mem. Amer. Math. Soc. No. 209 **16** (1978).
27. ———, *On the point spectrum of the Schrödinger operators of multiparticle systems*, Comm. Math. Phys. **48** (1976), 155–164.
28. ———, *On quantum mechanics of many-body systems with dilation-analytic potentials*, Bull. Amer. Math. Soc. **84** (1978), 152–154.
29. B. Simon, *Quantum mechanics for Hamiltonians defined as quadratic forms*, Princeton University Press, 1971.
30. ———, *Quadratic form techniques and the Balslev-Combes theorem*, Comm. Math. Phys. **27** (1972), 1–9.
31. ———, *Resonances in n -body quantum systems with dilation analytic potentials and the foundations of time-dependent perturbation theory*, Ann. of Math. **97** (1973), 247–274.
32. ———, *Absence of positive eigenvalues in a class of multiparticle quantum systems*, Math. Ann. **207** (1974), 133–138.
33. ———, *Scattering theory and quadratic forms: On a theorem of Schechter*, Comm. Math. Phys. **53** (1977), 151–153.
34. ———, *Geometric methods in multiparticle quantum systems*, Comm. Math. Phys. **55** (1977), 259–274.
35. E. M. Stein, *Interpolation of linear operators*, Trans. Amer. Math. Soc. **83** (1956), 482–492.
36. L. E. Thomas, *Asymptotic completeness in two- and three-particle quantum mechanical scattering*, Ann. Physics **90** (1975), 127–165.
37. E. C. Titchmarsh, *Introduction to the theory of Fourier integrals*, Oxford Press, London, 1950.
38. C. van Winter, *Complex dynamical variables for multiparticle systems and analytic interactions. I*, J. Math. Anal. Appl. **47** (1974), 633–670.
39. ———, *Complex dynamical variables for multiparticle systems with analytic interactions. II*, J. Math. Anal. Appl. **48** (1974), 368–399.
40. S. Weinberg, *Systematic solution of multiparticle scattering problems*, Phys. Rev. B **133** (1964), 232–256.
41. O. A. Yakubovskii, *On the integral equations in the theory of N particle scattering*, Soviet J. Nuclear Phys. **5** (1967), 937–942.
42. V. Enss, *Asymptotic completeness for quantum mechanical potential scattering*, Comm. Math. Phys. (to appear).
43. C. van Winter, *Projection operators associated with analytic multiparticle Hamiltonians*, Notices Amer. Math. Soc. **25** (1978), A-320.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY 08540

Current address: Department of Mathematical Physics, The Rockefeller University, New York, New York 10021