

CONVEXITY AND TIGHTNESS FOR RESTRICTIONS OF HAMILTONIAN FUNCTIONS TO FIXED POINT SETS OF AN ANTISYMPLECTIC INVOLUTION

BY

J. J. DUISTERMAAT¹

ABSTRACT. The Kostant convexity theorem for real flag manifolds is generalized to a Hamiltonian framework. More precisely, it is proved that if f is the momentum mapping for a Hamiltonian torus action on a symplectic manifold M and Q is the fixed point set of an antisymplectic involution of M leaving f invariant, then $f(Q) = f(M)$ is a convex polytope. Also it is proved that the coordinate functions of f are tight, using "half-turn" involutions of Q .

1. Introduction. Recently the theorem of Kostant [10], on convexity of certain projections of complex flag manifolds, has been generalized to a Hamiltonian framework by Guillemin and Sternberg [7], and, independently, by Atiyah [2]. However, Kostant proved his theorem for the real flag manifolds as well, and it is the first purpose of this paper to show that also this real version has a generalization in a Hamiltonian setting. More specifically, let M be a compact connected smooth manifold of dimension $2n$, provided with a symplectic form σ . Let T be a torus acting on M in a Hamiltonian way, with Hamiltonian functions f_X , $X \in \mathfrak{t}$, and momentum mapping $f: M \rightarrow \mathfrak{t}^*$. Here \mathfrak{t} is the Lie algebra of T and \mathfrak{t}^* its dual, see §2 for more details. Furthermore, let τ be a smooth involution of M such that $\tau^*\sigma = -\sigma$ and such that $f_X \circ \tau = f_X$ for all $X \in \mathfrak{t}$. (One can always arrange the latter condition by passing to a suitable subtorus T_0 of T , the new momentum mapping then being equal to f followed by the natural projection $\mathfrak{t}^* \rightarrow \mathfrak{t}_0^*$.) Let Q be the fixed point set of τ , which we assume to be nonvoid. Then $f(Q) = f(M)$, the convex hull of finitely many points in \mathfrak{t}^* . For a more detailed description of the extremal points of $f(Q) = f(M)$, see Theorem 2.5 and formula (2.32). The proof follows the pattern of Guillemin and Sternberg [7], which in turn was inspired by Heckman [8].

If M is a Kähler manifold, then a more refined result is true in terms of gradient flows, this will be discussed in §4.

Secondly, Atiyah [2], following Frankel [6], observed that the Hamiltonian functions f_X , $X \in \mathfrak{t}$, are tight in the sense that the sum of the Betti numbers of M is equal to the sum of the Betti numbers of the critical set of f_X . In the case of isolated critical points this implies that f_X is a Morse function on M with the minimal number of critical points. This too generalizes a known result for complex flag manifolds, but

Received by the editors December 7, 1981 and, in revised form, January 21, 1982.

1980 *Mathematics Subject Classification*. Primary 53C15; Secondary 53C55, 58F05, 55M20.

¹ Partially supported by NSF Grant MCS 80-23356, while visiting the University of California at Berkeley.

© 1983 American Mathematical Society
0002-9947/82/0000-0437/\$04.50

which actually is true for the real flag manifolds as well, due to Takeuchi and Kobayashi [13], see also Duistermaat, Kolk and Varadarajan [3, §4]. In §3 it will be shown that also the tightness generalizes to our setting. The theorem is that $\dim H^*(Q; \mathbb{Z}/2\mathbb{Z}) = \dim H^*(C_X; \mathbb{Z}/2\mathbb{Z})$ if C_X is the critical set of the function $f_X|_Q$, $X \in \mathfrak{t}$.

Finally in §5 we describe how the real flag manifolds fit into the framework described above.

It is a pleasure for me to thank Michael Atiyah for his suggestion, made to me at the Arbeitstagung in Bonn in June 1981 (and in [2]), that the real flag manifolds really should be treated as the fixed point set of an involution in a symplectic manifold. I also thank John Millson, resp. Alan Weinstein for some discussions at UCLA, resp. Berkeley, which stimulated me further.

2. Convexity. We recall that (M, σ) is a compact connected symplectic manifold with a Hamiltonian action of a torus T on it. That is, there is a linear map $X \mapsto f_X$ from the Lie algebra \mathfrak{t} of T to the space of smooth functions on M , such that

(2.1) For each $X \in \mathfrak{t}$, the infinitesimal action \tilde{X} of X on M is equal to the Hamilton vector field of the function f_X , and

(2.2) The functions f_X , $X \in \mathfrak{t}$, are in involution.

In formula, (2.1) reads

$$(2.1') \quad \tilde{X} \lrcorner \sigma = -df_X, \quad X \in \mathfrak{t},$$

whereas assuming this, (2.2) is equivalent to

$$(2.2') \quad \tilde{X} \lrcorner df_Y = 0 \quad \text{for all } X, Y \in \mathfrak{t},$$

that is, f_Y is constant along the T -orbits in M .

The mapping $f: M \rightarrow \mathfrak{t}^*$, defined by

$$(2.3) \quad \langle X, f(m) \rangle = f_X(m), \quad m \in M, X \in \mathfrak{t},$$

is called the *momentum mapping* of the Hamiltonian T -action.

The next ingredient which we introduce is a smooth map $\tau: M \rightarrow M$ which is an *involution* of M , that is

$$(2.4) \quad \tau \circ \tau = \text{identity on } M,$$

and which is *antisymplectic*, that is

$$(2.5) \quad \tau^* \sigma = -\sigma.$$

We will assume that the Hamiltonian T -action and the involution τ are related to each other by the condition that the functions f_X are τ -invariant, that is

$$(2.6) \quad \tau^* f_X = f_X \quad \text{for all } X \in \mathfrak{t}.$$

Assuming (2.5), this is equivalent to

$$(2.7) \quad \tau^* \tilde{X} = -\tilde{X}, \quad X \in \mathfrak{t},$$

in view of (2.1'). In turn, (2.7) is equivalent to

$$(2.8) \quad \tau \tilde{g} \tau^{-1} = \tilde{g}^{-1}, \quad g \in T,$$

if \tilde{g} denotes the action of g on M . That is, τ maps T -orbits to T -orbits, but at the same time reverses the time on the orbits of the 1-parameter subgroups of T .

If G is a compact group acting by smooth mappings on a manifold M , then averaging over G of an arbitrary Riemannian metric on M leads to a G -invariant Riemannian metric β on M . If m is a fixed point for the G -action, then the exponential map, centered at m , with respect to β , intertwines the linear action of G on $T_m M$ with the local action of G around m . That is, the G -action is linear orthogonal on suitable local coordinates. In particular the fixed point set for the action of any subset of G has finitely many components, each of which is a closed smooth submanifold of M .

We assume that the fixed point set of τ in M ,

$$(2.9) \quad Q = \{m \in M; \tau(m) = m\},$$

is nonvoid. It has finitely many smooth, compact, connected components. From (2.5) we obtain that if $m \in Q$, then both $T_m M = \text{Ker}(D\tau(m) - I)$ and

$$(2.10) \quad P_m = \text{Ker}(D\tau(m) + I)$$

are isotropic subspaces for σ_m . Because also

$$(2.11) \quad T_m M = T_m Q \oplus P_m,$$

the conclusion is that $T_m Q$ and P_m are Lagrange subspaces of $T_m M$. That is, Q is a Lagrange submanifold of M , and the P_m , $m \in Q$, form a Lagrange subbundle of $T_Q M$, complementary to TQ . In fact, Meyer [11] showed that there is an open τ -invariant neighborhood U of Q in M and a symplectic diffeomorphism Φ from U onto an open neighborhood V of the zero section in T^*Q , such that $\Phi\tau\Phi^{-1}$ maps $p \in (T_q Q)^* \cap V$ to $-p$, for all $q \in Q$.

Another rigidity property of compact Lie group actions on a compact manifold M is that there are only finitely many orbit types (Mostow [12], Yang [15]). In the case of the action of a torus T this means that there are only finitely many possibilities for the stabilizer groups

$$(2.12) \quad T_m = \{g \in T; g(m) = m\},$$

as m ranges over M . This fact will be used both in the proof of the convexity and of the tightness theorem mentioned in the introduction.

2.1. LEMMA. *Let $m \in Q$, $X \in \mathfrak{t}$. Then $d(f_X|Q)(m) = 0$ implies that $df_X(m) = 0$, which in turn is equivalent to the condition that m is a fixed point for the action of $\exp tX$, $t \in \mathbf{R}$.*

PROOF. $T_m Q = \text{Ker}(D\tau(m) - I) = \{u + D\tau(m)(u); u \in T_m M\}$. So using (2.6), we get $0 = df_X(m)(u + D\tau(m)(u)) = 2df_X(m)(u)$ for all $u \in T_m M$.

2.2. PROPOSITION. *Let $m \in Q$, $D(f|Q)(m) = 0$. Then there exist smooth symplectic local coordinates q_j, p_j for a neighborhood U of m (zero at m), such that, for suitable $\omega_j \in \mathfrak{t}^*$,*

$$(2.13) \quad f_X = f_X(m) + \sum_{j=1}^n \omega_j(X) \cdot (q_j^2 + p_j^2)/2,$$

and moreover

$$(2.14) \quad \tau: (q, p) \mapsto (q, -p)$$

in U . In particular $U \cap Q = \{p = 0\}$.

PROOF. Although the proof is a combination of the by now standard arguments in Meyer [11] and Guillemin-Sternberg [7, §4], we will give it in some detail for the convenience of the reader. We begin with the result on the tangent level.

In view of Lemma 2.1, m is not only a fixed point for τ , but also for the T -action. Then the

$$(2.15) \quad X_m = D\tilde{X}(m): T_m M \rightarrow T_m M, \quad X \in \mathfrak{t},$$

form a commuting family of infinitesimally symplectic transformations, antisymmetric with respect to β_m . In particular they are simultaneously diagonalizable over \mathbb{C} with purely imaginary eigenvalues. That is,

$$(2.16) \quad T_m M \otimes \mathbb{C} = \sum_{\lambda}^{\oplus} E_{\lambda}, \quad X_m|_{E_{\lambda}} = \lambda(X) \cdot \text{identity on } E_{\lambda};$$

here X_m is extended to a complex linear endomorphism of $T_m M \otimes \mathbb{C}$ and λ is real linear: $\mathfrak{t} \rightarrow \mathbb{C}$, taking only purely imaginary values. The complex conjugation maps E_{λ} to $E_{\bar{\lambda}} = E_{-\lambda}$. Because the X_m are infinitesimally symplectic, the spaces

$$(2.17) \quad E_{\lambda, -\lambda} = (E_{\lambda} + E_{-\lambda}) \cap T_m M$$

are mutually σ_m -orthogonal. Because they span $T_m M$, they form a symplectic vector space decomposition of $T_m M$.

Now (2.7) implies that

$$(2.18) \quad D\tau(m) \circ X_m = -X_m \circ D\tau(m),$$

showing that $D\tau(m)$ maps E_{λ} to $E_{-\lambda}$, so it leaves $E_{\lambda, -\lambda}$ invariant. This reduces the problem to the case that all X_m are real multiples of one infinitesimally symplectic mapping J , the square of which we can take equal to $-I$. Moreover, J maps $T_m Q$ to P_m and P_m to $T_m Q$. Now

$$(2.19) \quad \langle u, v \rangle = \sigma_m(u, Jv), \quad u, v \in T_m Q,$$

defines a nondegenerate symmetric bilinear form on $T_m Q$. We can write $T_m Q = Q_m^+ \oplus Q_m^-$, $\langle Q_m^+, Q_m^- \rangle = 0$, with (2.19) being positive, resp. negative definite, on Q_m^+ , resp. Q_m^- . The spaces $Q_m^+ + J(Q_m^+)$ and $Q_m^- + J(Q_m^-)$ are σ_m -orthogonal to each other, leading to a $D\tau(m)$ -invariant symplectic vector space decomposition. This reduces the problem to the case that (2.19) is definite.

In the positive case, let e_1, \dots, e_n be an orthonormal basis of $T_m Q$ with respect to (2.19). Then

$$(2.20) \quad (q, p) \mapsto \sum_j (q_j \cdot e_j - p_j \cdot J e_j)$$

is a symplectic mapping, with respect to the symplectic form $\sum_j dp_j \wedge dq_j$ on (q, p) -space. It intertwines J with the mapping $(q, p) \mapsto (p, -q)$, which, as a linear vector field, is Hamiltonian with Hamilton function

$$(2.21) \quad \sum_j (q_j^2 + p_j^2)/2.$$

In the negative definite case we get (2.21) with a minus sign in front. This proves the proposition on the tangent level.

For the *local* normal form we already know that there are local coordinates linearizing τ and the T -action, hence also the vector fields \tilde{X} , $X \in \mathfrak{t}$. However, in these coordinates the symplectic form σ will in general not be equal to the constant symplectic form $\sigma_m = \sum_j dp_j \wedge dq_j$. The proof is finished by applying the following version (also known to Weinstein) of the equivariant Darboux lemma of Weinstein [14].

2.3. LEMMA. *Let G be a compact group acting by diffeomorphisms on a symplectic manifold (M, σ) , such that $\tilde{g}^*\sigma = \epsilon(g) \cdot \sigma$, ϵ a continuous homomorphism: $G \rightarrow \{-1, +1\}$. Let m be a fixed point for the G -action, σ' another smooth symplectic form defined on a neighborhood of m , such that $\tilde{g}^*\sigma' = \epsilon(g) \cdot \sigma'$ for all $g \in G$, and $\sigma'_m = \sigma_m$. Then there exists a G -equivariant local diffeomorphism Ψ around m , such that $\Psi(m) = m$, $D\Psi(m) = I$, and $\Psi^*\sigma' = \sigma$.*

PROOF. Write $\sigma_t = \sigma + t \cdot (\sigma' - \sigma)$, so that $\sigma_0 = \sigma$, $\sigma_1 = \sigma'$, $\tilde{g}^*\sigma_t = \epsilon(g) \cdot \sigma_t$ for all $g \in G$, $t \in [0, 1]$. One attempts to find local diffeomorphisms Ψ_t , G -equivariant, depending smoothly on t , such that $\Psi_0 = \text{identity}$, and $\Psi_t(m) = m$, $D\Psi_t(m) = I$, $\Psi_t^*\sigma_t = \sigma$ for all $t \in [0, 1]$. Differentiating with respect to t one finds that the velocity field v_t has to satisfy the equations

$$(2.22) \quad d(v_t \lrcorner \sigma_t) = \sigma - \sigma', \quad v_t(m) = 0, \quad Dv_t(m) = 0,$$

for all $t \in [0, 1]$. By the Poincaré lemma there is a smooth 1-form α such that $d\alpha = \sigma - \sigma'$, $\alpha(m) = 0$, $D\alpha(m) = 0$. Let v_t be the unique vector field such that $v_t \lrcorner \sigma_t = \alpha$. Then v_t depends smoothly on t and satisfies (2.22). Each \tilde{g}^*v_t , $g \in G$, will again satisfy (2.22), because

$$\begin{aligned} \epsilon(g) \cdot (\sigma - \sigma') &= \tilde{g}^*(\sigma - \sigma') = \tilde{g}^*d(v_t \lrcorner \sigma_t) \\ &= d\tilde{g}^*(v_t \lrcorner \sigma_t) = d(\tilde{g}^*v_t \lrcorner \tilde{g}^*\sigma_t) \\ &= d(\tilde{g}^*v_t \lrcorner \epsilon(g) \cdot \sigma_t) = \epsilon(g) \cdot d(\tilde{g}^*v_t \lrcorner \sigma_t). \end{aligned}$$

So averaging over G we get a smooth G -invariant vector field v_t , depending smoothly on t , and satisfying (2.22). Integrating it we get a G -equivariant 1-parameter family of local diffeomorphisms Ψ_t , such that $\Psi_t(m) = m$, $D\Psi_t(m) = I$, and $t \mapsto \Psi_t^*\sigma_t$ is constant, hence equal to σ . Taking $\Psi = \Psi_1$, the lemma is proved.

Ignoring the T -action in Proposition 2.2, we have, locally, recovered the theorem of Meyer mentioned before. Moreover, the set

$$(2.23) \quad C = \{m \in Q; D(f|Q)(m) = 0\}$$

is equal to $F \cap Q$, where F is the fixed point set of the T -action in M . Each connected component C_k of C , which in the local coordinates of Proposition 2.2 reads as

$$(2.24) \quad \{(q, p); p = 0 \text{ and } q_j = 0 \text{ whenever } \omega_j \neq 0\},$$

is a Lagrange submanifold of some connected component $F_{j(k)}$ of F , the connected components F_j of F being symplectic submanifolds of M . F_j is τ -invariant if it meets

Q , and then the C_k with $j(k) = j$ are the connected components of the fixed point set for the involution τ in F_j .

We now want to prove that $f(Q)$ is equal to the convex hull of $f(C)$, which consists of only finitely many points because f is obviously constant on each C_k . As in Guillemin-Sternberg [7, §5], the first step is

2.4. LEMMA. *For each $X \in \mathfrak{t}$, the function $f_X|Q$ has a unique local maximal value.*

PROOF. Fix $X \in \mathfrak{t}$. Let $m \in Q$ be a critical point of $f_X|Q$. From Lemma 2.1 we know that m is a fixed point for the action of $\exp tX$, $t \in \mathbf{R}$. Replacing, in this proof, T by the closure of $\{\exp tX, t \in \mathbf{R}\}$ in T , we get that $D(f|Q)(m) = 0$. A glance at (2.13) and the local characterization of Q shows that m is a local maximum for $f_X|Q$ if and only if m is a local maximum for f_X , both conditions being equivalent to $\omega_j(X) \leq 0$ for all j . A Morse theoretic argument gives that the set of points in M where f_X has a local maximum is connected, see Atiyah [2] and Guillemin-Sternberg [7]. So f_X on M has only one local maximal value, and therefore the same must hold for $f_X|Q$.

Now, let $\xi \in \mathfrak{t}^*$ be a boundary point of $f(Q)$, and let $m \in Q$ be such that $f(m) = \xi$. Then

$$(2.25) \quad \mathfrak{t}_m = \{X \in \mathfrak{t}; d(f_X|Q)(m) = 0\}$$

is nonzero, otherwise $D(f|Q)(m)$ would be surjective, contradicting that ξ is a boundary point. In view of Lemma 2.1 we may replace T by the subtorus $\exp \mathfrak{t}_m$, fixing m , and apply Proposition 2.2. Write

$$(2.26) \quad \gamma_m = \sum_{j=1}^n \vartheta_j \cdot \omega_j, \quad \vartheta_j \geq 0,$$

for the convex cone in \mathfrak{t}_m^* generated by the vectors ω_j of (2.13), with \mathfrak{t} replaced by \mathfrak{t}_m . Then (2.13) shows that there is a neighborhood U of m in Q and a neighborhood V of $\pi_m(\xi)$ in \mathfrak{t}_m^* , such that

$$(2.27) \quad \pi_m(f(U)) = V \cap (\pi_m(\xi) + \gamma_m).$$

Here π_m is the projection: $\mathfrak{t}^* \rightarrow \mathfrak{t}_m^*$ obtained by restriction of linear forms to \mathfrak{t}_m . This implies of course that

$$(2.28) \quad f(U) \subset \xi + \pi_m^{-1}(\gamma_m).$$

Note that $\text{Ker } \pi_m = \text{Im } D(f|Q)(m)$. The fact that there are only finitely many possibilities for the T_m in (2.12), and that \mathfrak{t}_m is equal to the Lie algebra of T_m , leads to $\text{Im } D(f|Q)(m') \supset \text{Im } D(f|Q)(m)$ for all m' in a neighborhood of m , so (2.27) can actually be strengthened to

$$(2.29) \quad f(U) = W \cap (\xi + \pi_m^{-1}(\gamma_m))$$

for some neighborhood W of ξ in \mathfrak{t}^* .

Because ξ is a boundary point of $f(Q)$, we see from (2.13) and (2.29) that $\gamma_m \neq \mathfrak{t}_m^*$, so the dual cone

$$(2.30) \quad \delta_m = \{X \in \mathfrak{t}_m; \omega_j(X) \leq 0 \text{ for all } j\}$$

must contain nonzero X . However, $X \in \mathfrak{t}_m$ means in view of (2.28) that $\langle X, f(U) \rangle \leq \langle X, \xi \rangle$, that is, $\langle X, \xi \rangle$ is a local maximal value for $f_X|Q$. Since Lemma 2.4 implies that local maximal values are global maximal values, $\langle X, f(Q) \rangle \leq \langle X, \xi \rangle$ for all $X \in \delta_m$. That is, (2.28) can be strengthened to

$$(2.31) \quad f(Q) \subset \xi + \pi_m^{-1}(\gamma_m).$$

In particular, since $\delta_m \neq \{0\}$, each boundary point of $f(Q)$ is on the boundary of a half-space containing $f(Q)$, so $f(Q)$ is convex. Because $f(Q)$ is compact, it is equal to the convex hull of its extremal points. But (2.29) shows that ξ can only be extremal if $\mathfrak{t}_m = \mathfrak{t}$, that is $m \in C$, and γ_m is a proper cone in \mathfrak{t}^* . Also, (2.13) shows that γ_m is constant, say equal to $\gamma(C_k)$, if m runs along a connected component C_k of C . Since a nonvoid compact convex set is equal to the intersection of the half-spaces containing it and supported at the extremal points, the local description in (2.29), (2.31) of $f(Q)$ leads to

$$(2.32) \quad f(Q) = \bigcap_{\gamma(C_k) \text{ a proper cone}} (f(C_k) + \gamma(C_k)).$$

Working on M and ignoring the involution τ , the above arguments also lead to

$$(2.33) \quad f(M) = \bigcap_{\gamma(F_j) \text{ a proper cone}} (f(F_j) + \gamma(F_j)),$$

where the F_j are the connected components of F , the fixed point set of the T -action on M , and $\gamma(F_j)$ is the convex cone in \mathfrak{t}^* generated by the $\omega_j \in \mathfrak{t}^*$ as in (2.13), in the analogue of Proposition 2.2. disregarding the involution τ . Now each C_k was contained in an $F_{j(k)}$, of course $f(C_k) = f(F_{j(k)})$. However, (2.13) shows that also $\gamma(C_k) = \gamma(F_{j(k)})$. So comparing (2.32) and (2.33) we get $f(Q) \supset f(M)$. Because the other inclusion is trivial, we have proved

2.5. THEOREM. $f(Q) = f(M)$, and is equal to the convex hull of the finitely many values $f(m)$ of f at points $m \in Q$ such that

$$(2.34) \quad D(f|Q)(m) = 0 \text{ and the convex cone } \gamma_m \text{ generated by the } \omega_j \in \mathfrak{t}^* \text{ in (2.13) is proper.}$$

Moreover, if $m' \in M$, $Df(m') = 0$ and the corresponding cone $\gamma_{m'}$ is proper, then there exists $m \in Q$ such that (2.34) holds, and $f(m') = f(m)$, $\gamma_{m'} = \gamma_m$.

REMARK. The proof shows that the same statements are true if Q is replaced by any connected component Q_0 of Q .

3. Tightness. We make the same assumptions as in §2, namely that (M, σ) is a compact symplectic manifold with on it a Hamiltonian action of a torus T , with Hamilton functions f_X , $X \in \mathfrak{t}$. Also, τ is a smooth involution of M such that $\tau^*\sigma = -\sigma$, $\tau^*f_X = f_X$ for all $X \in \mathfrak{t}$. In this section we prove

3.1. THEOREM. Let C_X be the critical set of $f_X|Q$, $X \in \mathfrak{t}$. Then

$$(3.1) \quad \dim H^*(Q; \mathbf{Z}/2\mathbf{Z}) = \dim H^*(C_X; \mathbf{Z}/2\mathbf{Z}).$$

We begin the proof with a reduction to the case of a circle action. By restricting to the closure of $\{\exp tX; t \in \mathbf{R}\}$ in T , we may assume that $\{\exp tX; t \in \mathbf{R}\}$ is dense in

T . In view of Lemma 2.1 we have now $C_X = F \cap Q$, where F is the fixed point set of the action of T on M . Because there are only finitely many possibilities for the stabilizer groups T_m in (2.12), it follows that the collection of $Y \in \mathfrak{t}$ such that $C_Y = C_X$ is open and dense in \mathfrak{t} . Because the set of Y , such that $t \mapsto \exp tY$ is periodic, is dense in \mathfrak{t} , we may assume that $t \mapsto \exp tX$ is periodic, and multiplying X by a suitable factor we get $\exp X = 1$. All this without changing the critical set.

Now (2.8) implies that τ maps T -orbits to T -orbits, reversing the time order on them. In particular the T -orbits through $m \in Q$ are τ -invariant, and

$$(3.2) \quad \tau(\exp t\tilde{X}(m)) = \exp -t\tilde{X}(m).$$

But then $\tau(\exp \frac{1}{2}\tilde{X}(m)) = \exp -\frac{1}{2}\tilde{X}(m) = \exp \frac{1}{2}\tilde{X}(m)$, that is

3.2. LEMMA. $\exp \frac{1}{2}\tilde{X}$ maps Q to itself,² and thereby defines a smooth involution of Q .

Examples (see the remark below) show that the fixed point set in Q of $\exp \frac{1}{2}\tilde{X}$, which contains C_X , is not necessarily equal to C_X (assuming that the minimal period is equal to 1). For this reason we now define, by induction over $k \in \mathbb{N}$,

$$(3.3) \quad Q_{(0)} = Q, \quad Q_{(k)} = \{m \in Q_{(k-1)}; \exp 2^{-k}\tilde{X}(m) = m\}.$$

By induction over k one proves

3.3. LEMMA. $\exp 2^{-k}X$ is an involution of $Q_{(k-1)}$. The $Q_{(k)}$ form a decreasing family of subspaces of Q . $Q_{(k)}$ has finitely many connected components, each of which is a smooth compact submanifold of $Q_{(k-1)}$, resp. of Q . There exists an $N \in \mathbb{N}$ such that $Q_{(N)} = C_X$.

The last statement follows because such decreasing sequences of submanifolds must stabilize, that is, there is an $N \in \mathbb{N}$ such that $Q_{(k)} = Q_{(N)}$ for all $k \geq N$. But $m \in Q_{(k)}$ for all k implies that $\tilde{X}(m) = 0$, hence $dJ_X(m) = 0$, or $m \in C_X$.

It is known (Floyd [4 or 5, §4]) that if B is the fixed point set of a periodic transformation of prime period p in a compact manifold A , then

$$\dim H^*(A; \mathbb{Z}/p\mathbb{Z}) \geq \dim H^*(B; \mathbb{Z}/p\mathbb{Z}).$$

Lemma 3.3 therefore implies

$$(3.4) \quad \dim H^*(Q; \mathbb{Z}/2\mathbb{Z}) \geq \dim H^*(Q_{(1)}; \mathbb{Z}/2\mathbb{Z}) \geq \cdots \geq \dim H^*(C_X; \mathbb{Z}/2\mathbb{Z}).$$

But the opposite inequality $\dim H^*(Q; R) \leq \dim H^*(C_X; R)$, valid for general coefficient rings R , follows from the Bott-Morse inequalities. This proves Theorem 3.1. The idea to conclude tightness by exhibiting the critical set as the fixed point set for a periodic map has been introduced by Frankel [6] in a Kähler framework.

REMARK. If m is an isolated fixed point for $\exp \frac{1}{2}\tilde{X}: Q \rightarrow Q$, then this “half-turn” is the Cartan involution around m with respect to any invariant Riemannian metric. Suppose now that the isometry group of the connected component Q_0 of Q acts transitively, as is the case for the real flag manifolds; see §5. Then the existence of isolated fixed points for the half-turn on Q_0 makes Q_0 into a Riemannian symmetric

² Note that Q is not invariant under the \tilde{X} -flow. In fact $\tilde{X}(m) \in P_m$ where P_m is the space complementary to $T_m Q$, defined in (2.10).

space. For symmetric real flag manifolds the tightness of $f_X|_{Q_0}$ was proved in Takeuchi [16, pp. 167–168] by identifying the critical set with the fixed point set of a Cartan involution. Conversely, if Q_0 is a real flag manifold which is not a symmetric space, and if $f_X|_{Q_0}$ has isolated critical points (as generically is the case), then the fixed point set of $\exp \frac{1}{2}\tilde{X}$ in Q cannot be equal to C_X .

4. Gradient flows. If β is a Riemannian metric on M which is T - and τ -invariant, then the gradient vector fields of the functions f_X , $X \in \mathfrak{t}$, are also T - and τ -invariant. In particular they are tangent to the fixed point set Q of τ , the gradient flows leaving Q invariant. The formula $\beta(u, v) = \sigma(u, Jv)$ defines a tensor field J on M which is T -invariant and τ -anti-invariant, and $\text{grad } f_X$ is equal to J times the Hamiltonian vector field X of the function f_X .

If J is an integrable almost complex structure, that is if M is a complex analytic manifold with complex structure equal to J , then $\beta + i\sigma$ is a Kähler metric on M . The T -action consists of the holomorphic mappings, whereas the involution τ is anti-holomorphic, making Q into a “real subspace” of M in a strong sense. Because the automorphism group of a compact complex manifold is a complex Lie group, the action of T extends to a holomorphic action of the complexification T_c of T . Its Lie algebra $\mathfrak{t} \otimes \mathbb{C}$ can be written as $\mathfrak{t} \oplus \mathfrak{a}$ where $\mathfrak{a} = i\mathfrak{t}$. The exponential map is taken to be injective on \mathfrak{a} , making $A = \exp \mathfrak{a}$ into a vector subgroup of T_c , and $(t, a) \mapsto t \cdot a$ is a diffeomorphism from $T \times A$ onto T_c . Now $\text{grad } f_X = J \cdot \tilde{X} = \widetilde{i \cdot X}$, so the gradient vector fields together make up the infinitesimal action of A . In particular the gradient flows commute with each other, in fact this is the only additional assumption which will be used in the sequel.

4.1. THEOREM. *Let Y be an A -orbit in M and $F_j, j = 1, \dots, p$, the components of the common critical points of the functions $f_X, X \in \mathfrak{t}$, which intersect the closure \bar{Y} of Y in M . Then $f(\bar{Y})$ is equal to the convex polytope P with extremal points equal to $c_j = f(F_j), j = 1, \dots, p$. For each open face φ of P the inverse image $f^{-1}(\varphi)$ in Y consists of a simple A -orbit, and f induces a homeomorphism of \bar{Y} onto P .*

This is Theorem 2 of Atiyah [2]. That theorem was phrased in terms of T_c -orbits in M , but using the invariance of f under the T -action, one readily translates it into the above statement. Because Q is A -invariant we can apply Theorem 4.1 to the A -orbits in Q . Then Theorem 4.1 is a generalization of the corresponding statement for the real flag manifolds, due to Heckman [8, Chapter 2, Theorem 3]. The following result shows that Theorem 4.1 can be regarded as a refinement of Theorem 2.5 in the case of commuting gradient flows.

4.2. PROPOSITION. *For all m' in an open dense subset Q' of Q , the set of $f(m)$ such that $m \in Q, D(f|_Q)(m) = 0$, and m is in the closure of the A -orbit through m' , is equal to the set of extremal points of $f(Q)$.*

PROOF. Without loss of generality we may assume that $f(Q) = f(M)$ has a nonempty interior, that is

$$(4.1) \quad Q^{\text{reg}} = \{m \in Q; D(f|_Q)(m) \text{ is surjective}\}$$

is nonvoid.

Recall that $f(Q_0) = f(M)$ for any connected component Q_0 of Q . Now $m \notin Q_0^{\text{reg}}$ means that there exists $X \in \mathfrak{t}$, $X \neq 0$, such that m is a critical point for f_X . For the description of the critical set C_X of $f_X|_{Q_0}$ we may, as in the proof of Theorem 3.1, assume that $t \mapsto \exp tX$ is periodic on M with minimal positive period equal to 1 when starting on Q_0 . Replacing T for the moment by the circle $\{\exp tX; t \in \mathbf{R}\}$, we read off from (2.13) that

$$(4.2) \quad C_X = \{q; q_j = 0 \text{ if } \omega_j(X) \neq 0\}, \quad \text{locally.}$$

Here $\omega_j(X) = 2\pi \cdot k_j$, $k_j \in \mathbf{Z}$, and not all k_j are even, noting that the half-turn $\exp \frac{1}{2}X$ of Lemma 3.2 is a nontrivial involution of Q_0 . If $\text{codim } C_X = 1$, locally, then only one of the $\omega_j(X)$ is nonzero, hence odd. The half-turn maps q_j to $-q_j$, so one side of C_X to the other. There are only finitely many C_X 's composing $Q_0 \setminus Q_0^{\text{reg}}$. Therefore, writing

$$(4.3) \quad T_{1/2} = \{g \in T; \tilde{g}^2(m) = m \text{ for all } m \in Q_0\},$$

we get that $Q_0^{\text{reg}}/T_{1/2}$ is connected. (This argument is reminiscent of the transitivity of the action of the Weyl group on the set of Weyl chambers.)

In particular, since f is T -invariant, $f(Q_0^{\text{reg}})$ is connected. It is open; and dense in $f(Q_0) = f(M)$, because Q_0^{reg} is dense in Q_0 , $Q_0 \setminus Q_0^{\text{reg}}$ being equal to the union of finitely many closed submanifolds of codimension ≥ 1 of Q_0 .

Since there are only finitely many possibilities for the $f(\bar{Y})$, Y an A -orbit in Q_0 , one has that for each $m' \in Q_0$ there is a neighborhood U of m' in Q_0 such that $f(A \cdot m'') \supset f(A \cdot m')$ for all $m'' \in U$. Let Q'_0 be the set of $m' \in Q_0$ such that $f(A \cdot m'') = f(A \cdot m')$ for all $m'' \in Q_0$, m'' near m' . Then Q'_0 is open and dense in Q_0 . Each connected component R of Q'_0 is A -invariant, and $f(\bar{R}) = f(A \cdot m')$ for each $m' \in R$. We shall show that $f(\bar{R}) = f(Q_0) = f(M)$, thus completing the proof of the proposition in view of Theorem 4.1. Indeed, if $m \in Q_0^{\text{reg}}$ then f is a diffeomorphism from $A \cdot m$ to an open subset of \mathfrak{t}^* , so $f(m) \notin \partial f(\bar{R})$. Because R meets the open dense subset Q_0^{reg} of Q_0 , the open set $f(Q_0^{\text{reg}})$ meets $f(\bar{R})^{\text{int}}$. Since $\partial f(\bar{R}) \in f(Q_0^{\text{reg}}) = \emptyset$, it follows that $f(\bar{R})^{\text{int}}$ contains the connected set $f(Q_0^{\text{reg}})$. But this implies that $f(\bar{R}) = f(Q_0)$.

Choose $X \in \mathfrak{t}$. The stable (or unstable) manifolds of the gradient vector field of f_X define a decomposition of M and of Q . These are cell decompositions if f_X is a Morse function rather than only Bott-Morse. The stable manifolds in Q are the connected components of the intersections with Q of the stable manifolds in M . *In the case of the flag manifolds*, and f_X a Morse function, the closures of the stable manifolds in M are complex algebraic varieties, defining cycles (the Schubert cycles) which form a basis for the homology of M . The closures of the stable manifolds in Q are real algebraic varieties, defining cycles modulo 2 which form a basis for the homology mod 2 of Q . Also, two critical points in Q are connected by a gradient curve in M only if they are connected by a gradient curve in Q . See [3, §4]. Finally it is known that the image under f of both the real and the complex Schubert cycles are convex polytopes, see Heckman [8, Chapter 2, Corollary 2], and Atiyah [2, §4].

It might be interesting to investigate which of these properties generalize to the present setting.

5. Flag manifolds. For any connected Lie group U with Lie algebra \mathfrak{u} , Kirillov [9]³ introduced a symplectic form on each orbit of the coadjoint action of U in \mathfrak{u}^* , as follows. For $\xi \in \mathfrak{u}^*$, the coadjoint orbit \mathcal{O} of ξ can be identified with U/U_ξ , where

$$(5.1) \quad U_\xi = \{g \in U; (\text{Ad } g)^*(\xi) = \xi\},$$

is the stabilizer of ξ in U , which has Lie algebra

$$(5.2) \quad \mathfrak{u}_\xi = \{X \in \mathfrak{u}; (\text{ad } X)^*(\xi) = 0\}.$$

The symplectic form on $T_\xi \mathcal{O} \cong \mathfrak{u}/\mathfrak{u}_\xi$ is defined by

$$(5.3) \quad \sigma_\xi(X, Y) = \xi([X, Y]), \quad X, Y \in \mathfrak{u}/\mathfrak{u}_\xi,$$

and it is then shown that σ_ξ extends to a unique U -invariant symplectic form σ on \mathcal{O} . The action of U on \mathcal{O} is Hamiltonian (though nonabelian, see Abraham and Marsden [1] for the definitions in this general case), and the momentum mapping is equal to the inclusion mapping: $\mathcal{O} \rightarrow \mathfrak{u}^*$. If T is a torus in U , then its action on \mathcal{O} is Hamiltonian with momentum mapping equal to the projection: $\mathfrak{u}^* \rightarrow \mathfrak{t}^*$, restricted to \mathcal{O} .

Up to coverings, these are the only symplectic manifolds with transitive Hamiltonian group actions. If U is compact, then the theorem of Atiyah [2] and Guillemin and Sternberg [7] gives that the projection of the coadjoint orbit of ξ in \mathfrak{u}^* to \mathfrak{s}^* , \mathfrak{s} = the Lie algebra of a maximal torus in U , is equal to the convex hull of the Weyl group orbit of ξ . As we shall see below, these coadjoint orbits are the complex flag manifolds and the convexity theorem is Kostant's for the complex case. Note that because the center of U is contained in U_ξ , one may assume here that U has trivial center.

Now we turn to a description of the real flag manifolds, see also [3, §2]. Let G be a real connected semisimple Lie group with trivial center, and let $G = KAN$ be its Iwasawa decomposition. We may think of $G = \text{Ad } G$ as a matrix group, then K, A, N are the groups of respectively the orthogonal, diagonal with positive eigenvalues, upper triangular unipotent elements of G . Let $\mathfrak{g}, \mathfrak{k}, \mathfrak{a}$ be the Lie algebras of G, K, A respectively. For any $H \in \mathfrak{a}$, the $\text{Ad } K$ -orbit of H in \mathfrak{g} (actually contained in a sphere in the orthogonal complement of \mathfrak{k}) can be identified with K/K_H , where

$$(5.4) \quad K_H = \{k \in K; \text{Ad } k(H) = H\}$$

is the centralizer of H in K . The functions

$$(5.5) \quad f_{H', H}(k) = \langle H', \text{Ad } k(H) \rangle, \quad k \in K/K_H, \quad H, H' \in \mathfrak{a}$$

(the bilinear form here is the Killing form), can be considered as testing the orthogonal projection of $\text{Ad } K(H)$ to \mathfrak{a} by linear forms on \mathfrak{a} .

This orthogonal projection is actually the infinitesimal version of the Iwasawa projection $\pi: G \rightarrow \mathfrak{a}$ defined by

$$(5.6) \quad x \in K \cdot \exp \pi(x) \cdot N, \quad x \in G.$$

³ The fact that the form is closed and the relation with general homogeneous symplectic manifolds were observed later by Kostant and Souriau.

This projection will be applied to the K -orbit

$$(5.7) \quad \{k \cdot \exp H \cdot k^{-1}; k \in K\} \cong K/K_H$$

of $\exp H$ in G . The full convexity theorem of Kostant now states that both the Iwasawa projection of (5.7), and its infinitesimal version applied to $\text{Ad } K(H)$, have their image equal to the convex hull of the Weyl group orbit of H in \mathfrak{a} . We shall only discuss the functions $f_{H',H}$ in (5.5), noting that Heckman [8] showed that the convexity theorem for the Iwasawa projection can be proved from its infinitesimal version by a homotopy argument.

As we shall show below, K/K_H is a connected component of Q and $f_{H',H} = f_X|_Q$, where Q, f_X are as in §§2, 3. The symplectic manifold M is equal to a complex flag manifold U/U_ξ as above, for a suitable U , resp. ξ , and X is related to H' by a linear isomorphism. This then puts the infinitesimal version of Kostant's convexity theorem in the framework of Theorem 2.5. Moreover, in Takeuchi and Kobayashi [13] and [3, §4], it is proved that for generic H' , $f_{H',H}$ is a tight Morse function on K/K_H . So Theorem 3.1 provides a new proof for this, and extends the tightness to arbitrary $H' \in \mathfrak{a}$.

Let $G_{\mathbb{C}}$ be the complexification of G , with Iwasawa decomposition

$$(5.8) \quad G_{\mathbb{C}} = UBV$$

(we are clearly running out of letters). Here U is the maximal compact subgroup of $G_{\mathbb{C}}$, which in fact is another real form of $G_{\mathbb{C}}$. If τ is the complex conjugation of $G_{\mathbb{C}}$ around G , then G is the connected component of 1 of the fixed point set of τ in $G_{\mathbb{C}}$. Moreover, we can arrange that U is τ -invariant and K is the connected component of 1 of the fixed point set of τ in U . Similarly B , resp. V , are τ -invariant and A , resp. N , are the fixed point sets of τ in B , resp. V . Of these groups only V is complex, in general. In fact, the complexification C of B is a Cartan subgroup of $G_{\mathbb{C}}$, $S = C \cap U$ is a maximal torus in u , its Lie algebra \mathfrak{s} is equal to $i \cdot \mathfrak{b}$ if \mathfrak{b} denotes the Lie algebra of B .

For $H \in \mathfrak{b}$, $U/U_H \xrightarrow{\sim} G_{\mathbb{C}}/U_H B V$, where $U_H B V$ turns out to be a complex closed subgroup of $G_{\mathbb{C}}$. It contains $C V$, which is a maximal solvable subgroup of $G_{\mathbb{C}}$, called a Borel subgroup. The Borel subgroups of $G_{\mathbb{C}}$ are all conjugate to each other. The subgroups $P_{\mathbb{C}}$ of $G_{\mathbb{C}}$ containing a Borel subgroup are called the parabolic subgroups. They are also characterized as those for which $G_{\mathbb{C}}/P_{\mathbb{C}}$ is a complex projective variety. The $G_{\mathbb{C}}/P_{\mathbb{C}}$ are the complex flag manifolds. Since up to conjugacy each parabolic subgroup of $G_{\mathbb{C}}$ is of the form $U_H B V$ for some $H \in \mathfrak{b}$, this exhibits the U/U_H as the general complex flag manifolds.

Now $U_H = U_{\xi}$ as in (5.1), if we define $\xi \in \mathfrak{s}^*$ by

$$(5.9) \quad \xi(Z) = \langle iH, Z \rangle, \quad Z \in \mathfrak{s}.$$

This identifies the coadjoint orbits of compact connected Lie groups with the complex flag manifolds.

If $H \in \mathfrak{a} \subset \mathfrak{b}$, then $U_H B V$ is equal to the complexification of $P = K_H A N$, which therefore is called a real parabolic subgroup of G . As a corollary, $K/K_H \xrightarrow{\sim} G/P$ has

$U/U_H \xrightarrow{\sim} G_C/P_C$ as its complexification. Conversely, K/K_H is equal to the connected component of $1 \cdot K_H$ of the fixed point set of τ in U/U_H . The K/K_H are called the real flag manifolds. For the classical groups they can be identified with spaces of flags of linear subspaces of a vector space, isotropic with respect to the bilinear form (not necessarily symmetric or nondegenerate) of which G is taken as the isometry group. In particular all (isotropic) Grassmann manifolds are included in the list of examples.

Since τ leaves the elements of \mathfrak{a} fixed, and maps $\xi = iH$ to $\overline{iH} = -iH = -\xi$, we see from (5.3) that τ is antisymplectic. On the other hand, taking for $T \subset S$ the torus in U generated by $\mathfrak{t} = i\mathfrak{a} \subset i\mathfrak{b} = \mathfrak{s}$, we get that T acts in a Hamiltonian way on U/U_H , with Hamilton function f_X of \tilde{X} , $X \in \mathfrak{t}$, equal to $f_{H',H}$, taking $H' = -iX$. In particular f_X is τ -invariant.

REFERENCES

1. R. Abraham and J. Marsden, *Foundations of mechanics*, 2nd ed., Benjamin, New York, 1978.
2. M. F. Atiyah, *Convexity and commuting Hamiltonians*, preprint, Oxford Univ. Press, London, 1981.
3. J. J. Duistermaat, J. A. C. Kolk and V. S. Varadarajan, *Functions, flows and oscillatory integrals on flag manifolds and conjugacy classes in real semisimple Lie groups*, preprint, Utrecht, 1981.
4. E. E. Floyd, *On periodic maps and the Euler characteristics of associated spaces*, Trans. Amer. Math. Soc. **72** (1952), 138–147.
5. ———, *Periodic maps via Smith theory*, Seminar on Transformation Groups (A. Borel, ed.), Ann. of Math. Studies, no. 46, Princeton Univ. Press, Princeton, N. J., 1960, pp. 35–47.
6. T. Frankel, *Fixed points on Kähler manifolds*, Ann. of Math. (2) **70** (1959), 1–8.
7. V. Guillemin and S. Sternberg, *Convexity properties of the moment mapping*, preprint, M.I.T., 1981.
8. G. J. Heckman, *Projections of orbits and asymptotic behaviour of multiplicities for compact Lie groups*, Thesis, Leiden, 1980.
9. A. A. Kirillov, *Unitary representations of nilpotent Lie groups*, Uspehi Mat. Nauk **17** (1962), 57–110 = Russian Math. Surveys **17** (1962), 53–104.
10. B. Kostant, *On convexity, the Weyl group and the Iwasawa decomposition*, Ann. Sci. École Norm. Sup. (4) **6** (1973), 413–455.
11. K. R. Meyer, *Hamiltonian systems with discrete symmetry*, J. Differential Equations **41** (1981), 228–238.
12. G. D. Mostow, *On a conjecture of Montgomery*, Ann. of Math. (2) **65** (1957), 513–516.
13. M. Takeuchi and S. Kobayashi, *Minimal embeddings of R-spaces*, J. Differential Geom. **2** (1968), 203–215.
14. A. Weinstein, *Symplectic manifolds and their Lagrangian submanifolds*, Adv. in Math. **6** (1971), 329–346.
15. C.-T. Yang, *On a problem of Montgomery*, Proc. Amer. Math. Soc. **8** (1957), 255–275.
16. M. Takeuchi, *Cell decomposition and Morse inequalities on certain symmetric spaces*, J. Fac. Sci. Univ. Tokyo **12** (1965), 81–192.

MATHEMATISCH INSTITUUT, RIJKSUNIVERSITEIT UTRECHT, BUDAPESTLAAN 6, DE UITHOF, 3508 TA UTRECHT, THE NETHERLANDS