

PROJECTIONS ON TENSOR PRODUCT SPACES

BY

E. J. HALTON¹ AND W. A. LIGHT

ABSTRACT. $(S, \Sigma, \mu), (T, \Theta, \nu)$ are finite, nonatomic measure spaces. G and H are finite-dimensional subspaces of $L_1(S)$ and $L_1(T)$ respectively. Both G and H contain the constant functions. It is shown that the relative projection constant of $L_1(S) \otimes H + G \otimes L_1(T)$ in $L_1(S \times T)$ is at least 3.

I. Introduction. Suppose X is a normed linear space with subspace Y . A projection $P: X \rightarrow Y$ is a linear transformation whose range is Y and which acts as the identity on Y . We set $\lambda(Y, X) = \inf\{\|P\|: P \text{ is a projection from } X \text{ onto } Y\}$. This number is called the relative projection constant of Y in X . If we can find a projection from X onto Y whose norm is $\lambda(Y, X)$, then this projection is called minimal. Computing $\lambda(Y, X)$ even for quite "small" Y is usually a very difficult task, and the best that can be done in most cases is to provide bounds.

In this area some recent work has focused attention on bivariate function spaces. In particular several authors have considered the cases $X = C(S \times T)$, $Y = C(S) + C(T)$ or $X = L_p(S \times T)$, $Y = L_p(S) + L_p(T)$ (see [3, 4 and 6] for details). Recently, Cheney and Franchetti [1] improved the results of [6] by showing that if $X = C(S \times T)$ and $Y = C(S) \otimes H + G \otimes C(T)$, where G and H are finite-dimensional subspaces of $C(S)$ and $C(T)$ respectively and each containing the constant functions, then $\lambda(Y, X) \geq 3$. Jameson and Pinkus [6] had previously showed that if G and H are just spanned by the constant functions, then $\lambda(Y, X) = 3$.

In this paper we shall take G as an n -dimensional subspace of $L_1(S)$ or $L_\infty(S)$ spanned by $\{g_1, \dots, g_n\}$. H will be an m -dimensional subspace of $L_1(T)$ or $L_\infty(T)$ spanned by $\{h_1, \dots, h_m\}$. Both G and H will be assumed to contain the functions which are constant almost everywhere with respect to the appropriate measure. S and T will be finite, nonatomic measure spaces. We shall show that $\lambda(Y, X) \geq 3$ for $X = L_1(S \times T)$, $Y = L_1(S) \otimes H + G \otimes L_1(T)$.

II. Preliminary results. In this section we shall deal entirely with $L_1(S \times T)$. However, we shall be able to derive the same results in $L_\infty(S \times T)$. We shall indicate at the end of this section the minor differences that arise in $L_\infty(S \times T)$. So let us take $X = L_1(S \times T)$, $U = L_1(S) \otimes H$, $V = G \otimes L_1(T)$ and $W = U + V$. We shall also assume without loss of generality that (S, Σ, μ) and (T, Θ, ν) are such that $\mu(S) = \nu(T) = 1$. For $x \in X$ the sections x_s, x_t are defined by $x_s(t) = x(s, t)$ and $x_t(s) = x(s, t)$ for almost all (s, t) belonging to $S \times T$. Unadorned norm symbols will denote L_1 -norms. Thus, for example,

$$\|u\| = \int |u(s)| d\mu, \quad u \in L_1(S).$$

Received by the editors December 19, 1983.

1980 *Mathematics Subject Classification.* Primary 41A65, 41A50.

¹Supported by the Science and Engineering Research Council of Great Britain.

©1985 American Mathematical Society
0002-9947/85 \$1.00 + \$.25 per page

The L_∞ -norms are written with a subscript, as

$$\|f\|_\infty = \text{ess sup } |f(s, t)|, \quad f \in L_\infty(S \times T).$$

We begin with an elementary lemma whose proof can be found in [5], for example.

LEMMA 2.1. (i) *There exists a constant β such that each element w of W has a representation $w = u + v$, where $u \in U$, $v \in V$ and $\|u\| + \|v\| \leq \beta\|w\|$.*

(ii) *There exists a constant γ such that for all $x_i \in L_1(T)$, $\|x_i\| \leq \gamma\|\sum_{i=1}^n x_i g_i\|$.*

Here the constant β depends only on U and V while the constant γ depends only on the choice of basis for G .

It is well known (see [2, p. 308]) that any finite nonatomic measure space can be written as the union of a finite number of disjoint sets each with measure no more than some prescribed tolerance ε . This fact can be used to show that S and T may be partitioned into sets S_i and T_i , where $1 \leq i \leq k$ and $\mu(S_i) = \mu(T_j) = 1/k$. In outline this is done as follows. Firstly, partition S into a finite number of disjoint sets of length at most $1/2k$. By taking unions if necessary we may assume that $1/2k \leq \mu(A_1) \leq 1/k$. Partition $S \setminus A_1$ into sets of measure at most $1/4k$. By taking the union of as many of these sets as necessary together with A_1 we obtain A_2 with $3/4k \leq \mu(A_2) \leq 1/k$. This defines the first two members of a sequence of sets A_1, A_2, A_3, \dots . Now setting $S_1 = \bigcup_{n=1}^\infty A_n$ gives S_1 with measure $1/k$ as required. Further members in the partition are produced by applying this technique to $S \setminus S_1$ and so on. With the aid of these partitions we define functionals $E_i \in [L_1(S)]^*$ and $F_j \in [L_1(T)]^*$ by

$$\begin{aligned} E_i u &= \int_{S_i} u d\mu, & u &\in L_1(S), \quad 1 \leq i \leq k, \\ F_j v &= \int_{T_j} v d\nu, & v &\in L_1(T), \quad 1 \leq j \leq k. \end{aligned}$$

Now define functions $x_i \in L_1(S)$ and $y_i \in L_1(T)$ by

$$x_i = k\chi_{S_i}, \quad y_i = k\chi_{T_i}, \quad 1 \leq i \leq k.$$

Here χ_{S_i} denotes the characteristic function of the set S_i . Since $\{S_i\}$ is a partition of S we have $\sum_{i=1}^k x_i = k$ and similarly $\sum_{i=1}^k y_i = k$. We shall set $z_{ij} = x_i y_j$, $1 \leq i, j \leq k$. It is clear that

$$\|E_i\| = \|F_i\| = \|x_i\| = \|y_i\| = 1, \quad 1 \leq i \leq k,$$

and

$$E_i(x_j) = F_i(y_j) = \delta_{ij}, \quad 1 \leq i, j \leq k.$$

We also define $D_{ab} \in [L_1(S \times T)]^*$ by $D_{ab}z = E_a(F_b z_s)$ for all $z \in L_1(S \times T)$ and $1 \leq a, b \leq k$.

LEMMA 2.2. *Let P be a projection from $L_1(S \times T)$ onto W . Then we have*

$$\|P\| \geq \frac{1}{k^2} \sum_{a,b=1}^k \sum_{i,j=1}^k |(D_{ab})(Pz_{ij})|.$$

PROOF. We have for any $z \in L_1(S \times T)$ with $\|z\| = 1$,

$$\|P\| \geq \|Pz\| = \sup \left\{ \left| \iint \phi \cdot Pz d\sigma \right| : \phi \text{ belongs to the unit ball in } L_\infty(S \times T) \right\}.$$

Now since the functionals D_{ab} , $1 \leq a, b \leq k$, have disjoint support and are of unit norm we may write

$$\|P\| \geq \sum_{a,b=1}^k |(D_{ab})(Pz)| \quad \text{or} \quad \|P\| \geq \sum_{a,b=1}^k |(D_{ab})(Pz_{ij})|.$$

Summing each side of this inequality over $1 \leq i, j \leq k$ gives

$$k^2 \|P\| \geq \sum_{a,b=1}^k \sum_{i,j=1}^k |(D_{ab})(Pz_{ij})|$$

which is equivalent to the required result. \square

LEMMA 2.3. *For any projection P from $L_1(S \times T)$ onto W we have*

$$\|P\| \geq 3 - \frac{2}{k^2} \sum_{a,b=1}^k (D_{ab})(Pz_{ab}),$$

PROOF. Consider first

$$\sum_{i=1}^k (D_{ab})(Pz_{ij}) = (D_{ab}) \left(P \sum_{i=1}^k z_{ij} \right) = (D_{ab})(P(ky_j)) = (D_{ab})(ky_j) = \delta_{bj}.$$

Similarly, $\sum_{j=1}^k (D_{ab})(Pz_{ij}) = \delta_{ai}$. This allows us to reason as follows:

$$\begin{aligned} k^2 \|P\| &\geq \sum_{a,b=1}^k \sum_{i,j=1}^k |(D_{ab})(Pz_{ij})| \\ &\geq \sum_{a,b=1}^k \left[- \sum_{\substack{i,j=1 \\ i \neq a, j \neq b}}^k (D_{ab})(Pz_{ij}) + \sum_{j=1}^k (D_{ab})(Pz_{aj}) \right. \\ &\quad \left. + \sum_{i=1}^k (D_{ab})(Pz_{ib}) - (D_{ab})(Pz_{ab}) \right] \\ &= \sum_{a,b=1}^k \left[- \sum_{i,j=1}^k (D_{ab})(Pz_{ij}) + 2 \sum_{j=1}^k (D_{ab})(Pz_{aj}) \right. \\ &\quad \left. + 2 \sum_{i=1}^k (D_{ab})(Pz_{ib}) - 2(D_{ab})(Pz_{ab}) \right] \\ &= \sum_{a,b=1}^k (-1 + 2\delta_{aa} + 2\delta_{bb} - 2(D_{ab})(Pz_{ab})) \\ &= 3k^2 - 2 \sum_{a,b=1}^k (D_{ab})(Pz_{ab}). \quad \square \end{aligned}$$

We conclude this section as promised, with a brief discussion of the L_∞ -case. Lemma 2.3 is really not dependent on the norm employed so much as the closedness of $U + V$. We define $E_i, F_i, x_i, y_i, 1 \leq i \leq k$, as before but now $\|E_i\| = \|F_i\| = 1/k$ and $\|x_i\| = \|y_i\| = k, 1 \leq i \leq k$. We continue to have the biorthogonality property. The change of scaling does not affect Lemma 2.2 and since Lemma 2.3 is an algebraic result resting only on the biorthogonality conditions we continue to have the following result.

LEMMA 2.4. *Let P be any projection from $L_\infty(S \times T)$ onto $L_\infty(S) \otimes H + G \otimes L_\infty(T)$, where H is finite dimensional in $L_\infty(T)$, G is finite dimensional in $L_\infty(S)$ and G and H contain the functions which are constant almost everywhere on S and T respectively. Then*

$$\|P\| \geq 3 - \frac{2}{k^2} \sum_{a,b=1}^k (D_{ab})(Pz_{ab}).$$

III. Main theorem. With the aid of the results in the previous section we can now deduce the required result in $L_1(S \times T)$.

THEOREM 3.1. *Let G, H be finite-dimensional subspaces of $L_1(S)$ and $L_1(T)$ respectively such that they each contain the functions which are constant almost everywhere with respect to the appropriate measure. Then*

$$\lambda(L_1(S) \otimes H + G \otimes L_1(T), L_1(S \times T)) \geq 3.$$

PROOF. By Lemma 2.1 we can express Pz_{ab} as

$$Pz_{ab} = u_{ab} + v_{ab} = \sum_{i=1}^n x_i^{a,b} g_i + \sum_{j=1}^m y_j^{a,b} h_j,$$

where $\|u_{ab}\| + \|v_{ab}\| \leq \beta \|Pz_{ab}\|$. Then

$$\begin{aligned} \sum_{a,b=1}^k (D_{ab})(Pz_{ab}) &= \sum_{a,b=1}^k \left[\sum_{i=1}^n E_a(g_i) F_b(x_i^{a,b}) + \sum_{j=1}^m E_a(y_j^{a,b}) F_b(h_j) \right] \\ &\leq \sum_{a,b=1}^k \left[\sum_{i=1}^n |E_a(g_i)| |F_b(x_i^{a,b})| + \sum_{j=1}^m |E_a(y_j^{a,b})| |F_b(h_j)| \right]. \end{aligned}$$

Now by Lemma 2.1 again we may argue as follows

$$|F_b(x_i^{a,b})| \leq \|x_i^{a,b}\| \leq \gamma \left\| \sum_{i=1}^n x_i^{a,b} g_i \right\| \leq \beta \gamma \|Pz_{ab}\| \leq \beta \gamma \|P\|.$$

Similarly, $|E_a(y_j^{a,b})| \leq \beta \delta \|P\|$, where δ is an appropriate constant depending only on the basis chosen for H . Hence, there exists a real number $R > 0$ such that

$$\sum_{a,b=1}^k (D_{ab})(Pz_{ab}) \leq R \|P\| \sum_{a,b=1}^k \left[\sum_{i=1}^n |E_a(g_i)| + \sum_{j=1}^m |F_b(h_j)| \right].$$

Now by our definition of $E_a, F_b, 1 \leq a, b \leq k$, we have

$$\sum_{a=1}^k |E_a(g_i)| \leq \|g_i\|, \quad \sum_{b=1}^k |F_b(h_j)| \leq \|h_j\|$$

and so

$$\sum_{a,b=1}^k (D_{ab})(Pz_{ab}) \leq kR\|P\| \left[\sum_{i=1}^n \|g_i\| + \sum_{j=1}^m \|h_j\| \right].$$

Hence,

$$\begin{aligned} \|P\| &\geq 3 - \frac{2}{k^2} \sum_{a,b=1}^k (D_{ab})(Pz_{ab}) \\ &\geq 3 - \frac{2}{k} R\|P\| \left[\sum_{i=1}^n \|g_i\| + \sum_{j=1}^m \|h_j\| \right]. \end{aligned}$$

Since this inequality holds for all natural numbers k , we have the required result. \square

REFERENCES

1. E. W. Cheney and C. Franchetti, *Minimal projections in tensor product spaces* (Center for Numerical Analysis Report No. 184, Univ. of Texas at Austin, Austin, Texas); J. Approx. Theory (to appear).
2. N. Dunford and J. T. Schwartz, *Linear operators*, Part I, Interscience, New York, 1957.
3. E. J. Halton and W. A. Light, *Minimal projections in bivariate function spaces*, J. Approx. Theory (to appear).
4. ———, *Minimal projections in L_p -spaces*, Univ. of Lancaster Math. Dept. Report, Lancaster, England, Sept. 1983.
5. S. M. Holland, W. A. Light and L. J. Sulley, *On proximality in $L_1(T \times S)$* , Proc. Amer. Math. Soc. **86** (1982), 279–282.
6. G. J. O. Jameson and A. Pinkus, *Positive and minimal projections in function spaces*, J. Approx. Theory **37** (1983), 182–195.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LANCASTER, LANCASTER LA1 4YL, ENGLAND