

SYMMETRY BREAKING FOR A CLASS OF SEMILINEAR ELLIPTIC PROBLEMS

MYTHILY RAMASWAMY AND P. N. SRIKANTH

ABSTRACT. We study positive solutions of the Dirichlet problem for $-\Delta u = u^p - \lambda$, $p > 1$, $\lambda > 0$, on the unit ball Ω . We show that there exists a positive solution (u_0, λ_0) of this problem which satisfies in addition $\partial u_0 / \partial n = 0$ on $\partial\Omega$. We prove also that at (u_0, λ_0) , the symmetry breaks, i.e. asymmetric solutions bifurcate from the positive radial solutions.

1. Introduction. We consider here the symmetry breaking for semilinear elliptic problems of the type

$$(1.1) \quad \begin{aligned} -\Delta u &= f(u, \lambda) && \text{in } B_1, \\ u &= 0 && \text{on } \partial B_1, \end{aligned}$$

where B_1 is the unit ball on \mathbf{R}^n , $n \geq 2$, and $f: \mathbf{R} \times \mathbf{R} \rightarrow \mathbf{R}$ is a C^2 function. We say that the symmetry breaks at λ if the radial solution (u_0, λ_0) is a nonradial bifurcation point, i.e. if every neighborhood of (u_0, λ_0) contains solutions of (1.1), (u, λ) with u nonradial.

Symmetry breaking problems for positive solutions of semilinear elliptic equations have been studied by Smoller and Wasserman [4, 5]. In fact they prove that for the function $f \in C^2$ satisfying

$$(1.2) \quad \begin{aligned} f(0) &< 0, && \left(\frac{f(t)}{t} \right)' < 0, \\ f''(t) &\leq 0 \quad \forall t > 0 && \text{and} \quad f(t) > 0 \quad \text{for some } t > 0, \end{aligned}$$

there is an interval $(R_1, R_2]$ such that for all $R \in (R_1, R_2]$ there exists a unique positive radial solution of

$$\begin{aligned} -\Delta u &= f(u) && \text{in } B_R \subset \mathbf{R}^n, \\ u &= 0 && \text{on } \partial B_R \end{aligned}$$

and only when $R = R_2$, the necessary condition for symmetry breaking, namely

$$\partial u / \partial n = 0 \quad \text{on } \partial B_R,$$

is satisfied. They conjecture that the symmetry breaks at R_2 . In both [5] and [1], it is shown that there exists a nonempty subclass of functions satisfying (1.2), for which this conjecture is true. However only for a "generic" subclass, the result is true, mainly due to the difficulty involved in verifying a certain nondegeneracy. Here we give an explicit class of functions depending on a parameter λ , namely

$$f(u, \lambda) = u^p - \lambda, \quad p > 1,$$

Received by the editors November 5, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 35J25; Secondary 35P30.

©1987 American Mathematical Society
 0002-9947/87 \$1.00 + \$.25 per page

for which the symmetry breaks when we consider the problem in the unit ball. However this function does not satisfy (1.2).

The one dimensional problem for the odd function $f(u, \lambda) = u|u|^{p-1} - \lambda$ was studied in [3]. The behavior of the solution branches described there leads to the conjecture that for the higher dimensional analogue of the problem, when a positive solution satisfies $\partial u / \partial n = 0$ on $\partial \Omega$, symmetry breaks at that point. We prove here that it does happen indeed.

2. Main results. Consider the equation

$$(2.1) \quad \begin{aligned} -\Delta u &= u^p - \lambda & \text{in } \Omega \equiv B_1 \subset \mathbf{R}^n, \\ u &= 0 & \text{on } \partial \Omega. \end{aligned}$$

Here $p < (n+2)/(n-2)$ in view of Pohozhaev's nonexistence result. From [2], we know that the positive solutions of (2.1) are radial. At $\lambda = 0$, the unique positive solution u_0 of (2.1) is in fact nondegenerate. Since a proof of this nondegeneracy could not be located in the literature, we indicate the proof here for the sake of completeness. Now onwards, we let

$$f(u) = u^p, \quad 1 < p < (n+2)/(n-2).$$

Lemma 1 will show that $(u_0, 0)$ is radially nondegenerate. We just remark here that nonradial degeneracy can also be ruled out at $(u_0, 0)$ and in fact all along the positive solution branch with $\partial u / \partial n < 0$ on $\partial \Omega$.

LEMMA 1. *Let u be a positive solution of (2.1) with $\partial u / \partial n \leq 0$ on $\partial \Omega$. Assume that there exists a radial eigenfunction v satisfying*

$$(2.2) \quad \begin{aligned} -\Delta v &= f'(u)v & \text{in } \Omega, \\ v &= 0 & \text{on } \partial \Omega. \end{aligned}$$

Then

$$(2.3) \quad v'(1)u'(1) = -\frac{2p\lambda}{p-1} \int_0^1 r^{n-1}v(r)dr.$$

PROOF. It is easy to verify

$$-\Delta(ru') = f'(u)(ru') + 2(f(u) - \lambda) \quad \text{in } \Omega.$$

Now using the equation for u , we have

$$(2.4) \quad \begin{aligned} -\Delta \left(ru' + \frac{2}{p-1}u \right) &= f'(u) \left(ru' + \frac{2}{p-1}u \right) - \frac{2p}{p-1}\lambda & \text{in } \Omega, \\ \left(ru' + \frac{2}{p-1}u \right) &= u'(1) & \text{on } \partial \Omega. \end{aligned}$$

We multiply (2.2) by $(ru' + 2u/(p-1))$ and (2.4) by v , subtract one from the other and integrate over Ω . then (2.3) follows. \square

At $(u_0, 0)$, if there were any radial degeneracy, then Lemma 1 yields an absurd equation $u'(1)v'(1) = 0$. Thus $(u_0, 0)$ has to be nondegenerate. Now the implicit function theorem gives the branch of positive solutions of (2.1) for λ close to 0.

We study the behavior of this branch for large positive λ . The following theorem is an extension of the nonexistence result in one dimension. (Cf. [3].)

THEOREM 1. *There exists no positive solution to equation (2.1) for large positive λ .*

PROOF. The proof follows by contradiction. Let if possible (u_j, λ_j) be a sequence of positive solutions, with $\lambda_j \rightarrow \infty$. Since u_j is radial, it satisfies

$$(2.5) \quad \begin{aligned} -u_j'' - (n-1)u_j'/r &= f(u_j) - \lambda_j \quad \text{in } (0, 1), \\ u_j'(0) &= 0; \quad u_j(1) = 0. \end{aligned}$$

Now $\Delta u_j < 0$ at 0, it being the maximum. At 1, $\Delta u_j > 0$. Thus there exists a point a where $f(u_j(a)) - \lambda_j = 0$. Now choose a point $b \in (a, 1)$ such that

$$(2.6) \quad f(u_j(b)) = \lambda_j/2.$$

We first prove that $b \rightarrow 1$ and hence a also, for $\lambda_j \rightarrow \infty$. We have in $[b, 1]$,

$$\frac{\lambda_j}{2} < \frac{1}{r^{n-1}} \left(\frac{d}{dr} (r^{n-1} u_j') \right).$$

Integration of this from b to 1 leads to

$$(2.7) \quad \frac{\lambda_j}{2n} (1 - b^n) < u_j'(1) - b^{n-1} u_j'(b) = b^{n-1} |u_j'(b)| - |u_j'(1)|.$$

Multiplying (2.5) by $u_j' r^{2(n-1)}$ and integrating between b and 1,

$$\begin{aligned} & \frac{1}{2} \left[u_j'^2(b) b^{2(n-1)} - u_j'^2(1) \right] \\ &= -2(n-1) \int_b^1 (F(u_j) - \lambda u_j) r^{2n-3} + [(F(u_j) - \lambda u) r^{2n-2}]_b^1 \\ &\leq 2(n-1) \int_b^1 (\lambda u_j - F(u_j)) r^{2n-3} + \left(2 - \frac{1}{p+1} \right) (u_j(b))^{p+1} b^{2n-2} \\ &\leq \left(u_j(b)^{p+1} \left(2(n-1) + 2 - \frac{1}{p+1} \right) \right). \end{aligned}$$

Here $F(u)$ stands for the primitive of f , namely $(u^{p+1})/(p+1)$. Further we have

$$\begin{aligned} |u_j'(b)| b^{n-1} - |u_j'(1)| &\leq \sqrt{u_j'(b) b^{2(n-1)} - u_j'^2(1)} \\ &\leq c |u_j(b)|^{(p+1)/2}. \end{aligned}$$

This along with (2.7) results in

$$(1 - b^n) < c_1 |u_j(b)|^{((p+1)/2)-p}.$$

As $p > 1$, this goes to 0 when $\lambda_j \rightarrow \infty$, in view of (2.6). Thus $b \rightarrow 1$.

The functions f and u_j being convex and f being monotone, the composition $f \circ u_j$ is convex in $[a, 1]$. Thus

$$\begin{aligned} f \circ u_j(1/2 + a/2) &\leq (f \circ u_j(a))/2 = f \circ u_j(b), \\ \Rightarrow (1+a)/2 &> b; \quad (1-b) > (b-a). \end{aligned}$$

Hence it follows that for λ_j large enough, $1 - a < \varepsilon$, for a preassigned ε . Let $\mu_1 = \mu_1(-\Delta, B_{1-\varepsilon})$ be the first eigenvalue of $-\Delta$ in $B_{1-\varepsilon}$, with Dirichlet boundary conditions. Define

$$v(x) = u_j(x) - u_j(a).$$

Choose λ_j large enough so that $f(u_j(a))/u_j(a) > \mu_1$. From the equation, we get

$$\begin{aligned} -\Delta v &> \mu_1(-\Delta, B_a)v \quad \text{in } B_a, \\ v &\geq 0 \quad \text{in } B_a \\ v &= 0 \quad \text{on } \partial B_a. \end{aligned}$$

This is easily shown to be impossible by comparing this with the eigenvalue equation in B_a . Thus the theorem holds. \square

Now it follows that the positive solution branch does not extend beyond a certain λ_0 . It has to encounter a singular point. Theorem 2 shows that this cannot be a radial singularity.

THEOREM 2. *There is no radial degeneracy all along the positive solution branch satisfying (2.1) with $\partial u/\partial n \leq 0$ on $\partial\Omega$.*

PROOF. Assume if possible, for some positive solution (u, λ) of (2.1), there exists a radial eigenfunction v satisfying (2.2). The idea is to get a contradiction by comparing the equation for a suitable test function with that of v . Define

$$w(r) = \int_0^r t^{n-1} v \, dt.$$

It is easy to verify

$$(2.8) \quad -\Delta w = -r^{n-1}v' - 2(n-1)r^{n-2}v \quad \text{in } \Omega.$$

Now multiplying (2.8) by v and (2.2) by w , integrating and subtracting one from the other,

$$v'(1)w(1) = - \int_0^1 r^{2(n-1)}vv' - 2(n-1) \int_0^1 r^{2n-3}v^2 - \int_0^1 f'(u)vw r^{n-1}.$$

After an integration by parts in the first and last terms, this leads to

$$(2.9) \quad v'(1)w(1) = -(n-1) \int_0^1 r^{2n-3}v^2 + \frac{1}{2} \int_0^1 f''(u)u'w^2.$$

Since $u > 0$ and $u' < 0$ in $(0, 1)$, the right-hand side is completely negative. v can always be chosen so as to have $v'(1) > 0$. Then it follows that

$$(2.10) \quad \int_0^1 r^{n-1}v < 0.$$

Now Lemma 1 gives

$$(2.11) \quad -((p-1)/2p\lambda)v'(1)u'(1) < 0.$$

This leads to a contradiction as $u'(1) \leq 0$. Thus the positive solution branch for (2.1) is radially nondegenerate. \square

From the above two theorems, we conclude that there exists a positive solution (u_0, λ_0) for (2.1) satisfying $\partial u_0/\partial n = 0$ on $\partial\Omega$. This is indeed a singular point since the linearized problem has at least n nonradial solutions $\{\partial u/\partial x_i\}_{1 \leq i \leq n}$. Lemma 2.2 of [1] (also cf. Theorem 20 of [4]) shows that there can be n nonradial solutions or $(n+1)$ solutions including one radial solution. Theorem 2 rules out the second possibility. Now to show the symmetry breaking at (u_0, λ_0) , we need the following definitions and Theorem 3.

We define

$$G(u, \lambda) \equiv -\Delta u - f(u) + \lambda$$

from the space

$$X = \{u \in C_0^{2,\alpha} : u(x_1, \dots, x_{n-1}, x_n) = u(-x_1, -x_2, \dots, -x_{n-1}, x_n)\}$$

into $C^\alpha(\Omega)$. We have

$$\begin{aligned} \ker G_u(u_0, \lambda_0) &= \text{span}\{\partial u_0 / \partial x_n\}, \\ \text{dimension}(\ker G_u^0) &= \text{codimension}(RaG_u^0) = 1. \end{aligned}$$

Let ϕ be the unique solution of the problem

$$(2.12) \quad G_u^0 \phi + G_\lambda^0 = 0; \quad \int_\Omega \phi \frac{\partial u_0}{\partial x_n} = 0.$$

Note that $G_\lambda^0 = 1$ is in $Ra(G_u^0)$ since

$$\int_\Omega 1 \cdot \frac{\partial u_0}{\partial x_n} = 0.$$

Now we prove the symmetry breaking.

THEOREM 3. (u_0, λ_0) is a nonradial bifurcation point.

PROOF. We use standard arguments to derive the bifurcation equation. This extends the proof for one dimension as in Proposition (5.1) in [3]. Assuming that the branch is given by $(u(s), \lambda(s))$, we have

$$(2.13) \quad G(u(s), \lambda(s)) = 0 \quad \text{for } s \in (s_0 - \varepsilon, s_0 + \varepsilon)$$

where $(u(s_0), \lambda(s_0)) = (u_0, \lambda_0)$. Differentiation with respect to s gives $G_u^0 u'(s) + G_\lambda^0 \lambda'(s) = 0$. This along with (2.12) yields,

$$u'(s) - \lambda'(s)\phi = \alpha_1 \phi_2,$$

for some scalar α_1 . Let

$$\lambda'(s) = \alpha_0, \quad u'(s) = \alpha_0 \phi + \alpha_1 \phi_2.$$

Second differentiation of (2.13) gives, at $s = s_0$,

$$G_u^0 u''(s) = -[G_{uu}^0(u'_0(s), u'_0(s)) + 2G_{u\lambda}^0(u'_0(s), \lambda'_0(s)) + G_{\lambda\lambda}^0(\lambda'_0(s))^2] - G_\lambda^0 \lambda''(s).$$

The bracketed term is in $Ra(G_u^0)$ and hence

$$\int \frac{\partial u_0}{\partial x_n} [- - -] = 0.$$

This after simplification results in

$$\begin{aligned} a\alpha_0^2 + 2b\alpha_0\alpha_1 + c\alpha_1^2 &= 0, \\ a &= \int \frac{\partial u_0}{\partial x_n} G_{uu}^0 \left(\frac{\partial u_0}{\partial x_n}, \frac{\partial u_0}{\partial x_n} \right), \\ b &= \int \frac{\partial u_0}{\partial x_n} G_{uu}^0 \left(\frac{\partial u_0}{\partial x_n}, \phi \right), \\ c &= \int \frac{\partial u_0}{\partial x_n} G_{uu}^0(\phi, \phi). \end{aligned}$$

Equation (2.12) can be rewritten as

$$(2.14) \quad \begin{aligned} -\Delta\phi &= f'(u)\phi - 1 && \text{in } \Omega, \\ \phi &= 0 && \text{on } \partial\Omega, \quad \int_{\Omega} \phi \frac{\partial u_0}{\partial x_n} = 0. \end{aligned}$$

Since there is no radial degeneracy, there exists a unique radial ϕ satisfying this equation. This ϕ satisfies the orthogonality condition also and hence by uniqueness this must be the solution of (2.14). Thus a and c have to vanish. We now show $b \neq 0$.

The uniqueness arguments given above and equation (2.4) give

$$(2.15) \quad \phi(r) = \frac{p-1}{2p\lambda} \left(ru' + \frac{2}{p-1}u \right), \quad \phi'(1) = \frac{p-1}{2p}.$$

By differentiating equation (2.1),

$$-\Delta(\Delta u) = f''(u)|\nabla u|^2 + f'(u)\Delta u \quad \text{in } \Omega.$$

Green's formula gives

$$\int_{\Omega} (\Delta u \cdot \Delta\phi - \phi\Delta(\Delta u)) = \int_{\partial\Omega} \frac{\partial\phi}{\partial n} \Delta u - \int_{\partial\Omega} \phi \frac{\partial}{\partial n} (\Delta u),$$

which after simplification results in

$$-\lambda\phi'(1)\text{vol}(\partial\Omega) = \int_{\Omega} f''(u)|\nabla u|^2 \phi.$$

(2.15) now shows $b \neq 0$, since at least for some i ,

$$\int_{\Omega} f''(u) \left(\frac{\partial u}{\partial x_i} \right)^2 \phi \neq 0.$$

Without loss of generality, we let $i = n$.

We have at (u_0, λ_0) :

- (i) G_u^0 is singular and $\dim(\ker G_u^0) = 1$; $\text{codimension}(Ra G_u^0) = 1$.
- (ii) $G_{\lambda}^0 \in Ra(G_u^0)$.
- (iii) The nondegeneracy condition $b \neq 0$ holds.

Now using standard arguments (cf., for example, [1]) it is easy to show that indeed (u_0, λ_0) is a bifurcation point. The two distinct curves cutting at (u_0, λ_0) have the tangent directions

$$(du_1, d\lambda_1) = (1, \phi), \quad (du_2, d\lambda_2) = \left(0, \frac{\partial u_0}{\partial x_n} \right)$$

the first being the radial solution branch and the latter being the branch of nonradial solutions.

3. Concluding remarks. After analyzing the proofs more carefully, the following remarks seem in order.

REMARK 1. In one dimension, for the odd nonlinearity $f(u) = u|u|^{p-1}$, a global picture was obtained in [2] by a scaling argument. When $n > 1$, this scaling argument works only for $B_{z(\lambda)}$ where $z(\lambda)$ is the first zero of $u(r)$. It fails outside because of the presence of the term $(n-1)u'/r$. But we observe that the proof

of Theorem 1 will go through for the nonlinearity $f(u) + \lambda$, in the annulus $\tilde{\Omega} = B_1 \setminus B_{a(\lambda)}$, provided $a(\lambda)$, the last zero of $u(r)$ before 1, tends to 0 as λ tends to ∞ . Thus we can conclude that each S_k^\pm , $k \geq 1$, can exist only in $[-\lambda_k, \tilde{\lambda}_k]$, where $S_k^\pm = \{(u, \lambda): \text{a solution of (2.1), } u \text{ has } (k-1) \text{ nodes in } (0, 1), u(0) > 0 \text{ (or } < 0)\}$. As in one dimension, we expect the positive solution branch S_1^+ to continue as S_3^+ , etc., by developing more and more nodes. Each time a new node is developed, we get a solution $(u, \lambda) \in S_k^\pm$ with $\partial u / \partial n = 0$ on $\partial\Omega$. These are potential symmetry breaking points. We conjecture that the radial solution branch has purely radial degeneracy (turning points) and purely nonradial degeneracy (symmetry breaking points) alternately.

REMARK 2. Even though this paper discusses a special class of functions, what seems to be interesting is the way we prove the nondegeneracy. Perhaps it is possible to prove the nondegeneracy (or uniqueness) using similar test functions, wherever the time map is used to prove such results. Hopefully this approach may lead to the settling of the conjecture of Smoller and Wasserman [5].

REMARK 3. Recall that in view of Theorem 1, we concluded that the positive solution branch has to encounter a singular point. This is a consequence of Theorem 1 and also the existence of a priori bounds for positive solutions. (See DeFigueiredo, Lions, Nussbaum, *A priori estimates and existence of positive solutions of semilinear elliptic equations*, J. Math. Pures Appl. **61** (1983)).

REMARK 4. Since the unique positive solution of

$$\begin{aligned} -\Delta u &= u^p \quad \text{in } B_1, \quad p < (n+2)/(n-2), \\ u &= 0 \quad \text{on } \partial B_1 \end{aligned}$$

is nondegenerate, we can get a branch of positive solutions, using implicit function theorem, for ε in $[0, \varepsilon_0)$, for equation

$$\begin{aligned} -\Delta u &= u^p + \varepsilon u^q \quad \text{in } B_1, \\ u &= 0 \quad \text{on } \partial B_1 \end{aligned}$$

for any q . This fact has already been observed in a more general set up by Rabinowitz for $q > (n+2)/(n-2)$ (*Variational methods for nonlinear elliptic eigenvalue problems*, Indiana Univ. Math. J. **23** (1974)) and by Brezis and Nirenberg (*Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*, Comm. Pure Appl. Math. **36** (1983)) for $q = (n+2)/(n-2)$, through different arguments.

REFERENCES

1. G. Cerami, *Symmetry breaking for a class of semilinear elliptic problems*, Nonlinear Anal. **10** (1986), 1-14.
2. B. Gidas, W. M. Ni and L. Nirenberg, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys. **68** (1979), 209-243.
3. M. Ramaswamy, *On the global set of solutions to a nonlinear ODE-theoretical and numerical description*, J. Differential Equations **65** (1986), 1-48.
4. J. A. Smoller and A. G. Wasserman, *Existence, uniqueness and nondegeneracy of positive solutions of semilinear elliptic equations*, Comm. Math. Phys. **95** (1984), 129-159.
5. —, *Symmetry breaking for positive solutions of semilinear elliptic equations*, Arch. Rational Mech. Anal. **95** (1986), 217-225.