

CONJUGACY CLASSES WHOSE SQUARE IS AN INFINITE SYMMETRIC GROUP

GADI MORAN

ABSTRACT. Let X_ν be the set of all permutations ξ of an infinite set A of cardinality \aleph_ν with the property: every permutation of A is a product of two conjugates of ξ . The set X_0 is shown to be the set of permutations ξ satisfying one of the following three conditions:

- (1) ξ has at least two infinite orbits.
- (2) ξ has at least one infinite orbit and infinitely many orbits of a fixed finite size n .
- (3) ξ has: no infinite orbit; infinitely many finite orbits of size k, l and $k + l$ for some positive integers k, l ; and infinitely many orbits of size > 2 . It follows that $\xi \in X_0$ iff some transposition is a product of two conjugates of ξ , and ξ is not a product σi , where σ has a finite support and i is an involution.

For $\nu > 0$, $\xi \in X_\nu$ iff ξ moves \aleph_ν elements, and satisfies (1), (2) or (3'), where (3') is obtained from (3) by omitting the requirement that ξ has infinitely many orbits of size > 2 . It follows that for $\nu > 0$, $\xi \in X_\nu$ iff ξ moves \aleph_ν elements and some transposition is the product of two conjugates of ξ .

The covering number of a subset X of a group G is the smallest power of X (if any) that equals G [AH]. These results complete the classification of conjugacy classes in infinite symmetric groups with respect to their covering number.

0. BACKGROUND AND RESULTS

Let A be a countably infinite set, and let $S = S_A$ denote the group of all permutations of A . For $X, Y \subseteq S$ let $X \cdot Y = \{\xi\eta : \xi \in X, \eta \in Y\}$ and let $X^1 = X$, $X^{n+1} = X^n \cdot X$. For $\xi \in S$ let $[\xi] = \{\eta\xi\eta^{-1} : \eta \in S\}$ denote the conjugacy class (coc) of ξ in S , and consider the equation

$$(1) \quad [\xi]^2 = S.$$

Let $X_0 \subseteq S$ be the set of all $\xi \in S$ satisfying (1). We shall establish

Theorem 1. *Let $\xi \in S$. Then $\xi \in X_0$ if and only if ξ satisfies one of the following three mutually exclusive conditions:*

- (1) ξ has at least two infinite orbits.

Received by the editors August 3, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 20B30, 20B07; Secondary 20E34, 05C25.

- (2) (a) ξ has precisely one infinite orbit.
- (b) ξ has infinitely many orbits of a fixed finite size n .
- (3) (a) ξ has no infinite orbit.
- (b) For some positive integers k, l , ξ has infinitely many orbits of size n , for $n = k, n = l$ and $n = k + l$.
- (c) ξ has infinitely many orbits of size greater than 2.

Much of the knowledge and technique accumulated in evaluating products of cocs in S is used in a full proof of Theorem 1. Recently Droste [D3] combined these to establish Theorem 1 for those $\xi \in S$ which have at least one infinite orbit, or fix infinitely many elements. To complete the argument one more new theorem is needed, namely, Theorem A, which we state below.

Let $\text{INF} \subseteq S$ be the set of all $\xi \in S$ that move infinitely many elements. Call $\xi \in \text{INF}$ *almost involution* if ξ has no infinite orbit and finitely many finite orbits of length > 2 (hence ξ has infinitely many orbits of size 2). Let $\text{AI} \subseteq \text{INF}$ be the set of all almost involutions.

Theorem A. *Let $\xi, \eta \in \text{INF}$ have no infinite orbit, but infinitely many finite orbits of length at least k , $k \in \{3, 4\}$.*

- (1) *If $k = 3$ then $\text{INF-AI} \subseteq [\xi] \cdot [\eta]$.*
- (2) *If $k = 4$ then $\text{INF} \subseteq [\xi] \cdot [\eta]$.*
- (3) *The value $k = 3$ ($k = 4$) in (1) (in (2)) cannot be reduced.*

Our tool for establishing Theorem A is the use of bicolored planar maps, first introduced in [M4] (see also [M5]). This technique is reviewed in §2.

A brief history about the search for X_0 is in order. In 1951 Ore [O] proved that every member of S is a commutator. Since $[\xi] = [\xi^{-1}]$ holds in S , $\eta = \zeta \xi \zeta^{-1} \xi^{-1}$ implies $\eta \in [\xi]^2$ and so Ore's result shows that S is a union of squares of conjugacy classes.

In 1960 Gray [G] showed that if $\xi \in S$ has infinitely many infinite orbits and no finite orbits, then $S = [\xi]^2$, and so $X_0 \neq \emptyset$. He also noted that the obvious necessary condition $\xi \in \text{INF}$ is not sufficient for $\xi \in X_0$.

In 1973 Bertram [B] showed that if $\xi \in S$ has infinitely many orbits of length n for $n = 1, 2, 3$ but no other orbit, then $\xi \in X_0$. He used it to show that $[\xi]^4 = S$ for every $\xi \in \text{INF}$ (see also [DG1]), and conjectured that actually $[\xi]^3 = S$ for $\xi \in \text{INF}$.

Shortly afterwards it was noted [M1; M2, (2), p. 76] (see also [DG1]) that if $\rho_0 \in S$ is a fixed-point-free involution, $R_0 = [\rho_0]$, then $R_0^3 \neq S$ (see [M1, M6] for the actual evaluation of R_0^3). However, twelve years later Droste [D2] showed that R_0 is the *only* coc in S for which Bertram's conjecture fails.

In 1980 we determined [M3] the set of $\xi \in S$ with parity features, i.e. with the property that $[\xi]^2$ contains no finitary odd permutations ($\xi \in S$ is called *finitary* if it moves only finitely many elements). We now formulate as Theorem

2 an unpublished observation made in the course of that investigation. This result turns out to have much significance in the present context.

Let $Y_0 \subseteq S$ be the set of η 's satisfying:

- (2) $[\eta]^2$ contains a transposition.

Theorem 2. *Let $\xi \in S$. Then $\xi \in Y_0$ if and only if ξ satisfies one of the three mutually exclusive conditions (1), (2), or (3)(a) and (3)(b) mentioned in Theorem 1.*

A proof of Theorem 2 is given in §1.

Comparing Theorems 1 and 2 we conclude:

Corollary 3. *Let $\xi \in \text{INF}$. Then $[\xi]^2 = S$ if and only if:*

- (i) $[\xi]^2$ contains a transposition.
- (ii) ξ is not almost involution.

The first indication that X_0 and Y_0 are closely related was given in 1982 in Droste's paper [D1]. Indeed, Theorem 1 of [D1], implies that if $\xi \in S$ has at least one infinite orbit, then $\xi \in X_0$ if and only if $\xi \in Y_0$. Other interesting results of this paper fit into the research on covering numbers of groups that was carried out intensively at the same time (see [AH]). If C is a coc in a group G , we write $\text{cn}(C) = n$ if n is the smallest positive integer such that $C^n = G$. $\text{cn}(C)$ is called the *covering number of G by C* and, when G is fixed in context, *the covering number of C* . If no such integer exists, we write $\text{cn}(C) = \omega$. During 1980–1983 these numbers were computed for various cocs in various groups, and the effort to obtain them in simple factors of the decomposition chain of infinite symmetric groups led us to the discovery and use of the method of bicolored planar maps [ACM, M4, M5]. Droste's work [D1] has immediate relevance to this work as well (see [D1, §5]).

Combining his old results on products of cocs in S with the fresh ones obtained in the course of research on the covering numbers, Droste [D2] obtained the remarkable result mentioned above, that R_0 is the *only* coc C in S with $\text{cn}(C) = 4$. Theorem 1 determines the cocs C in S with $\text{cn}(C) = 2$. The remaining cocs in INF —including those with Parity Features (see [M3])—all have covering number 3. Since $\text{cn}(C) = \omega$ for every finitary coc C , we know now $\text{cn}(C)$ for every coc C in S .

Applying this same body of knowledge and some more recent acquaintance with products of cocs in finite symmetric groups due independently to Boccara [Bo] and Dvir [Dv], Droste then found that X_0 and Y_0 are indeed very close. In [D3, Theorem 2], he shows that if ξ has *no* infinite orbit, and infinitely many fixed points, then $\xi \in X_0$ iff $\xi \in Y_0$ and ξ is not an almost-involution. The complete determination of X_0 is reduced, then, to familiarity with those ξ 's in X_0 that have no infinite orbit, and only finitely many fixed points. Now, $X_0 \subseteq Y_0$ and Theorem 2 further reduce the search to the ones that satisfy

conditions (3)(a) and (3)(b), stated in Theorem 1. Theorem A provides us with the missing link, and closes the argument for Theorem 1.

Let us review some relevant results. For an arbitrary group G , let X_G denote the set of $g \in G$ satisfying $[g]^2 = G$. For an arbitrary ordinal ν , let S_ν denote the group of all permutations of a set of cardinality \aleph_ν , and let $X_\nu = X_{S_\nu}$. Let $S_\nu^\tau \subseteq S_\nu$ be the normal subgroup of S_ν consisting of permutations moving less than \aleph_τ elements and let $H_\tau^\nu = S_\nu^{\tau+1}/S_\nu^\tau$, $\tau = 0, \dots, \nu$. Then we have

Proposition 4 [M4, ACM]. *Let $\nu > 0$. Then $X_G \cup \{1\} = G$ if $G = H_\tau^\nu$, $\tau = 0, \dots, \nu + 1$. Also, $X_G \cup \{1\} = G$ if $G = J_1$, where J_1 is the smallest Janko group.*

J_1 is the only finite group G known to satisfy $X_G \cup \{1\} = G$. Infinite groups G satisfying $X_G \cup \{1\} = G$ were noted earlier (see e.g. [H]).

Fix an ordinal ν . For any ordinal τ let Y_ν^τ denote the set of all $\xi \in S_\nu^\tau$ satisfying (2), that is, $[\xi]^2$ contains a transposition. Obviously, Theorem 2 implies its generalization obtained by replacing “ $\xi \in S$ ” by “ $\xi \in S_\nu$ ”.

Corollary 5. *Let $\nu > 0$, let $0 \leq \tau \leq \nu$, and let $\xi \in S_\nu^{\tau+1}$. The following are equivalent:*

- (a) $\xi \in X_\nu$.
- (b) $\xi \in Y_\nu^\tau$ and ξ moves \aleph_τ elements.

This corollary is established by Droste [D3, Corollary 3] for the case $0 \leq \tau < \nu$ from the restricted formulation of Theorem 1. Theorem 1 in full easily implies it for the case $0 < \tau = \nu$, by the same argument.

The paper is organized as follows: In §1 we introduce some convenient notation, and use it to reformulate Theorems 1, 2 and A as Theorems 1', 2' and A'. We then prove Theorem 1' using Theorem 2', and Theorem A'. A proof of Theorem 2' is also given there.

Theorem A', the main new result of this paper, is established in §§2–4. In §2 we review the method of using bicolored planar maps. In §3 Theorem A' is reduced to two theorems, Theorem B and Theorem C, whose proof relies on the use of bicolored planar maps.

Theorem B provides the inclusion claimed in Theorem A(1) and (2) for permutations ζ in INF-AI (in case (1)) or in INF (in case (2)) that have no infinite orbit, and is proved in §3.

Theorem C takes care of the most difficult case—when ζ has at least one infinite orbit (where one easily sees that precisely one infinite orbit is the crucial case). It is proved in §4.

1. VOCABULARY

In this section we introduce our notation and use it to reformulate our results.

\mathbb{Z} denotes the set of integers, \mathbb{N} denotes the set of positive integers, and $\mathbb{N}_0 = \{0\} \cup \mathbb{N}$, $\mathbb{N}^+ = \mathbb{N} \cup \{\aleph_0\}$. $|A|$ denotes the cardinality of the set A and S_A

the group of all permutation of A . For $a \in A$ and $\xi \in S_A$ we let $\xi(a)$ denote the value taken by ξ at a . Hence the composition $\xi\zeta$ is to be computed by acting first with ζ , then with ξ . $1_A \in S_A$ is the identity permutation: $1_A(a) = a$ for all $a \in A$. If $B \subseteq A$, $\xi|B$ denotes the restriction of ξ to B . If $\xi|B \in S_B$ then B is called ξ -invariant. If $A = B_1 \cup B_2$, $B_1 \cap B_2 = \emptyset$, B_1 and B_2 are ξ -invariant and $\xi_i \in S_{B_i}$ is $\xi|B_i$, $i = 1, 2$, then we write $\xi = \xi_1 + \xi_2$. More generally, if $\{B_i; i \in I\}$ is a partition of A into ξ -invariant sets, $\xi_i = \xi|B_i$, $i \in I$, then we write $\xi = \sum_{i \in I} \xi_i$.

For $\xi \in S_A$ and $a \in A$ let $(a)_\xi = \{\xi^n(a); n \in \mathbb{Z}\}$. $(a)_\xi$ is called the ξ -orbit of a . Let $(A)_\xi = \{(a)_\xi; a \in A\}$ be the partition of A into ξ -orbits, and for $n \in \mathbb{N}^+$ let $(A)_{\xi,n} = \{B \in (A)_\xi; |B| = n\}$. We define the cardinal number $\bar{\xi}(n)$ by

$$\bar{\xi}(n) = |(A)_{\xi,n}| = \text{cardinality of the set of } \xi\text{-orbits of cardinality } n.$$

By a *type* we mean a cardinal valued function t defined on \mathbb{N}^+ . Thus, for any permutation $\xi \in S_A$, $\bar{\xi}$ is a type. Moreover, for $\xi, \eta \in S_A$ we have $[\xi] = [\eta]$ iff $\bar{\xi} = \bar{\eta}$ (see e.g. [S]). For types s, t and a cardinal number k we define the types $s + t$ and kt by $(s + t)(n) = s(n) + t(n)$ and $(kt)(n) = k \cdot t(n)$ ($n \in \mathbb{N}^+$). More generally, if t_i is a type for all $i \in I$, then a type $\sum_{i \in I} t_i$ is defined by

$$\left(\sum_{i \in I} t_i \right) (n) = \sum_{i \in I} t_i(n) \quad (n \in \mathbb{N}^+).$$

Obviously we have

$$\overline{\xi + \eta} = \bar{\xi} + \bar{\eta}, \quad \overline{\sum_{i \in I} \xi_i} = \sum_{i \in I} \bar{\xi}_i.$$

Given three types r, s, t we write $P(r, s, t)$ for the following statement:

There is a set A and $\xi, \eta, \zeta \in S_A$ such that $\bar{\xi} = r$, $\bar{\eta} = s$, $\bar{\zeta} = t$ and $\xi\eta\zeta = 1_A$.

We have $[\xi] \subseteq [\eta][\zeta]$ iff $P(\bar{\xi}, \bar{\eta}, \bar{\zeta})$.

The main properties of P are (see [M4]):

Symmetry. $P(t_1, t_2, t_3)$ iff $P(t_{\theta(1)}, t_{\theta(2)}, t_{\theta(3)})$ for some $\theta \in S_{\{1, 2, 3\}}$,
iff $P(t_{\theta(1)}, t_{\theta(2)}, t_{\theta(3)})$ for all $\theta \in S_{\{1, 2, 3\}}$.

Superadditivity. If $P(r_i, s_i, t_i)$ for all $i \in I$, then

$$P\left(\sum_{i \in I} r_i, \sum_{i \in I} s_i, \sum_{i \in I} t_i\right).$$

From superadditivity follows also

Homogeneity. $P(r, s, t)$ implies $P(kr, ks, kt)$ for any cardinal number k .

For $n \in \mathbb{N}^+$ define the types n^*, \tilde{n} by:

$$n^*(n) = 1, \quad n^*(m) = 0 \quad \text{for } m \neq n, \quad \tilde{n} = \aleph_0 \cdot n^*.$$

Then for any type t we have $t = \sum_{n \in \mathbb{N}^+} t(n) \cdot n^*$. Define also $|t| = \sum_{n \in \mathbb{N}^+} t(n) \cdot n$. Then for $\xi \in S_A$ we have $|\bar{\xi}| = |A|$.

If $K \subseteq \mathbb{N}^+$, t a type, we set $t[K] = \sum_{n \in K} t(n)$. We adopt the usual conventions for intervals in \mathbb{N}^+ : if $k, l \in \mathbb{N}^+$ then:

$$(k, l) = \{n \in \mathbb{N}^+ : k < n < l\}, \quad [k, l) = \{n \in \mathbb{N}^+ : k \leq n < l\},$$

$$(k, l] = \{n \in \mathbb{N}^+ : k < n \leq l\}, \quad [k, l] = \{n \in \mathbb{N}^+ : k \leq n \leq l\}.$$

Let $\xi \in S_A$. We define $M(\xi) \subseteq A$ by $M(\xi) = \{a \in A : \xi(a) \neq a\}$.

A permutation $\xi \in S_A$ is *finitary* if $M(\xi)$ is finite, and a type t is *finitary* if $t = \bar{\xi}$ for some finitary ξ . So t is finitary iff $t(\aleph_0) = 0$ and $t[(1, \aleph_0)] < \aleph_0$.

Let $\mathbf{x}_0 = \{\bar{\xi} : \xi \in X_0\}$. Thus, $t \in \mathbf{x}_0$ iff $P(t, s, t)$ holds true for every type s with $|s| = \aleph_0$.

We formulate Theorem 1, Theorem A and Theorem 2 as follows:

Theorem 1'. *Let t be a type. The following are equivalent:*

- (i) $t \in \mathbf{x}_0$.
- (ii) $|t| = \aleph_0$ and t satisfies one of the following three mutually exclusive conditions:

$$(1') \quad t(\aleph_0) \geq 2.$$

$$(2') \quad (a) \quad t(\aleph_0) = 1.$$

$$(b) \quad t(n) = \aleph_0 \text{ for some } n \in \mathbb{N}.$$

$$(3') \quad (a) \quad t(\aleph_0) = 0.$$

$$(b) \quad t(k) = t(l) = t(k+l) = \aleph_0 \text{ for some } k, l \in \mathbb{N}.$$

$$(c) \quad t[(2, \aleph_0)] = \aleph_0.$$

Let $\mathbf{inf} = \{\bar{\xi} : \xi \in \text{INF}\}$. Thus $t \in \mathbf{inf}$ iff $|t| = \aleph_0$ and $t(\aleph_0) > 0$ or $t[(1, \aleph_0)] = \aleph_0$. Define $\mathbf{ai} \subseteq \mathbf{inf}$ by $\mathbf{ai} = \{\bar{\xi} : \xi \in \text{AI}\}$. Thus $t \in \mathbf{ai}$ iff $t(2) = \aleph_0$, $t[(2, \aleph_0)] < \aleph_0$ and $t(\aleph_0) = 0$.

Theorem A'. *Let $r, t \in \mathbf{inf}$ satisfy $r(\aleph_0) = t(\aleph_0) = 0$.*

- (1) *If $r[[3, \aleph_0]] = t[[3, \alpha_0]] = \aleph_0$ and $s \in \mathbf{inf-ai}$, then $P(r, s, t)$.*
- (2) *If $r[[4, \aleph_0]] = t[[4, \aleph_0]] = \aleph_0$ and $s \in \mathbf{inf}$, then $P(r, s, t)$.*
- (3) *Statement (1) fails if we require only $r[[2, \aleph_0]] = t[[2, \aleph_0]] = \aleph_0$. Statement (2) fails if we require only $r[[3, \aleph_0]] = t[[3, \aleph_0]] = \aleph_0$.*

Let $\mathbf{y}_0 = \{\bar{\xi} : \xi \in Y_0\}$. Thus, $t \in \mathbf{y}_0$ iff $P(t, \tilde{1} + 2^*, t)$. Obviously, $\mathbf{x}_0 \subseteq \mathbf{y}_0$.

Theorem 2'. *Let t be a type. The following are equivalent:*

- (i) $t \in \mathbf{y}_0$.
- (ii) $|t| = \aleph_0$ and t satisfies one of the following three mutually exclusive conditions:

$$(1'') \quad t(\aleph_0) \geq 2.$$

$$(2'') \quad t(\aleph_0) = 1 \text{ and } t(n) = \aleph_0 \text{ for some } n \in \mathbb{N}.$$

$$(3'') \quad t(\aleph_0) = 0 \text{ and } t(k) = t(l) = t(k+l) = \aleph_0 \text{ for some } k, l \in \mathbb{N}.$$

We now reformulate two earlier results and use them, together with Theorem A' and Theorem 2', to prove Theorem 1'. The first, due to Droste, is a restricted form of Theorem 1'.

Proposition 1.1 (Droste [D1, Theorem 1; D3, Theorem 2]). *Let t be a type, $|t| = \aleph_0$, and $t(\aleph_0) \geq 1$ or $t(1) = \aleph_0$. Then the following are equivalent:*

- (i) $t \in \mathbf{x}_0$.
- (ii) t satisfies one of the three mutually exclusive conditions (1'), (2'), (3') of Theorem 1'.

(The fact that if t satisfies (1') of Theorem 1' then $t \in \mathbf{x}_0$ was noted already in 1981 by Droste and Göbel; see [DG2].)

Proposition 1.2 (see [M2, Corollary 2.5; D2, Theorem 4.1]). *Let $t \in \mathbf{ai}$ and let $P(t, s, t)$. Then $s[\mathbb{N}^+] = \aleph_0$. Hence $t \notin \mathbf{x}_0$, as $P(t, \aleph_0^*, t)$ does not hold, for instance.*

Proof of Theorem 1' (assuming Theorems 2' and A').

(i) \rightarrow (ii) Let $t \in \mathbf{x}_0$. By $\mathbf{x}_0 \subseteq \mathbf{y}_0$, $t \in \mathbf{y}_0$. By Theorem 2', t satisfies one of the conditions (1'), (2'), or (3''). But (3'') is equivalent to (3')(a) and (3')(b). Thus, it is left to show that if t satisfies (3'') it satisfies also (3')(c), i.e., $t \notin \mathbf{ai}$. But $t \notin \mathbf{ai}$ follows from Proposition 1.2.

(ii) \rightarrow (i) Let t satisfy (ii), i.e., $|t| = \aleph_0$ and t satisfies (1'), (2') or (3'). If t satisfies (1') or (2') then $t \in \mathbf{x}_0$ by Proposition 1.1. If t satisfies (3') and $t(1) = \aleph_0$ then again $t \in \mathbf{x}_0$ by Proposition 1.1, so assume t satisfies (3') but $t(1) < \aleph_0$. By (3')(b) $t(k) = t(l) = t(k+l) = \aleph_0$ for some $k, l \in \mathbb{N}$, and by $t(1) < \aleph_0$, $2 \leq k, 2 \leq l$ so $k+l \geq 4$. Thus, $t[[4, \aleph_0]] = \aleph_0$. By Theorem A'(2) and [D3, Theorem 1], $t \in \mathbf{x}_0$. \square

Proof of Theorem 2'. We first make the following observation: Let $A = B \cup B'$, $B \cap B' = \emptyset$, $b \in B$, and $b' \in B'$. Let $\xi \in S_A$ and let $B = (b)_\xi$, $B' = (b')_\xi$ (so B, B' are ξ -orbits). Let (b, b') be the transposition interchanging b and b' . Let $\zeta = (b, b')\xi$. Then we have:

- (1) If $|B| = |B'| = \aleph_0$ then $\bar{\zeta} = 2 \cdot \aleph_0^*$, i.e. ζ has two infinite orbits.
- (2) If $|B| = \aleph_0$ and $|B'| = n < \aleph_0$ then $\bar{\zeta} = \aleph_0^*$, i.e. ζ has one infinite orbit.
- (3) If $|B| = k$ and $|B'| = l$, $k, l < \aleph_0$, then $\bar{\zeta} = (k+l)^*$, i.e. ζ has one orbit, of size $k+l$.

Thus we have

- (1) $P(2 \cdot \aleph_0^*, \hat{1} + 2^*, 2 \cdot \aleph_0^*)$.
- (2) $P(\aleph_0^*, \hat{1} + 2^*, n^* + \aleph_0^*)$, $n \in \mathbb{N}$.
- (3) $P((k+l)^*, (k+l-2) \cdot 1^* + 2^*, k^* + l^*)$.

Also:

- (4) $P(t, |t| \cdot 1^*, t)$ holds for any type t .

(ii) \rightarrow (i) Assume that $|t| = \aleph_0$ and t satisfies (1'), (2') or (3'). We show that $t \in \mathbf{y}_0$, i.e., $P(t, \tilde{1} + 2^*, t)$.

In case (1') we have $t = 2 \cdot \aleph_0^* + t'$ so by (1) and (4) $P(t, \tilde{1} + 2^*, t)$.

In case (2') we have $t = \aleph_0^* + \tilde{n} + t' = n^* + \aleph_0^* + \tilde{n} + t'$, so by (2) and (4) $P(t, \tilde{1} + 2^*, t)$.

In case (3') we have $t = \tilde{k} + \tilde{l} + (\widetilde{k+l}) + t'$, and so we have $t = (k+l)^* + t' = k^* + l^* + t'$, and so by (3) and (4) $P(t, \tilde{1} + 2^*, t)$.

(i) \rightarrow (ii) Let $t \in \mathbf{y}_0$. Then there are $\xi, \zeta \in S_A$ and a transposition $(b, b') \in S_A$ such that $\zeta = (b, b')\xi$, and $\bar{\xi} = \bar{\zeta} = t$. We may further assume that b and b' belong to distinct ξ -orbits B and B' (else b and b' belong to distinct ζ -orbits and we have $\xi = (b, b')\zeta$). Let $\xi = \xi' + \xi''$, $\zeta = \zeta' + \zeta''$ where $\xi', \zeta' \in S_{B \cup B'}$. Then we have $\zeta' = (b, b')\xi'$ (where here $(b, b') \in S_{B \cup B'}$), $\xi'' = (\zeta'')^{-1}$. Hence $\bar{\xi}' = |B|^* + |B'|^*$, and $\bar{\xi}'' = \bar{\zeta}''$. Now $t = \bar{\xi}' + \bar{\xi}'' = \bar{\zeta}' + \bar{\zeta}''$, and $P(\bar{\xi}', \overline{(b, b')}, \bar{\xi}'')$ imply that t satisfies (1'), (2') or (3') as follows.

We know that $\bar{\xi}' = 2 \cdot \aleph_0^*$, or $\bar{\xi}' = \aleph_0^* + n^*$, or $\bar{\xi}' = k^* + l^*$ ($n, k, l \in \mathbb{N}$). If $\bar{\xi}' = 2 \cdot \aleph_0^*$ then $t(\aleph_0) \geq 2$ so (1') holds. If $\bar{\xi}' = \aleph_0^* + n^*$ then $\bar{\zeta}' = \aleph_0^*$ and so setting $t'' = \bar{\xi}'' = \bar{\zeta}''$ we have $t = \aleph_0^* + n^* + t'' = \aleph_0^* + t''$, so $t(\aleph_0) \geq 1$, and by $t(n) = t''(n) + 1 = t''(n)$, we have $t(n) = \aleph_0$. Thus t satisfies (1') or (2').

If $\bar{\xi}' = k^* + l^*$ then $\bar{\zeta}' = (k+l)^*$ and we obtain as before $t = k^* + l^* + t'' = (k+l)^* + t''$, so $t(k) = 1 + t''(k) = t''(k)$, $t(l) = 1 + t''(l) = t''(l)$, $t(k+l) = 1 + t''(k+l) = t''(k+l)$.

It follows that $t(k), t(l), t(k+l) \geq \aleph_0$, and t satisfy (1'), (2') or (3'). \square

2. BICOLORED PLANAR MAPS REVISITED

In this section we review briefly the bicolored-planar-map method introduced in [M4], and use it to establish Theorem A' in subsequent sections. The following treatment is self-contained and slightly different from the one offered in [M4].

By a *planar graph* we mean a pair $G = (V_G, E_G)$ where $V = V_G$ is a set of points in the plane, called the *vertices* of G , and $E = E_G$ is the set of *edges* of G . Each edge $e \in E$ is the range of a continuous mapping f of the closed unit interval $[0, 1]$ into the plane, whose restriction to the open unit interval $(0, 1)$ is a homeomorphism. $f(0)$ and $f(1)$ are called the *endpoints* of e . They should be vertices, and may be nondistinct. If $f(0) = f(1)$ we call e a *loop*. Two distinct edges may have the same set of endpoints (*multiplicity of edges* (including loops) is allowed). An edge e meets V by the set of its endpoints, and the intersection of two distinct edges is a set of vertices (so two edges meet at common endpoints). In addition to those usual conditions, we impose two more assumptions as follows:

(a) every bounded set meets only finitely many edges.

The *degree* $d(v)$ of $v \in V$ is defined as the number of times v occurs as an endpoint of an edge (so a loop contributes two to this count). It follows from (a) that every vertex has a finite degree.

We let $G_* = V \cup (\bigcup E)$ denote the set of all points of the plane in V or in e for some $e \in E$. Our second assumption is:

(b) G_* is connected.

It follows from (a) that the complement G_*^c of G_* is open. A connected component of G_*^c is called a *G-region*. It further follows from (b) that every *G-region* is simply connected, and so homeomorphic to an open disc. Let $F = F_G$ denote the set of all *G-regions*.

By a *map* we mean a triple (V, E, F) , where $G = (V, E)$ is a planar graph and $F = F_G$ is the set of *G-regions*.

By a *bicolored planar map* (*bpm*) we mean a quadruple (V, E, B, W) , where $(V, E, B \cup W)$ is a map, and for distinct $b, b' \in B$ ($w, w' \in W$) we have $\text{cl}(b) \cap \text{cl}(b') \subseteq V$ ($\text{cl}(w) \cap \text{cl}(w') \subseteq V$). Here $\text{cl}(X)$ is the closure of the set X in the plane. B (W) is the set of black (white) regions of the map.

Let (V, E, B, W) be a bpm, $e \in E$. Then there is a unique $b_e \in B$ and a unique $w_e \in W$ such that $e \in \text{cl}(b_e) \cap \text{cl}(w_e)$. For a region $f \in B \cup W$, let $F_f = \{e \in E: f = f_e\}$ be the set of edges in its closure, and let $s(f) = |E_f|$. We say that f is of *size* $s(f)$.

We now use the orientation of the plane to define three permutations $\beta, v, \omega \in S_E$ as follows. Orient $e \in E$ by requiring that a traveller moving on e as oriented sees b_e to his left and w_e to his right. (Thus, the boundary of a bounded $b \in B$ is oriented counterclockwise.) Define the *tail* $t(e)$, and that *head* $h(e)$, of $e \in E$ to be the endpoint travelled from and to by this traveller on e .

Let us say that e' follows e around $b \in B$ ($w \in W$) iff $b = b_e = b_{e'}$ ($w = w_e = w_{e'}$) and $h(e) = t(e')$.

Let $v \in V$, and let E_v denote the set of edges $e \in E$ satisfying $t_e = v$ (so E_v is the set of edges emanating from v). By (a) E_v is a finite set. Let $s(v) = |E_v|$. Thus, $s(v)$ the out-degree of v . We have $d(v) = 2s(v)$, as our orientation makes (V, E) into an Eulerian graph. We say that v is of *size* $s(v)$.

Let $e, e' \in E_v$. We say that e' follows e around v if for any small enough circle C centered at v , $P_{e'}$ follows P_e in the clockwise sense on C , where P_e ($P_{e'}$) is the first point on C met by a traveller on e (e') moving from $t(e) = v$ to $h(e)$ ($t(e') = v$ to $h(e')$).

Define β, v and ω to be the black, vertex and white permutations respectively acting on the set of edges E . That is:

$$\begin{aligned} \beta(e) &= e' \quad \text{iff } e' \text{ follows } e \text{ around } b_e = b_{e'}, \\ v(e) &= e' \quad \text{iff } e' \text{ follows } e \text{ around } t_e = t_{e'}, \\ \omega(e) &= e' \quad \text{iff } e' \text{ follows } e \text{ around } w_e = w_{e'}. \end{aligned}$$

One readily verifies:

Proposition 2.1. *Let (V, E, B, W) be a bicolored planar map. Then $\beta, v, \omega \in S_E$ defined above satisfy:*

- (1) $\{E_b : b \in B\}$ is the partition of E into β -orbits.
- (2) $\{E_v : v \in V\}$ is the partition of E into v -orbits.
- (3) $\{E_w : w \in W\}$ is the partition of E into ω -orbits.
- (4) $\beta = v\omega$; hence, $P(\bar{\beta}, \bar{v}, \bar{\omega})$.
- (5) Let $G = \langle \beta, v \rangle = \langle v, \omega \rangle = \langle \omega, \beta \rangle$ be the group of permutations of E generated by any two of the three permutations β, v, ω . Then G acts transitively on E .




FIGURE 1

The edges of this bicolored planar map—marked here 1 to 12—oriented as marked define the associated three permutations:

The black permutation β :

$$\beta = (1)(2)(3)(4, 5, 6)(7, 8, 9)(10, 11, 12)$$

The vertex permutation v :

$$v = (1, 5, 7)(2, 8, 10)(3, 11, 4)(6, 12, 9)$$

The white permutation ω :

$$\omega = (1, 7, 2, 10, 3, 4)(5, 9)(6, 11)(8, 12)$$

which satisfy $\beta = v\omega$.

The relation $P(3 \cdot (1^* + 3^*), 4 \cdot 3^*, 6^* + 3 \cdot 2^*)$ follows.

Figure 1 displays Proposition 2.1 in a particular case, where the bicolored map M has 12 edges, oriented as required to define the permutations β, v, ω .

Establishing P relations on countable sets by means of bpms proved useful in the past. Here we shall use such maps to demonstrate that such relations can

be established, while some extra side conditions are imposed. Figure 2 is an example. One concludes from it that permutations β, v, ω of a countable set exist such that:

- (1) $\beta = v\omega$.
- (2) $\bar{\beta} = \tilde{3}$, $\bar{v} = \tilde{1} + 3 \cdot 2^* + \tilde{3}$, $\bar{\omega} = \tilde{5}$.
- (3) Every ω -orbit contains precisely one v -fixed point and v has no other fixed points.




FIGURE 2

3. PROOF OF THEOREM A

In this section we reduce the proof of Theorem A to the establishment of Theorem C, below. Theorem C is proved—using bicolored planar maps—in §4.

We first establish Theorem A(3), which asserts that in a sense Theorem A(1) and (2) is a sharp result.

Let $\xi, \eta \in \text{INF}$ have no infinite orbit but infinitely many orbits of length at least k .

(a) If $k = 2$, $\text{INF-AI} \subseteq [\xi] \cdot [\eta]$ does not necessarily follow.

Indeed, let $\xi = \eta$ be a fixed-point-free involution. Then $[\xi] = R_0$, and so $[\xi] \cdot [\eta] = R_0^2$ is the set of nicely even permutations, i.e. permutations $\zeta \in S$ with $\bar{\zeta}(n)$ an even cardinal for all $n \in \mathbb{N}^+$ (see [M2]); and so, if $\zeta \in \text{INF}$ has a single infinite orbit, then $\zeta \notin R_0^2 = [\xi] \cdot [\eta]$.

(b) If $k = 3$, $\text{INF} \subseteq [\xi] \cdot [\eta]$ does not necessarily follow.

Indeed, if $\xi, \eta \in \text{INF}$ satisfy $\xi^3 = \eta^3 = 1$ and $[\xi] \neq [\eta]$, then $[\xi] \cdot [\eta]$ contains no involution (see [M5, Proposition 3.4]). (Even if $[\xi] = [\eta]$ then

$[\xi][\eta] = [\xi]^2$ may omit some nonfinitary involutions; see [D3, Example 4.6].) Theorem A(3) is proved.

Theorem A(1) and (2) follow from the following two theorems.

Theorem B. *Let $\xi, \eta \in \text{INF}$ have no infinite orbit, but infinitely many orbits of length at least k , and let $\zeta \in \text{INF}$ have no infinite orbit. Then:*

- (1) *If $k = 3$ and ζ has infinitely many orbits of length at least 3, then $\zeta \in [\xi][\eta]$.*
- (2) *If $k = 4$, then $\zeta \in [\xi][\eta]$.*

Theorem C. *Let $\xi, \eta \in \text{INF}$ have no infinite orbit, but infinitely many orbits of length at least 3. Let $\zeta \in \text{INF}$ have at least one infinite orbit. Then $\zeta \in [\xi] \cdot [\eta]$.*

Let us restate these theorems, using the terminology developed in §2.

Theorem B'. *Let r, s, t be types satisfying*

- (i) $|r| = |s| = |t| = \aleph_0$.
- (ii) $r(\aleph_0) = s(\aleph_0) = t(\aleph_0) = 0$.
- (iii) $r[[k, \aleph_0]] = s[[k, \aleph_0]] = t[[l, \aleph_0]] = \aleph_0$.

Then

- (1) *If $k = l = 3$, then $P(r, s, t)$.*
- (2) *If $k = 4$ and $l = 2$, then $P(r, s, t)$.*

Theorem C'. *Let r, s, t be types satisfying:*

- (i) $|r| = |s| = \aleph_0$.
- (ii) $r(\aleph_0) = s(\aleph_0) = 0$.
- (iii) $r[[3, \aleph_0]] = s[[3, \aleph_0]] = \aleph_0$.
- (iv) $|t| = \aleph_0$, $t(\aleph_0) \geq 1$.

Then $P(r, s, t)$.

Theorem C' is the main technical result here, and will be proved in §4. The next two lemmas are special cases of Theorem B', from which Theorem B' easily follows.

Lemma B1. *Let $m \in \mathbb{N}$, and let r, s, t be types satisfying:*

- (i) $|r| = |s| = |t| = \aleph_0$.
- (ii) $r(\aleph_0) = s(\aleph_0) = t(\aleph_0) = 0$.
- (iii) $r[[3, \aleph_0]] = s[[3, \aleph_0]] = t[[3, \aleph_0]] = \aleph_0$.
- (iv) $r(1) = r(2) = s(1) = s(2) = t(1) = t(2) = 0$.

Then $P(m^ + r, s, t)$, $P(r, m^* + s, t)$, $P(r, s, m^* + t)$.*

Lemma B2. *Let $m \in \mathbb{N}$, and let r, s be types satisfying:*

- (i) $|r| = |s| = \aleph_0$.
- (ii) $r(\aleph_0) = s(\aleph_0) = 0$.
- (iii) $r[[4, \aleph_0]] = s[[4, \aleph_0]] = \aleph_0$.
- (iv) $r(1) = r(2) = r(3) = s(1) = s(2) = s(3) = 0$.

Then $P(m^* + r, s, \tilde{2})$, $P(r, m^* + s, \tilde{2})$ and $P(r, s, m^* + \tilde{2})$.

Proof of Theorem B' from Lemmas B1 and B2.

Let r, s, t , be types satisfying (i)–(iii) of Theorem B'.

Case 1. Assume that $k = l = 3$. By (iii) we can select three disjoint sets $I, J, K \subseteq \mathbb{N}$, and positive integers l_i for $i \in I$, m_j for $j \in J$, n_k for $k \in K$; and types r_p, s_p, t_p for $p \in I \cup J \cup K$ such that

(a) r_p, s_p, t_p satisfy (i)–(iv) of Lemma B1 for $p \in I \cup J \cup K$.

(b)

$$\begin{aligned} r &= \sum_{i \in I} (l_i^* + r_i) + \sum_{j \in J} r_j + \sum_{k \in K} r_k, \\ s &= \sum_{i \in I} s_i + \sum_{j \in J} (m_j^* + s_j) + \sum_{k \in K} s_k, \\ t &= \sum_{i \in I} t_i + \sum_{j \in J} t_j + \sum_{k \in K} (n_k^* + t_k). \end{aligned}$$

By Lemma B1 and (a), $P(l_i^* + r_i, s_i, t_i)$, $P(r_j, m_j^* + s_j, t_j)$, $P(r_k, s_k, n_k^* + t_k)$ hold for $i \in I, j \in J, k \in K$. Hence by superadditivity of P , $P(r, s, t)$ follows from (b).

Case 2. Assume now that $k = 4$ and $l = 2$. If $t(2) < \aleph_0$ then by (iii) $t[[3, \aleph_0]] = \aleph_0$ and so, $P(r, s, t)$ follows from Case 1.

If $t(2) = \aleph_0$ then we select again disjoint subset $I, J, K \subseteq \mathbb{N}$, positive integers l_i, m_j, n_k for $i \in I, j \in J, k \in K$ and types r_p, s_p , for $p \in I \cup J \cup K$ so that:

(a) r_p, s_p satisfy (i)–(iv) of Lemma B2.

(b)

$$\begin{aligned} r &= \sum_{i \in I} (l_i^* + r_i) + \sum_{j \in J} r_j + \sum_{k \in K} r_k, \\ s &= \sum_{i \in I} s_i + \sum_{j \in J} (m_j^* + s_j) + \sum_{k \in K} s_k, \\ t &= |I| \cdot \tilde{2} + |J| \cdot \tilde{2} + \sum_{k \in K} (n_k^* + \tilde{2}). \end{aligned}$$

By Lemma B2, (a), (b) and the superadditivity of P , $P(r, s, t)$ follows. \square

Lemmas B1 and B2 are proved by constructing an appropriate bicolored planar map and using Proposition 2.1. In fact, they follow from the following proposition on planar bicolored maps, which only slightly generalizes Proposition 5.1 of [M4]. (By the symmetry of P and the conditions of these lemmas, only three P -relations need to be verified.)

Proposition B. Let $m \in \mathbb{N}$ and let $(l_i)_{i \in \mathbb{N}}, (m_j)_{j \in \mathbb{N}}$ and $(n_k)_{k \in \mathbb{N}}$ be sequences of integers, satisfying at least one of the following three conditions:

- (1) $m_1 = m$; $2 < l_i, m_j, n_k$ for $i, j, k \in \mathbb{N}$, $j > 1$.
- (2) $m_1 = m$; $3 < l_i, n_k$, $1 < m_j$ for $i, j, k \in \mathbb{N}$, $j > 1$.
- (3) $l_1 = m$; $3 < l_i, n_k$, $1 < m_j$ for $i, j, k \in \mathbb{N}$, $i > 1$.

Then there is a bicolored planar map M and enumeration $(b_i)_{i \in \mathbb{N}}$, $(v_j)_{j \in \mathbb{N}}$, $(w_k)_{k \in \mathbb{N}}$ of its black regions, vertices and white regions respectively, so that $|E_{b_i}| = l_i$, $|E_{v_j}| = m_j$, $|E_{w_k}| = n_k$ for $i, j, k \in \mathbb{N}$.

Hence $P(\sum_{i \in \mathbb{N}} l_i^*, \sum_{i \in \mathbb{N}} m_i^*, \sum_{i \in \mathbb{N}} n_i^*)$ holds, by Proposition 2.1.

(Recall that E_{b_i} (E_{w_k}) is the set of edges in the closure of the black region b_i (the white region w_k) and $2|E_{v_i}|$ is the degree of v_i .)

Proof. See Proposition 5.1 in [M4], which is Proposition B for $m = 1$, and cases (1), (3). To obtain a proof for Proposition B in general we need only to modify the base of inductive construction of the bicolored map given there. In cases (1) and (2), the base consists of the unit disc, partitioned radially into m white and m black alternating regions, sharing the origin as a common vertex (Figure 3(a)), while in case (3) it consists of the unit disc as a black region, with m vertices on its boundary (Figure 3(b)).




FIGURE 3. Base of induction for proof of Proposition B.

(a) Cases (1), (2), with: $m = 3, (l_1, l_2, l_3) = (5, 4, 3)$,

$(n_1, n_2, n_3) = (3, 4, 5)$.

(b) Case (3), with $m = 3$.

4. PROOF OF THEOREM C'

First notice that we may assume in Theorem C that ζ has precisely one infinite orbit, i.e., enough to prove the following special case of Theorem C'.

Theorem C''. Let r, s, t , be types satisfying:

- (i) $|r| = |s| = \aleph_0$.
- (ii) $r(\aleph_0) = s(\aleph_0) = 0$.
- (iii) $r[[3, \aleph_0]] = s[[3, \aleph_0]] = \aleph_0$.
- (iv) $|t| = \aleph_0, t(\aleph_0) = 1$.

Then $P(r, s, t)$.

Indeed, assume Theorem C'' and let r, s, t satisfy assumptions (i)–(iv) of Theorem C' , with $t(\aleph_0) = k \geq 1$. Let $K \subseteq \mathbb{N}$ be a set of cardinality k . Then one can associate with each $i \in K$ types r_i, s_i, t_i satisfying (i)–(iv) of Theorem C'' and $r = \sum_{i \in K} r_i, s = \sum_{i \in K} s_i, t = \sum_{i \in K} t_i$. By Theorem C'' , $P(r_i, s_i, t_i)$ holds for each $i \in K$, and so by superadditivity of P , $P(r, s, t)$ holds as well.

The rest of this section is devoted to the proof of Theorem C'' . We assume that r, s, t are types satisfying (i)–(iv). In particular, $t = \aleph_0^* + t''$, where $t''(\aleph_0) = 0$. Let A be a set of cardinality \aleph_0 . Our goal is to define $\xi, \eta, \zeta \in S_A$ such that $\bar{\xi} = r, \bar{\eta} = s, \bar{\zeta} = t$ and $\xi = \eta\zeta$. This is achieved in four steps as follows.

Step 1. Select types $r', r'', s', s'', t', t''$ so that:

- (1.1) $r = r' + r'', s = s' + s'', t = t' + t''$.
- (1.2) $r'[[3, \aleph_0]] = r''[[3, \aleph_0]] = s'[[3, \aleph_0]] = s''[[3, \aleph_0]] = \aleph_0$.
- (1.3) $t' = \aleph_0^*$.

Let $A_1, A_2, A_3, A_4 \subseteq A$ satisfy:

- (1.4) $A = \bigcup_{i=1}^4 A_i, A_i \cap A_j = \emptyset$ for $1 \leq i < j \leq 4$.
- (1.5) $|A_1| = |A_2| = |A_3| = \aleph_0, |A_4| = |t''|$.

We let $A_{i,j} = A_i \cup A_j$ and $A_{ijk} = A_i \cup A_j \cup A_k$, $1 \leq i, j, k \leq 4$. Also, let $1_i = 1_{A_i}, 1_{ij} = 1_{A_{ij}}, 1_{ijk} = 1_{A_{ijk}}$, $1 \leq i, j, k \leq 4$.

Step 2. Define $\xi' \in S_{A_1}, \eta' \in S_{A_{12}}, \zeta' \in S_{A_{123}}, \theta' \in A_{23}$ so that:

- (2.1) $\bar{\xi}' = r', \bar{\eta}' = s', \bar{\zeta}' = \aleph_0^*, \bar{\theta}' = \tilde{\gamma}$.
- (2.2) $\xi' + \theta' = (\eta' + 1_3)\zeta'$.

Step 3. Define $\xi'' \in S_{A_{234}}, \eta'' \in S_{A_{34}}, \zeta'' \in S_{A_4}$ so that:

- (3.1) $\bar{\xi}'' = r'', \bar{\eta}'' = s'', \bar{\zeta}'' = t''$.
- (3.2) $\xi'' = (1_2 + \eta'')(\theta' + \zeta'')$.

Step 4. Let $\xi = \xi' + \xi'', \eta = \eta' + \eta'', \zeta = \zeta' + \zeta''$. Then

- (4.1) $\bar{\xi} = r, \bar{\eta} = s, \bar{\zeta} = t$.
- (4.2) $\xi = \eta\zeta$.

Theorem C'' follows from (4.1) and (4.2).

Details. Step 1 is obviously possible, and needs no further comment. Step 4 follows from the previous steps as follows: (4.1) follows from (1.1), (2.1) and

(3.1). We establish (4.2):

$$\begin{aligned}
 \eta\zeta &= (\eta'' + \eta')(\zeta' + \zeta'') = [(1_{12} + \eta'')(\eta' + 1_{34})][(\zeta' + 1_4)(1_{123} + \zeta'')] \\
 &= (1_{12} + \eta'')[(\eta' + 1_3) + 1_4](\zeta' + 1_4)(1_{123} + \zeta'') \\
 &= (1_{12} + \eta'')[(\eta' + 1_3)\zeta' + 1_4 1_4](1_{123} + \zeta'') \\
 &= [(1_1 + 1_2 + \eta'')(\zeta' + \theta' + 1_4)](1_1 + 1_{23} + \zeta'') \\
 &= (1_1 \zeta' + (1_2 + \eta'')(\theta' + 1_4))(1_1 + 1_{23} + \zeta'') \\
 &= 1_1 \zeta' 1_1 + (1_2 + \eta'')[(\theta' + 1_4)(1_{23} + \zeta'')] \\
 &= \zeta' + (1_2 + \eta'')(\theta' + \zeta'') = \zeta' + \zeta'' = \xi.
 \end{aligned}$$

Here we used (2.2) for the fifth equality and (3.2) for the last-but-one equality.

Steps 2 and 3 form the core of the argument, and follow from two slightly more general Propositions 4.1 and 4.2 that we formulate next.

Proposition 4.1. *Let r^+, s^+, t^+ be types satisfying:*

- (i) $|r^+| = |s^+| = |t^+| = \aleph_0$.
- (ii) $r^+(\aleph_0) = s^+(\aleph_0) = t^+(\aleph_0) = 0$.
- (iii) $r^+[[3, \aleph_0]] = s^+[[2, \aleph_0]] = t^+[[2, \aleph_0]] = \aleph_0$.

Then there are $\xi' \in S_{A_1}, \eta' \in S_{A_{12}}, \zeta' \in S_{A_{123}}, \theta' \in S_{A_{23}}$ such that:

$$(4.1.1) \quad \overline{\xi'} = r^+, \overline{\eta'} = s^+, \overline{\zeta'} = \aleph_0^*, \overline{\theta'} = t^+.$$

$$(4.1.2) \quad \xi' + \theta' = (\eta' + 1_3)\zeta'.$$

(4.1.3) *If D is any θ' -orbit then $D \cap A_2$ contains a single element a_D (hence $A_2 = \{a_D : D \text{ is a } \theta'\text{-orbit}\}$).*

Proposition 4.2. *Let $r^{++}, s^{++}, t_0^{++}, t_1^{++}$ be types satisfying:*

- (i) $|r^{++}| = |s^{++}| = |t_0^{++}| = \aleph_0, |t_1^{++}| = |A_4| \leq \aleph_0$.
- (ii) $r^{++}(\aleph_0) = s^{++}(\aleph_0) = t_0^{++}(\aleph_0) = t_1^{++}(\aleph_0) = 0$.
- (iii) $r^{++}[[3, \aleph_0]] = s^{++}[[3, \aleph_0]] = t_0^{++}[[7, \aleph_0]] = \aleph_0, t_0^{++}[[1, 6]] = 0$.

Let further $\theta' \in S_{A_{23}}$ be given so that:

$$(iv) \quad \overline{\theta'} = t_0^{++}.$$

(v) *If D is any θ' -orbit then $D \cap A_2$ contains a single element a_D (hence $A_2 = \{a_D : D \text{ is a } \theta'\text{-orbit}\}$).*

Then there are $\xi'' \in S_{A_{234}}, \eta'' \in S_{A_{34}}, \zeta'' \in S_{A_4}$ such that

$$(4.2.1) \quad \overline{\xi''} = r^{++}, \overline{\eta''} = s^{++}, \overline{\zeta''} = t_1^{++}.$$

$$(4.2.2) \quad \xi'' = (1_2 + \eta'')(\theta' + \zeta'').$$

Let $M = (V, E, B, W)$ be a bicolored planar map. For $B' \subseteq B, V' \subseteq V, W' \subseteq W$, the types $t_{B'} = t_{B'}^M, t_{V'} = t_{V'}^M, t_{W'} = t_{W'}^M$ are defined as follows.

Let $n \in \mathbb{N}^+$. Then

$t_{B'}(n)$ = number of black regions $b \in B'$ of size n ,

$t_{V'}(n)$ = number of vertices $v \in V'$ of size n ,

$t_{W'}(n)$ = number of white regions $w \in W'$ of size n .

(Recall that the *size of a region* f is $|E_f|$, where E_f is the set of edges in its closure, and the *size of a vertex* v is $|E_v|$, which equals half its degree.) $t_B(t_V, t_W)$ is called the black (the vertex, the white) type of M .

Propositions 4.1 and 4.2 follow from their respective counterparts, Propositions 4.1' and 4.2', that provide the necessary bicolored maps, using Proposition 2.1. We start with:

Proposition 4.1'. *Let r^+, s^+, t^+ be types satisfying (i)–(iii) of Proposition 4.1. Then there is a bicolored planar map $M = (V, E, B, W)$, and partitions $\{E_1, E_2, E_3\}$ of E , $\{B', B''\}$ of B , $\{V', V''\}$ of V such that:*

(4.1'.1) $E_1 = \bigcup_{b \in B'} E_b$, $E_2 \cup E_3 = \bigcup_{b \in B''} E_b$, $E_1 \cup E_2 = \bigcup_{v \in V'} E_v$, $E_3 = \bigcup_{v \in V''} E_v$.

(4.1'.2) $t_{B'} = r^+$, $t_{B''} = t^+$, $t_{V'} = s^+$, $t_{V''} = \tilde{1}$, $t_W = \aleph_0^*$.

(4.1'.3) If $b \in B''$ then $E_b \cap E_2$ contains a single element.

Proof of Proposition 4.1 from 4.1'. Let $\delta: E \rightarrow A_{123}$ be any bijection that maps E_i onto A_i , $i = 1, 2, 3$. Let β, v and ω be the black, vertex and white permutations of M . Let $\beta' = \beta|E_1$, $\beta'' = \beta|E_2 \cup E_3$, $v' = v|E_1 \cup E_2$, $v'' = v|E_3$, and define $\xi', \eta', \zeta', \theta'$ by $\xi' = \delta\beta'\delta^{-1}$, $\eta' = \delta v'\delta^{-1}$, $\zeta' = \delta\omega\delta^{-1}$, $\theta' = \delta\beta''\delta^{-1}$. Then (4.1.1)–(4.1.3) hold. (See Proposition 2.1 for (4.1.2).)

Proof of Proposition 4.1'—The Caterpillar Map. A bicolored planar map $M = (V, E, B, W)$ is called a *one way infinite caterpillar map*—or, briefly, a *caterpillar map*—if it satisfies the following conditions:

All black regions have finite size (hence are bounded) and the set B of black regions is a disjoint union $B = B^+ \cup B^{++}$ where:

(C.1) $B^+ = \{b_n : n \in \mathbb{N}\}$ where $\text{cl}(b_n) \cap \text{cl}(b_m) = \emptyset$ if $|n - m| > 1$, while $\text{cl}(b_n) \cap \text{cl}(b_{n+1}) = \{v_n\}$, $v_n \in V$ ($n \in \mathbb{N}$). The regions $b_n \in B^+$ are called *links* and the vertices v_n are called *joints*. Other vertices of b_n are called *nonjoints*.

(C.2) Every $b'' \in B^{++}$ has a unique vertex $v_{b''}$ such that $\text{cl}(b'') \cap \text{cl}(b_n) = \{v_{b''}\}$ for some $n \in \mathbb{N}$ ($v_{b''}$ may be a joint or a nonjoint). The regions $b'' \in B^{++}$ are called *toes*.

It follows that $W = \{w\}$, i.e. there is only one (unbounded) white region.

Assume now that r^+, s^+, t^+ are types satisfying (i)–(iii) of Proposition 4.1. We shall show that a caterpillar map $M = (V, E, B, W)$ exists with partitions $\{E_1, E_2, E_3\}$ of E , $\{B', B''\}$ of B and $\{V', V''\}$ of V such that (4.1'.1)–(4.1'.3) hold.

(a) Let $\{b_n : n \in \mathbb{N}\}$ be a sequence of disjoint translation of the unit disc in the plane such that (C.1) holds. Let C_n denote the boundary of b_n and let v_n be the point of intersection of C_n with C_{n+1} .




FIGURE 4. A caterpillar map

The black faces b_1, \dots, b_5, \dots are links.
 The black faces b'_1, \dots, b'_{10} are toes.
 The vertices v_1, \dots, v_4, \dots are joints.
 The infinite region w is the only white face.

(b) Let $(r_n)_{n \in \mathbb{N}}$ be a sequence of integers satisfying $r_n > 1$ and $r^+ = r^+(1) \cdot 1^* + \sum_{n \in \mathbb{N}} r_n^*$. Make b_n into a black region of size r_n by adding, if necessary, extra vertices on its boundary C_n . Notice that, for $n > 1$, b_n already has v_{n-1} and v_n as vertices in its closure. We add more vertices if $r_n > 2$, and this is true of infinitely many n 's by $r^+[[3, \aleph_0]] = \aleph_0$. Let $B'_0 = \{b_n : n \in \mathbb{N}\}$ be the set of links of M .

(c) Let $(s_k)_{k \in \mathbb{N}}$ be a sequence of integers satisfying $s_k > 0$ and $s^+ = \sum_{k \in \mathbb{N}} s_k^*$. Let $(u_k)_{k \in \mathbb{N}}$ be an enumeration of all vertices defined so far, such that:

1. If $s_k = 1$ then u_k is not a joint.
2. $s_k > 1$ for infinitely many nonjoint vertices u_k .

(d) Add loops lying in the unbounded region, each enclosing a black region and pending at a vertex u_k as necessary to guarantee that the degree of u_k is $2s_k$ (and so its size is s_k). Select a coinfinite subset B'_1 of black regions enclosed by these loops of cardinality $|B'_1| = r^+(1)$, and set $B' = B'_0 \cup B'_1$ (so $t_{B'} = r^+$). Let $(b''_l)_{l \in \mathbb{N}}$ be an enumeration of the black regions enclosed by the other loops.

(e) Let $(t_l)_{l \in \mathbb{N}}$ be a sequence of integers satisfying $t^+ = \sum_{l \in \mathbb{N}} t_l^*$. Add vertices on the loop enclosing b''_l so as to make the size of b''_l equal t_l . Notice that by $t^+[[2, \infty]] = \aleph_0$, $t_l > 1$ for infinitely many l 's. Let $B'' = \{b''_l : l \in \mathbb{N}\}$ (so $t_{B''} = t^+$).

The caterpillar map $M = (V, E, B, W)$ is now complete. (Its "body" consists of the links b_n of B'_0 ; its "toes" form $B'_1 \cup B''$.) We let $V' = \{u_k : k \in \mathbb{N}\}$ be the set of vertices that lie in the closure of some link $b' \in B'$ and $V'' =$

$V - V'$. Then E_1, E_2, E_3 are defined by (4.1'.1). A straightforward checking verifies (4.1'.1)–(4.1'.3).

Proposition 4.2'. *Let $r^{++}, s^{++}, t_0^{++}, t_1^{++}$ be types satisfying (i)–(iii) of Proposition 4.2. Then there is a bicolored planar map $M = (V, E, B, W)$ and partitions $\{E_2, E_3, E_4\}$ of E , $\{V', V''\}$ of V , $\{W', W''\}$ of W such that:*

(4.2'.1) $E_2 \cup E_3 = \bigcup_{w \in W'} E_w$, $E_4 = \bigcup_{w \in W''} E_w$, $E_3 \cup E_4 = \bigcup_{v \in V'} E_v$, $E_2 = \bigcup_{v \in V''} E_v$.

(4.2'.2) $t_B = r^{++}$, $t_{V'} = s^{++}$, $t_{V''} = \tilde{1}$, $t_{W'} = t_0^{++}$, $t_{W''} = t_1^{++}$.

(4.2'.3) If $w \in W'$ then $\text{cl}(w)$ contains a unique distinguished vertex $v_w \in V''$ of degree 2.

(4.2'.4) Let $w \in W'$ and let $E_{v_w} = \{e_w\}$.¹ Then $E_{v_w} = E_w \cap E_2$ (hence $E_2 = \{e_w : w \in W'\}$).

Proof of Proposition 4.2 from 4.2'. Let M be the map provided by Proposition 4.2', and let $\beta, v, \omega \in S_E$ be its black, vertex and white permutations, respectively (see §2). Let $\theta' \in S_{A_{23}}$ satisfy (iv) and (v) of Proposition 4.2. Since $\bar{\theta}' = t_0^{++} = t_{W'}$, there is a bijection ε from W' onto the set $(A_{23})_{\theta'}$ of all θ' -orbits, satisfying $s(w) = |\varepsilon(w)|$ (i.e., ε preserves sizes).

Define a bijection $\delta : E \rightarrow A_{234}$ as follows: For each $w \in W'$, set

$$(*) \quad \delta(e_w) = a_{\varepsilon(w)},$$

where $e_w \in E$ and $a_{\varepsilon(w)} \in A$ are guaranteed by (4.2'.4) and 4.2(v), respectively. Thus, (*) defines $\delta|_{E_2}$ as a bijection of E_2 onto A_2 .

Extend δ to $E_2 \cup E_3 = \bigcup_{w \in W'} E_w$ by the requirement:

$$(**) \quad \theta' \delta(e) = \delta \omega(e), \quad e \in E_2 \cup E_3.$$

By (**), $\delta' = \delta|_{E_2 \cup E_3}$ is a bijection of $E_2 \cup E_3$ with $A_{23} = \bigcup_{D \in (A_{23})_{\theta'}} D$, and letting $\omega' = \omega|_{E_2 \cup E_3}$ we have $\theta' = \delta' \omega' (\delta')^{-1}$.

By (4.2'.1) and (4.2'.2) we have $|E_4| = |t_{W''}| = |t_1^{++}| = |A_4|$. Let $\delta|_{E_4}$ be an arbitrary bijection of E_4 with A_4 . Define $\xi'' \in A_{234}$, $\eta'' \in A_{34}$, $\zeta'' \in A_4$ by

$$\begin{aligned} \zeta'' &= \delta \omega'' \delta^{-1}, \quad \text{where } \omega'' = \omega|_{E_4}, \\ \eta'' &= \delta v'' \delta^{-1}, \quad \text{where } v'' = v|_{E_3 \cup E_4}, \\ \xi'' &= (1_2 + \eta'')(\theta' + \zeta''). \end{aligned}$$

Since $E_2 = \{e_w : w \in W'\}$ we have $v|_{E_2} = 1_{E_2}$ by (4.2'.3), (4.2'.4) and Proposition 2.1. Thus $\delta v \delta^{-1} = \delta(1_{E_2} + v'') \delta^{-1} = 1_2 + \eta''$. Similarly, $\delta \omega \delta^{-1} = \delta(\omega' + \omega'') \delta^{-1} = \theta' + \zeta''$. Hence we have

$$\xi'' = \delta v \omega \delta^{-1} = \delta \beta \delta^{-1}.$$

¹ By §2, whenever v has degree 2, E_v is a singleton $\{e\}$, where e is the unique edge along which one can travel from v with a white region w on the right-hand side.

Thus, $\overline{\xi''} = \overline{\beta} = t_B = r^{++}$, $\overline{\eta''} = \overline{v''} = t_{V'} = s^{++}$ and $\overline{\zeta''} = \overline{\omega''} = t_{W''} t_1^{++}$. (4.2.1) and (4.2.2) are established and Proposition 4.2 follows.

The rest of this paper is dedicated to the

Proof of Proposition 4.2'. Let $r^{++}, s^{++}, t_0^{++}, t_1^{++}$ satisfy (i)–(iii) of Proposition 4.2, and let

$$\begin{aligned} r_0^{++} &= \sum_{3 \leq n} r(n) \cdot n^*, & r_1^{++} &= r(1) \cdot 1^* + r(2) \cdot 2^*, \\ s_0^{++} &= \sum_{3 \leq n} s(n) \cdot n^*, & s_1^{++} &= s(1) \cdot 1^* + s(2) \cdot 2^*, \end{aligned}$$

so that

$$r^{++} = r_0^{++} + r_1^{++}, \quad s^{++} = s_0^{++} + s_1^{++}.$$

By (ii) and (iii) of Proposition 4.2 there are three sequences of integers:

$$b = (l_i)_{i \in \mathbb{N}}, \quad v = (m_j)_{j \in \mathbb{N}}, \quad w = (n_k)_{k \in \mathbb{N}}$$

satisfying $3 \leq l_i, m_j$ ($i, j \in \mathbb{N}$) and $7 \leq n_k$ ($k \in \mathbb{N}$) such that

$$r_0^{++} = \sum_{i \in \mathbb{N}} l_i^*, \quad s_0^{++} = \sum_{j \in \mathbb{N}} m_j^*, \quad t_0^{++} = \sum_{k \in \mathbb{N}} n_k^*.$$

Also, there are disjoint (possibly empty) subsets $T_B, T_V, T_W \subseteq \mathbb{N}$ and a T_B -sequence $b' = (l'_i)_{i \in T_B}$, a T_V -sequence $v' = (m'_j)_{j \in T_V}$ and a T_W -sequence $w' = (n'_k)_{k \in T_W}$ of positive integers such that

$$r_1^{++} = \sum_{i \in T_B} l_i'^*, \quad s_1^{++} = \sum_{j \in T_V} m_j'^*, \quad t_1^{++} = \sum_{k \in T_W} n_k'^*$$

(so $l'_i, m'_j \in \{1, 2\}$ for $i \in T_B, j \in T_V$).

We shall construct a bicolored planar map $M = (V, E, B, W)$ with partitions $\{V'_0, V'_1, V''\}$ of V , $\{B_0, B_1\}$ of B , $\{W', W''\}$ of W and a bijection $f: W' \rightarrow V''$ such that:

(4.2'' .1) $t_{B_0} = r_0^{++}$, $t_{B_1} = r_1^{++}$, $t_{V'_0} = s_0^{++}$, $t_{V'_1} = s_1^{++}$, $t_{W'} = t_0^{++}$, $t_{W''} = t_1^{++}$.

(4.2'' .2) If $w \in W'$ then $f(w) \in V''$ is a vertex of degree 2 that belongs to $\text{cl}(w)$.

Proposition 4.2' then follows. (Define $E_2 = \bigcup_{v \in V''} E_v$, $E_3 = (\bigcup_{w \in W'} E_w) - E_2$, $E_4 = E - (E_2 \cup E_3)$, $V' = V'_0 \cup V'_1$.)

Outline of the construction of M . M will be obtained as the increasing union of maps M_n , where M_n is drawn on the closed disk D_n of radius n centered at the origin. The inductive extension of M_n to M_{n+1} involves the addition of extra black regions, extra vertices and extra white regions. The sizes of black and white regions are dictated mostly by the sequences b, w , and the fresh vertices lie mostly on the circle C_{n+1} of radius $n + 1$ centered at the origin.

In the process of extending M_n to M_{n+1} , vertices on C_n obtain their degrees as dictated by the list v . As a rule, a white region w introduced during the extension is provided with the vertex $f(w) = v_w$ which lies strictly between the circles C_n and C_{n+1} , and supports a black triangular region with apex v_w with a circular arc—a segment of C_{n+1} —as an opposite side (see Figure 2).

For the above inductive construction to proceed smoothly, the occurrences of regions of small size and vertices of small degree should be limited. Such occurrences are forced upon us by the need to construct the black regions in B_1 , the vertices in V'_1 (which are of size 1 or 2) and possibly some of the white regions in W'' (which are of arbitrary finite size). We overcome this difficulty by constructing these map-components one at a time—that is, at most one at an inductive extension of M_n to M_{n+1} .

The construction of a map-element from $B_1 \cup V'_1 \cup W''$ is referred to in the sequel as *accomplishing a task*. The tasks are in one-to-one correspondence with the set $T = T_B \cup T_V \cup T_W \subseteq \mathbb{N}$. We take care of the first task in proceeding from M_1 to M_2 , and if a task is accomplished in the transition from M_n to M_{n+1} , the next one is accomplished in the transition from M_{n+1} to M_{n+2} or from M_{n+2} to M_{n+3} . Thus, the n th task is already accomplished in M_{2n} , hence all tasks are eventually accomplished in M .

Details. We identify for the sake of brevity the plane with the set of complex numbers \mathbb{C} . For $r > 0$, let $U_r = \{z \in \mathbb{C} : |z| < r\}$, $C_r = \{z \in \mathbb{C} : |z| = r\}$ and $D_r = U_r \cup C_r$. For a subset $X \subseteq \mathbb{C}$ let $[X]$ denote its convex closure. For $x_1, \dots, x_n \in \mathbb{C}$ let $[x_1, \dots, x_n] = [\{x_1, \dots, x_n\}]$. Thus $[x, y]$ is the line segment connecting x and y .

Let $x, y \in C_r$, say $x = re^{i\varphi}$, $y = re^{i\psi}$, $0 \leq \psi - \varphi < 2\pi$. Then $[x, y] = \{re^{ix} : \varphi \leq x \leq \psi\}$ denote the arc on C_r leading from x to y in the positive sense, i.e., counterclockwise. $\text{cl}(X)$ and $\text{Int}(X)$ denote as usual the closure and the interior of $X \subseteq \mathbb{C}$.

Let $M = (V, E, B, W)$ be any bicolored planar map, and let r be a positive real number. Define $V^r = V \cap D_r$, $E^r = \{e \cap D_r : e \in E\}$, $B^r = \{b \cap D_r : b \in B\}$, $W^r = \{w \cap D_r : w \in W\}$ and set $M^r = (V^r, E^r, B^r, W^r)$. We have $C_r \subseteq \bigcup E$ if and only if $C_r \subseteq \bigcup E^r$. If $C_r \subseteq \bigcup E^r$ we say that M^r is an r -disk-map or, briefly, that M^r is an r -map.

Assume that M^r is an r -map. Then (V^r, E^r) is obviously a planar graph, $B^r \subseteq B$, $W^r \subseteq W$. We call $v \in V^r$ an *inner vertex* iff $|v| < r$, and a *rim vertex* if $|v| = r$. Inner vertices have the same (even) degree in M and in M^r , and the degree of a rim vertex is at least 2.

The required map $M = (V, E, B, W)$ will be defined as the increasing union of disk maps, namely, its restrictions $M^n = (V^n, E^n, B^n, W^n)$ to the disks D_n . To help verify (4.2''.1) and (4.2''.2) we shall further specify partitions $\{^nV'_0, ^nV'_1, ^nV''\}$ of $V^n - C_n$, $\{^nB_0, ^nB_1\}$ of B^n and $\{^nW', ^nW''\}$ of W^n , which eventually define the partitions $\{V'_0, V'_1, V''\}$, $\{B_0, B_1\}$ and $\{W', W''\}$

of V, B, W respectively. In fact, ${}^nV'_1, {}^nB_1, {}^nW''$ will be obtained as established *tasks*, as follows.

For $p \in T = T_B \cup T_V \cup T_W$, we shall define in the course of the construction the p th task $T(p)$ to be a particular map element—a black region of size l'_p if $p \in T_B$, a vertex of size m'_p (i.e., degree $2m'_p$) if $p \in T_V$, and a white region of size n'_p if $p \in T_W$. The task $T(p)$ will always have specified *task vertex* $v(p)$ associated with it. If $p \in T_B \cup T_W$ then $v(p)$ belongs to $\text{cl } T(p)$, and if $p \in T_V$ then actually $T(p) = v(p)$.

It follows from the fact that M^n is a disk-map, that for every $n \in \mathbb{N}$, C_n is a union of edges in E , in fact in E^n . In addition, the set of positive integers $\mathbb{N} = \{1, 2, \dots\}$ will be a subset of V , and its convex closure, the ray $[1, \infty)$, will also occur as a union of edges in E . In general task vertices $v(p)$ will occur on this ray as half-integers, $v(p) = n_p + \frac{1}{2}$ for some $n_p \in \mathbb{N}$. (The only exceptions are some of the tasks $T(p)$ with $p \in T_V$, $m'_p = 1$; i.e., some vertices in V'_1 of degree 2.)

We say that a black region b (a vertex v ; a white region w) *accomplishes* the integer k if $s(b) = k$ ($s(v) = k; s(w) = k$).

Base of the inductive construction. Let $M^1 = (V^1, E^1, B^1, W^1)$ where

$$\begin{aligned} V^1 &= \{v_1 = 1, v_2, \dots, v_{l_1}\}, \quad \text{with } v_k = e^{(2\pi i/l_1) \cdot (k-1)}, \quad k = 1, \dots, l_1, \\ E^1 &= \{e_1, \dots, e_{l_1}\}, \quad \text{with } e_k = [v_k, v_{k+1}], \quad k = 1, \dots, l_1, \quad (v_{l_1+1} = v_1), \\ B^1 &= \{U_1\}, \quad W^1 = \emptyset. \end{aligned}$$




FIGURE 5. M_1 with $l_1 = 3$

We further let:

$${}^1V'_0 = V^1, \quad {}^1V'_1 = {}^1V'' = \emptyset, \quad {}^1B_0 = B^1, \quad {}^1B_1 = \emptyset, \quad W' = W'' = \emptyset.$$

Thus, U_1 accomplishes l_1 and let $b^{(1)}$ denote the sequence $(l_i)_{i \in \mathbb{N} - \{1\}}$. Let also $v^{(1)} = v, w^{(1)} = w$. Let $t^{(1)} = t = (t_p)_{p \in T}$ denote the task-sequence $b' \cup v' \cup w'$ (i.e., $t_p = l'_p$ if $p \in T_B$, $t_p = m'_p$ if $p \in T_V$, $t_p = n'_p$ if $p \in T_W$).

Induction hypothesis. Assume that $M^n = (V^n, E^n, B^n, W^n)$ is already defined, and so are the sequences of positive integers $b^{(n)}, v^{(n)}, w^{(n)}, t^{(n)}$. These are obtained from the sequences b, v, w, t by omitting a finite number of terms from each, namely, the terms accomplished in M_n ; that is, the terms omitted from the sequence b (v, w, t) stand in one-to-one correspondence with the regions in nB_0 (the set ${}^nV'_0$ of inner vertices of V_n ; the set ${}^nW'$ of white region of M_n ; the task-set ${}^nB_1 \cup {}^nV'_1 \cup {}^nW''$) that accomplish them. Also, each $w \in W'$ has an inner vertex $v_w \in {}^nV''$ of degree two in its closure, and ${}^nV'' = \{v_w : w \in W'\}$.

Let $v_1 = n, v_2, \dots, v_q \in V^n$ enumerate the rim vertices of M_n in the positive sense, and let $2 \leq d_n(v_i)$ denote the degree of v_i in M_n . We assume that one of the following two cases hold:

Case a. $d_n(v_i) \leq 4$ for $i = 1, \dots, q$.

Case b. $d_n(v_1) = 5$ and $d_n(v_i) \leq 4$ for $i = 2, \dots, q$.

Induction Step. The extension of M^n to M^{n+1} is made in several steps. In general, the 1-skeleton of the extension (i.e., the union of its edges) is defined, being divided by inserted vertices into edges later on, to obtain the required sizes for fresh map elements. The actual extension depends on whether M_n obeys Case a or Case b and whether any tasks are left. Thus, we distinguish two cases:

Case 1. Case b occurs, or no tasks are left.

Case 2. Case a occurs, and some tasks are left.

In Case 1 a "smooth" extension leads to M_{n+1} , where Case a occurs. In Case 2, the first nonaccomplished task is accomplished in the course of the extension.

We turn to the details. Recall that $v_1 = n, v_2, \dots, v_q \in C_n$ are the rim vertices of M^n , and $d_n(v_i)$ is the degree of v_i in M^n , $i = 1, \dots, q$. We let $v_{q+1} = v_1$ and $e_i = [v_i, v_{i+1}]$, $i = 1, \dots, q$, $e_{q+1} = e_1$. For $i = 1, \dots, q$ we define $X_i \in \{\underline{B}, \underline{W}\}$ as follows: $X_i = \underline{B}$ ($X_i = \underline{W}$) iff e_i lies in the closure of a white (black) region of M_n . Thus, $X_i = \underline{B}$ ($X_i = \underline{W}$) iff the region of M touching e_i and lying outside D_n should be black (white). We now proceed by the two cases as follows:

Case 1. $d_n(v_1) = 5$, $d_n(v_i) \leq 4$ for $i = 1, 2, \dots, q$ or $t^{(n)}$ is an empty sequence.

Step 1.1 (assigning sizes to new map elements supporting edges on C_n). Declare $s(v_1), \dots, s(v_q)$ to be the first q members of the sequence $v^{(n)}$ (so $s(v_i) \geq 3$, $i = 1, \dots, q$). Let f_i denote the M^{n+1} region lying outside D_n and having e_i in its closure (to be determined in full later), $i = 1, \dots, q$. Let $g_1, \dots, g_{q'}$ ($h_1, \dots, h_{q''}$) be the subsequence of f_1, \dots, f_q consisting of all f_i with $X_i = \underline{B}$ ($X_i = \underline{W}$). Let $s(g_1), \dots, s(g_{q'})$ be the first q' members of the sequence $b^{(n)}$, and let $s(h_1), \dots, s(h_{q''})$ be the first q'' members of the sequence $w^{(n)}$ (so $s(g_i) \geq 3$, $i = 1, \dots, q'$, and $s(h_i) \geq 6$,² $i = 1, \dots, q''$).

Step 1.2 (determination of all edges connecting C_n with C_{n+1}). Declare C_{n+1} to be part of the 1-skeleton of M_{n+1} . Declare $u_i = (1 + \frac{1}{n})v_i \in C_{n+1}$ to be a vertex in V^{n+1} , and $[v_i, u_i]$ to be an edge in E^{n+1} . Let $1 \leq i \leq q$, and consider the number $d'_i = 2s(v_i) - d_n(v_i)$ (which may be called “the degree of v_i outside D_n ”). Since $s(v_i) \geq 3$, $d_n(v_i) \leq 4$ for $i = 2, \dots, q$, we have $d'_i \geq 2$ for $i = 2, \dots, q$. In general, $d'(v_1) \geq 2$ as well. If, however, Case b holds, then $d_n(v_1) = 5$, so if in addition $s(v_1) = 3$, we have $d'(v_1) = 1$. We proceed according to two possible options.

Case 1.2.1. $d'_1 \geq 2$, or $s(f_1) > 3$.

(1.2.1.1) For $i = 1, \dots, q$ choose a vertex $u_i^+ \in [u_i, u_{i+1}]$ (where $u_{q+1} = u_1$) and an edge $e'_i \in E^{n+1}$ connecting v_i to u_i^+ as follows: If $s(f_i) > 3$, let $u_i^+ \neq u_i, u_{i+1}$ be an internal point of the arc $[u_i, u_{i+1}]$ such that $[v_i, u_i^+] \cap U_n = \emptyset$.

Let $e'_i = [v_i, u_i^+]$. If $s(f_i) = 3$ (in which case $X_i = \underline{B}$, i.e., f_i is to be a black region) let $u_i^+ = u_{i+1}$, and let e'_i be a straight line segment or any circular arc connecting v_i to u_i^+ meeting D_n at v_i and lying in $D_{n+1} \cap [\{v_i, v_{i+1}\} \cup [u_i, u_{i+1}]]$, that meets $[v_i, u_i]$ at nonzero angle.

(1.2.1.2) Declare $d'_i - 2$ distinct points other than u_i, u_i^+ on the arc $[u_i, u_i^+]$ to be vertices in V^{n+1} , and the line segments connecting them to v_i to be edges in E^{n+1} (these new points $u \in [u_i, u_i^+]$ are chosen so that the line segment $[v_i, u]$ does not intersect e'_i).

(1.2.1.3) Let \tilde{f}_i denote the open region bounded by the curves $e'_i, e_i = [v_i, v_{i+1}]$, $[v_{i+1}, u_{i+1}]$ and $[u_i^+, u_{i+1}]$ (if $u_i^+ \neq u_{i+1}$).

Case 1.2.2. $d'_1 = 1$ and $s(f_1) = 3$.

(1.2.2.1) For $i = 2, \dots, q+1$ choose a vertex $u_i^- \in [u_{i-1}, u_i]$ (where $u_{q+1} = u_1$) and an edge e'_i connecting v_i to u_i^- as follows: If $s(f_{i-1}) > 3$, let $u_i^- \neq u_{i-1}, u_i$ be an internal point of $[u_{i-1}, u_i]$ such that $[v_i, u_i^-] \cap U_n = \emptyset$. Let $e'_i = [v_i, u_i^-]$. If $s(f_{i-1}) = 3$ (in which case $X_i = \underline{B}$) let $u_i^- = u_{i-1}$ and let e'_i be a straight line segment or any circular arc connecting v_i to u_i^- meeting D_n at v_i and lying in $D_{n+1} \cap [\{v_i, v_{i-1}\} \cup [u_{i-1}, u_i]]$, that meet $[v_i, u_i]$ at nonzero angle.

² Actually, $s(h_i) \geq 7$; but in Case 1 $s(h_i) \geq 6$ is sufficient.




FIGURE 6. Step 1, Case 1.2.1, with $q = 5$, $X_1 = X_2 = X_5 = \underline{W}$, $X_3 = X_4 = \underline{B}$, $d'_1 = d'_2 = 2$, $d'_3 = d'_5 = 3$, $d'_4 = 4$.

(1.2.2.2) Declare $d'_i - 2$ distinct points of the arc $[u_i^-, u_i]$ other than u_i^- , u_i to be vertices in V^n and the line segments connecting them to be edges in E^{n+1} . (The new points $u \in [u_i^-, u_i]$ are chosen so that $[v_i, u] \cap e'_i = \emptyset$.)

(1.2.2.3) Let \tilde{f}_i denote the open region bounded by the curves e'_{i+1} , $e_i = [v_i, v_{i+1}]$, $[v_i, u_{i+1}^-]$ and $[u_i, u_{i+1}^-]$ (if $u_i \neq u_{i+1}^-$).

Case 1.2.3. We now establish notation and colour for the regions obtained so far in $D_{n+1} - D_n$. For $i = 1, \dots, q$ and $j = 1, \dots, d'_i$ we define the region \tilde{f}_{ij} and colour $X_{ij} \in \{\underline{B}, \underline{W}\}$ as follows. For $X \in \{\underline{B}, \underline{W}\}$ define X' by $\underline{B}' = \underline{W}$, $\underline{W}' = \underline{B}$.

In Case 1.2.1, let $\tilde{f}_{id'_i} = \tilde{f}_i$, $X_{id'_i} = X_i$. Let \tilde{f}_{ij-1} be the neighboring region to \tilde{f}_{ij} to the right. Let $X_{ij-1} = X'_{ij}$ (see Figure 6).

In Case 1.2.2, let $\tilde{f}_{i1} = \tilde{f}_i$, $X_{i1} = X_i$. Let \tilde{f}_{ij+1} be the region to the left of \tilde{f}_{ij} , $X_{ij+1} = X'_{ij}$ (see Figure 7).

Remarks. 1.2.4. The parity of the degrees $d(v_i) = 2s(v_i)$ and the fact that M_n is properly colored provide us with the fact that

$$X_{11}, \dots, X_{1d'_1}, \dots, X_{21}, \dots, X_{2d'_2}, \dots, X_{q1}, \dots, X_{qd'_q}$$

is an alternating sequence of \underline{B} and \underline{W} .

1.2.5. All vertices on C_{n+1} defined so far have degree ≤ 4 . The only vertices of degree 4 are some of the vertices u_1, \dots, u_q . u_i has degree 4 only in the following two cases: Case 1.2.1 and $s(f_{i-1}) = 3$, or Case 1.2.2 and $s(f_i) = 3$. Note that in both cases $X_i = \underline{B}$.

Let $f_{ij} \in W^{n+1}$. Thus we have

$$\begin{aligned} B^{n+1} - B^n &= \{b_k : k = 1, \dots, r\} \cup \{f_{ij} : X_{ij} = \underline{B}\}, \\ W^{n+1} - W^n &= \{f_{ij} : X_{ij} = \underline{W}\}. \end{aligned}$$

We declare $B^{n+1} - B^n \subseteq B_0^{n+1}$, $W^{n+1} - W^n \subseteq W_0^{n+1}$.

Step 1.4 (completion of M^{n+1} -additional vertices on C_{n+1}).

1.4.1. Assign a size $s(b)$ ($s(w)$) to each region of $B^{n+1} - B^n$ ($W^{n+1} - W^n$) other than f_1, \dots, f_q , whose sizes were already determined in Step 1.1. For this purpose use the first members of the sequence $b^{(n)}$ ($w^{(n)}$) not assigned to map elements in 1.1. Thus, $s(b) \geq 3$ for all $b \in B^{n+1} - B^n$ and $s(w) \geq 7$ for all $w \in W^{n+1} - W^n$.

1.4.2. Add vertices on C_{n+1} so that we have $s(b) = |E_b|$ and $s(w) = |E_w|$ for each $b \in B^{n+1} - B^n$ and $w \in W^{n+1} - W^n$. Declare the arcs into which C_{n+1} is divided edges in E_{n+1} . Let

$$\begin{aligned} {}^{n+1}V'' - V^n &= \{z_k : k = 1, \dots, r\} \subseteq U^{n+1} - D_n, \\ {}^{n+1}V'_0 - V^n &= V^{n+1} \cap C_{n+1}, \quad {}^{n+1}V'_1 = \emptyset. \end{aligned}$$

The induction step is complete. Notice that every vertex on C_{n+1} has degree 2, 3 or 4, i.e., Case a occurs.

Case 2. $d_n(v_i) \leq 4$, $i = 1, \dots, q$, and $t^{(n)}$ is not an empty sequence. Let t_p be the first member of $t^{(n)}$. The p th task $T(p)$ will be accomplished in $D_{n+1} - D_n$, in the course of this inductive step. We shall denote the possible tasks to be considered as $T(p)$ as follows:

Task B.k.: $p \in T_B$ and $t_p = k$ ($k \in \{1, 2\}$),

Task V.k.: $p \in T_V$ and $t_p = k$ ($k \in \{1, 2\}$),

Task W.k.: $p \in T_W$ and $t_p = k$ ($k \in \mathbb{N}$).

We proceed in steps:

Step 2.1 (assigning sizes to map elements supporting edges in C_{n+1}). Same as Step 1.1. Notice that $d'_i \geq 2$, $i = 1, \dots, q$.

Step 2.2 (division of the ring $D_{n+1} - D_n$ into compartments). Declare C_{n+1} to be part of the 1-skeleton of M_{n+1} . Declare $u_i = (1 + 1/n_i)v_i \in C_{n+1}$ to be a vertex in V^{n+1} , and $[v_i, u_i]$ to be edges in E^{n+1} , $i = 2, \dots, q$. Declare $[v_1, u_1]$ to be part of the 1-skeleton of M_{n+1} .

Let us call the region enclosed by the curves $[v_q, v_2], [v_2, u_2], [u_q, u_2], [v_q, u_q]$ the *task-region*.

Step 2.3 (accomplishment of the task). This step splits into cases dictated by the following parameters:

- (i) The task to be accomplished.




FIGURE 8

(ii) The values $X_1, X_q \in \{\underline{B}, \underline{W}\}$.

(ii) Sizes preassigned already to map elements in the task region, mainly $d'_1 = 2s(v_1) - d_n(v_1), s(f_1), s(f_q)$.

We shall describe this step by displaying the actual solution for each of the possible tasks, under the most tight assumption about (iii), namely: The size of a nontask black region is 3. The size of a nontask vertex is 3 (i.e., its degree is 6) and the size of a nontask white region is at most 7 (nontask white regions always possess an arc (possibly degenerated to a point) on C_{n+1} as part of their boundary, so their size is always augmentable by adding vertices on that arc). These cases are displayed in Figure 8. The other (easier) cases can be safely left to the reader.

With the mentioned assumptions about sizes, we denote the cases as follows: Let $X, Y \in \{\underline{B}, \underline{W}\}$. $B.k.X.Y$ ($V.k.X.Y$, $W.k.X.Y$) will stand for task $B.k$ ($V.k$, $W.k$) with $X_1 = X, X_q = Y$. In addition, the assumption $d'_1 = 1$ or $d'_1 = 2$ is utilized for the demonstration. In fact, $d'_1 = 1$ is ruled out, as we are in Case a; but obviously, $d'_1 > 1$ only facilitates the construction. In each figure, the letter "T" denotes the task. We note that once we display a figure for $U.k.B, \underline{W}$ ($U = B, V$ or W), a figure for $U.k.W.B$ will be obtained by reflecting on the real axis.

Step 2.4 (completion of M_{n+1}). Proceed to define M_{n+1} outside the task region as in Case 1, by suitably modifying Steps 1.2–1.4.

BIBLIOGRAPHY

- [ACM] Z. Arad, D. Chillag and G. Moran, *Groups with a small covering number*, Chapter 4 of [AH].
- [AH] Z. Arad and M. Herzog, *Products of conjugacy classes in groups*, Lecture Notes in Math., vol. 1112, Springer, 1985.
- [B] E. Bertram, *On a theorem of Schreier and Ulam for permutations*, J. Algebra **24** (1973), 316–322.
- [Bo] G. Boccara, *Cycles comme produit de deux permutations de classes donnees*, Discrete Math. **38** (1982), 129–142.
- [D1] M. Droste, *Products of conjugacy classes of the infinite symmetric groups*, Discrete Math. **47** (1983), 35–48.
- [D2] —, *Cubes of conjugacy classes covering the infinite symmetric group*, Trans. Amer. Math. Soc. **288** (1985), 381–393.
- [D3] —, *Squares of conjugacy classes in the infinite symmetric group*, Trans. Amer. Math. Soc. **303** (1987), 503–515.
- [DG1] M. Droste and R. Gobel, *On a theorem of Baer, Schreier and Ulam for permutations*, J. Algebra **58** (1979), 282–290.
- [DG2] —, *Products of conjugate permutations*, Pacific J. Math. **92** (1981), 47–60.
- [Dv] Y. Dvir, *Covering properties of permutation groups*, Chapter 4 of [AH].
- [G] A. B. Gray, *Infinite symmetric and monomial groups*, Ph.D. Thesis, New Mexico State University, Las Cruces, New Mexico, 1960.
- [H] P. Hall, *Some constructions for locally finite groups*, J. London Math. Soc. **34** (1959), 305–319.
- [M1] G. Moran, *The algebra of reflections of an infinite set*, Notices Amer. Math. Soc. **73T** (1973), A193.

- [M2] —, *The product of two reflection classes of the symmetric group*, Discrete Math. **15** (1976), 63–77.
- [M3] —, *Parity features for classes of the infinite symmetric group*, J. Combin. Theory Ser A **33** (1982), 82–98.
- [M4] —, *Of planar Eulerian graphs and permutations*, Trans. Amer. Math. Soc. **287** (1985), 323–341.
- [M5] —, *The product of conjugacy classes in some infinite simple groups*, Israel J. Math. **50** (1985), 54–74.
- [M6] —, *Products of involution classes in infinite symmetric groups*, Trans. Amer. Math. Soc. **307** (1988), 745–762.
- [O] O. Ore, *Some remarks on commutators*, Proc. Amer. Math. Soc. **2** (1951), 307–314.
- [S] W. R. Scott, *Group theory*, Prentice-Hall, Englewood Cliffs, N.J., 1964.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, HAIFA 31999, ISRAEL