

DISTAL FUNCTIONS AND UNIQUE ERGODICITY

EBRAHIM SALEHI

ABSTRACT. A. Knapp [5] has shown that the set, $D(S)$, of all distal functions on a group S is a norm closed subalgebra of $l^\infty(S)$ that contains the constants and is closed under the complex conjugation and left translation by elements of S . Also it is proved that [7] for any $k \in \mathbb{N}$ and any $\lambda \in \mathbb{R}$ the function $f: \mathbb{Z} \rightarrow \mathbb{C}$ defined by $f(n) = e^{i\lambda n^k}$ is distal on \mathbb{Z} . Now let W be the norm closure of the algebra generated by the set of functions

$$\{n \mapsto e^{i\lambda n^k} : k \in \mathbb{N}, \lambda \in \mathbb{R}\},$$

which will be called the *Weyl algebra*. According to the facts mentioned above, all members of the Weyl Algebra are distal functions on \mathbb{Z} . In this paper, we will show that any element of W is uniquely ergodic (Theorem 2.13) and that the set W does not exhaust all the distal functions on \mathbb{Z} (Theorem 2.14). The latter will answer the question that has been asked (to the best of my knowledge) by P. Milnes [6].

The term Weyl algebra is suggested by S. Glasner. I would like to express my warmest gratitude to S. Glasner for his helpful advise, and to my advisor Professor Namioka for his enormous helps and contributions.

1. PRELIMINARIES

Let S be a semigroup and $X = l^\infty(S)$ be the algebra of all bounded complex functions on S , equipped with the topology of pointwise convergence on S . Then $(S, l^\infty(S))$ forms a flow, where the action of S on $X = l^\infty(S)$ is defined by

$$(s, f) \mapsto sf = R_s f,$$

where $(sf)(t) = (R_s f)(t) = f(ts)$ for all $f \in X$ and $s, t \in S$. A member $f \in l^\infty(S)$ is called a *distal function* on S if it is, distal point relative to the flow $(S, l^\infty(S))$. For this flow and its distal functions we have the following results [7, 8]:

1.1. Theorem. *Let S be a semigroup and let $\Sigma(X)$ be the enveloping semigroup of the flow $(S, l^\infty(S))$. Then $\Sigma(X)$ is compact. Furthermore, each $\sigma \in \Sigma(X)$ is linear, multiplicative, and preserves the complex conjugation and the constant functions. Also $\|\sigma f\| \leq \|f\|$ holds for all $f \in l^\infty(S)$.*

Received by the editors June 20, 1988 and, in revised form, January 31, 1989. Presented to the Society April 12, 1986 in Indianapolis, Indiana.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 54H20; Secondary 28D99.

1.2. Theorem. Let D be the space of all distal functions on a semigroup S . Then D is a norm closed subalgebra of $l^\infty(S)$ containing constants and is closed under complex conjugation. Furthermore, D is invariant under each member σ of the enveloping semigroup of the flow $(S, l^\infty(S))$.

Given two flows (S, X) , and (S, Y) a continuous function $\varphi: X \rightarrow Y$ is called a *flow homomorphism* if $\varphi(sx) = s\varphi(x)$ holds for all $s \in S$ and $x \in X$. If, in addition, φ is onto (or 1-1), then it is called a *flow epimorphism* (or *monomorphism*). The following lemma facilitates a way to generate distal functions on S from distal points of a flow (S, X) .

1.3. Lemma. Let (S, X) be a flow, $F \in C(X)$, and $x_0 \in X$ be a distal point, and let the function $f: S \rightarrow \mathbb{C}$ be defined by $f(s) = F(sx_0)$. Then f is a distal on S .

Proof. Consider the mapping $\varphi: X \rightarrow l^\infty(S)$ defined by

$$\varphi(x)(t) = F(tx) \quad \text{for all } x \in X, t \in S,$$

which is a flow homomorphism. Since the homomorphic image of a distal flow is again distal [2], φ maps any distal point of X to a distal point of $l^\infty(S)$. Thus, $f = \varphi(x_0)$ is a distal function on S .

A (classical) *dynamical system* is a pair (X, T) , where X is a nonempty compact T_2 topological space, and T is a continuous map of X into itself. Note that this is precisely the flow (\mathbb{N}, X) with the action

$$(n, x) \mapsto T^n x,$$

and if $T: X \rightarrow X$ is a homeomorphism, then it is the flow (\mathbb{Z}, X) with the same action. Let (X, T) and (Y, S) be two dynamical systems. Then a continuous function $\varphi: X \rightarrow Y$ is called a *homomorphism* if $\varphi(Tx) = S\varphi(x)$. If, in addition, φ is onto, then it is called an *epimorphism*. A measure μ on X is said to be *invariant* under T if $\mu(f) = \mu(f \circ T)$ holds for all $f \in C(X)$. Following H. Furstenberg [3] we shall refer to the triple (X, T, μ) as a *process*, where (X, T) is a dynamical system, and μ is an invariant probability measure on the Borel subsets of X . For any $f \in C(X)$ we define an associated sequence $\{f_n\}$ of functions on X by

$$f_n(x) = \frac{1}{n+1} \sum_{m=0}^n f(T^m x).$$

A point $x \in X$ is said to be *generic* for the process (X, T, μ) if for any $f \in C(X)$ the sequence $\{f_n(x)\}$ converges to $\mu(f)$. A dynamical system (X, T) is said to be *uniquely ergodic* if there is a unique invariant probability measure on the Borel subsets of X ; if in addition, X is minimal, then (X, T) is called *strictly ergodic*.

The following theorem is a consequence of the Markov-Kakutani fixed point theorem [1, p. 45].

1.4. Theorem. *For any dynamical system (X, T) there is a probability measure μ on the Borel subsets of X that is invariant under T .*

And, for uniquely ergodic processes we have the following theorem [3, 9].

1.5. Theorem. *Let (X, T, μ) be a process. Then the following are equivalent:*

- (a) (X, T) is uniquely ergodic with invariant measure μ ;
- (b) For any $f \in C(X)$ its associated sequence $\{f_n\}$ converges to $\mu(f)$ uniformly on X ;
- (c) Every point of X is generic for the process (X, T, μ) .

1.6. Corollary. *Let (X, T) be a dynamical system in which for any $f \in C(X)$ its associated sequence $\{f_n\}$ converges pointwise to a constant on X . Then (X, T) is uniquely ergodic.*

1.7. Corollary. *Let A be a dense linear subspace of $C(X)$, and assume that for any $f \in A$ its associated sequence $\{f_n\}$ converges pointwise to a constant. Then (X, T) is uniquely ergodic.*

1.8. Corollary. *If the process (X, T, μ) is uniquely ergodic and $\varphi: (X, T) \rightarrow (Y, S)$ is an epimorphism, then (Y, S) is also uniquely ergodic.*

2. ERGODICITY OF ELEMENTS OF THE WEYL ALGEBRA

In this section we would like to study the dynamical system (X, S) , where $X = l^\infty(\mathbb{Z})$, and $S: X \rightarrow X$ is the shift operator defined by $Sf(n) = f(n+1)$. We note that this is exactly the flow $(\mathbb{Z}, l^\infty(\mathbb{Z}))$. For a function $f \in X$, let X_f be the orbit closure of f , that is, the closure of the set $\{S^n f; n \in \mathbb{Z}\}$. Since Σ , the enveloping semigroup of $(\mathbb{Z}, l^\infty(\mathbb{Z}))$, is compact, we have

$$X_f = \Sigma f = \{\sigma(f): \sigma \in \Sigma\}.$$

Also, if f is distal, then the X_f is minimal. Clearly, $S: X \rightarrow X$ is a homeomorphism, and $S(X_f) = X_f$. Thus the restriction of S to X_f , which will be denoted by S_f , is still a homeomorphism, and we can consider the dynamical system (X_f, S_f) . By Theorem 1.4, there is a probability measure on the Borel subsets of X_f , which is invariant under S_f , and therefore invariant under all S^n ($n \in \mathbb{Z}$).

2.1. Definition. A function $f \in X$ is said to be *uniquely ergodic* if the system (X_f, S_f) is uniquely ergodic. If in addition X_f is also minimal, then we call f *strictly ergodic*.

For a distal function f , since X_f is minimal unique ergodicity is the same as strict ergodicity; therefore we do not make any distinction between them.

2.2. Theorem. *Suppose that $f \in X$ is uniquely ergodic and that the function $F: \mathbb{C} \rightarrow \mathbb{C}$ is continuous. Then $F \circ f$ is uniquely ergodic.*

Proof. Let $h = F \circ f$ and X_h be the orbit closure of h , and let $\varphi: X_f \rightarrow X_h$ be defined by $\varphi(g) = F \circ g$ ($g \in X_f$). Then φ is an epimorphism, and thus by Corollary 1.8, h is uniquely ergodic.

2.3. Theorem. For a finite subset $\Omega = \{\omega_1, \omega_2, \dots, \omega_j\}$ of \mathbb{Z} let the functions

$$E(\Omega): X \rightarrow \mathbb{C}, \quad \overline{E}(\Omega): X \rightarrow \mathbb{C},$$

defined by

$$E(\Omega)(f) = f(\omega_1)f(\omega_2)\cdots f(\omega_j), \quad \text{and} \quad E(\emptyset)(f) = 1, \\ \overline{E}(\Omega)(f) = \overline{E(\Omega)(f)},$$

where $\overline{E(\Omega)(f)}$ is the complex conjugate of $E(\Omega)(f)$. Then f is uniquely ergodic if and only if for any finite subsets Ω, Δ of \mathbb{Z} ,

$$\frac{1}{n+1} \sum_{m=0}^n E(\Omega)\overline{E}(\Delta)(S^m(\sigma f))$$

converges, as $n \rightarrow \infty$, to a constant $c(\Omega, \Delta)$ that is independent of $\sigma \in \Sigma$.

Proof. \Rightarrow Let $\varphi = E(\Omega)\overline{E}(\Delta)$. Then $\varphi' \in C(X_f)$, and by 1.5(b), its associated sequence $\{\varphi_n\}$ converges to $\mu(\varphi)$ uniformly on X_f .

\Leftarrow Let A be the linear subspace of $C(X_f)$ generated by all the functions of the form $E(\Omega)\overline{E}(\Delta)$ with Ω and Δ being any finite subsets of \mathbb{Z} . Then by the Stone-Weierstrass theorem, A is dense in $C(X_f)$, and the theorem follows from Corollary 1.7.

2.4. Corollary. If f is uniquely ergodic, then the limit

$$\lim_n \frac{1}{n+1} \sum_{m=0}^n [\sigma(f)(m)]$$

exists and is a constant independent of σ .

Proof. With the notation of the Theorem 2.3, let $\Omega = \{0\}$, and $\Delta = \emptyset$.

2.5. Corollary. If the function f is uniquely ergodic, then so is τf for any $\tau \in \Sigma$.

Proof. Let $\sigma \in \Sigma$, and $\nu = \sigma \circ \tau$. Then by Theorem 2.3,

$$\frac{1}{n+1} \sum_{m=0}^n E(\Omega)\overline{E}(\Delta)(S^m(\sigma(\tau f))) = \frac{1}{n+1} \sum_{m=0}^n E(\Omega)\overline{E}(\Delta)(S^m(\nu f)),$$

which converges uniformly to a constant independent of ν .

2.6. Lemma. Let $p(x) = a_k x^k + a_{k-1} x^{k-1} + \cdots + a_1 x + a_0$ be a real polynomial, and the function $f: \mathbb{Z} \rightarrow \mathbb{C}$ be defined by

$$f(x) = e^{ip(x)} \quad (x \in \mathbb{Z}).$$

Then for each $\sigma \in \Sigma$ there is a real polynomial p_σ whose leading coefficient is a_k , and $\sigma f(n) = e^{ip_\sigma(n)}$.

Proof. The conclusion follows obviously if $k = \deg p(x) = 0$. So let $k > 0$, and note that for any $n, s \in \mathbb{Z}$

$$p(n+s) = p(n) + \frac{s}{1!} p'(n) + \frac{s^2}{2!} p''(n) + \cdots + \frac{s^k}{k!} p^{(k)}(n).$$

Now suppose that $\{s_\alpha\}$ is a net in \mathbb{Z} such that $\lim_\alpha s_\alpha = \sigma \in \Sigma$. By taking a subnet, if necessary, we may assume that

$$\lim_\alpha \exp\left(i \frac{s_\alpha^l}{l!}\right) = \exp(i\theta_l) \quad (l = 1, 2, \dots, k).$$

Consequently, for each $n \in \mathbb{Z}$

$$\sigma f(n) = \lim_\alpha f(n + s_\alpha) = \lim_\alpha e^{ip(n+s_\alpha)} = e^{ip_\sigma(n)},$$

where $p_\sigma(n) = p(n) + \theta_1 p'(n) + \theta_2 p''(n) + \dots + \theta_k p^{(k)}(n)$ is a real polynomial in n , whose leading coefficient is the same as that of $p(n)$, i.e., a_k .

2.7. Lemma. Let $F: \mathbb{Z} \rightarrow \mathbb{C}$ be a periodic function with period $u > 0$, and $p(x) = a_q x^q + a_{q-1} x^{q-1} + \dots + a_1 x + a_0$ be a real polynomial for which, if $q > 0$, the leading coefficient a_q is an irrational multiple of π . Then

$$\lim_{n \rightarrow \infty} \left[\frac{1}{n+1} \sum_{m=0}^n F(m) e^{ip(m)} \right] = \begin{cases} 0, & \text{if } q > 0, \\ \frac{e^{ia_0}}{u} [F(0) + \dots + F(u-1)], & \text{if } q = 0. \end{cases}$$

Proof. Observe that, for any $n \in \mathbb{N}$, there is a nonnegative integer k with $n = ku + r$ ($0 \leq r < u$). Hence

$$\begin{aligned} \frac{1}{n+1} \sum_{m=0}^n F(m) e^{ip(m)} &= \frac{ku}{ku+r+1} \cdot \frac{1}{ku} \sum_{m=0}^{ku-1} F(m) e^{ip(m)} \\ &\quad + \frac{1}{n+1} \sum_{m=ku}^n F(m) e^{ip(m)}. \end{aligned}$$

If $F(m) e^{ip(m)}$ is bounded by M , then $|\sum_{m=ku}^n F(m) e^{ip(m)}| \leq uM$. Therefore,

$$\lim_{n \rightarrow \infty} \left[\frac{1}{n+1} \sum_{m=ku}^n F(m) e^{ip(m)} \right] = 0.$$

Also, $\lim_{k \rightarrow \infty} ku/(ku+r+1) = 1$. Thus it is sufficient to calculate

$$\lim_{k \rightarrow \infty} \left[\frac{1}{ku} \sum_{m=0}^{ku-1} F(m) e^{ip(m)} \right].$$

But

$$\begin{aligned}
 \sum_{m=0}^{ku-1} F(m)e^{ip(m)} &= \sum_{j=0}^{k-1} \sum_{m=ju}^{(j+1)u-1} F(m)e^{ip(m)} \\
 &= \sum_{j=0}^{k-1} \sum_{m=0}^{u-1} F(m+ju)e^{ip(m+ju)} \\
 &= \sum_{j=0}^{k-1} \sum_{m=0}^{u-1} F(m)e^{ip(m+ju)} \\
 &= \sum_{m=0}^{u-1} \sum_{j=0}^{k-1} F(m)e^{ip(m+ju)} \\
 &= \sum_{m=0}^{u-1} F(m) \sum_{j=0}^{k-1} e^{ip(m+ju)}.
 \end{aligned}$$

If $q = \deg p(x) = 0$, then $e^{ip(m+ju)} = e^{ia_0}$, and

$$\begin{aligned}
 \frac{1}{ku} \sum_{m=0}^{ku-1} F(m)e^{ip(m)} &= \frac{1}{ku} \sum_{m=0}^{u-1} F(m) \sum_{j=0}^{k-1} e^{ip(m+ju)} \\
 &= \frac{1}{ku} \sum_{m=0}^{u-1} F(m) \sum_{j=0}^{k-1} e^{ia_0} \\
 &= \frac{e^{ia_0}}{u} [F(0) + \cdots + F(u-1)].
 \end{aligned}$$

For $q > 0$, note that $p(m+ju)$ is a polynomial in j , whose leading coefficient, $a_q u^q$, is an irrational multiple of π . Then, by the theorem of Weyl [10, Satz 9]

$$\lim_{k \rightarrow \infty} \left[\frac{1}{k} \sum_{j=0}^{k-1} e^{ip(m+ju)} \right] = 0.$$

Consequently

$$\lim_{k \rightarrow \infty} \left[\frac{1}{ku} \sum_{m=0}^{ku-1} F(m)e^{ip(m)} \right] = \sum_{m=0}^{u-1} F(m) \left\{ \lim_{k \rightarrow \infty} \left[\frac{1}{ku} \sum_{j=0}^{k-1} e^{ip(m+ju)} \right] \right\} = 0,$$

and this proves the lemma.

2.8. Theorem. Let $k \in \mathbb{N}$ be fixed and let

$$p(x) = a_k x^k + a_{k-1} x^{k-1} + \cdots + a_1 x + a_0$$

be a real polynomial of degree k . Then the function $h: \mathbb{Z} \rightarrow \mathbb{C}$ defined by $h(x) = e^{ip(x)}$ is uniquely ergodic.

Proof. If a_1, a_2, \dots, a_k , are all rational multiplies of π , then h is periodic and X_h is finite and minimal. Hence it is uniquely ergodic. So, we may assume

that at least one of the coefficients a_1, a_2, \dots, a_k , is an irrational multiple of π . Suppose that a_q ($0 < q \leq k$) is the last such number, and let

$$\begin{aligned} Q(n) &= a_k x^k + \dots + a_{q+1} x^{q+1}, \\ R(n) &= a_q x^q + \dots + a_1 x + a_0. \end{aligned}$$

If $f(n) = e^{iQ(n)}$ and $g(n) = e^{iR(n)}$, then $h(n) = f(n)g(n)$, and the function f is a periodic function (say, with period u). By Lemma 2.6, for any σ , there is a real polynomial R_σ with the same leading coefficient as that of R (i.e., a_q), such that $\sigma g(n) = e^{iR_\sigma(n)}$. Now let $\Omega = \{\omega_1, \omega_2, \dots, \omega_j\}$, and $\Delta = \{\delta_1, \delta_2, \dots, \delta_l\}$ be any two finite subsets of \mathbb{Z} . Then by Theorem 2.3, we must show that

$$\lim_{n \rightarrow \infty} \left[\frac{1}{n+1} \sum_{m=0}^n E(\Omega) \bar{E}(\Delta) (S^m(\sigma h)) \right]$$

exists and its value is independent of σ . Using the properties of σ , S , E , and \bar{E} we note that

$$\begin{aligned} E(\Omega) \bar{E}(\Delta) (S^m \sigma h) &= E(\Omega) \bar{E}(\Delta) (S^m(\sigma f \sigma g)) \\ &= [E(\Omega) \bar{E}(\Delta) (S^m(\sigma f))] \cdot [E(m + \Omega) \bar{E}(m + \Delta) (\sigma g)] \\ &= [E(\Omega) \bar{E}(\Delta) (S^m(\sigma f))] \cdot [E(m + \Omega) \bar{E}(m + \Delta) e^{iR_\sigma}] \\ &= E(\Omega) \bar{E}(\Delta) (S^m(\sigma f)) \cdot e^{iG_\sigma(m)}, \end{aligned}$$

where $G_\sigma(m) = \sum_{\omega \in \Omega} R_\sigma(m + \omega) - \sum_{\delta \in \Delta} R_\sigma(m + \delta)$ is a polynomial in m . Now let $F: \mathbb{Z} \rightarrow \mathbb{C}$ be defined by

$$F(m) = E(\Omega) \bar{E}(\Delta) (S^m f).$$

Then F is periodic function of period u , and a straightforward calculation shows that

$$E(\Omega) \bar{E}(\Delta) (S^m \sigma f) = (\sigma F)(m).$$

Consequently,

$$\frac{1}{n+1} \sum_{m=0}^n E(\Omega) \bar{E}(\Delta) (S^m(\sigma h)) = \frac{1}{n+1} \sum_{m=0}^n (\sigma F)(m) e^{iG_\sigma(m)}.$$

Also, if $\{s_\alpha\}$ is a net in \mathbb{Z} such that $\lim_\alpha s_\alpha = \sigma \in \Sigma$, then

$$\begin{aligned} \sigma F(0) + \dots + \sigma F(u-1) &= \lim_\alpha [F(0 + s_\alpha) + \dots + F(u-1 + s_\alpha)] \\ &= \lim_\alpha [F(0) + \dots + F(u-1)] = F(0) + \dots + F(u-1), \end{aligned}$$

which shows that $\sigma F(0) + \dots + \sigma F(u-1)$ is a constant independent of $\sigma \in \Sigma$. Similarly, one can show that σF is also a periodic function of period u . On the other hand, by Taylor's theorem

$$\begin{aligned} G_\sigma(m) &= \sum_{\omega \in \Omega} R_\sigma(m + \omega) - \sum_{\delta \in \Delta} R_\sigma(m + \delta) \\ &= \lambda_0 R_\sigma(m) + \lambda_1 R'_\sigma(m) + \dots + \lambda_q R_\sigma^{(q)}(m), \end{aligned}$$

where $\lambda_\nu = \frac{1}{\nu!}(\omega_1^\nu + \cdots + \omega_j^\nu - \delta_1^\nu - \cdots - \delta_l^\nu)$ ($\nu = 0, 1, \dots, q$) is a rational number independent of σ .

Now, if $\lambda_0 = \lambda_1 = \cdots = \lambda_{q-1} = 0$, then $G_\sigma(m) = \lambda_q a_q q!$ and, by Lemma 2.7,

$$\begin{aligned} \lim_{n \rightarrow \infty} \left[\frac{1}{n+1} \sum_{m=0}^n (\sigma F)(m) e^{iG_\sigma(m)} \right] &= \frac{e^{i\lambda_q a_q q!}}{u} [\sigma F(0) + \cdots + \sigma F(u-1)] \\ &= \frac{e^{i\lambda_q a_q q!}}{u} [F(0) + \cdots + F(u-1)], \end{aligned}$$

which is independent of σ . If one of $\lambda_0, \lambda_1, \dots, \lambda_{q-1}$ is nonzero, say $\lambda_r \neq 0$, but $\lambda_0 = \lambda_1 = \cdots = \lambda_{r-1} = 0$ ($0 \leq r < q$), then $\deg G_\sigma(m) = r$, and its leading coefficient is the same as that of $\lambda_r R_\sigma^{(r)}(m)$, which is an irrational multiple of π . Therefore, by Lemma 2.7,

$$\lim_{n \rightarrow \infty} \left[\frac{1}{n+1} \sum_{m=0}^n (\sigma F)(m) e^{iG_\sigma(m)} \right] = 0.$$

Thus, in either case, the limit

$$\lim_{n \rightarrow \infty} \left[\frac{1}{n+1} \sum_{m=0}^n E(\Omega) \bar{E}(\Delta) (S^m \sigma h) \right] = \lim_{n \rightarrow \infty} \left[\frac{1}{n+1} \sum_{m=0}^n (\sigma F)(m) e^{iG(m)} \right]$$

exists and is independent of $\sigma \in \Sigma$. This completes the proof.

2.9. Theorem. Let the system (X, T) be strictly ergodic, with $T: X \rightarrow X$ a homeomorphism, and let the function $\Lambda: C(X) \times X \rightarrow l^\infty(\mathbb{Z})$ be defined by $\Lambda(F, x)(n) = F(T^n x)$. Then any element in the range of Λ is strictly ergodic.

Proof. Let $f = \Lambda(F, x_0)$ be fixed, and define the mapping $\varphi: X \rightarrow l^\infty(\mathbb{Z})$ by $\varphi(x)(n) = F(T^n x)$. Then $\varphi(x_0) = f$, and

$$\varphi(Tx)(n) = F(T^n(Tx)) = F(T^{n+1}x) = \varphi(x)(n+1) = (S\varphi(x))(n)$$

or $\varphi(Tx) = S\varphi(x)$, which shows that φ is a flow homomorphism. Since X is minimal, $\varphi[X] = X_f$, the orbit closure of f . Therefore, by Corollary 1.7, f is strictly ergodic.

Next we would like to show that any function $h: \mathbb{Z} \rightarrow \mathbb{C}$ of the form $h(n) = \sum_{j=1}^r A_j e^{iP_j(n)}$ is uniquely ergodic, where $A_j \in \mathbb{C}$ and $P_j(n)$ is a real polynomial. The proof is presented for the case $r = 2$, but it can be applied to any $r \in \mathbb{N}$. First we need the following theorem.

2.10. Theorem. Let f, g be two uniquely ergodic functions on \mathbb{Z} such that for any four finite subsets $\Omega_1, \Omega_2, \Delta_1$, and Δ_2 of \mathbb{Z} , the function

$$F(\Omega_1, \Omega_2, \Delta_1, \Delta_2) = \prod_{\omega \in \Omega_1} S^\omega f \prod_{\omega \in \Omega_2} S^\omega g \prod_{\delta \in \Delta_1} S^{\delta} \bar{f} \prod_{\delta \in \Delta_2} S^{\delta} \bar{g}$$

is uniquely ergodic. Then $f + g$ is uniquely ergodic.

Proof. First we consider $Y = \Sigma(f, g)$, the orbit closure of (f, g) , and we wish to show that it is uniquely ergodic. By Corollary 1.6, we must prove that for any $\varphi \in C(\Sigma(f, g))$, and any element $\sigma(f, g) = (\sigma f, \sigma g) \in \Sigma(f, g)$ the limit

$$\lim_{n \rightarrow \infty} \left[\frac{1}{n+1} \sum_{m=0}^n \varphi(S^m(\sigma f, \sigma g)) \right]$$

exists and is a constant independent of σ . For this we use the fact that $\Sigma(f, g) \subset \Sigma f \times \Sigma g = X_f \times X_g$. Following the notation of Theorem 2.3, for a finite subset Ω of \mathbb{Z} , let the functions

$$E_1(\Omega): \Sigma f \times \Sigma g \rightarrow \mathbb{C}, \quad E_2(\Omega): \Sigma f \times \Sigma g \rightarrow \mathbb{C}$$

be defined by $E_1(\Omega)(\sigma f, \tau g) = E(\Omega)(\sigma f)$, and $E_2(\Omega)(\sigma f, \tau g) = E(\Omega)(\tau g)$, and let $\overline{E_j}(\Omega)$ be the complex conjugate of $E_j(\Omega)$, $j = 1, 2$. Then $E_j(\Omega)$ is continuous and the set of linear combinations of the functions of the form

$$E_1(\Omega_1)E_2(\Omega_2)\overline{E_1}(\Delta_1)\overline{E_2}(\Delta_2)$$

is dense in $C(\Sigma f \times \Sigma g) = C(X_f \times X_g)$. Thus by Corollary 1.7, is enough to consider $\varphi = E_1(\Omega_1)E_2(\Omega_2)\overline{E_1}(\Delta_1)\overline{E_2}(\Delta_2)$. With this choice

$$\varphi(S^m(\sigma f, \sigma g)) = F(\Omega_1, \Omega_2, \Delta_1, \Delta_2)(m).$$

Since $F(\Omega_1, \Omega_2, \Delta_1, \Delta_2)(m)$ is uniquely ergodic, by Theorem 2.3, the limit

$$\lim_n \left[\frac{1}{n+1} \sum_{m=0}^n \varphi(S^m(\sigma f, \sigma g)) \right]$$

exists and is independent of σ . This proves that $Y = \Sigma(f, g)$ is uniquely ergodic. Next by Theorem 2.9, the mapping $\Lambda: C(Y) \times Y \rightarrow l^\infty(\mathbb{Z})$ defined by $\Lambda(G, y)(n) = G(S^n y)$ produces uniquely ergodic functions. Now let $y_0 = (f, g)$, and $G = E_1(0) + E_2(0)$. Then

$$\Lambda(G, y_0)(n) = G(S^n y_0) = G(S^n(f, g)) = G(S^n f, S^n g) = f(n) + g(n)$$

is uniquely ergodic.

2.11. Corollary. Let $p(n)$, $q(n)$ be any two real polynomials, and $f(n) = e^{ip(n)}$, and $g(n) = e^{iq(n)}$. Then $f + g$ is uniquely ergodic.

Proof. It is sufficient to show that, for any four finite subsets Ω_1 , Ω_2 , Δ_1 , and Δ_2 of integers the function

$$F(\Omega_1, \Omega_2, \Delta_1, \Delta_2) = \prod_{\omega \in \Omega_1} S^\omega f \prod_{\omega \in \Omega_2} S^\omega g \prod_{\delta \in \Delta_1} S^\delta \overline{f} \prod_{\delta \in \Delta_2} S^\delta \overline{g}$$

is uniquely ergodic. But $F(\Omega_1, \Omega_2, \Delta_1, \Delta_2)(n) = e^{iQ(n)}$, where

$$Q(n) = \sum_{\omega \in \Omega_1} p(n + \omega) + \sum_{\omega \in \Omega_2} q(n + \omega) - \sum_{\delta \in \Delta_1} p(n + \delta) - \sum_{\delta \in \Delta_2} q(n + \delta)$$

is a real polynomial in n . Therefore, by Theorem 2.8, $F(\Omega_1, \Omega_2, \Delta_1, \Delta_2)$ is uniquely ergodic.

So far, it has been shown that any element of the algebra A generated by the set $\{n \mapsto e^{i\lambda n^k} : \lambda \in \mathbb{R}, k \in \mathbb{N}\}$ is strictly ergodic. Next, we would like to demonstrate the main result of this paper; that is, any element of the Weyl algebra W is strictly ergodic. To accomplish this result, we need the following theorem.

2.12. Theorem. *The set of all uniquely ergodic functions in $X = l^\infty(\mathbb{Z})$ is closed under uniform convergence.*

Proof. Let $f_\alpha \in X$ be a sequence of uniquely ergodic functions that converge uniformly to f . We choose Ω, Δ to be any two finite subsets of \mathbb{Z} , and define the function $F: l^\infty(\mathbb{Z}) \rightarrow \mathbb{C}$ by $F(f) = E(\Omega)\overline{E}(\Delta)(f)$. For each α , the function f_α is uniquely ergodic, and therefore there is a number c_α independent of $\sigma \in \Sigma$ such that

$$\begin{aligned} \lim_{n \rightarrow \infty} \left[\frac{1}{n+1} \sum_{m=0}^n E(\Omega)\overline{E}(\Delta)(S^m(\sigma f_\alpha)) \right] \\ = \lim_{n \rightarrow \infty} \left[\frac{1}{n+1} \sum_{m=0}^n F(S^m \sigma f_\alpha) \right] = c_\alpha. \end{aligned}$$

Let $M = \sup_\alpha \|f_\alpha\|_\infty$. Then $S^m(\sigma f_\alpha)$ and $S^m(\sigma f)$ are in the set

$$B = \{h \in l^\infty(\mathbb{Z}) : \|h\|_\infty \leq M\}.$$

Since F is uniformly continuous on B , $\{c_\alpha\}$ is Cauchy, and it converges to a number $c \in \mathbb{C}$. Now we would like to show that for any $\sigma \in \Sigma$

$$\lim_{n \rightarrow \infty} \left[\frac{1}{n+1} \sum_{m=0}^n F(S^m(\sigma f)) \right] = c.$$

Given $\varepsilon > 0$, by uniform continuity of F on B , there is a $\delta > 0$ such that

$$h_1, h_2 \in B \text{ and } \|h_1 - h_2\| < \delta \Rightarrow |F(h_1) - F(h_2)| < \varepsilon/3.$$

Let α be chosen so that $|c_\alpha - c| < \varepsilon/3$, and $\|f_\alpha - f\|_\infty < \delta$, which implies that $|F(f_\alpha) - F(f)| < \varepsilon/3$. Therefore

$$\begin{aligned} \left| \frac{1}{n+1} \sum_{m=0}^n F(S^m(\sigma f)) - c \right| &\leq \frac{1}{n+1} \sum_{m=0}^n |F(S^m(\sigma f)) - F(S^m(\sigma f_\alpha))| \\ &\quad + \left| \frac{1}{n+1} \sum_{m=0}^n F(S^m(\sigma f_\alpha)) - c_\alpha \right| + |c_\alpha - c|. \end{aligned}$$

Now, since f_α is uniquely ergodic, there is a number $M_0 \in \mathbb{N}$ such that

$$n \geq M_0 \Rightarrow \left| \frac{1}{n+1} \sum_{m=0}^n F(S^m(\sigma f_\alpha)) - c_\alpha \right| < \frac{\varepsilon}{3};$$

therefore

$$n \geq M_0 \Rightarrow \left| \frac{1}{n+1} \sum_{m=0}^n F(S^m(\sigma f)) - c \right| < \varepsilon,$$

which completes the proof.

2.13. Theorem. *Any element of the Weyl algebra is uniquely ergodic.*

Proof. This follows from Theorem 2.12 and the fact that the Weyl algebra is the uniform closure of a set of uniquely ergodic functions.

2.14. Theorem. *The Weyl algebra does not exhaust all distal functions.*

Proof. By Theorem 2.13, it is enough to introduce a distal function on \mathbb{Z} that is not uniquely ergodic. Let $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$ be the unit circle, $\lambda \in \mathbb{T}$, and $g: \mathbb{T} \rightarrow \mathbb{T}$ be any continuous function, and let $\varphi: \mathbb{T}^2 \rightarrow \mathbb{T}^2$ be defined by

$$\varphi(u, v) = (\lambda u, g(u)v).$$

Then the flow $(\mathbb{Z}, \mathbb{T}^2)$ is distal, where the action is $(n, \xi) \mapsto \varphi^n(\xi)$ [4]. In [3], Furstenberg has introduced a continuous function $g: \mathbb{T} \rightarrow \mathbb{T}$ and an element $\lambda \in \mathbb{T}$ such that not all ergodic averages do exist; that is, there is a $\xi \in \mathbb{T}^2$ and $h \in C(\mathbb{T}^2)$ such that

$$\lim_{n \rightarrow \infty} \left[\frac{1}{n+1} \sum_{m=0}^n h(\varphi^m(\xi)) \right]$$

does not exist. Now if we define $f: \mathbb{Z} \rightarrow \mathbb{C}$ by $f(m) = h(\varphi^m(\xi))$, then, by Lemma 1.3, f is distal, while, by Corollary 2.4 (with $\sigma = R_0$, the right translation by 0); it is not uniquely ergodic ($f \notin \mathbf{W}$).

REFERENCES

1. J. R. Brown, *Ergodic theory and topological dynamics*, Academic Press, New York, 1976.
2. Robert Ellis, *Lectures on topological dynamics*, Benjamin, New York, 1969.
3. H. Furstenberg, *Strict ergodicity and transformation of the torus*, Amer. J. Math. **83** (1961), 573–601.
4. —, *The structure of distal flow*, Amer. J. Math. **85** (1963), 477–515.
5. A. W. Knap, *Distal functions on groups*, Trans. Amer. Math. Soc. **128** (1967), 1–40.
6. Paul Milnes, *Minimal and distal functions on a semidirect products of groups*, Preprint.
7. Isaac Namioka, *Affine flows and distal points*, Math. Z. **184** (1983), 259–269.
8. —, *Right topological groups, distal flows and a fixed point theorem*, Math. Systems Theory **6** (1972), 193–209.
9. J. C. Oxtoby, *Ergodic sets*, Bull. Amer. Math. Soc. **58** (1952), 116–136.
10. H. Weyl, *Über die Gleichverteilung von Zahlen mod eins*, Math. Ann. **77** (1916), 313–352.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEVADA, LAS VEGAS, NEVADA 89154