

MULTIVARIATE ORTHOGONAL POLYNOMIALS AND OPERATOR THEORY

YUAN XU

ABSTRACT. The multivariate orthogonal polynomials are related to a family of commuting selfadjoint operators. The spectral theorem for these operators is used to prove that a polynomial sequence satisfying a vector-matrix form of the three-term relation is orthonormal with a determinate measure.

1. INTRODUCTION

The purpose of this paper is to use the operator theory to study the multivariate orthogonal polynomials. First let us recall the relevant one variable theory of orthogonal polynomials (cf. [5, 16]).

Let $\{p_n(x)\}_{n=0}^{\infty}$ be a sequence of polynomials satisfying the three-term relations

$$xp_n(x) = a_np_{n+1}(x) + b_np_n(x) + a_{n-1}p_{n-1}(x).$$

Then Favard's Theorem states that the sequence p_n is orthonormal with respect to a measure μ if and only if a_n are positive real numbers. It is known that we can relate the measure of orthogonality to the spectral measure of a selfadjoint operator; the matrix representation of this operator is a Jacobi matrix (cf. [6, 15]). Therefore it is possible to use operator theory to study the nature of the measure (cf. [6, 7]). In particular, one can prove that the measure is of compact support if and only if $\{a_n\}$ and $\{b_n\}$ are bounded. However, the usual way of proving the latter result (cf. [5]) does not use operator theory, but a more elementary method which requires the knowledge of the zeros of orthogonal polynomials.

The theory of multivariate orthogonal polynomials is far from complete. There are only a few papers in the literature dealing with the general multivariable theory (cf. [4, 9, 10, 17]). In a recent paper [17], we have proved an extension of Favard's Theorem, where the orthogonality is with respect to a quasi-inner product (see §2). Earlier results can be found in [9]. The three-terms relation now takes the vector-matrix form

$$x_i \mathbb{P}_n(\mathbf{x}) = A_{n,i} \mathbb{P}_{n+1}(\mathbf{x}) + B_{n,i} \mathbb{P}_n(\mathbf{x}) + A_{n-1,i}^T \mathbb{P}_{n-1}(\mathbf{x}), \quad 1 \leq i \leq d,$$

where $\mathbf{x} = (x_1, \dots, x_d)^T \in \mathbf{R}^d$, \mathbb{P}_n is a polynomial vector, and $A_{n,i}$ and $B_{n,i}$ are matrices (see §2). One problem for the multivariate orthogonal polynomials

Received by the editors May 14, 1992.

1991 *Mathematics Subject Classification.* Primary 42C05, 47B15.

Key words and phrases. Multivariate orthogonal polynomials, recurrence relation, commuting selfadjoint operators, determinate measure.

is that the zeros are actually common zeros of a family of polynomials, thus, the elegant theorems for zeros in the one variable case are no longer available. However, as we shall show, the multivariate orthogonal polynomials are related to a family of commuting selfadjoint operators. The purpose of this paper is to establish this relation. In particular, we shall prove that if the norms of matrices in the three-terms relation are bounded, then the set of polynomials is orthonormal with respect to a nonnegative measure with compact support. In a subsequent paper the nature of the measure will be studied via the operator approach.

The paper is organized as follows. In §2 we introduce the notation and present the basic results of multivariate orthogonal polynomials. The main results are stated in §3. The operator approach and the proof of the main theorem are in §4.

2. MULTIVARIATE ORTHOGONAL POLYNOMIALS

Let \mathbf{N}_0 be the set of nonnegative integers. For $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbf{N}_0^d$ and $\mathbf{x} = (x_1, \dots, x_d) \in \mathbf{R}^d$ we write $\mathbf{x}^\alpha = x_1^{\alpha_1} \cdots x_d^{\alpha_d}$. The number $|\alpha| = \alpha_1 + \cdots + \alpha_d \in \mathbf{N}_0$ is called the total degree of \mathbf{x}^α . For $n \in \mathbf{N}_0$ we denote by Π_n^d the set of polynomials of total degree at most n in d variables, and Π^d the set of all polynomials in d variables.

A real valued linear functional \mathcal{L} is said to induce a quasi-inner product on Π^d by the formula $\langle P, Q \rangle = \mathcal{L}(PQ)$ provided there exists a basis B of Π^d such that

$$(2.1) \quad \mathcal{L}(PQ) \begin{cases} = 0, & \text{if } P \neq Q, \\ \neq 0, & \text{if } P = Q, \forall P, Q \in B, \end{cases}$$

and $B = \{P_j^k\}_{k=0}^\infty_{j=1}^{r_k^d}$, where the superscript k means that P_j^k is of total degree k and $r_k^d = \dim \Pi_k^d - \dim \Pi_{k-1}^d$. We take $\mathcal{L}(P^2) = 1$ in (2.1) when \mathcal{L} induces an inner product. Two polynomials P and Q are said to be orthogonal to each other with respect to \mathcal{L} if $\mathcal{L}(PQ) = 0$. For each $k \in \mathbf{N}_0$, let $V_k^d \subset \Pi_k^d$ denote the set of polynomials of degree exactly equal to k , together with zero, that are orthogonal to all polynomials in Π_{k-1}^d . Then V_k is a vector space of dimension $r_k^d = \binom{k+d-1}{k}$, and V_k 's are mutually orthogonal. Throughout this paper, the letter d is reserved for the number of variables or the dimension. It is fixed and will be omitted sometimes. For example, we write r_k for r_k^d in the following. For a sequence of polynomials $\{P_j^k\}_{j=1}^{r_k}$ of total degree k , we introduce the vector notation

$$(2.2) \quad \mathbb{P}_k(\mathbf{x}) = [P_1^k(\mathbf{x}), P_2^k(\mathbf{x}), \dots, P_{r_k}^k(\mathbf{x})]^T.$$

If a matrix $P = (p_{ij})$ is given whose elements p_{ij} are polynomials in Π^d , we denote by $\mathcal{L}(P)$ the matrix whose elements are the values $\mathcal{L}(p_{ij})$. For our convenience, if $\{P_j^k\}_{j=1}^{r_k}$ is a basis for V_k^d , we shall say that \mathbb{P}_k is a basis for V_k^d and that $\{\mathbb{P}_k\}_{k=0}^\infty$ is an orthogonal basis for Π^d . If \mathcal{L} is an inner product, $\{\mathbb{P}_k\}_{k=0}^\infty$ is orthogonal with respect to \mathcal{L} and $\mathcal{L}(\mathbb{P}_k \mathbb{P}_k^T) = I_{r_k}$, then we say $\{\mathbb{P}_n\}_{n=0}^\infty$ is orthonormal. Throughout this paper, the $n \times n$ identity matrix is denoted by I_n , or simply I . The notation $A : i \times j$ means that A is a matrix of size $i \times j$. For $\mathbf{x} \in \mathbf{R}^d$ we write $\mathbf{x} = (x_1, \dots, x_d)^T$.

In [17], we studied the multivariate orthogonal polynomials from the point of view that the orthogonality is given in terms of V_n rather than a particular basis for V_n . One important characteristic of the orthogonal polynomials is the three-terms recurrence relation, as seen in the following multivariable version of the Favard's Theorem [17].

Theorem 2.1. *Let $B = \{P_j^k\}_{k=0}^\infty_{j=1}^{r_k}$, $P_1^0 \neq 0$, be an arbitrary sequence in Π^d . Then the following statements are equivalent:*

(1) *There exists a linear functional \mathcal{L} which induces a quasi-inner product in Π^d and makes $\{P_k\}_{k=0}^\infty$ an orthogonal basis in Π^d .*

(2) *For $k \geq 1$, $1 \leq i \leq d$, there exist matrices $A_{k,i} : r_k \times r_{k+1}$, $B_{k,i} : r_k \times r_k$, and $C_{k,i} : r_k \times r_{k-1}$, such that*

$$(a) \quad x_i P_k = A_{k,i} P_{k+1} + B_{k,i} P_k + C_{k,i} P_{k-1}, \quad 1 \leq i \leq d.$$

$$(b) \quad \text{rank } A_k = r_{k+1}, \quad \text{rank } C_{k+1} = r_{k+1},$$

where $A_k : dr_k \times r_{k+1}$ and $C_k : r_k \times dr_{k-1}$ are defined by

$$A_k = [A_{k,1}^T | A_{k,2}^T | \cdots | A_{k,d}^T]^T \quad \text{and} \quad C_k = [C_{k,1} | C_{k,2} | \cdots | C_{k,d}].$$

A weaker result appears in [9]. In this paper we study the recurrence relation with $C_{n,i} = A_{n-1,i}^T$. From Theorem 2.1 we have

Theorem 2.2. *Let $\{P_n\}_{n=0}^\infty$, $P_0 = 1$, be a sequence in Π^d . Then the following statements are equivalent:*

(1) *There exists a linear functional which induces an inner product in Π^d and makes $\{P_n\}_{n=0}^\infty$ an orthonormal basis in Π^d .*

(2) *For $k \geq 0$, $1 \leq i \leq d$, there exist matrices $A_{k,i} : r_k \times r_{k+1}$ and $B_{k,i} : r_k \times r_k$, such that*

$$(a) \quad x_i P_k = A_{k,i} P_{k+1} + B_{k,i} P_k + A_{k-1,i}^T P_{k-1}, \quad 1 \leq i \leq d,$$

$$(b) \quad \text{rank } A_k = r_{k+1}, \text{ where } A_{-1,i} \text{ is taken to be zero.}$$

Proof. If $\{P_n\}_{n=0}^\infty$ is orthonormal with respect to an inner product \mathcal{L} , then $\mathcal{L}(P_n P_n^T) = I$ and $\{P_n\}$ satisfies (a) by (2) in Theorem 2.1. On the other hand, if (2) holds, then from Theorem 2.1 there exists a quasi-inner product \mathcal{L} which makes $\{P_n\}_{n=0}^\infty$ orthogonal. We only need to show that the matrix $H_n = \mathcal{L}(P_n P_n^T)$ is the identity matrix for every $n \in \mathbb{N}_0$. Multiplying (a) by P_{k+1} and (a) with k in place of $k-1$ by P_k respectively, and applying \mathcal{L} leads to two formulas for $\mathcal{L}(x_i P_k P_{k+1}^T)$, which gives

$$A_{k,i} H_{k+1} = H_k A_{k,i}, \quad 1 \leq i \leq d.$$

Therefore,

$$(2.3) \quad A_k H_{k+1} = G_k A_k, \quad G_k = \text{diag}(H_k, \dots, H_k).$$

Since $P_0 = 1$, we have $H_0 = [1]$. Suppose H_k is an identity matrix. Then G_k is also an identity matrix. Thus, it follows from (b) and (2.3) that H_{k+1} is an identity matrix. The proof is completed by induction. \square

We note that if \mathcal{L} defines an inner product and $\{P_n\}_{n=0}^\infty$ is an orthogonal basis in Π^d , then $H_n = \mathcal{L}(P_n P_n^T)$ is a positive definite matrix. Hence, there exists a nonsingular matrix S_n such that $H_n = S_n S_n^T$. It is easy to verify that $Q_n = S_n^{-1} P_n$ satisfies the recurrence relation (a) in Theorem 2.2, and Q_n

is an orthonormal basis for V_n . Therefore, from the point of view that the orthogonality is in terms of V_n , it is enough to study the simpler recurrence relation (a) in Theorem 2.2. In particular, it is enough for the purpose of analyzing the measure of orthogonality.

We note that $\{\mathbb{P}_n\}$ satisfying (a) in Theorem 2.2 is uniquely determined by the assumption $\mathbb{P}_0 = 1$ and the rank condition (b). The coefficient matrices in the recurrence relation satisfy the following matrix equations.

Corollary 2.3. *In the statement (2) of Theorem 2.2, the matrices $A_{k,i}$ and $B_{k,i}$ satisfy the equations*

$$(2.4) \quad A_{k,i}A_{k+1,j} = A_{k,j}A_{k+1,i},$$

$$(2.5) \quad A_{k,i}B_{k+1,j} + B_{k,i}A_{k,j} = B_{k,j}A_{k,i} + A_{k,j}B_{k+1,i},$$

$$(2.6) \quad \begin{aligned} & A_{k-1,i}^T, A_{k-1,j} + B_{k,i}B_{k,j} + A_{k,i}A_{k,j}^T \\ & = A_{k-1,j}^T A_{k-1,i} + B_{k,i}B_{k,j} + A_{k,j}A_{k,i}^T, \end{aligned}$$

for $i \neq j$, $1 \leq i, j \leq d$, and $k \geq 0$, where $A_{-1,i} = 0$. Moreover, $B_{k,i}$ are symmetric.

Proof. By Theorem 2.2, there is an inner product \mathcal{L} that makes \mathbb{P}_k orthonormal. From the recurrence relation, we have two different ways of calculating the matrices $\mathcal{L}(x_i x_j \mathbb{P}_k \mathbb{P}_{k+2}^T)$, $\mathcal{L}(x_i x_j \mathbb{P}_k \mathbb{P}_k^T)$, and $\mathcal{L}(x_i x_j \mathbb{P}_k \mathbb{P}_{k+1}^T)$. These calculations lead to the desired matrices equations. For examples, by the recurrence relation,

$$\begin{aligned} \mathcal{L}(x_i x_j \mathbb{P}_k \mathbb{P}_{k+2}^T) &= \mathcal{L}(x_i \mathbb{P}_k x_j \mathbb{P}_{k+2}^T) \\ &= \mathcal{L}[(A_{k,i} \mathbb{P}_{k+1} + \cdots)(\cdots + A_{k+1,j}^T \mathbb{P}_{k+1})^T] = A_{k,i} A_{k+1,j}, \end{aligned}$$

and

$$\mathcal{L}(x_i x_j \mathbb{P}_k \mathbb{P}_{k+2}^T) = \mathcal{L}(x_j \mathbb{P}_k x_i \mathbb{P}_{k+2}^T) = A_{k,j} A_{k+1,i},$$

which leads to equation (2.4). \square

The condition $\text{rank } A_k = r_{k+1}$ implies that there exist matrices $D_{k,i} : r_{k+1} \times r_k$, $1 \leq i \leq d$, $k \geq 0$, such that

$$(2.7) \quad \sum_{i=1}^d D_{k,i} A_{k,i} = I.$$

3. MAIN RESULTS

Let $\mathcal{M} = \mathcal{M}(\mathbf{R}^d)$ denote the set of nonnegative Borel measures μ on \mathbf{R}^d , defined on the σ -algebra of Borel sets, such that

$$\int_{\mathbf{R}^d} |\mathbf{x}^\alpha| d\mu(\mathbf{x}) < +\infty, \quad \forall \alpha \in \mathbf{N}_0^d.$$

We are especially interested in the linear functional that has an integral representation

$$(3.1) \quad \mathcal{L}(f) = \int_{\mathbf{R}^d} f(\mathbf{x}) d\mu(\mathbf{x}), \quad \mu \in \mathcal{M}.$$

Such a \mathcal{L} induces an inner product $\langle \cdot, \cdot \rangle$ by $\langle P, Q \rangle = \mathcal{L}(PQ)$. The measure μ is called a representation of \mathcal{L} . If $\{\mathbb{P}_n\}_{n=0}^\infty$ is orthogonal with respect to \mathcal{L} expressible as in (3.1) we say that $\{\mathbb{P}_n\}_{n=0}^\infty$ is orthogonal with respect to the measure μ . By the moments of $\mu \in \mathcal{M}$ we mean the numbers $\mu_\alpha = \int \mathbf{x}^\alpha d\mu(\mathbf{x})$, $\forall \alpha \in \mathbb{N}_0^d$. Two measures are called equivalent if they have the same moments. If the equivalent class of measures having the same moments as μ consists of μ only, the measure μ is called determinate, a terminology from the theory of moments, see [3, 7].

In contrast to one variable theory, Theorem 2.2 does not say whether the inner-product has a measure of representation. Our main result shows that such a measure exists and is unique, when the coefficient matrices satisfy a boundedness condition.

Let $\|\cdot\|_2$ be the spectral norm for matrices. It is induced by the Euclidean norm for vectors.

$$\|A\|_2 = \max\{\sqrt{\lambda} : \lambda \text{ is an eigenvalue of } A^T A\}.$$

Our main result is as follows.

Theorem 3.1. *Let $\{\mathbb{P}_n\}_{n=0}^\infty$, $\mathbb{P}_0 = 1$, be a sequence in Π^d . Then the following statements are equivalent:*

(i) *There exists a determinate measure $\mu \in \mathcal{M}$ with compact support in \mathbb{R}^d such that $\{\mathbb{P}_n\}_{n=0}^\infty$ is orthonormal with respect to μ .*

(ii) *The statement (2) in Theorem 2.2 together with*

$$(3.2) \quad \sup_{k \geq 2} \|A_{k,i}\|_2 < +\infty, \quad \sup_{k \geq 0} \|B_{k,i}\|_2 < +\infty, \quad 1 \leq i \leq d.$$

The proof of this theorem is in §4. It is based on the spectral theorem of a family of commuting selfadjoint operators. The connection between such a family of operators and multivariate orthogonal polynomials is the main establishment of the paper, and is of interest in itself. The operator approach is also used in the study of the moment problem (cf. [3, 7]). Our results in §4 indicate that there is a close connection between the multivariable moment problem and orthogonal polynomials. We intend to study this connection in our future investigation. Let us also mention that the family of commuting selfadjoint operators plays an important role in the quantum mechanics (cf. [12]). It is interesting to see the possible applications of multivariate orthogonal polynomials in that direction.

For $d = 1$; Theorem 3.1 is well known (cf. [5, p. 75 and p. 109]). In [9], an integral representation for the inner product is given, but the measure is not shown to be nonnegative nor determinate.

4. THE OPERATOR APPROACH

4.1. Spectral theory for commuting selfadjoint operators. We recall the part of spectral theory that will be needed, see [1, 2, 12, 13, 14]. Let \mathcal{H} be a separable Hilbert space. Each selfadjoint operator T in \mathcal{H} has a unique spectral representation $T = \int x dE(x)$ where E is the spectral measure, which is a projection valued measure defined for Borel sets of \mathbb{R} such that $E(\mathbb{R})$ is the identity operator in \mathcal{H} and $E(B \cap C) = E(B) \cap E(C)$ for Borel sets $B, C \subseteq \mathbb{R}$. For any $f \in \mathcal{H}$ the mapping $B \rightarrow \langle E(B)f, f \rangle$ is an ordinary measure defined for the Borel sets $B \subseteq \mathbb{R}$ and denoted $\langle Ef, f \rangle$.

The selfadjoint operators T_1, \dots, T_d in \mathcal{H} with spectral measure E_1, \dots, E_d , respectively, are said to be mutually commuting (mutually permutable in [13]) if their spectral measures commute, i.e.,

$$(4.1) \quad E_i(B)E_j(C) = E_i(C)E_j(B)$$

for any $i, j = 1, \dots, d$ and any two Borel sets $B, C \subseteq \mathbf{R}$. If T_1, \dots, T_d commute, then

$$E = E_1 \otimes \cdots \otimes E_d$$

is a spectral measure on \mathbf{R}^d with values that are selfadjoint projections in \mathcal{H} . In particular, E is the unique measure such that

$$E(B_1 \times \cdots \times B_d) = E_1(B_1) \cdots E_d(B_d)$$

for any Borel sets $B_1, \dots, B_d \subseteq \mathbf{R}$. The measure E is called the spectral measure of the commuting family T_1, \dots, T_d .

We shall consider only bounded operators, see Remark 4.2 below. A vector $\Phi_0 \in \mathcal{H}$ is a cyclic vector in \mathcal{H} with respect to the commuting family of bounded selfadjoint operators T_1, \dots, T_d in \mathcal{H} if the linear manifold $\{P(T_1, \dots, T_d)\Phi_0, P \in \Pi^d\}$ is dense in \mathcal{H} . We summarize the spectral theorem for T_1, \dots, T_d in the following.

Theorem 4.1. *Let \mathcal{H} be a separable Hilbert space and T_1, \dots, T_d be a commuting family of bounded selfadjoint operators in \mathcal{H} . Let S_i denote the spectrum of T_i , $1 \leq i \leq d$. If Φ_0 is a cyclic vector in \mathcal{H} with respect to T_1, \dots, T_d , then T_1, \dots, T_d are unitarily equivalent to the multiplication operators X_1, \dots, X_d ,*

$$(4.2) \quad (X_i f)(\mathbf{x}) = x_i f(\mathbf{x}), \quad 1 \leq i \leq d,$$

defined on $L^2(\mathbf{R}^d, \mu)$, where the measure μ is defined by $\mu(B) = \langle E(B)\Phi_0, \Phi_0 \rangle$ for the Borel set $B \subset \mathbf{R}^d$ with support $S \subset S_1 \times \cdots \times S_d$.

The unitary equivalence means that there exists a unitary mapping $U: \mathcal{H} \rightarrow L^2(\mathbf{R}^d, \mu)$ such that

$$(4.3) \quad UT_i U^{-1} = X_i, \quad 1 \leq i \leq d.$$

We note that the measure μ satisfies the relation

$$(4.4) \quad \langle \Phi_0, P(T_1, \dots, T_d)\Phi_0 \rangle = \int_S P(\mathbf{x}) d\mu(\mathbf{x})$$

for all polynomials $P \in \Pi^d$. Since T_1, \dots, T_d are bounded, S_1, \dots, S_d are compact. Therefore S is compact and μ is of compact support.

Remark. For bounded selfadjoint operators T_1 and T_2 , (4.1) is a sufficient and necessary condition for $T_1 T_2 = T_2 T_1$. However, for unbounded operators it has been shown by Nelson [11] that there are selfadjoint operators T_1 and T_2 with a common dense domain such that $T_1 T_2 = T_2 T_1$, but their spectral measures do not commute.

4.2. The definition of the operators. Let \mathcal{H} be a separable Hilbert space with fixed orthonormal basis $\{\psi_n\}_{n=0}^\infty$. For our purpose we shall rewrite $\{\psi_n\}_{n=0}^\infty$ as $\{\psi_n\}_{n=0}^\infty = \{\phi_j^k\}_{j=1}^{r_k} \}_{k=0}^\infty$, where $r_k = r_k^d = \binom{k+d-1}{k}$ for a fixed positive integer d as before. We introduce the formal vector notation

$$(4.5) \quad \Phi_k = [\phi_1^k, \dots, \phi_{r_k}^k]^T, \quad k \in \mathbf{N}_0.$$

For our convenience we shall say that $\{\Phi_n\}_{n=0}^\infty$ is orthonormal, formally

$$\langle \Phi_k, \Phi_m^T \rangle = (\langle \phi_i^k, \phi_j^m \rangle)_{i=1}^{r_k} {}_{j=1}^{r_m} = \delta_{ij} \delta_{km} I_{r_k}.$$

For every $f \in \mathcal{H}$ we can write in the vector-matrix notation that

$$(4.6) \quad f = \sum_{k=0}^{\infty} \mathbf{a}_k^T \Phi_k, \quad \mathbf{a}_k \in \mathbf{R}^{r_k}.$$

If $T : \mathcal{H} \rightarrow \mathcal{H}$ is a linear operator, we mean by $T\Phi_k$ the vector

$$T\Phi_k = [T\phi_1^k, \dots, T\phi_{r_k}^k]^T.$$

We denote by $F_{k,m}$ the matrix

$$F_{k,m} = (\langle T\Phi_k, \Phi_m^T \rangle) = (\langle T\phi_i^k, \phi_j^m \rangle)_{i=1}^{r_k} {}_{j=1}^{r_m}.$$

Then we can write

$$Tf = \sum_{k=0}^{\infty} \mathbf{b}_k^T \Phi_k, \quad \mathbf{b}_k^T = \sum_{j=0}^{\infty} \mathbf{a}_j^T F_{j,k}.$$

The usual matrix representation of the linear operator now takes the form $T = (F_{ij})_{i,j=1}^\infty$. The elements of T are matrices $F_{ij} : r_i \times r_j$, whose sizes are different for different pairs of (i, j) . Note that we have used the same symbol for both the operator and its matrix representation.

We now define the linear operators associated with the recurrence relation in Theorem 2.2. Let $A_{k,i} : r_k \times r_{k+1}$ and $B_{k,i} : r_k \times r_k$ be given matrices such that the rank condition (b) in Theorem 2.2 is satisfied. Furthermore, assume that $B_{k,i}$ are symmetric, and $A_{k,i}$ and $B_{k,i}$ satisfy equations (2.4), (2.5), and (2.6). We then define $T_i : \mathcal{H} \rightarrow \mathcal{H}$, $1 \leq i \leq d$, to be the linear operator whose matrix representation with respect to $\{\Phi_k\}_{k=0}^\infty$ is given by

$$(4.7) \quad T_i = \begin{bmatrix} B_{0,i} & A_{0,i} & & \bigcirc \\ A_{0,i}^T & B_{1,i} & A_{1,i} & \\ & A_{1,i}^T & B_{2,i} & \ddots \\ \bigcirc & & \ddots & \ddots \end{bmatrix}, \quad 1 \leq i \leq d.$$

We can consider T_i as matrix operator which acts on sequences in l^2 via matrix multiplication. For $d = 1$, we have $r_k = 1$ for all $k \in \mathbf{N}_0$ and the matrix T_1 is the Jacobi matrix (cf. [15]).

4.3. Properties of T_i . We restrict ourselves to the bounded operators.

Lemma 4.3. *The operator T_i is bounded if and only if $\sup_{k \geq 0} \|A_{k,i}\|_2 < \infty$ and $\sup_{k \geq 0} \|B_{k,i}\|_2 < \infty$.*

Proof. For any $f \in \mathcal{H}$, $f = \sum \mathbf{a}_k^T \Phi_k$, we have $\|f\|_{\mathcal{H}}^2 = \langle f, f \rangle = \sum \mathbf{a}_k^T \mathbf{a}_k$. It follows easily from the definition that

$$\begin{aligned} T_i f &= \sum_{k=0}^{\infty} \mathbf{a}_k^T [A_{k,i} \Phi_{k+1} + B_{k,i} \Phi_k + A_{k-1,i}^T \Phi_{k-1}] \\ &= \sum_{k=0}^{\infty} [\mathbf{a}_{k-1}^T A_{k-1,i} + \mathbf{a}_k^T B_{k,i} + \mathbf{a}_{k+1}^T A_{k+1,i}^T] \Phi_k \end{aligned}$$

where we define $A_{-1,i} = 0$. Therefore, if $\sup_{k \geq 0} \|A_{k,i}\|_2 < +\infty$ and $\sup_{k \geq 0} \|B_{k,i}\|_2 < +\infty$, then

$$\begin{aligned} \|T_i f\|_{\mathcal{H}}^2 &= \sum_{k=0}^{\infty} \|\mathbf{a}_{k-1}^T A_{k-1,i} + \mathbf{a}_k^T B_{k,i} + \mathbf{a}_{k+1}^T A_{k,i}\|_2^2 \\ &\leq 3 \left(2 \sup_{k \geq 0} \|A_{k,i}\|_2^2 + \sup_{k \geq 0} \|B_{k,i}\|_2^2 \right) \|f\|_{\mathcal{H}}^2. \end{aligned}$$

Thus, T_i is a bounded operator. Conversely, suppose that $\|A_{k,i}\|_2$, say, goes to infinity for a subsequence of \mathbf{N}_0 . Let \mathbf{a}_k be vectors such that $\|\mathbf{a}_k\|_2 = 1$ and $\|\mathbf{a}_k^T A_{k,i}\|_2 = \|A_{k,i}\|_2$. Then we have $\|\mathbf{a}_k \Phi_k\| = \|\mathbf{a}_k\|_2 = 1$. Therefore, it follows from

$$\|T_i\|_{\mathcal{H}}^2 \geq \|T_i \mathbf{a}_k^T \Phi_k\|_{\mathcal{H}}^2 = \|\mathbf{a}_k^T A_{k,i}\|_2^2 + \|\mathbf{a}_k^T B_{k,i}\|_2^2 + \|\mathbf{a}_k^T A_{k-1,i}\|_2^2 \geq \|A_{k,i}\|_2^2$$

that T_i is unbounded. \square

Lemma 4.4. Suppose T_i , $1 \leq i \leq d$, are bounded. Then T_i , $1 \leq i \leq d$, are selfadjoint operators, and T_1, \dots, T_d commute.

Proof. Since T_i is bounded, it is selfadjoint if it is symmetric, i.e., $\langle T_i f, g \rangle = \langle f, T_i g \rangle$. But this follows from the obvious symmetry of the matrix representation (4.7). To prove that T_1, \dots, T_d commute, we only need to show

$$T_k T_j f = T_j T_k f, \quad \forall f \in \mathcal{H},$$

since T_i 's are bounded. A simple calculation shows that this is equivalent to the matrix equations (2.4), (2.5), and (2.6), which are assumed. \square

Lemma 4.5. Suppose T_1, \dots, T_d are bounded operators. Then $\Phi_0 \in \mathcal{H}$ is a cyclic vector with respect to T_1, \dots, T_d , and

$$(4.8) \quad \Phi_n = \mathbb{P}_n(T_1, \dots, T_d) \Phi_0$$

where $\mathbb{P}_n(x_1, \dots, x_d)$ is of the form (2.2).

Proof. If (4.8) is true, then Φ_0 is a cyclic vector by definition. To prove (4.8), we use induction. Clearly $\mathbb{P}_0 = 1$. From the definition of T_i we have

$$T_i \Phi_0 = A_{0,i} \Phi_1 + B_{0,i} \Phi_0, \quad 1 \leq i \leq d.$$

Therefore,

$$A_{0,i} \Phi_1 = T_i \Phi_0 - B_{0,i} \Phi_0, \quad 1 \leq i \leq d.$$

Multiply this equation by $D_{0,i}$ in (2.7) and sum for $i = 1, \dots, d$, we get

$$\Phi_1 = \sum_{i=1}^d D_{0,i} T_i \Phi_0 - \sum_{i=1}^d D_{0,i} B_{0,i} \Phi_0 = \left(\sum_{i=1}^d T_i D_{0,i} - E_0 \right) \Phi_0$$

where $E_0 = \sum D_{0,i} B_{0,i}$. Therefore

$$\mathbb{P}_1(\mathbf{x}) = \sum_{i=1}^d \lambda_i D_{0,i} - E_0.$$

Since $D_{0,i}$ is of the size $r_1 \times r_0 = d \times 1$, \mathbb{P}_1 is of the desired form. Likewise, for $k \geq 1$,

$$T_i \Phi_k = A_{k,i} \Phi_{k+1} + B_{k,i} \Phi_k + A_{k-1,i}^T \Phi_{k-1}, \quad 1 \leq i \leq d,$$

therefore

$$\Phi_{k+1} = \sum_{i=1}^d T_i D_{k,i} \Phi_k - E_k \Phi_k - F_k \Phi_{k-1}$$

where

$$E_k = \sum_{i=1}^d D_{k,i} B_{k,i} \quad \text{and} \quad F_k = \sum_{i=1}^d D_{k,i} A_{k-1,i}^T.$$

By induction we then have

$$\Phi_{k+1} = \left[\sum_{k=1}^d T_i D_{k,i} \mathbb{P}_k(T) - E_k \mathbb{P}_k(T) - F_k \mathbb{P}_{k-1}(T) \right] \Phi_0,$$

where $T = (T_1, \dots, T_d)^T$. Therefore

$$\mathbb{P}_{k+1}(\mathbf{x}) = \sum_{k=1}^d \lambda_i D_{k,i} \mathbb{P}_k(\mathbf{x}) - E_k \mathbb{P}_k(\mathbf{x}) - F_k \mathbb{P}_{k-1}(\mathbf{x}).$$

Clearly, every component of \mathbb{P}_{k+1} is a polynomial in Π_{k+1} . \square

From these lemmas and Theorem 4.1, we have proved

Theorem 4.6. *If $\sup_{k \geq 0} \|A_{k,i}\|_2 < \infty$ and $\sup_{k \geq 0} \|B_{k,i}\|_2 < \infty$ for $1 \leq i \leq d$, then there exists a measure $\mu \in \mathcal{M}$ with compact support such that T_1, \dots, T_d are unitarily equivalent to the multiplication operators X_1, \dots, X_d in $L^2(\mathbf{R}^d, \mu)$.*

4.4. Proof of the main theorem. The unitary equivalence in Theorem 4.6 associates the cyclic vector Φ_0 with the function $f(\mathbf{x}) = 1$ and $(T_1^{\alpha_1} \dots T_d^{\alpha_d})\Phi_0$ with $f(\mathbf{x}) = \mathbf{x}^\alpha$. Moreover, the orthonormal basis $\{\Phi_n\}_{n=0}^\infty$ in \mathcal{H} corresponds to $\{\mathbb{P}_n\}_{n=0}^\infty$ in $L^2(\mathbf{R}^d, \mu)$ as shown in (4.8). We have

Lemma 4.7. *The polynomials $\{\mathbb{P}_n\}_{n=0}^\infty$ in Lemma 4.5 are orthonormal with respect to μ , and they satisfy the recurrence relation (a) in Theorem 2.2 with the matrices in (4.7) as the coefficients.*

Proof. Since $\mu(B) = \langle E(B)\Phi_0, \Phi_0 \rangle$ in Theorem 4.1, we have by (4.4) that

$$\int \mathbb{P}_n(\mathbf{x}) \mathbb{P}_m^T(\mathbf{x}) d\mu(\mathbf{x}) = \langle \mathbf{P}_n \Phi_0, \mathbb{P}_m^T \Phi_0 \rangle = \langle \Phi_n, \Phi_m^T \rangle.$$

This proves that $\{\mathbb{P}_n\}$ are orthonormal. From Theorem 2.2 the polynomials $\{\mathbb{P}_n\}$ satisfy a recurrence relation of the form (a) in that theorem. By the unitary equivalence in Theorem 4.6, the multiplication operators X_1, \dots, X_d at (4.2) have the same matrix representation (4.7) with respect to $\{\mathbb{P}_n\}$ in $L^2(\mathbf{R}^d, \mu)$. Since $X_i \mathbb{P}_n = x_i \mathbb{P}_n$, the coefficient matrices in the recurrence relation that $\{\mathbb{P}_n\}$ satisfy are the same matrices in (4.7). \square

Proof of Theorem 3.1. (i) \Rightarrow (ii). From Theorem 2.2, we only need to prove (3.2). However, since μ has compact support, the multiplication operators X_1, \dots, X_d in $L^2(\mathbf{R}^d, \mu)$ are bounded. Since X_1, \dots, X_d have the matrix representation (4.7) with respect to $\{\mathbb{P}_n\}$, (3.2) follows from Lemma 4.3.

(ii) \Rightarrow (i). If $\{\mathbb{P}_n\}$ satisfies the recurrence relation in the theorem, then we can use the coefficient matrices to define operators T_i through (4.7). By Lemma 4.7, the polynomials in Lemma 4.5 satisfy the same recurrence relation. Since

the recurrence relation determinates uniquely the polynomials under the consideration, the $\{\mathbb{P}_n\}$ are the polynomials in Lemma 4.5. Therefore, the existence of the measure μ with compact support has been established by our previous results. That μ is determinate follows from the boundedness of the multiplication operators and [3, Theorem 4]. \square

ACKNOWLEDGMENT

The author thanks one referee for clarifying one inaccurate point in the manuscript.

REFERENCES

1. N. I. Akhiezer and I. M. Glazman, *Theory of linear operators in Hilbert space*, Ungar, New York, 1961.
2. S. K. Berberian, *Notes on spectral theory*, Van Nostrand, Princeton, N.J., 1966.
3. C. Berg, *The multidimensional moment problem and semigroups*, Moments in Mathematics, Proc. Sympos. Appl. Math., Vol. 37, Amer. Math. Soc., Providence, R.I., 1987, pp. 110–124.
4. M. Bertran, *Notes on orthogonal polynomials in ν -variables*, SIAM J. Math. Anal. **6** (1975), 250–257.
5. T. S. Chihara, *An introduction to orthogonal polynomials*, Math. and its Appl., Vol. 13, Gordon and Breach, New York, 1978.
6. J. Dombrowski, *Orthogonal polynomials and functional analysis*, Orthogonal Polynomials (P. Nevai, ed.), Kluwer Academic, Amsterdam, 1990, pp. 147–161.
7. ———, *Tridiagonal matrix representations of cyclic selfadjoint operators*, I, Pacific J. Math. **114** (1984), 325–334; II **120** (1985), 47–53.
8. B. Fuglede, *The multidimensional moment problem*, Math. Expositions **1** (1983), 47–65.
9. M. A. Kowalski, *Orthogonality and recursion formulas for polynomials in n variables*, SIAM J. Math. Anal. **13** (1982), 316–323.
10. H. L. Krall and I. M. Sheffer, *Orthogonal polynomials in two variables*, Ann. Mat. Pura Appl. (4) **76** (1967), 325–376.
11. E. Nelson, *Analytic vectors*, Ann. of Math. (2) **70** (1959), 572–615.
12. E. Prugovecki, *Quantum mechanics in Hilbert space*, 2nd ed., New York, 1981.
13. F. Riesz and B. Sz.-Nagy, *Functional analysis*, Ungar, New York, 1955.
14. B. D. Sleeman, *Multiparameter spectral theory in Hilbert Spaces*, Pitman, London, 1978.
15. M. Stone, *Linear transformations in Hilbert space and their applications to analysis*, Amer. Math. Soc. Colloq. Publ., Vol. 15, Amer. Math. Soc., Providence, R.I., 1932; reprinted 1983.
16. G. Szegő, *Orthogonal polynomials*, Amer. Math. Soc. Colloq. Publ., Vol. 23, Amer. Math. Soc., Providence, R.I., 4th ed., 1975.
17. Y. Xu, *On multivariate orthogonal polynomials*, SIAM J. Math. Anal. **24** (1993), 783–794.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF ARKANSAS AT LITTLE ROCK,
LITTLE ROCK, ARKANSAS 72204

Current address: Department of Mathematics, University of Oregon, Eugene, Oregon 97403

E-mail address: yuan@bright.uoregon.edu