

ON CR -MAPPINGS BETWEEN ALGEBRAIC CAUCHY-RIEMANN MANIFOLDS AND SEPARATE ALGEBRAICITY FOR HOLOMORPHIC FUNCTIONS

RUSLAN SHARIPOV AND ALEXANDER SUKHOV

ABSTRACT. We prove the algebraicity of smooth CR -mappings between algebraic Cauchy-Riemann manifolds. A generalization of separate algebraicity principle is established.

1. INTRODUCTION

One of the most surprising phenomena of geometric theory of several complex variables is the following one: every local biholomorphism of the (real) unit sphere of \mathbb{C}^n , $n > 1$, extends to a global automorphism of the unit ball and, particularly, to a complex rational mapping of the whole space \mathbb{C}^n (of course, this is not true in one variable). Poincaré [Po] was the first who obtained such result for $n = 2$; later the general case was considered independently by Tanaka [Ta], Pelles [Pe] and Alexander [Al]. We mention also a quite elementary and very elegant proof of the result of this type (in fact, more general) in Rudin's book [Ru].

Now there exist several generalizations of this classical result in different directions. Thus, Tumanov and Henkin [TH], Tumanov [Tu], Forstneric [Fo] and Sukhov [Su1, Su3] investigated the rational extendability of (locally defined) CR -mappings between quadric Cauchy - Riemann manifolds of higher codimensions. On the other hand, for the case of hypersurfaces Webster in his famous paper [W1] discovered a very natural extension of the Poincaré phenomenon in the algebraic category: a local biholomorphism between two real algebraic Levi - nondegenerate hypersurfaces extends to an algebraic mapping on all \mathbb{C}^n . In the present paper we generalize this result to CR -mappings between real algebraic Cauchy - Riemann manifolds of higher codimensions (Theorem 1) (another generalization was obtained in [Su4]). Our technique is based on the Webster reflection principle modified in the spirit of [Su1] and a generalization of the classical separate algebraicity theorem, where we replace the parallel lines by families of algebraic curves (Theorem 2). We hope that this second result is of self-interest.

The paper is organized as follows. In section 2 we give the precise definitions and statements of our results. In sections 3 and 4 we prove Theorem 1 provided Theorem 2 holds. Section 5 is devoted to the proof of Theorem 2.

Received by the editors January 26, 1995.

1991 *Mathematics Subject Classification*. Primary 32D15, 32D99, 32F25, 32H99.

Key words and phrases. CR -mapping, reflection principle, algebraic CR -manifold, separate algebraicity.

Paper written under the financial support of the International Scientific Foundation of Soros, project #RK4000

2. THE RESULTS

Let Ω be a domain in \mathbb{C}^n . A closed subset M of Ω is called a generic real algebraic manifold of codimension $d \geq 1$ if

$$(2.1) \quad M = \{z \in \Omega : \rho_j(z, \bar{z}) = 0, j = 1, \dots, d\},$$

where ρ_j are real polynomials and $\bar{\partial}\rho_1 \wedge \dots \wedge \bar{\partial}\rho_d \neq 0$ in Ω . They are called the defining functions of M . We denote by $T_p M$ and $T_p^c M$ the real and complex tangent spaces of M at a point $p \in M$ (recall that $T_p^c M = T_p M \cap J(T_p M)$, where J is the complex structure operator in \mathbb{C}^n). This is well known that for a generic manifold the complex dimension of $T_p^c M$ does not depend on $p \in M$ and is equal to $n - d$; it is called the CR -dimension of M .

We denote by $H_p(\rho_j, u, v)$ the value of the Levi form (complex hessian) of the function ρ_j on vectors u, v at the point $p \in M$, i.e.

$$(2.2) \quad H_p(\rho_j, u, v) = \sum_{\nu, \mu=1}^n \frac{\partial^2 \rho_j}{\partial z_\nu \partial \bar{z}_\mu}(p) u_\nu \bar{v}_\mu.$$

The Levi cone (at $p \in M$) of the manifold M of the form (2.1) is said to be the convex hull of the set

$$(2.3) \quad \{\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{R}^d : \alpha_j = H_p(\rho_j, u, u), u \in T_p^c M\}.$$

If the Levi cone of M has a non-empty interior in \mathbb{R}^d , then we say that M possess a non-degenerate Levi cone at p . Evidently this condition does not depend on the choice of defining functions and is invariant with respect to changes of coordinates.

The vector valued hermitian form

$$L_p(u, v) = (H_p(\rho_1, u, v), \dots, H_p(\rho_d, u, v))$$

is called the Levi form of M at p .

Along with M we consider a domain Ω' in $\mathbb{C}^{n'}$ and a generic real algebraic manifold M' of codimension $d' \geq 1$ of the form

$$(2.4) \quad M' = \{z' \in \Omega' : \rho'_j(z', \bar{z}') = 0, j = 1, \dots, d'\},$$

where ρ'_j are real polynomials and $\bar{\partial}\rho'_1 \wedge \dots \wedge \bar{\partial}\rho'_{d'} \neq 0$ in Ω' . Fix a hermitian scalar product in $\mathbb{C}^{n'}$ that is denoted by $\langle \cdot, \cdot \rangle'$. Fix a point $p' \in M'$. With every defining function ρ'_j we associate the Levi operator $L_{p'}^j : T_{p'}^c M' \longrightarrow T_{p'}^c M'$ determined by $H_{p'}(\rho'_j, u, v) = \langle L_{p'}^j(u), v \rangle'$ for all $u, v \in T_{p'}^c M'$. Of course, this definition of the Levi operator depends on the choice of hermitian scalar product. But as we shall see further this dependence is inessential. In what follows we use the similar notation for the Levi operators of M (without primes).

Let U be an open connected subset of the manifold M . The mapping $F : U \longrightarrow M'$ of the smoothness class C^1 is called a CR -mapping if for any point $p \in U$ the tangent map dF_p is \mathbb{C} -linear after restriction to the complex tangent space $T_p^c M$ (in this case $dF_p(T_p^c M) \subset T_{p'}^c M'$). We say that F extends to an algebraic mapping of all \mathbb{C}^n if the graph of F is a part of n -dimensional complex algebraic manifold in $\mathbb{C}^{n+n'}$.

Now we can formulate the first main result of our paper. It is given by the following theorem.

Theorem 1. *Let $\Omega \in \mathbb{C}^n$ and $\Omega' \in \mathbb{C}^{n'}$ be domains, $M \subset \Omega$ and $M' \subset \Omega'$ be generic real algebraic manifolds of the form (2.1) and (2.4) respectively, M having the non-degenerate Levi cone at some point $p \in M$. Suppose $U \subset M$ is an open connected subset of M containing p and let $F : U \longrightarrow M'$ be a CR-mapping of class C^1 satisfying the following condition*

$$(2.5) \quad \sum_{j=1}^{d'} L_{p'}^j(dF_p(T_p^c M)) = T_{p'}^c M', \text{ where } p' = F(p).$$

Then F extends to an algebraic mapping of the whole space \mathbb{C}^n .

We note first that (2.5) does not depend on the choice of the hermitian scalar product in $\mathbb{C}^{n'}$ by means of which we defined the Levi operators $L_{p'}^j$. If $\tilde{L}_{p'}^j$ is defined by another scalar product, then it is connected with $L_{p'}^j$ by the equality $\tilde{L}_{p'}^j = A L_{p'}^j$, where A is a non-degenerate \mathbb{C} -linear operator in $T_{p'}^c M'$. Therefore (2.5) holds for operators $\tilde{L}_{p'}^j$ as well.

Recall that the Levi form of M is called non-degenerate if $L_p(u, v) = 0$ for any $v \in T_v^c M$ implies $u = 0$ [W2].

Corollary. *Let $F : M \longrightarrow M'$ be a CR diffeomorphism of class C^1 between two real algebraic manifolds in \mathbb{C}^n with non-degenerate Levi forms and non-degenerate Levi cones. Then F extends to an algebraic mapping on all \mathbb{C}^n .*

This assertion follows from Theorem 1 quite similarly to [Su1]. We emphasize that Theorem 1 treats a considerably more general situation, since M and M' are allowed to have different CR dimensions. Our proof of Theorem 1 is based on the modification of the Webster reflection principle [W1]. The crucial technical tool here is a “curved” version of the following classical separate algebraicity principle.

Claim. Let $f(\mathbf{z}) = f(z_1, \dots, z_n)$ be a function in some domain $D \subset \mathbb{C}^n$. If $f(\mathbf{z})$ is algebraic in each separate variable z_i for any fixed values of other variables, then $f(\mathbf{z})$ is an algebraic function in D .

Proof of this classical theorem can be found in [BM]. Function $f(\mathbf{z})$ in this assertion is algebraic along the straight lines parallel to the coordinate axes. The domain D foliates into n families of such lines. In order to generalize this result for our purposes we introduce n families of algebraic curves in D

[illegible]

Here $m = 1, \dots, n$ is the number of the family, t_m is a parameter on each particular curve of the m -th family ($R_i^{(m)}$ depends on t_m algebraically). Parameters $c_1^{(m)}, \dots, c_{n-1}^{(m)}$ identify curves of the m -th family.

Definition 1. The family of algebraic curves (2.6) is called *algebraically depending on parameters* if each of the defining functions $R_i^{(m)}$, $i = 1, \dots, n$, in (2.6) is algebraic in $c_1^{(m)}, \dots, c_{n-1}^{(m)}$.

In this case because of the above-mentioned classical separate algebraicity principle functions $R_i^{(m)}$ in (2.6) are algebraic in the whole set of their arguments $t_m, c_1^{(m)}, \dots, c_{n-1}^{(m)}$.

Definition 2. The family of curves (2.6) is called nonsingular in D if the curves of this family fill the whole domain D and the mapping $R^{(m)} : (t_m, c_1^{(m)}, \dots, c_{n-1}^{(m)}) \rightarrow \mathbf{z}$ is a local diffeomorphism.

Definition 3. Let (2.6) define n nonsingular families of algebraic curves in D . Then we have n tangent vectors to the curves at each point of D . We shall say that n families of curves (2.6) are in general position if these vectors are linearly independent at any point $\mathbf{z} \in D$.

Now we can state the “curved” separate algebraicity principle which partially generalizes the classical one and forms the second main result of our paper.

Theorem 2. Let $D \subset \mathbb{C}^n$ be a domain equipped with n families (2.6) of nonsingular algebraic curves algebraically depending on parameters and being in general position. Then each holomorphic function $f(\mathbf{z})$ in D , which is algebraic in t_m after restriction to any particular curve from any one of these families, extends to an algebraic function on \mathbb{C}^n .

3. TANGENT CR -FIELDS AND THE MAIN EQUATIONS

First we shall recall briefly some facts of the theory of CR -structures (reader can find more details in [Ch]). Let $(T_p^c M)_{\mathbb{R}}$ be the complex tangent space $T_p^c M$ considered as a vector space over real numbers \mathbb{R} . Then $(T_p^c M)_{\mathbb{R}} \otimes^{\mathbb{R}} \mathbb{C}$ is a complexification for $(T_p^c M)_{\mathbb{R}}$. The operator J of canonical complex structure in this complexification is \mathbb{C} -linear with respect to the own complex structure of $T_p^c M$. Therefore it has two eigenvalues $+i$ and $-i$. Then we have the decomposition $(T_p^c M)_{\mathbb{R}} \otimes \mathbb{C} = T_p^c M^{1,0} \oplus T_p^c M^{0,1}$, where

$$T_p^c M^{1,0} = \{(v, -iv) : v \in T_p^c M\} \text{ and } T_p^c M^{0,1} = \{(v, +iv) : v \in T_p^c M\}$$

are the eigenspaces for J corresponding to the eigenvalues $+i$ and $-i$ respectively. The following two maps $v \mapsto (v, -iv)$ and $v \mapsto (v, +iv)$ realize the canonical isomorphisms $T_p^c M \equiv T_p^c M^{1,0}$ and $\overline{T_p^c M} \equiv T_p^c M^{0,1}$. Thus we have $T_p^c M_{\mathbb{R}} \otimes \mathbb{C} = T_p^c M \oplus \overline{T_p^c M}$. Because of the last decomposition the sections of the vector bundles $T^c M$ and $\overline{T^c M}$ are called the CR vector fields of the types $(1, 0)$ and $(0, 1)$ respectively. It is easy to check that the field $V = \sum_{j=1}^n v_j(z) \partial / \partial z_j$ in \mathbb{C}^n is the vector field of the type $(1, 0)$ on M if and only if the vector $(v_1(a), \dots, v_n(a))$ is in $T_a^c M$ for any $a \in M$.

Let us go back to Theorem 1. Note that without loss of generality we can take $p = 0$ and $F(p) = 0$. Taking $z = (x, y)$ and $z' = (x', y')$ and denoting $\mathbb{C}^n = \mathbb{C}_x^k \times \mathbb{C}_y^d$ and $\mathbb{C}^{n'} = \mathbb{C}_{x'}^{k'} \times \mathbb{C}_{y'}^{d'}$ one can bring the defining polynomials for M and M' to the form

$$(3.1) \quad \begin{aligned} p_j &= y_j + \bar{y}_j + o(|z|), & j &= 1, \dots, d, \\ p'_j &= y'_j + \bar{y}'_j + o(|z|), & j &= 1, \dots, d'. \end{aligned}$$

For this choice of coordinates we have $T_0^c M = \mathbb{C}_x^k = \{(x, y) : y = 0\}$ and $T_0^c M' = \mathbb{C}_{x'}^{k'} = \{(x', y') : y' = 0\}$. Now let us consider vector fields T_q , $q = 1, \dots, k$, of the form

$$(3.2) \quad T_q = \Delta(z, \bar{z}) \frac{\partial}{\partial x_q} - \sum_{j=1}^d a_{jq}(z, \bar{z}) \frac{\partial}{\partial y_j},$$

where Δ is the determinant of the matrix

$$(3.3) \quad \Delta_{sj} = \left(\frac{\partial \rho_s}{\partial y_j} \right)_{s=1, \dots, d}^{j=1, \dots, d}.$$

Everywhere in this paper we shall obey the following rule for denoting the matrix elements: lower index outside the right bracket is the row number and upper index is the column number. The coefficients a_{jq} in (3.4) we define as follows

$$(3.4) \quad a_{jq} = \sum_{s=1}^d \Delta b_{js} \frac{\partial \rho_s}{\partial x_q},$$

where b_{js} is the inverse of the matrix (3.3). According to elementary facts from linear algebra the matrix with the elements Δb_{js} is a conjugate matrix for (3.3), i.e. its elements are the algebraic cofactors for the elements of the transpose of (3.3). Therefore the coefficients of the vector fields in (3.2) are polynomials in x_i and y_i .

Clearly, the restrictions of the fields T_q , $q = 1, \dots, k$, form a base of the bundle $T^c M$ over a neighborhood of the origin in M . Moreover, it is obvious that

$$(3.5) \quad \Delta(0) = 1, \quad a_{jq}(0) = 0, \quad j = 1, \dots, d, \quad q = 1, \dots, k.$$

Note that because of $dF_0(T_0^c M) \subset T_0^c M'$, we get by (3.1)

$$(3.6) \quad \partial F_j / \partial x_q(0) = 0, \quad j = k' + 1, \dots, n', \quad q = 1, \dots, k.$$

Together with the fields (3.2) let us consider the conjugate fields $\overline{T_q}$. Recall that a C^1 -function h defined on an open connected subset $U \subset M$ is called a CR -function if for any $z = (x, y) \in U$ one has

$$(3.7) \quad \overline{T_q} h = \Delta \frac{\partial h}{\partial \bar{x}_q} - \sum_{j=1}^d \bar{a}_{jq} \frac{\partial h}{\partial \bar{y}_j} = 0, \quad q = 1, \dots, k.$$

These are the tangent Cauchy-Riemann equations. It is well known that a C^1 -mapping $F : M \rightarrow M'$ is a CR -mapping (in the sense of section 2) if and only if each its component is a CR -function on M . Indeed, (3.7) means that $\bar{\partial} h(V^q) = 0$, $q = 1, \dots, k$, (where $\{V^q\}_{q=1}^k$ is a base of $T_p^c M$) for any point $p \in M$. Therefore $\bar{\partial} h|_{T_p^c M} = 0$ and (3.7) means that the restriction $dh|_{T_p^c M}$ is a \mathbb{C} -linear function for any $p \in M$.

According to the Boggess-Polking theorem [BP] from the non-degeneracy of the Levi cone of M at the origin $0 \in M$ we derive that the mapping F extends holomorphically to a wedge with the edge M . Using the results of [Su1, Su2] we obtain that F extends holomorphically to a neighborhood of the origin in \mathbb{C}^n . Therefore everywhere below we take F being holomorphic in a neighborhood $\Omega \ni 0$ in \mathbb{C}^n and suppose $U = M \cap \Omega$.

The condition $F(U) \subset M'$ means that $\rho'_j(F, \bar{F}) = 0$ for $z \in U$ and $j = 1, \dots, d'$. Applying the tangent operators (3.2) to both sides of these equalities we obtain

$$(3.8) \quad T_q \rho'_j(F, \bar{F}) = 0, \quad q = 1, \dots, k, \quad j = 1, \dots, d' \text{ for } z \in U.$$

Now we introduce the vector-function $D(F)$ holomorphic in Ω

$$D(F) = \left(\frac{\partial F_1}{\partial z_1}, \dots, \frac{\partial F_{n'}}{\partial z_1}, \dots, \frac{\partial F_1}{\partial z_n}, \dots, \frac{\partial F_{n'}}{\partial z_n} \right).$$

Its values are in $\mathbb{C}^{nn'}$.

The left-hand sides of (3.8) can be considered as polynomials in z, \bar{z}, F, \bar{F} and $\partial F_j / \partial z_s, \partial \bar{F}_j / \partial \bar{z}_s$. But since F is holomorphic, we have $\partial \bar{F} / \partial z_s = 0$. Thus, the following expressions

$$(3.9) \quad \Phi_{qj}(z, \bar{z}, F, \bar{F}, D(F)) = T_q \rho'_j(F, \bar{F})$$

are polynomials in $z, \bar{z}, F, \bar{F}, D(F)$. By $\tilde{0}$ we denote the point $(0, 0, 0, 0, D(F)(0))$ in $\mathbb{C}^n \times \mathbb{C}^n \times \mathbb{C}^{n'} \times \mathbb{C}^{n'} \times \mathbb{C}^{nn'}$.

Lemma 3.1. *In the set of functions (3.9) with $q = 1, \dots, k, j = 1, \dots, d'$ one can choose a subset $\Phi_1 = \Phi_{q(1)j(1)}, \dots, \Phi_{k'} = \Phi_{q(k')j(k')}$ such that the following Jacobian matrix is of rank k'*

$$(3.10) \quad \left(\frac{\partial \Phi_j}{\partial \bar{F}_s}(\tilde{0}) \right)_{j=1, \dots, k'}^{s=1, \dots, k'}.$$

Proof. Let us fix j , consider the set of functions Φ_{qj} , $q = 1, \dots, k$, and form the Jacobian matrix

$$\left(\frac{\partial \Phi_{qj}}{\partial \bar{F}_s}(\tilde{0}) \right)_{j=1, \dots, k'}^{s=1, \dots, k}.$$

It follows from (3.1), (3.2), (3.5), (3.6), (3.8), (3.9) that this matrix can be rewritten in the form

$$(3.11) \quad \left(\sum_{r=1}^{k'} \frac{\partial^2 \rho'_j}{\partial z'_r \partial \bar{z}'_s}(0) \frac{\partial F_r}{\partial z_q}(0) \right)_{j=1, \dots, k'}^{s=1, \dots, k}.$$

The condition (2.5) does not depend on the choice of scalar product in $\mathbb{C}^{n'}$. Therefore one can take the hermitian scalar product, which defines Levi operators

L_0^j , being canonical in the coordinates (3.1). Then each operator L_0^j in the standard basis e'_r , $r = 1, \dots, k'$, of $\mathbb{C}^{k'} = T_0^c M'$ has the matrix of the following form

$$\left(\frac{\partial^2 \rho'_j}{\partial z'_j \partial \bar{z}'_s}(0) \right)_{j=1, \dots, k'}^{s=1, \dots, k'}.$$

Therefore the q -th row of the matrix (3.11) consists of the coordinates of the vector $L_0^j(dF_0(e_q))$, where e_q , $q = 1, \dots, k$, is the standard basis of $\mathbb{C}^{k'} = T_0^c M'$, i.e.

$${}^t(\partial \Phi_{qj} / \partial \bar{F}_1(\tilde{0}), \dots, \partial \Phi_{qj} / \partial \bar{F}_k(\tilde{0})) = L_0^j(dF_0(e_q))$$

where ${}^t(\cdot)$ denotes the transpose. According to (2.5) the rank of the set of vectors $L_0^j(dF_0(e_q))$, $j = 1, \dots, d'$, $q = 1, \dots, k$, is equal to k . Thus we get the proof of the lemma. \square

Let Φ_j , $j = 1, \dots, k'$, be the functions chosen according to Lemma 3.1. For $z \in U$ we have

$$(3.12) \quad \begin{aligned} \Phi_j(z, \bar{z}, F, \bar{F}, D(F)) &= 0, & j &= 1, \dots, k', \\ \rho'_s(F, \bar{F}) &= 0, & s &= 1, \dots, d'. \end{aligned}$$

Lemma 3.2. *The rank of the Jacobian matrix for (3.12) with respect to \bar{F} at the point $\tilde{0}$ is equal to n' .*

Proof. The Jacobian matrix mentioned in the lemma has the form

$$(3.13) \quad \left(\begin{array}{c|c} \frac{\partial \Phi_j(\tilde{0})}{\partial F_s} & * \\ \hline \tilde{0}_{d'}^{k'} & I_{d'} \end{array} \right).$$

The upper left block of the matrix (3.13) is formed by the matrix (3.10), $\mathbf{0}$ is the zero $d' \times k'$ matrix and I is a unit $d' \times d'$ matrix. Therefore by Lemma 3.1 the rank of (3.13) is equal to $k' + d' = n'$. \square

Applying the complex conjugation to the first k' equations (3.13), we get

$$(3.14) \quad \begin{aligned} P_j(z, \bar{z}, F, \bar{F}, \overline{D(F)}) &= 0, & j &= 1, \dots, k', \\ \rho'_s(F, \bar{F}) &= 0, & s &= 1, \dots, d', \end{aligned}$$

where P_j are polynomials in $z, \bar{z}, F, \bar{F}, \overline{D(F)}$. This system of equations is of crucial importance in what follows.

4. GEOMETRY OF SEGRE SURFACES

For a real algebraic manifold M of the form (2.1) the Segre surface associated to a fixed point $z \in \mathbb{C}^n$ is a complex algebraic set in \mathbb{C}^n of the form

$$(4.1) \quad Q(z) = \{w \in \mathbb{C}^n : \rho_j(w, \bar{z}) = 0, j = 1, \dots, d\}.$$

We denote by A the graph of the mapping F over a neighborhood $\Omega \ni 0$ in \mathbb{C}^n . Also let

$$(4.2) \quad A_\zeta = \{(z, z') \in \mathbb{C}^{n+n'} : z' = F(z), \rho_j(z, \bar{\zeta}) = 0, j = 1, \dots, d\}$$

denote the graph of the restriction of F to the Segre surface $Q(\zeta)$. Evidently, every A_ζ is a $k(n-d)$ -dimensional complex manifold in $\Omega \times \mathbb{C}^{n'}$.

Lemma 4.1. *For any point $\zeta \in \Omega$ the complex manifold A_ζ is a piece of a complex k -dimensional algebraic variety \tilde{A}_ζ in $\mathbb{C}^{n+n'}$.*

Proof. It follows from Lemma 3.2 that one can apply the implicit function theorem to the system (3.14). We get $F(z) = R(z, \bar{z})$ for $z \in M \cap \Omega$ (where R is a real analytic function in Ω algebraic in z). By (3.1) and the implicit function theorem we get $M \cap \Omega = \{z = (x, y) \in \Omega : y = \phi(x, \bar{z})\}$. Therefore,

$$(4.3) \quad F(x, \phi(x, \bar{z})) = R(x, \phi(x, \bar{z}), \bar{z}),$$

for $z = (x, y) \in M \cap \Omega$. Consider antiholomorphic functions $F^*(\theta, \xi) = F(\bar{\theta}, \phi(\bar{\theta}, \bar{\xi}))$ and $R^*(\theta, \xi) = R(\bar{\theta}, \phi(\bar{\theta}, \bar{\xi}), \bar{\xi})$, where $\theta \in \mathbb{C}^k$, $\xi \in \mathbb{C}^n$. Then (4.3) means that these functions coincide on the manifold

$$\hat{M} = \{(\theta, \xi) : \bar{\theta} = (\xi_1, \dots, \xi_k), \xi \in M\},$$

which is obviously generic in a neighborhood of the origin in \mathbb{C}^{n+k} . Now it follows from the uniqueness theorem [Pi] that $F(\theta, \phi(\theta, \bar{\xi})) = R(\theta, \phi(\theta, \bar{\xi}), \bar{\xi})$ in a neighborhood of the origin in \mathbb{C}^{n+k} . Hence,

$$F(x, \phi(x, \bar{\zeta})) = R(x, \phi(x, \bar{\zeta}), \bar{\zeta})$$

for any fixed ζ in a neighborhood of the origin in \mathbb{C}^n . But the set

$$\{(x, \phi(x, \bar{\zeta})) : x \text{ runs over a neighborhood of the origin}\}$$

coincides with the Segre surface $Q(\zeta) = \{z : \rho_j(z, \bar{\zeta}) = 0, j = 1, \dots, d\}$. Thus we get $F(z) = R(z, \bar{\zeta})$ for $z \in Q(\zeta)$. Since R was obtained by (3.14), the set

$$A_\zeta = \{(z, z') : z \in Q(\zeta), z' = F(z)\}$$

is contained in an $(n - d)$ -dimensional complex algebraic variety of the form

$$(4.5) \quad \begin{aligned} P_j(z, \bar{\zeta}, z', \overline{F(\zeta)}, \overline{DF(\zeta)}) &= 0, & j &= 1, \dots, k', \\ \rho'_s(z', \overline{F(\zeta)}) &= 0, & s &= 1, \dots, d', \\ \rho_l(z, \bar{\zeta}) &= 0, & l &= 1, \dots, d, \end{aligned}$$

proves the desired assertion. \square

Fix $\theta \in \mathbb{C}^k$ and consider the d -parametric family of Segre surfaces $Q(\theta, \tau)$, $\tau \in \mathbb{C}^d$.

Lemma 4.2. *There exists a neighborhood $U \ni 0$ in \mathbb{C}^n of the form $U = U_x \times U_y$, $U_x \subset \mathbb{C}^k$, $U_y \subset \mathbb{C}^d$ such that for any fixed $\theta \in U_x$, the family of Segre surfaces $Q(\theta, \tau)$, $\tau \in U_y$, has the following properties:*

- (1) *for any $\tau', \tau'' \in U_y$ the intersection $Q(\theta, \tau') \cap Q(\theta, \tau'')$ is empty;*
- (2) *for any $z = (x, y) \in U$ there exists a unique $\tau \in U_y$ such that $(x, y) \in Q(\theta, \tau)$.*

Proof. One can represent the Segre surface as $Q(\theta, \tau) = \{z \in U : \bar{\tau} = S(z, \bar{\theta})\}$, where S is an analytic function and U is a neighborhood of the origin. Now, if $z = (x, y)$ is in $Q(\theta, \tau') \cap Q(\theta, \tau'')$, then $\bar{\tau}' = S(z, \bar{\theta}) = \bar{\tau}''$; this implies (1). For $z = (x, y)$ we set $\tau = \bar{S}(z, \bar{\theta})$. Then $z \in Q(\theta, \tau)$ and we get (2). \square

By the implicit function theorem

$$Q(\theta, \tau) \cap U = \{(x, y) \in U : y = R(x, \bar{\theta}, \bar{\tau})\},$$

where R is an algebraic function, i.e. locally $Q(\theta, \tau)$ is the graph over the coordinate plane $\mathbb{C}_x^k = \mathbb{C}_{z_1 \dots z_k}^k$. Let $X_{(\theta, \tau)}^j$ be holomorphic vector fields on $Q(\theta, \tau)$, the natural liftings to $Q(\theta, \tau)$ of the coordinate vector fields $\partial/\partial z_j$, $j = 1, \dots, k$ in $\mathbb{C}_{z_1 \dots z_k}^k$. It follows from Lemma 4.2 that for any point $(x, y) \in U$ there exists the unique surface $Q(\theta, \tau)$, $\tau = \bar{S}(z, \bar{\theta})$ passing through (x, y) . Hence, one can consider the holomorphic vector fields $Y_{(\theta)}^j(z)$ (depending on the parameter θ) in U defined as follows: given z in U and fixed θ we set $\tau = \bar{S}(z, \bar{\theta})$ and $Y_{(\theta)}^j(z) = X_{(\theta, \tau)}^j(z)$. Their integral curves evidently are linear sections of the Segre surfaces by parallel planes and, therefore, form families of complex algebraic curves in \mathbb{C}^n algebraically depending on parameters.

Lemma 4.3. *The set of vectors $Y_{(\theta)}^j(0)$, $j = 1, \dots, k$, θ runs over a neighborhood of the origin in \mathbb{C}^k , spans \mathbb{C}^n .*

Proof. If $(\theta, \tau) \in Q(0)$, i.e. $\rho_j(\theta, \tau, 0, 0) = 0$, $j = 1, \dots, d$, then it follows from (3.1) and the implicit function theorem that $\tau = o(|\theta|)$. Now let $0 \in Q(\theta, \tau)$ (recall that this is equivalent to $(\theta, \tau) \in Q(0)$). By the implicit function theorem we get

$$Q(\theta, \tau) = \{(x, y) : y + \bar{\tau} = \phi(x, \bar{\theta}, y, \bar{\tau})\} = \{(x, y) : y + \bar{\tau} = \psi(x, \bar{\theta}, \bar{\tau})\}.$$

Since $\tau = o(|\theta|)$, we have

$$\psi = \langle L(x), \theta + o(|\theta|) \rangle + o(|\theta|) + o(|x|),$$

where

$$\langle L(\xi), \eta \rangle = (\langle L_1(\xi), \eta \rangle, \dots, \langle L_d(\xi), \eta \rangle)$$

is the Levi form of M , L_j being the Levi operators of M at the origin. Hence

$$Y_{(\theta)}^j(0) = (e_j | \langle L(e_j), \theta \rangle + o(|\theta|)) = (0, \dots, 1, \dots, 0, \langle L(e_j), \theta \rangle + o(|\theta|)),$$

where 1 is on the j -th position and $e_j, j = 1, \dots, k$, is the standard basis of \mathbb{C}^k .

Assume there exists $\alpha \in \mathbb{C}^n \setminus \{0\}$ such that

$$\langle \alpha, Y_{(\theta)}^j(0) \rangle = 0, \quad j = 1, \dots, k,$$

for any $\theta \in U_x$. Then

$$\begin{aligned} \langle \alpha, Y_{(\theta)}^j(0) \rangle &= \alpha_j + \sum_{\nu=1}^d (\langle L_\nu(e_j), \theta \rangle + o(|\theta|)) \alpha_{k+\nu} \\ &= \alpha_j + \sum_{\nu=1}^d \alpha_{k+\nu} \langle L_\nu(e_j), \theta \rangle + o(|\theta|) \\ &= \alpha_j + \left\langle \sum_{\nu=1}^d \alpha_{k+\nu} L_\nu(e_j), \theta \right\rangle + o(|\theta|) \equiv 0 \end{aligned}$$

are algebraic in t and holomorphic in other arguments. In order to prove Theorem 2 it suffices to show the algebraicity of these functions in all their arguments. We shall proceed by induction on i (the number of the function f_i in (5.4)). However, first we need some preliminaries.

Let $\alpha = (\alpha_1, \dots, \alpha_n)$ be an integer multi-index and $|\alpha| = \alpha_1 + \dots + \alpha_n$. We denote by $f_\alpha(z)$ the derivative

$$(5.5) \quad f_\alpha = \frac{\partial^{|\alpha|} f}{\partial z_1^{\alpha_1} \dots \partial z_n^{\alpha_n}}.$$

Lemma 5.1. *Under the assumptions of Theorem 2 one can find a smaller subdomain $D' \subset D$ such that all derivatives $f_\alpha(z)$ are algebraic in t_m after restriction to each curve of any family.*

Proof. The family of curves (2.6) is non-singular, therefore the transformation (2.6) from z_1, \dots, z_n to $t_m, c_1^{(m)}, \dots, c_{n-1}^{(m)}$ and the inverse are implemented by algebraic functions. Hence in place of (5.5) we can consider the derivatives

$$(5.6) \quad f_\alpha = \frac{\partial^{\alpha_1 + \dots + \alpha_{n-1}} f}{\partial c_1^{(m)\alpha_1} \dots \partial c_{n-1}^{(m)\alpha_{n-1}}}$$

and prove their algebraicity in t_m . Differentiation by t_m and transformation to the original variables z_1, \dots, z_n do not destroy their algebraicity.

In order to prove the algebraicity of the derivatives (5.6) we shall use the algebraicity of the function $f(t_m, c_1^{(m)}, \dots, c_{n-1}^{(m)})$ in t_m for fixed values of the other arguments. This means that we have an irreducible polynomial $P(f, t)$ in the ring $\mathbb{C}[f, t]$ of complex polynomials in the variables f, t such that $f(t_m)$ satisfies the equation

$$(5.7) \quad P(f(t_m), t_m) \equiv 0$$

(see [VW, L]). Note that the coefficients of the polynomial (5.7) and even its degrees in f and t depend on the parameters $(c_1^{(m)}, \dots, c_{n-1}^{(m)})$. Let us define the following subsets in the range of values of these parameters:

$$(5.8) \quad C_{qk} = \{(c_1^{(m)}, \dots, c_{n-1}^{(m)}) : \deg_f P = q, \deg_t P = k\}.$$

The union of the countable number of C_{qk} coincides with the whole range of values of the parameters $c_1^{(m)}, \dots, c_{n-1}^{(m)}$. This allows us to use the following well-known Baire theorem.

Baire's Theorem. *A complete metric space cannot be a countable union of nowhere dense subsets.*

The proof can be found in [RS]. We apply this fact to the of range of parameters $c_1^{(m)}, \dots, c_{n-1}^{(m)}$, and conclude that the closure of at least one of the sets C_{qk} has non-empty interior. Choose a domain D' whose natural projection lies in the interior of such C_{qk} . Also, let us choose in D' a curve of the family (2.6) with parameters in C_{qk} . Without loss of generality one can assume that this curve corresponds to

the parameters $c_i^{(m)} = 0$ and the point $t_m = 0$ on this curve is in D' . For the polynomial (5.7) we have

$$(5.9) \quad P(f, t) = \sum_{i=0}^q \sum_{j=0}^k a_{ij} f^i t^j.$$

We normalize the polynomial (5.9) by setting some of its nonzero coefficients a_{rs} to be equal to 1. This polynomial vanishes after the substitution $f = f(t_m)$ and $t = t_m$. Let us consider the functions

$$(5.10) \quad \varphi_{ij} = f(t)^i t^j, \text{ where } i = 0, \dots, q \text{ and } j = 0, \dots, k.$$

They are algebraic in t and depend holomorphically on the parameters $c_i^{(m)}$. If these parameters vanish, these functions (as functions of t) are linearly dependent. But elimination of the function φ_{rs} with $a_{rs} = 1$ makes the rest of the functions linearly independent. Otherwise we would have another nonzero polynomial $\tilde{P}(f, t)$ of the form (5.9) for which the equality (5.7) holds. Since P is irreducible, we have $\tilde{P}(f, t) = uP(f, t)$, where $u \in \mathbb{C}[f, t]$. But $\deg_f \tilde{P} \leq \deg_f P$ and $\deg_t \tilde{P} \leq \deg_t P$, therefore $u \in \mathbb{C} \subset \mathbb{C}[f, t]$. Comparing the coefficients $\tilde{a}_{rs} = 0$ and $a_{rs} = 1$ we find that the equality $\tilde{P}(f, t) = uP(f, t)$ cannot be true for $u \neq 0$.

We denote by X the set of all functions in (5.10), and by X' this set without φ_{rs} . Let us consider the Taylor expansions in t of the functions (5.10). One can treat their coefficients as infinitely-dimensional vectors (columns) of the linear space \mathbb{C}^∞ . Such vectors corresponding to the functions from X' form the $\infty \times N$ -matrix A , where N denote the number of elements of X' . The columns of A are linearly independent if $c_i^{(m)} = 0$. Therefore, there is an $N \times N$ -submatrix \tilde{A} of A with non-zero determinant. This minor is holomorphic in $c_i^{(m)}$ and therefore does not vanish in a neighborhood of the origin. Hence, the columns of A and the functions in X' are linearly independent for $c_i^{(m)}$ in a neighborhood of the origin.

Let us add the last column B corresponding to the function φ_{rs} to the matrix A , and consider the minors of order $(N + 1)$ of the extended matrix $A|B$. They vanish for $c_i^{(m)} = 0$ and for the parameters from the dense set C_{qk} . Therefore, they vanish identically. Thus, the functions in X' are linearly independent and the functions in X are linearly dependent for every $c_i^{(m)}$ in a neighborhood of the origin. Thus, φ_{rs} is a linear combination of the functions in X' . Its coefficients up to the sign coincide with the coefficients of the polynomial (5.9). They are defined uniquely by the linear system with the extended matrix $(\tilde{A}|B)$, where the N -th column \tilde{B} is formed by the elements of B lying in the rows defining \tilde{A} . Thus, the coefficients of the polynomial (5.9) are holomorphic in $c_i^{(m)}$, on a neighborhood of the origin. Now one can differentiate the equation (5.7) with respect to $c_i^{(m)}$, and we easily complete the proof. \square

Let us consider the functions (5.4). One can shrink D to $D' \subset D$ according to Lemma 5.1. Also, one can assume that the degrees of the polynomials (5.7) in f and t_m depend only on m in D' . Choose a point z^0 from (5.3) in the domain D' . This determines the functions (5.4). For the function f_1 Lemma 5.1 gives the algebraicity in t of the derivatives

$$(5.11) \quad \left. \frac{\partial^s f}{\partial t_2^s} \right|_{(t, 0, \dots, 0)}.$$

The derivatives (5.11) coincide with the derivatives of the function f_2 from (5.4) for $t = 0$. In fact,

$$(5.12) \quad \left. \frac{\partial^s f_2(t_1, t, 0, \dots, 0)}{\partial t^s} \right|_{t=0} = \left. \frac{\partial^s f}{\partial t_2^s} \right|_{(t_1, 0, \dots, 0)}.$$

We need the following

Lemma 5.2. *An algebraic function $f(t)$ is defined uniquely by its value and the values of a finite number of its derivatives at a regular point. If these values depend algebraically on a parameter τ , then $f = f(t, \tau)$ is an algebraic function of both variables t and τ .*

Assume the defining irreducible polynomial of the algebraic function $f(t)$ has the form (5.9). Repeating the above arguments, we again consider the functions (5.10) and their Taylor expansions at a regular point (one can assume it to be $t = 0$). The coefficients of these expansions depend algebraically on f and its derivatives at $t = 0$. Considering as above a nonsingular submatrix \tilde{A} and the corresponding linear system, we apply the Cramer rule and get the coefficients of the polynomial (5.9). In the second hypothesis of our claim they are algebraic in τ . By the separate algebraicity principle we conclude the argument.

We apply Lemma 5.2 to the function $f_2(t_1, t, 0, \dots, 0)$, taking into account its algebraicity in t and the algebraicity of derivatives (5.12) in t_1 . Therefore, the function $f_2(t_1, t, 0, \dots, 0)$ is algebraic in both variables.

This is the first induction step.

Assume that the functions f_1, \dots, f_m in (5.4) are algebraic. It follows from Lemma 5.1 that the derivatives

$$(5.13) \quad \left. \frac{\partial^s f}{\partial t_{m+1}^s} \right|_{(t_1, \dots, t_m, 0, \dots, 0)} = \left. \frac{\partial^s f_{m+1}(t_1, \dots, t_{m+1}, 0, \dots, 0)}{\partial t_{m+1}^s} \right|_{t_{m+1}=0}$$

are algebraic as well. Lemma 5.2 and the algebraicity of the derivatives (5.13) in t_1, \dots, t_m give the induction step from m to $m + 1$. This completes the proof of Theorem 2.

REFERENCES

- [Al] H.Alexander, *Holomorphic mappings from the ball and polydisk*, Math. Ann. **209** (1974), 249–256. MR **50**:5018
- [BM] S.Bochner and W.T.Martin, *Several complex variables*, Princeton University Press, Princeton, 1948. MR **10**:366a
- [BP] A.Boggress and J.Polking, *Holomorphic extension of CR functions*, Duke Math. J. **49** (1982), 757–784. MR **84j**:32018
- [Ch] E.Chirka, *Introduction to the geometry of CR manifolds*, Uspehi Mat. Nauk **46** (1991), 81–164. MR **92m**:32012
- [Fo] F.Forstneric, *Mappings of quadric Cauchy - Riemann manifolds*, Math. Ann. **292** (1992), 163–180. MR **93e**:32033
- [L] S.Lang, *Algebra*, Addison-Wesley, Reading, MA, 1965.
- [Pe] D.Pelles, *Proper holomorphic self-maps of the unit ball*, Math. Ann. **190** (1971), 298–305. MR **43**:3501
- [Pi] S.I.Pinchuk, *Boundary uniqueness theorem for holomorphic functions of several complex variables*, Mat. Zametki **15** (1974), 205–215. MR **50**:2558

- [Po] H.Poincaré, *Les fonctions analytiques de deux variables et la representation conforme*, Rend. Circ. Mat. Palermo **23** (1907), 185–220.
- [RS] M.Reed and B.Simon, *Methods of modern mathematical physics. V 1. Functional analysis*, Academic Press, New York and London, 1972. MR **58**:12429a
- [Ru] W.Rudin, *Function Theory on the Unit Ball of \mathbb{C}^n* , Springer, New York, 1980. MR **82i**:32002
- [Su1] A.Sukhov, *On CR mappings of real quadric manifolds*, Michigan Math. J. **41** (1994), 143–150. MR **94**:08
- [Su2] ———, *On holomorphic mappings of domains of the type of “wedge”*, Mat. Zametki **52** (1992), 141–145. MR **94e**:32047
- [Su3] ———, *On the mapping problem for quadric Cauchy - Riemann manifolds*, Indiana Univ. Math. J. **42** (1993), 27–36. MR **94e**:32039
- [Su4] ———, *On algebraicity of complex analytic sets*, Math. USSR Sbornik **74** (1991), 419–426. MR **93e**:32014
- [Ta] N.Tanaka, *On the pseudo-conformal geometry of hypersurfaces of the space of n complex variables*, J. Math. Soc. Japan **14** (1962), 397–429. MR **26**:3086
- [TH] A.Tumanov and G.Henkin, *Local characterization of holomorphic automorphisms of Siegel domains*, Funct. Anal. and Appl. **17** (1983), 49–61.
- [Tu] A.Tumanov, *Finite dimensionality of the group of CR automorphisms of standard CR manifolds and proper holomorphic mappings of Siegel domains*, Izv. Akad. Nauk USSR Ser. Mat. **52** (1988), 651–659. MR **89f**:32061
- [VW] B.L.Van der Waerden, *Algebra 1*, Springer Verlag, Berlin–Heidelberg–New York, 1966. MR **41**:8186
- [W1] S.Webster, *On the mapping problem for algebraic real hypersurfaces*, Invent. Math. **43** (1977), 53–68. MR **57**:3431
- [W2] ———, *Holomorphic mappings of domains with generic corners*, Proc. Amer. Math. Soc. **86** (1982), 236–240. MR **83k**:32041

DEPARTMENT OF MATHEMATICS, BASHKIR STATE UNIVERSITY, FRUNZE STREET 32, 450074 UFA, RUSSIA

E-mail address: yavdat@bgu.bashkiria.su

CMI, UNIVERSITE DE PROVENCE, 39 RUE F. JOLIOT-CURIE, 13453 MARSEILLE CEDEX 13, FRANCE